

New York
State College of Agriculture
At Cornell University
Ithaca, N. Y.

Library

Cornell University Library
QK 117.G75 1867

Manual of the botany of the northern Uni



3 1924 000 408 330

mann



Cornell University
Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

<http://www.archive.org/details/cu31924000408330>

B O T A N Y

OF

THE NORTHERN UNITED STATES.

MANUAL

THE BOTANY

THE NORTHERN UNITED STATES,

INCLUDING THE DISTRICT EAST OF THE MISSISSIPPI AND
NORTH OF NORTH CAROLINA AND TENNESSEE.

ARRANGED ACCORDING TO THE NATURAL SYSTEM.

BY ASA GRAY,

FISHER PROFESSOR OF NATURAL HISTORY IN HARVARD UNIVERSITY.

Fifth Edition.

WITH TWENTY PLATES,

ILLUSTRATING THE SEDGES, GRASSES, FERNS, ETC.

NEW YORK:
IVISON, PHINNEY, BLAKEMAN, & CO.
CHICAGO: S. C. GRIGGS & CO.

1867.

QK:
117
675
1867

Entered according to Act of Congress, in the year 1867, by
IVISON, PHINNEY, BLAKEMAN, & CO.,
in the Clerk's Office of the District Court of the United States for the Southern District of
New York.

②

UNIVERSITY PRESS : WELCH, BIGELOW, & Co.,
CAMBRIDGE.

TO

JOHN TORREY, LL. D.

ALMOST twenty years have passed since the first edition of this work was dedicated to you,— more than thirty, since, as your pupil, I began to enjoy the advantage of being associated with you in botanical pursuits, and in a lasting friendship. The flow of time has only deepened the sense of gratitude due to you from your attached friend,

ASA GRAY.

CAMBRIDGE, May 30, 1867.

CONTENTS.

| | |
|---|-----------|
| EXPLANATION OF SIGNS USED IN THIS WORK | Page 8 |
| EXPLANATION OF ABBREVIATION OF AUTHORS' NAMES | 9 |
| PREFACE | 11 |
| ANALYTICAL KEY TO THE ORDERS | 21 |
| FLORA. — PHENOGAMOUS OR FLOWERING PLANTS. | 33 |
| Dicotyledonous or Exogenous Plants | 33 |
| Angiospermous, Polypetalous | 33 |
| Monopetalous | 202 |
| Apetalous | 403 |
| Gymnospermous Plants | 468 |
| Monocotyledonous or Endogenous Plants | 475 |
| CRYPTOGAMOUS OR FLOWERLESS PLANTS. | 653 |
| Acrogenous Plants (Equisetaceæ, Ferns, &c.) | 653 |
| ADDENDA | 679 |
| INDEX | 681 |
| EXPLANATION OF THE PLATES | 695 |

SIGNS USED IN THIS WORK.

① An annual plant.

② A biennial plant.

⌚ A perennial plant.

? A mark of doubt.

! A mark of affirmation or authentication.

These signs are, however, very rarely employed in this volume.

1°, 2', 3". To save space, the sign of degrees (°) is used for feet; of minutes ('), for inches; of seconds ("), for lines,—the (English) line being the twelfth part of an inch.

The dash — between two figures, as “5-10,” means from 5 to 10, &c.

PRINCIPAL ABBREVIATIONS

OF THE NAMES OF AUTHORS CITED IN THIS VOLUME.

| | |
|--|--|
| <p><i>Adans.</i> = Adanson. <i>Ait.</i> Aiton. <i>All.</i> Allione. <i>Anders.</i> Andersson. <i>Andr.</i> Andrews. <i>Arn.</i> Arnott. <i>Aubl.</i> Aublet. <i>Bart.</i> Barton. <i>Bartl.</i> Bartling. <i>Beauv.</i> Palisot de Beauvois. <i>Benth.</i> Bentham. <i>Bernh.</i> Bernhardi. <i>Bieb.</i> Bieberstein. <i>Bigel.</i> Bigelow. <i>Boiss.</i> Boissier. <i>Bong.</i> Bongard. <i>Borkh.</i> Borkhausen. <i>Brong.</i> Brongniart. <i>Cass.</i> Cassini. <i>Cav.</i> Cavanilles. <i>Cham.</i> Chamisso. <i>Chav.</i> Chavannes. <i>Darlingt.</i> Darlington. <i>DC.</i> De Candolle. <i>A. DC.</i> Alphonse De Candolle. <i>Desf.</i> Desfontaines. <i>Dew.</i> Dewey. <i>Dill.</i> Dillenius. <i>Desv.</i> Desvaux. <i>Dougl.</i> Douglas. <i>Ehrh.</i> Ehrhart. <i>Ell.</i> Elliott. <i>Endl.</i> Endlicher. <i>Engelm.</i> Engelmann. <i>Fisch.</i> Fischer. <i>Gærtn.</i> Gärtner.</p> | <p><i>Gmel.</i> = Gmelin. <i>Good.</i> Goodenough. <i>Grev.</i> Greville. <i>Griseb.</i> Grisebach. <i>Gronov.</i> Gronovius. <i>Hartm.</i> Hartmann. <i>Hoffm.</i> Hoffmann. <i>Hook.</i> W. J. Hooker. <i>Hook. f. (filius)</i> J. D. Hooker. <i>Hornem.</i> Hornemann. <i>Huds.</i> Hudson. <i>H. B. K.</i> Humboldt, Bonpland, & <i>Jacq.</i> Jacquin. [Kunth. <i>Juss.</i> JUSSIEU. <i>A. Juss.</i> Adrien Jussieu. <i>L. or Linn.</i> LINNÆUS. <i>Lag.</i> Lagasca. <i>Lam.</i> Lamarck. <i>Lamb.</i> Lambert. <i>Ledeb.</i> Ledebour. <i>L'Her.</i> L'Heritier. <i>Lehm.</i> Lehmann. <i>Lesqx.</i> Lesquereux. <i>Lestib.</i> Lestibudois. <i>Lindl.</i> Lindley. <i>Lodd.</i> Loddiges. <i>Lour.</i> Loureiro. <i>Mart.</i> Martins. <i>Mich.</i> Micheli. <i>Michx.</i> Michaux (the elder). <i>Michx. f.</i> F. A. Michaux (the <i>Mill.</i> Miller. [younger.) <i>Mitch.</i> Mitchell. <i>Muhl.</i> Muhlenberg. <i>Nees,</i> Nees von Esenbeck. <i>Nutt.</i> Nuttall.</p> |
|--|--|

| | | | | | |
|--------------------------|---|----------------|------------------------|---|------------------|
| <i>Pav.</i> | = | Pavon. | <i>Steud.</i> | = | Stendel. |
| <i>Pers.</i> | | Persoon. | <i>Sulliv.</i> | | Sullivant. |
| <i>Pluk.</i> | | Plukenet. | <i>Thunb.</i> | | Thunberg. |
| <i>Plum.</i> | | Plumier. | <i>Torr.</i> | | Torrey. |
| <i>Poir.</i> | | Poiret. | <i>Torr. & Gr.</i> | | Torrey and Gray. |
| <i>R. Br.</i> | | ROBERT BROWN. | <i>Tourn.</i> | | Tournefort. |
| <i>Raf.</i> | | Rafinesque. | <i>Trautv.</i> | | Trautvetter. |
| <i>Reichenb.</i> | | Reichenbach. | <i>Trev.</i> | | Treviranus. |
| <i>Rich.</i> | | Richard. | <i>Trin.</i> | | Trinius. |
| <i>Richards.</i> | | Richardson. | <i>Tuckerm.</i> | | Tuckerman. |
| <i>Rœm.</i> | | Rœmer. | <i>Turcz.</i> | | Turczaninow. |
| <i>Rottb.</i> | | Rottböll. | <i>Vaill.</i> | | Vaillant. |
| <i>Salisb.</i> | | Salisbury. | <i>Vent.</i> | | Ventenat. |
| <i>Schk.</i> | | Schkuhr. | <i>Vill.</i> | | Villars. |
| <i>Schlecht.</i> | | Schlechtendal. | <i>Wahl.</i> | | Wahlenberg. |
| <i>Schrad.</i> | | Schrader. | <i>Walp.</i> | | Walpers. |
| <i>Schreb.</i> | | Schreber. | <i>Walt.</i> | | Walter. |
| <i>Schult.</i> | | Schultes. | <i>Wangh.</i> | | Wangenheim. |
| <i>Schw. or Schwein.</i> | | Schweinitz. | <i>Wedd.</i> | | Weddell. |
| <i>Scop.</i> | | Scopoli. | <i>Willd.</i> | | Willdenow. |
| <i>Ser.</i> | | Seringe. | <i>With.</i> | | Withering. |
| <i>Soland.</i> | | Solander. | <i>Wulf.</i> | | Wulfen. |
| <i>Spreng.</i> | | Sprengel. | <i>Zucc.</i> | | Zuccarini. |

P R E F A C E .

THIS work is designed as a compendious Flora of the Northern portion of the United States, for the use of students and of practical botanists.

The first edition (published in 1848) was hastily prepared to supply a pressing want. Its plan, having been generally approved, has not been altered, although the work has been to a great extent twice rewritten, and the geographical range extended. The second edition, much altered, appeared in 1856. The third and fourth were merely revised upon the stereotype plates, and some pages added, especially to the latter.

The *Garden Botany*, an Introduction to a Knowledge of the Common Cultivated Plants, which was prefixed to this fourth edition in 1863, is excluded from the present edition, and is to be incorporated into a simpler and more elementary work, but of wider scope, designed especially for school instruction, and for those interested in cultivation, — entitled *Field, Forest, and Garden Botany*.

In the present edition it has been found also expedient to remand to a supplementary volume the *Mosses* and *Liverworts*, so carefully and generously elaborated for the previous editions of this work by my friend, W. M. SULLIVANT, Esq. It is hoped that the *Lichenes*, if not all the other orders of the Lower Cryptogamia, may be added to this supplementary volume, so that our students may extend their studies into these more recondite and difficult departments of Botany.*

* The following important aids, moreover, are already provided, viz. *The Icones Muscorum*, or *Figures and Descriptions of most of those Mosses peculiar to*

Six plates, illustrating the genera of the *Cyperaceæ* or Sedge Family, are now added to the eight which illustrate the *Gramineæ* or Grasses, and the six which illustrate the *Filices* or Ferns and their allies: all are from original drawings by Mr. Isaac Sprague; and they should render the study of these families comparatively easy, even to the beginner.

In other respects the changes in this edition are only in details, and such as the progress of botanical knowledge, and the longer experience of the author and his associates or correspondents in teaching, have seemed to render necessary or advisable.

I am newly indebted to DR. GEORGE ENGELMANN, of St. Louis, for a revision of the account of *Cuscuta* and *Sagittaria*, &c., formerly prepared by him, and for the complete re-elaboration of the genera *Callitriche*, *Euphorbia*, *Pinus*, *Juncus*, and *Isoetes*. I have also to express my special acknowledgments to my friends, DR. J. W. ROBBINS, of Uxbridge, Massachusetts, who contributed the whole article on the difficult genus *Potamogeton*; — to CHARLES F. AUSTIN, of Closter, New Jersey, who furnished that on the *Lemnaceæ*; — and to Prof. DANIEL CADY EATON, of Yale College, who has entirely re-elaborated the Ferns for the present edition. The *Salicaceæ* and the genus *Carex*, as is well known, were contributed to the first edition by my old friend and associate, JOHN CAREY, Esq., now of London. Deprived of his further and important assistance, I have

Eastern North America which have not been heretofore figured, by WM. S. SULLIVANT, LL. D. Imp. 8vo, with 129 copper-plates.

Musci-Boreali-Americana, sive Specimina Exsiccata, etc. — A second and enlarged edition of the arranged collection of Mosses of the United States, published by Messrs. SULLIVANT and LESQUEREUX, of which the first issue was noticed in the preface to former editions of the Manual. The present edition comprises 536 species or varieties of Mosses, and is supplied by Mr. Leo Lesquereux, of Columbus, Ohio, for \$35 in gold, or £7 sterling.

Lichenes Exsiccati, by Professor EDWARD TUCKERMAN, of Amherst College; — of which four vols. (small 4to) have already been issued. A small volume on the *Genera of North American Lichenes* is now in preparation by the same author.

Nereis Boreali-Americana, — an account of the Marine Algæ of the United States, by the late Professor WM. H. HARVEY, a large quarto volume with fifty colored plates, — published by the Smithsonian Institution.

myself revised these articles as well as I could, in advance of the publication of Andersson's work on the *Salicaceæ* in the forthcoming volume of De Candolle's *Prodromus*, and of the posthumous volume of the late Dr. Boott's *Illustrations of Carex*. In the latter genus, however, I have been essentially aided by WILLIAM BOOTT, Esq., of Boston, and S. T. OLNEY, Esq., of Providence, who have made the *Carices* a special study. To render due acknowledgments to the correspondents who have contributed to the value of the *Manual* by the communication of specimens, notes, and corrections, would require me to enumerate all the cultivators and numerous amateurs of botany in this country. In special instances their names will be found scattered throughout the pages of the work. The necessity of economizing space to the utmost, so as to keep the volume within the dimensions of a *manual*, alone has debarred me from fuller citations of the names of collectors and of particular stations of rare or local plants. For the same reason I have generally omitted synonymes, except in case of some original or recent changes in nomenclature.

There is abundant reason, I doubt not, for me to renew the request that those who use this book will kindly furnish information of all corrections or additions that may appear to be necessary, so that it may be made more accurate and complete hereafter, and maintain the high character which it has earned.

Some explanations are needful in respect to details of typography, reference, and arrangement.

GEOGRAPHICAL LIMITATION, DISTRIBUTION, &c. As is stated on the title-page, this work is intended to comprise the plants which grow spontaneously in the United States north of North Carolina and Tennessee and east of the Mississippi. A *Flora* of the whole national domain, upon a similar plan (the issue of which I may now hope will not be delayed many years longer), would be much too bulky and expensive for the main purpose which this *Manual* fulfils.

For its purpose, the present geographical limitation is, on the whole, the best, — especially since the botany of the States south of our district has been so well provided for by my friend Dr. Chapman's *Flora of the Southern States*, issued by the same publishers. The southern boundary here adopted coincides better than any other geographical line with the natural division between the cooler-temperate and the warm-temperate vegetation of the United States; very few characteristically Southern plants occurring north of it, and those only on the low coast of Virginia, in the Dismal Swamp, &c. Our western limit, also, while it includes a considerable prairie vegetation, excludes nearly all the plants peculiar to the great Western woodless plains, which approach our borders in Iowa and Missouri. Our northern boundary, being that of the United States, varies through about five degrees of latitude, and nearly embraces Canada proper on the east and on the west, so that nearly all the plants of Canada East on this side of the St. Lawrence, as well as those of the deep peninsula of Canada West, will be found in this volume.*

I have here endeavored to indicate, briefly and generally, the district in which each species occurs, or in which it most abounds, in the following manner: 1. When the principal area of a species is southward rather than northward, I generally give first its northern limit, so far as known to me, if within the United States, and then its southern limit if within our boundaries, or add that it extends *southward*, meaning thereby that the species in question occurs in the States south of Virginia or Kentucky. Thus *Magnolia glauca*, p. 49, a prevailing Southern species, but which is sparingly found as far north as Massachusetts, is recorded as growing "near Cape Ann and New York southward, near the coast"; *M. acuminata*, "W. New York to Ohio and southward," &c. While in species of northern

* For the geographical statistics of our botany, see three articles in *The American Journal of Science and Arts*, Second Series, Vol. XXII. and Vol. XXIII. 1856 - 57.

range, the southern limits are mentioned; as *Anemone Pennsylvanica*, "W. New England to Illinois and northwestward." And so of Western plants; e. g. *Isopyrum biternatum*, p. 44, "Ohio to Illinois, Kentucky, and westward"; *Amorpha canescens*, p. 130, "Michigan to Wisconsin and southwestward." But this rule has not always been closely adhered to. 2. Where no habitat or range is mentioned, the species is supposed to be diffused over our whole area; or nearly so; and usually beyond it. 3. When the species is quite local or rare, the special habitat is given; e. g. *Vesicaria Shortii*, p. 73, and *Alyssum Lescurii*, p. 72; *Sullivantia Ohionis*, p. 169, &c. Except in such cases, or when the known geographical range of a species has been recently extended, the want of space has generally demanded the omission of particular stations, which are so appropriately given in local Floras and in more detailed works, but for which there is no room in a manual like this.

For the same reason, I could not here undertake to specify the range of those species which extend beyond the geographical limits of this work, or beyond the United States. Nevertheless, to facilitate the comparison of our American flora with that of Europe, I have appended the mark (Eu.) to those species which are indigenous to both.

DISTINCTION BETWEEN INTRODUCED AND INDIGENOUS SPECIES. Foreign plants which have become denizens of the soil have to be described along with the genuine indigenous members of our flora; but the *introduced* species are distinguished by the specific name being printed in a different type, namely, in small capitals,* while the names of the indigenous species are in heavier, *antique* letter.† Moreover, the country from which they came is specified (mostly Europe), as well as the nature of the denizenship. That is, following the suggestions of M. Alphonse De Candolle, I have classified our introduced plants as well as I could into two sorts, the

* For example, under *Ranunculus*, **R. BULBOSUS**, **ACRIS**, &c., p. 43.

† For example, **R. repens**, on the same page.

thoroughly *naturalized*, and the *adventive*; the first comprising those species which have made themselves perfectly at home in this country, propagating themselves freely by seed beyond the limits of cultivated grounds; the second, those which are only locally spontaneous, and perhaps precarious, or which are spontaneous only in cultivated fields, around dwellings, or in manured soil, and which, still directly or indirectly dependent upon civilized man, would probably soon disappear if he were to abandon the country. (I here rank with the *adventive* plants those weeds of cultivation which De Candolle terms plants cultivated without or against man's will.) Accordingly the species *naturalized from Europe* are indicated, at the close of the paragraph, by the phrase "(Nat. from Eu.)": those *adventive*, or less established, by the phrase "(Adv. from Eu.)," &c.

DISTINCTION OF GRADE OF VARIETIES. Vain is the attempt to draw an absolute line between varieties and species. Yet in systematic works the distinction has to be made absolute, and each particular form to be regarded as a species or a variety, according to the botanist's best judgment. Varieties, too, exhibit all degrees of distinctness. Such as are marked and definite enough to require names are distinguished here into two sorts, according to their grade: 1. Those which, I think, cannot be doubted to be varieties of the species they are referred to, have the name printed in small capitals.* These varieties make part of the common paragraph. 2. Those so distinct and peculiar that they have been, or readily may be, taken for species, and are some of them not unlikely to establish the claim: of these the name is printed in the same type as that of the species; and they are allowed the distinction of a separate paragraph,† — except where the variety itself is the only form in the country, as in the first species of *Anemone*.‡

* As, for instance, the three varieties of *Lespedeza violacea*, p. 137, viz. *DIVERGENS*, *SESSILIFOLIA*, and *ANGUSTIFOLIA*. See also, under *Ranunculus Flammula*, var. *INTERMEDIUS*, p. 41.

† As, Var. *reptans*, of the above-mentioned species.

‡ *A. patens*, L., var. *Nuttalliana*, p. 36.

ACCENTUATION OF NAMES. As a guide to correct pronunciation of botanical names (in which great carelessness prevails), I have marked the accented syllable; and have also (following Loudon's convenient mode) indicated what is called the long sound of the vowel by the grave (`) and the short sound by the acute (´) accent-mark.

INDICATION OF PROMINENT CHARACTERS is made by the use of *Italic type*, for the leading distinctions of the orders, and for those points in the specific descriptions by which two or more species of the same division may be most readily or surely discriminated, — the latter a plan adopted from *Koch's Flora Germanica*.

The ready discrimination of the genera is provided for by a *Synopsis*, in small type, of the leading characters of all the genera, when more than two, under each order. In this the genera are analytically disposed under their proper sub-orders, tribes, or other such natural groups, of whatever rank, properly characterized; and then, to save room, all these subordinal or tribal names and characters are left out of the body of the order, the genera following each other without a break.

Whenever a genus comprises several species, pains have been taken to render important differences conspicuous, and to abridge the labor of analysis, by proper grouping, and when needed by a series of rightly subordinated divisions and subdivisions. Divisions of the highest rank, or *Subgenera*, have the sectional mark (§) followed by the subgeneric name.* Those less important are indicated by the §, without a name; subsections or divisions of lower grade are marked by stars (*); their divisions, if any, by the +, and theirs again by the ++, &c.

Having in view the needs of students rather than of learned botanists, I have throughout endeavored to smooth the beginner's way by discarding many an unnecessary technical word or phrase, and

* As § 1. ATRAGENE, under Clematis, p. 35, and § 1. PULSATILLA, under Anemone, p. 36.

by casting the language somewhat in a vernacular mould, — perhaps at some sacrifice of brevity, but not, I trust, of the precision for which botanical language is distinguished.

ARRANGEMENT OF THE ORDERS. The Natural Orders are disposed in a series which nearly corresponds, in a general way, with De Candolle's arrangement (varied somewhat more in this edition, to come nearer to that adopted thus far in Bentham and Hooker's new *Genera Plantarum*), beginning with the highest class and ending with the lowest; and commencing this first and far the largest class (of *Dicotyledonous* or *Exogenous Plants*) with those orders in which the flowers are mostly provided with double floral envelopes, viz. with both calyx and corolla, and in which the corolla consists of separate petals (the *Polypetalous* division); beginning this series with those orders in which the several organs of the flower are most distinct and separate (*hypogynous*), and proceeding to those which have the parts most combined among themselves and consolidated with each other (*perigynous* and *epigynous*); then follow those with the petals combined into a monopetalous corolla (the *Monopetalous* division; and finally, those destitute of a corolla or destitute of all floral envelopes (the *Apetalous* division). The class of *Monocotyledonous* or *Endogenous Plants* opens with orders exhibiting one form of simplified flowers, passes to those with the organs most combined and consolidated, then to those less combined by adnation of parts, and closes with other simplified and reduced forms. The present problem in Botany is to group the numerous Natural Orders in each class into natural alliances. But this has not yet been done in such a manner as to be available to the ordinary student. I do not here attempt, therefore, to group the orders naturally, but let them follow one another in what seems to be on the whole a natural and manageable sequence. And, by means of an artificial

ANALYTICAL KEY TO THE ORDERS (p. 21), I enable the student to refer readily to its proper order any of our plants, upon taking the pains to ascertain the structure of its flowers, and sometimes of

the fruit, and following out a series of easy steps in the analysis. This key is founded upon the most obvious distinctions which will well answer the purpose, and is so contrived as to provide for all the exceptional instances and variant cases I could think of. I shall be disappointed if the attentive student is not able by it to refer to its proper order any to him unknown plant of the Northern States of which he has flowering specimens in hand. Referring to the Order which the Key leads him to, the student will find its most distinctive points, — which he has chiefly to consider, — brought together and printed in italics in the first sentence of the ordinal description, and thus can verify his results.

The Synopsis which follows will then lead him to the genus, to be verified in turn by the full generic description in its place; and the progress thence to the species, when there are several to choose from, is facilitated by the arrangement under divisions and subdivisions, as already explained.

It will be seen that the Key directs the inquirer to ascertain, first, the Class of the plant under consideration, — which, even without the seeds, is revealed at once by the plan of the stem, as seen in a cross-section, and usually by the veining of the leaves, and is commonly confirmed by the numerical plan of the flower; — then, if of the first class, the sub-class is at once determined by the pistil, whether of the ordinary kind, or an open scale bearing naked ovules. If the former, then the choice between the three divisions is determined by the presence or absence of the petals, and whether separate or united. Each division is subdivided by equally obvious characters, as, p. 21, first the number of stamens, then, whether the calyx is free from or connected with the surface of the ovary. And, finally, a series of successively subordinated propositions, — each set more indented upon the page than the preceding, leads to the name of the order sought for, followed by the number of the page upon which that order is described in the body of the work.

More particular instructions for the use of this book in the study of our plants are here superfluous; as these, as well as the needful preliminary knowledge, will be acquired from the author's *Introduction to Structural and Systematic Botany (Botanical Text-Book)*, or from the simpler *First Lessons in Botany*, — one or the other of which must needs be previously studied, and be the inseparable companion of *The Manual*.

ANALYTICAL KEY

TO THE ORDERS OF ALL THE PLANTS DESCRIBED IN THIS WORK.

SERIES I. PHÆNOGAMOUS OR FLOWERING PLANTS, those producing real flowers and seeds.

CLASS I. DICOTYLEDONOUS OR EXOGENOUS PLANTS.

Stems formed of bark, wood, and pith; the wood forming a zone between the other two, and increasing, when the stem continues from year to year, by the annual addition of a new layer to the outside, next the bark. Leaves netted-veined. Embryo with a pair of opposite cotyledons, or in Subclass II. often three or more in a whorl. Parts of the flower mostly in fours or fives.

SUBCLASS I. ANGIOSPERMÆ. Pistil consisting of a closed ovary which contains the ovules and the seeds.

DIVISION I. POLYPETALOUS: the calyx and corolla both present; the latter of *separate* petals.

A. *Stamens numerous, at least more than 10, and more than twice the sepals.*

1. *Calyx entirely free and separate from the pistil or pistils.*

- Pistils numerous but cohering over each other in a solid mass on an elongated receptacle. Page
MAGNOLIACEÆ, 48
- Pistils numerous, separate, but concealed in a hollow receptacle.
Leaves opposite, entire; no stipules. CALYCANTHACEÆ, 162
Leaves alternate, with stipules. Rosa, in ROSACEÆ, 146
- Pistils several, immersed in hollows of the upper surface of a large top-shaped receptacle. Nelumbium, in NYMPHÆACEÆ, 54
- Pistils more than one, separate, not enclosed in the receptacle.
Stamens inserted on the calyx, distinct. ROSACEÆ, 146
Stamens united with the base of the petals, monadelphous. MALVACEÆ, 98
Stamens inserted on the receptacle.
Filaments much shorter than the anther: trees. ANONACEÆ, 50
Filaments longer than the anther.
Flowers diœcious: twiners with alternate leaves. MENISPERMACEÆ, 51
Flowers perfect: if climbers, the leaves opposite.
Leaves not peltate: petals deciduous. RANUNCULACEÆ, 34
Leaves peltate: petals persistent. Brasenia, in NYMPHÆACEÆ, 54

- Pistils more than one, or several-lobed, the ovaries united below
the middle. RESEDACEÆ, 76
- Pistils several, their ovaries cohering in a ring around an axis,
but separating in fruit. MALVACEÆ, 98
- Pistils strictly one as to the ovary: the styles or stigmas may be several.
- Leaves punctate under a lens with transparent dots. HYPERICACEÆ, 83
- Leaves not punctate with transparent dots.
- Ovary simple, 1-celled, with one parietal placenta.
- Leaves 2-3-ternately compound or dissected. RANUNCULACEÆ, 34
- Leaves peltate, simply lobed. Podophyllum, in BERBERIDACEÆ, 52
- Ovary compound, 1-celled, with a central placenta. PORTULACACEÆ, 97
- Ovary compound, 1-celled, with 2 or more parietal placenta.
- Calyx caducous. Juice milky or colored. PAPAVERACEÆ, 58
- Calyx deciduous, of 4 sepals. CAPPARIDACEÆ, 75
- Calyx persistent, of 3 or 5 sepals. CISTACEÆ, 80
- Ovary compound, several-celled.
- Calyx valvate in the bud, and
- Persistent: stamens monadelphous: anthers 1-celled. MALVACEÆ, 98
- Deciduous: anthers 2-celled. TILIACEÆ, 103
- Calyx imbricated in the bud, persistent.
- Shrubs: stamens borne on the base of the petals. CAMELLIACEÆ, 103
- Aquatic or marsh herbs: ovules many,
- On 5 placenta in the axis. SARRACENIACEÆ, 57
- On the 8-24 partitions. NYMPHÆACEÆ, 54

2. *Calyx more or less coherent with the surface of the (compound) ovary.*

- Ovary 10-30-celled: ovules many, on the partitions: aquatic. NYMPHÆACEÆ, 54
- Ovary 2-5-celled.
- Leaves alternate, with stipules. Pomeæ, in ROSACEÆ, 146
- Leaves opposite, without stipules. Philadelphus, in SAXIFRAGACEÆ, 163
- Leaves alternate, without stipules. STYRACACEÆ, 309
- Ovary one-celled, with the ovules parietal.
- Fleshy plants with no true foliage: petals many. CACTACEÆ, 184
- Rough-leaved plants: petals 5 or 10. LOASACEÆ, 184
- Ovary one-celled, with the ovules rising from the base. PORTULACACEÆ, 97

B. *Stamens of the same number as the petals and opposite them.*

- Pistils 3-6, separate. Flowers dioecious. Woody vines. MENISPERMACEÆ, 51
- Pistil only one.
- Ovary one-celled: anthers opening by uplifted valves. BERBERIDACEÆ, 52
- Ovary one-celled: anthers not opening by uplifted valves.
- Style and stigma one: ovules more than one. PRIMULACEÆ, 313
- Style one: stigmas 3: sepals 2: ovules several. PORTULACACEÆ, 97
- Styles 5: ovule and seed only one. PLUMBAGINACEÆ, 312
- Ovary 2-4-celled.
- Calyx-lobes minute or obsolete: petals valvate. VITACEÆ, 112
- Calyx 4-5-cleft, valvate in the bud: petals involute. RHAMNACEÆ, 113

C. *Stamens not more than twice as many as the petals, when of just the number of the petals then alternate with them.*

1. *Calyx free from the ovary, i. e. the ovary wholly superior.*

* *Ovaries 2 or more, separate.*

Stamens united with each other and with a large and thick stigma common to the two ovaries. ASCLEPIADACEÆ, 394

Stamens unconnected, on the receptacle, free from the calyx.

Leaves punctate with pellucid dots. RUTACEÆ, 109

Leaves not pellucid-punctate.

Tree, with pinnate leaves. Ailanthus, in SIMARUBACEÆ, 110

Low shrub with pinnate leaves. Zanthorhiza, in RANUNCULACEÆ, 34

Herbs, not fleshy. RANUNCULACEÆ, 34

Herbs, with thick fleshy leaves. CRASSULACEÆ, 171

Stamens unconnected, inserted on the calyx,

Just twice as many as the pistils (flower symmetrical). CRASSULACEÆ, 171

Not just the number or twice the number of the pistils.

Leaves without stipules. SAXIFRAGACEÆ, 163

Leaves with stipules. ROSACEÆ, 146

* * *Ovaries 2-5, somewhat united at the base, separate above.*

Leaves punctate with pellucid dots. RUTACEÆ, 109

Leaves not pellucid-punctate.

Shrubs or trees with opposite leaves. SAPINDACEÆ, 116

* Terrestrial herbs: the carpels fewer than the petals. SAXIFRAGACEÆ, 163

* * * *Ovaries or lobes of ovary 3 to 5, with a common style.* GERANIACEÆ, 105

* * * * *Ovary only one, and*

+ *Simple, with one parietal placenta.* LEGUMINOSÆ, 123

+ + *Compound, as shown by the number of cells, placenta, styles, or stigmas.*

Ovary one-celled.

Corolla irregular: petals 4: stamens 6. FUMARIACEÆ, 60

Corolla irregular: petals and stamens 5. VIOLACEÆ, 76

Corolla regular or nearly so.

Ovule solitary: shrubs or trees: stigmas 3. ANACARDIACEÆ, 111

Ovules solitary or few: herbs. Some anomalous CRUCIFERÆ, 62

Ovules more than one, in the centre or bottom of the cell.

Petals not inserted on the calyx. CARYOPHYLLACEÆ, 87

Petals on the throat of a bell-shaped or tubular calyx. LYTHRACEÆ, 182

Ovules several or many, on two or more parietal placenta.

Leaves punctate with pellucid and dark dots. HYPERICACEÆ, 83

Leaves beset with reddish gland-tipped bristles. DROSERACEÆ, 82

Leaves neither punctate nor bristly-glandular.

Sepals 5, very unequal or only 3. CISTACEÆ, 80

Sepals and petals 4: stamens 6. ANOMALOUS CRUCIFERÆ, 62

Sepals and petals 5: stamens 5 or 10.

Ovary and stamens raised on a stalk. PASSIFLORACEÆ, 185

Ovary sessile. SAXIFRAGACEÆ, 163

Ovary 2 - several-celled.

Flowers irregular.

Anthers opening at the top,

Six or eight and 1-celled: ovary 2-celled. POLYGALACEÆ, 120

Ten and 2-celled: ovary 5-celled. Rhodora, in ERICACEÆ, 286

Anthers opening lengthwise.

Stamens 12 and petals 6 on the throat of a tubu-

lar inflated or gibbous calyx. Cuphea, in LYTHRACEÆ, 182

Stamens 5 - 8 or 10, and petals hypogynous, or nearly so.

Ovary 3-celled. SAPINDACEÆ, 116

Ovary 5-celled. Impatiens, &c. in GERANIACEÆ, 105

Flowers regular or nearly so.

Stamens neither just as many nor twice as many as petals,

Triadelphous: petals 5. HYPERICACEÆ, 83

Tetradynamous (or rarely only 2 or 4): petals 4:

pungent herbs. CRUCIFERÆ, 62

Distinct and fewer than the 4 petals. OLEACEÆ, 400

Distinct and more numerous than the petals. SAPINDACEÆ, 116

Stamens just as many or twice as many as the petals.

Ovules and seeds only 1 or 2 in each cell.

Herbs: flowers monœcious or diœcious. EUPHORBIACEÆ, 430

Herbs: flowers perfect and symmetrical.

Cells of the ovary as many as the sepals, &c. GERANIACEÆ, 105

Cells of the ovary (divided) twice as many as
the styles, sepals, &c. LINACEÆ, 104

Shrubs or trees.

Leaves 3-foliolate, pellucid-punctate. Ptelea, in RUTACEÆ, 109

Leaves palmately veined: fruit 2-winged. SAPINDACEÆ, 116

Leaves pinnately veined, simple, not punctate.

Calyx not minute: pod colored, dehiscent:

seeds enclosed in a pulpy aril. CELASTRACEÆ, 115

Calyx minute: fruit a berry-like drupe. AQUIFOLIACEÆ, 305

Ovules (and usually seeds) several or many in each cell.

Stipules between the opposite and simple leaves. ELATINACEÆ, 86

Stipules between the opposite and compound leaves

(but they are caducous). Staphylea, in SAPINDACEÆ, 116

Stipules none when the leaves are opposite.

Stamens 5, monadelphous in a 10-toothed tube

or cup: leaves simple, all radical. GALACINEÆ, 305

Stamens 10, monadelphous at the base. Leaf-

lets 3, inversely heart-shaped. Oxalis, in GERANIACEÆ, 105

Stamens distinct, free from the calyx.

Style 1, undivided. ERICACEÆ, 286

Styles 2 - 5, separate. CARYOPHYLLACEÆ, 87

Stamens distinct, inserted on the calyx.

Styles 2 (or 3), or splitting into 2 in fruit. SAXIFRAGACEÆ, 163

Style 1: pod in the calyx, 1-celled at maturity. LYTHRACEÆ, 182

2. *Calyx-tube adherent to the ovary, at least to its lower half.*

Tendrill-bearing and often succulent herbs. CUCURBITACEÆ, 186
 Not tendrill-bearing.

Ovules and seeds more than one in each cell.

Ovary 1-celled, many-ovuled from the base. PORTULACACEÆ, 97

Ovary 1-celled, with 2 or 3 parietal placentæ. SAXIFRAGACEÆ, 163

Ovary 2-several-celled.

Anthers opening by pores at the apex : style 1. MELASTOMACEÆ, 181

Anthers not opening by pores.

Stamens on a flat disk which covers the ovary. CELASTRACEÆ, 115

Stamens inserted on the calyx,

Eight or four (rarely five) : style 1. ONAGRACEÆ, 176

Five or ten : styles 2-3, distinct. SAXIFRAGACEÆ, 163

Ovules and seeds only one in each cell.

Stamens 10 or 5 (instead of many), — rarely in *Cratægus*, in ROSACEÆ, 146

Stamens 2 or 8 ; style 1 : stigma 2-4-lobed : herbs. ONAGRACEÆ, 176

Stamens 4 or 8 : aquatics : styles or sessile stigmas 4. HALORAGACEÆ, 174

Stamens 8 : styles 2 : shrub. HAMAMELACEÆ, 173

Stamens 4 : style and stigma 1 : chiefly shrubs. CORNACEÆ, 199

Stamens 5 : flowers in umbels, or rarely in heads.

Fruit dry, splitting in two at maturity : styles 2. UMBELLIFERÆ, 187

Fruit berry-like : styles 2-5, separate, or united. ARALIACEÆ, 198

DIVISION II. MONOPETALOUS : calyx and corolla both present ; the latter with its petals united more or less into one piece.

A. *Stamens more numerous than the lobes of the corolla.*

Ovary 1-celled with one parietal placenta. LEGUMINOSÆ, 123

Ovary 1-celled, with 2 parietal placentæ. *Adlumia*, &c. in FUMARIACEÆ, 60

Ovary 1-celled with the ovules at the centre or base. STYRACACEÆ, 309

Ovary 2-celled with a single ovule in each cell. POLYGALACEÆ, 120

Ovary 3-many-celled.

Stamens free or nearly free from the corolla : style single. ERICACEÆ, 286

Stamens free from the corolla : styles 5. *Oxalis*, in GERANIACEÆ, 105

Stamens inserted on the base or tube of the corolla.

Filaments monadelphous : anthers 1-celled, kidney-shaped. MALVACEÆ, 98

Filaments 1-5-adelphous at base : anthers 2-celled.

Calyx free from the ovary. CAMELLIACEÆ, 103

Calyx coherent with the ovary or with its base. STYRACACEÆ, 309

Filaments wholly distinct : calyx free, persistent. EBENACEÆ, 307

B. *Stamens (fertile ones) as many as the lobes of the corolla and opposite them.*

Ovary 5-celled : corolla appendaged with scales inside. SAPOTACEÆ, 308

Ovary 1-celled : pod several-many-seeded : style 1. PRIMULACEÆ, 313

Ovary 1-celled : utricle 1-seeded : styles 5. PLUMBAGINACEÆ, 312

C. *Stamens as many as the lobes of the corolla and alternate with them, or fewer.***1.** *Ovary adherent to the calyx-tube (inferior).*

Tendrils-bearing herbs : anthers often united. CUCURBITACEÆ, 186
 Tendrils none.

Stamens united by their anthers into a ring or tube.

Flowers in an involucrate head. COMPOSITÆ, 215

Flowers separate, not involucrate : corolla irregular. LOBELIACEÆ, 282

Stamens separate, free from the corolla or nearly so, as

many as its lobes : stipules none : juice milky. CAMPANULACEÆ, 285

Stamens separate, inserted on the corolla.

One to three, always fewer than lobes of the corolla. VALERIANACEÆ, 213

Four or five : leaves opposite or whorled.

Flowers in a dense head, with an involucre : no stipules. DIPSACEÆ, 215

Flowers if in heads not involucrate.

Leaves whorled and without stipules.

Leaves opposite or whorled, and with stipules. } RUBIACEÆ, 208

Leaves opposite without stipules (but some-
times with appendages to the petioles imitat-
ing them).

CAPRIFOLIACEÆ, 202

2. *Ovary free from the calyx (superior).*

* *Corolla irregular : stamens (with anthers) 4 and didynamous, or only 2.*

Ovules and seeds solitary in the (1-4) cells.

Ovary 4-lobed, the style rising from between the lobes. LABIATÆ, 341

Ovary not lobed, the style from its apex. VERBENACEÆ, 339

Ovules numerous or at least as many as 2 in each cell.

Ovary and pod 1-celled,

With a free central placenta : stamens 2 : aquatics. LENTIBULACEÆ, 317

With 2 or more parietal very many-seeded placentæ :

stamens 4. OROBANCHACEÆ, 322

Ovary and fruit more or less 4-5-celled. *Martynia*, }

Ovary and pod 2-celled, but the 2 placentæ parietal. } BIGNONIACEÆ, 320

Ovary and pod 2-celled : placentæ in the axis.

Seeds rarely few, not on hooks, with albumen. SCROPHULARIACEÆ, 324

Seeds few, borne on hook-like or other projections
of the placentæ : no albumen. ACANTHACEÆ, 338

* * *Corolla somewhat irregular : stamens (with anthers) 5.*

Stamens free from the corolla : anthers with their cells

opening by a hole or chink at the top. *Azalea*, in ERICACEÆ, 286

Stamens inserted on the corolla.

Filaments or some of them woolly. *Verbascum*, in SCROPHULARIACEÆ, 324

Filaments not woolly. *Hyoscyamus*, in SOLANACEÆ, 380

* * * *Corolla regular.*

← *Stamens as many as the lobes of the corolla.*

Ovaries 2, separate; their

Styles and stigmas also wholly separate. *Dichondra*, in CONVULVULACEÆ, 374

Stigmas and sometimes styles united into one.

Filaments distinct: pollen in ordinary grains. APOCYNACEÆ, 392

Filaments monadelphous: pollen in masses. ASCLEPIADACEÆ, 394

Ovary one, but deeply 4-lobed around the style.

Leaves alternate. BORRAGINACEÆ, 360

Leaves opposite. *Mentha*, in LABIATÆ, 341

Ovary one: pod 2-lobed or 2-horned at the summit. LOGANIACEÆ, 391

Ovary one, not divided nor deeply lobed,

One-celled, with ovules parietal or on 2 parietal placenta.

Leaves (or in *Menyanthes* three leaflets) entire. GENTIANACEÆ, 384

Leaves toothed, lobed or pinnately compound. HYDROPHYLLACEÆ, 367

Two- to ten-celled.

Leafless parasitic twining plants. *Cuscuta*, in CONVULVULACEÆ, 374

Leaves opposite, their bases or petioles connected

by stipules or a stipular line LOGANIACEÆ, 391

Leaves when opposite without stipules.

Stamens free from the corolla or nearly so: style 1. ERICACEÆ, 286

Stamens almost free from the corolla: style none. AQUIFOLIACEÆ, 305

Stamens inserted on the tube of the corolla,

Four: pod 2-celled, circumcissile. PLANTAGINACEÆ, 310

Four: ovary 2-4-celled: ovules solitary. VERBENACEÆ, 339

Five or rarely more.

Fruit of two or four seed-like nutlets. BORRAGINACEÆ, 360

Fruit a few-seeded pod.

Style 3-cleft: seeds small. POLEMONIACEÆ, 370

Style single or 2-cleft, or again 2-cleft: seeds

large, only one or two in a cell. CONVULVULACEÆ, 374

Fruit a very many-seeded pod or berry.

Styles 2. *Hydrolea*, in HYDROPHYLLACEÆ, 367

Style single. SOLANACEÆ, 380

← ← *Stamens fewer than the lobes of the corolla.*

Stamens 4, didynamous.

Ovary 2-celled; the cells several-seeded. ACANTHACEÆ, 338

Ovary 2-4-celled; the cells 1-seeded. VERBENACEÆ, 339

Stamens only 2 with anthers: ovary 4-lobed. *Lycopus*, in LABIATÆ, 342

Stamens 2, rarely 3: ovary 2-celled.

Low herbs: corolla scarious, withering on the pod. PLANTAGINACEÆ, 310

Herbs: corolla rotate, or somewhat funnelform, and

slightly irregular. *Veronica*, in SCROPHULARIACEÆ, 324

Shrubs or trees: corolla perfectly regular. OLEACEÆ, 400

DIVISION III. APETALOUS: corolla (and sometimes calyx) wanting.

A. *Flowers not in catkins.*1. *Ovary or its cells containing many ovules.*

Ovary and pod inferior (i. e. calyx-tube adherent to the ovary),

Six-celled: stamens 6-12. ARISTOLOCHIACEÆ, 403

Four-celled: stamens 4. Ludwigia, in ONAGRACEÆ, 176

One-celled, with parietal placentæ. Chrysosplenium, in SAXIFRAGACEÆ, 163

Ovary and pod wholly naked (there being no calyx),

Two-celled, 2-beaked: flowers capitate: tree. . . . HAMAMELACEÆ, 173

Two-celled, many-ribbed: aquatic herb. PODOSTEMACEÆ, 429

Ovary and pod superior, i. e. free from the calyx,

Five-celled and 5-beaked, opening across the beaks, which

fall off at maturity: stamens 10. Penthorum, in CRASSULACEÆ, 171

Three-5-celled, opening round the middle. Sesuvium, in PORTULACACEÆ, 97

Three-celled and 3-valved. . . . Mollugo, in CARYOPHYLLACEÆ, 87

Two-celled or one-celled: placentæ central.

Stamens inserted on the throat or tube of the calyx. LYTHRACEÆ, 182

Stamens inserted on the receptacle or the base of the calyx,

Alternate with the 5 sepals. . . . Glaux, in PRIMULACEÆ, 313

Opposite the sepals when of the same number. CARYOPHYLLACEÆ, 87

One-celled, with one parietal placenta. } RANUNCULACEÆ, 34

Ovaries 2 or more, separate, simple. }

2. *Ovary or its cells containing only 1 or 2, rarely 3 or 4, ovules.** *Pistils more than one, and distinct or nearly so.*

Stamens inserted on the calyx. Leaves with stipules. . . . ROSACEÆ, 146

Stamens inserted on the receptacle.

Leaves punctate with pellucid dots. . . . Zanthoxylum, in RUTACEÆ, 109

Leaves not dotted.

Calyx present, and usually colored or petal-like. RANUNCULACEÆ, 34

Calyx absent. Flowers entirely naked, perfect, spiked. SAURURACEÆ, 427

* * *Pistil one, either simple or compound.*

Ovary partly inferior, the calyx coherent to its lower half,

2-celled: styles 2: stamens many. . . . HAMAMELACEÆ, 173

Ovary wholly inferior (in perfect or pistillate flowers).

Aquatic herbs: ovary 3-4 celled, or (Hippuris) 1-celled. HALORAGEÆ, 174

Woody plants: style or stigma one, entire: ovary 1-celled.

Stigma running down one side of the style. Nyssa, in CORNACEÆ, 199

Stigma terminal, with or without a style.

Parasitic on the branches of trees: anthers sessile. LORANTHACEÆ, 426

Not parasitic above ground: anthers on filaments. SANTALACEÆ, 425

- Ovary really free from the calyx, but permanently invested by its tube, or the base of it, so as to seem inferior.
- Shrubs, with scurfy leaves : flowers mostly dioecious. ELÆAGNACEÆ, 424
- Herbs : with the calyx colored like a corolla.
- Leaves opposite, simple. NYCTAGINACEÆ, 404
- Leaves alternate, pinnate. Poterium, in ROSACEÆ, 146
- Ovary plainly free from the calyx, which is sometimes wanting.
- Stipules (ochreæ) sheathing the stem at the nodes.
- Tree : the calyx none : flowers monœcious, in heads. PLATANACEÆ, 446
- Herbs : the calyx present and commonly petal-like. POLYGONACEÆ, 414
- Stipules not sheathing the stem, or none.
- Aquatic herbs, submersed or nearly so.
- Leaves whorled and dissected : style single. CERATOPHYLLACEÆ, 427
- Leaves opposite, entire : styles 2 : ovary 4-celled. CALLITRICHACEÆ, 427
- Not aquatics, herbs.
- Ovary 10-celled : berry 10-seeded. PHYTOLACCACEÆ, 405
- Ovary 3- (rarely 1-2-) celled : juice usually milky. EUPHORBIACEÆ, 430
- Ovary one-celled : juice not milky.
- Style, if any, and stigma only one : leaves simple :
- no scarious bracts around the flowers. URTICACEÆ, 440
- Style or stigmas 2 or 3 : embryo coiled or curved.
- Stipules not scarious ; leaves palmately cleft or palmately compound. Cannabineæ, in URTICACEÆ, 440
- Stipules scarious. Illecebreæ, in CARYOPHYLLACEÆ, 87
- Stipules and scarious bracts none : stamens inserted high up on the tube of the calyx. Scleranthus, in CARYOPHYLLACEÆ, 87
- Stipules none : but flowers with scarious bracts. AMARANTACEÆ, 411
- Stipules and scarious bracts none CHENOPODIACEÆ, 405
- Shrubs or trees.
- Ovules a pair in each cell of the ovary.
- Fruit 2-celled, a double samara. Acerineæ, in SAPINDACEÆ, 116
- Fruit a 1-celled and 1-seeded samara or a drupe. OLEACEÆ, 400
- Ovules single in each cell of the
- Three - nine-celled ovary : leaves heath-like. EMPETRACEÆ, 440
- Three-celled ovary : leaves broad. RHAMNACEÆ, 113
- One - two-celled ovary : styles or stigmas 2-cleft. URTICACEÆ, 440
- One-celled ovary : style and stigma single and entire.
- Anthers opening longitudinally THYMELEACEÆ, 424
- Anthers opening by uplifted valves. LAURACEÆ, 422
- B.** *Flowers (monœcious or dioecious) one or both sorts in catkins.*
1. *Only one sort of flowers in catkins or catkin-like heads.*
- Fertile flowers in a short catkin, head, or strobile. URTICACEÆ, 440
- Fertile flowers single or clustered : sterile ones in slender catkins.
- Leaves pinnate : fertile flowers and fruit naked. JUGLANDACEÆ, 447
- Leaves simple : fertile flowers 1-3 in an involucre or cup. CUPULIFERÆ, 449

2. *Both sterile and fertile flowers in catkins or catkin-like heads.*

- Ovary and pod 2-celled, many-seeded. Liquidambar, in HAMAMELACEÆ, 173
 Ovary and pod 1-celled, many-seeded; seeds furnished with
 a downy tuft at one end. SALICACEÆ, 461
 Ovary 1 - 2-celled, only one ovule in each cell: fruit 1-seeded.
 Parasitic on trees: fruit a berry. LORANTHACEÆ, 426
 Trees or shrubs, not parasitic.
 Calyx regular, in the fertile flower succulent in fruit. URTICACEÆ, 440
 Calyx none, or rudimentary and scale-like.
 Style and stigma one, simple: the flowers in heads. PLATANACEÆ, 446
 Styles or long stigmas 2.
 Fertile flowers 2 or 3 under each scale of the catkin. BETULACEÆ, 458
 Fertile flowers single under each scale: nutlets
 naked, waxy-coated or drupe-like. MYRICACEÆ, 457
 Fertile flowers in pairs at each scale, each in a mem-
 branous sac or with leafy bractlets. Carpinæ, in CUPULIFERÆ, 449

SUBCLASS II. GYMNOSPERMÆ. Pistil an open scale or altered leaf, bearing naked ovules on its margin or its upper surface, or in *Taxus* entirely wanting. Flowers monœcious or diœcious.

Represented in the Northern United States only by the order CONIFERÆ, 468

CLASS II. MONOCOTYLEDONOUS OR ENDOGENOUS PLANTS.

Stems with the wood collected into separate bundles or threads, which are irregularly dispersed throughout the whole diameter, leaving no distinct pith in the centre; not forming annual layers. (A transverse slice of the stem therefore exhibits the woody threads as dots scattered throughout the cellular tissue.) Leaves mostly parallel-veined (occasionally more or less reticulated). Embryo with a single cotyledon, and the first leaves in germination alternate. Parts of the flower generally in threes, never in fives.

A. SPADICEOUS DIVISION. *Flowers aggregated on a spadix or fleshy axis, or sometimes scattered, destitute of calyx and corolla (excepting some Aracæ and Naiadacæ, where, however, they are on a spadix), and also without glumes (husky scales). Leaves sometimes with netted veins.*

- Little floating aquatics, with no distinction of stem and foliage. LEMNACEÆ, 478
 Immersed aquatics, branching and leafy. NAIDACEÆ, 482
 Reed-like or Flag-like marsh herbs, with linear and sessile
 nerved leaves: flowers in spikes or heads.
 Flowers monœcious, and quite destitute of floral envelopes. TYPHACEÆ, 480
 Flowers perfect, on a lateral spadix: sepals 6. Acorus, }
 Terrestrial or marsh plants: leaves mostly with a distinct }
 netted-veined blade, petioled. ARACEÆ, 475

B. PETALOIDEOUS DIVISION. *Flowers not collected on a spadix, furnished with floral envelopes (perianth) answering to calyx or to both calyx and corolla, either herbaceous or colored and petal-like.*

1. *Perianth adherent to the whole surface of the ovary.*

Flowers diœcious or polygamous, regular.

Aquatics : ovules and seeds several or numerous. HYDROCHARIDACEÆ, 494

Twiners : ovules and seeds one or two in each cell. DIOSCOREACEÆ, 518

Flowers perfect : ovules and seeds usually numerous.

Stamens only one or two : flower irregular, gynandrous. ORCHIDACEÆ, 497

Stamens three.

Anthers introrse, opening transversely. . . . BURMANNIACEÆ, 496

Anthers introrse or versatile, opening lengthwise. HEMODORACEÆ, 514

Anthers extrorse, opening lengthwise. . . . IRIDACEÆ, 515

Stamens 6 : flowers usually on a scape from a bulb. AMARYLLIDACEÆ, 512

2. *Perianth adherent only to the base or lower half of the ovary.*

Perianth woolly or roughish-mealy : the leaves equitant. HEMODORACEÆ, 514

Perianth smooth : the leaves grass-like. Stenanthium, &c., in LILIACEÆ, 520

3. *Perianth wholly free from the ovary.*

Pistils numerous or few in a head or ring. . . . ALISMACEÆ, 490

Pistil one : anthers 1-celled : flowers diœcious. Tendril-bearing. SMILACEÆ, 518

Pistil one, compound (cells or placentæ mostly 3) : anthers 2-celled.

Perianth not glumaceous or chaffy : flowers not in dense heads.

Stamens 6 (in one Smilacina 4), all alike and perfect.

Scurfy-leaved epiphyte : seeds hairy-tufted. . . . BROMELIACEÆ, 515

Rush-like marsh-herbs : carpels separating closed from the axis : seed without albumen. Triglochin, in ALISMACEÆ, 490

Terrestrial, not rush-like : seeds with albumen.

Perianth of similar divisions or lobes, mostly colored }
 Perianth of 3 foliaceous and green sepals and 3 col- } LILIACEÆ, 520
 ored withering-persistent petals. Trillium in }

Perianth of 3 persistent green sepals, and 3 ephemeral deliquescent petals COMMELYNACEÆ, 546

Stamens 6, dissimilar, or only three with perfect anthers.

Perianth of 3 herbaceous sepals and 3 unequal and ephemeral petals. . . . COMMELYNACEÆ, 546

Perianth tubular, 6-lobed. . . . PONTEDERIACEÆ, 544

Stamens 3, similar. Moss-like aquatic. Mayaca, under XYRIDACEÆ, 547

Perianth wholly glumaceous, of 6 similar divisions. Rushes. JUNCACEÆ, 536

Perianth partly glumaceous or chaff-like : flowers in very dense heads. Rush-like or aquatic.

Flowers perfect : inner perianth of three yellow petals :

perfect stamens and plumose sterile filaments each

3 : pod 1-celled, many-seeded on 3 parietal placentæ. XYRIDACEÆ, 547

Flowers monœcious or diœcious, whitish-bearded : sta-

mens 4 or 3 : pod 2-3-celled, 2-3-seeded. ERIOCAULONACEÆ, 549

C. GLUMACEOUS DIVISION. *Flowers destitute of any proper perianth, except sometimes small scales or bristles, but covered by glumes, i. e. husk-like or scale-like bracts.*

Glume a single scale-like bract with a flower in its axil. CYPERACEÆ, 550
 Glumes in pairs, of two sorts. GRAMINEÆ, 602

SERIES II. CRYPTOGAMOUS OR FLOWERLESS PLANTS:
 those destitute of stamens and pistils, in fructification producing *spores*
 instead of seeds.

CLASS III. ACROGENOUS PLANTS.

Plants with a stem containing woody tissue and vessels, as does the foliage when there is any (in the form of veins).

Fructification of several spore-cases borne on the under side of the shield-shaped stalked scales of a terminal spike or cone. Leaves none, except a whorl of teeth at each joint of the stem. EQUISETACEÆ, 653

Fructification borne on the leaves (fronds), commonly on their backs or margins. FILICES, 655

Fructification of spore-cases in the axil of simple leaves or bracts. LYCOPODIACEÆ, 672

Fructification on the branches or petioles. HYDROPTERIDES, 677

B O T A N Y

OF THE

NORTHERN UNITED STATES.

SERIES I.

PHÆNÓGAMOUS OR FLOWERING PLANTS.

VEGETABLES bearing proper flowers, that is, having stamens and pistils, and producing seeds, which contain an embryo.

CLASS I. DICOTYLÉDONOUS OR EXÓGE- NOUS PLANTS.

Stems formed of bark, wood, and pith; the wood forming a layer between the other two, increasing, when the stem continues from year to year, by the annual addition of a new layer to the outside, next the bark. Leaves netted-veined. Embryo with a pair of opposite cotyledons, or rarely several in a whorl. Flowers having their parts usually in fives or fours.

SUBCLASS I. ANGIOSPÉRMÆ.

Pistil consisting of a closed ovary, which contains the ovules and forms the fruit. Cotyledons only two.

DIVISION I. POLYPÉTALOUS EXÓGENOUS PLANTS.

Floral envelopes consisting of both calyx and corolla; the petals not united with each other. (Several genera or species belonging to Polypetalous Orders are destitute of petals.)

ORDER 1. RANUNCULÀCÆÆ. (CROWFOOT FAMILY.)

Herbs or woody climbers, rarely undershrubs, with a colorless acrid juice, polypetalous, or apetalous with the calyx often colored like a corolla, hypogynous; the sepals, petals, numerous stamens, and many or few (rarely single) pistils all distinct and unconnected. — Flowers regular or irregular. Sepals 3–15. Petals 3–15, or wanting. Stamens indefinite, rarely few: anthers short. Fruits either dry pods, or seed-like (achenia), or berries. Seeds anatropous (when solitary and suspended the rhapshe dorsal), with fleshy albumen and a minute embryo. — Stipules none. Leaves often dissected, their stalks dilated at the base. (A large family, mostly of acrid plants, some of them acrid-narcotic poisons.)

Synopsis of the Genera.

Tribe I. CLEMATIDÆÆ. Sepals valvate in the bud, or with the edges bent inwards. Petals none, or small. Achenia numerous, tailed with the feathery or hairy styles. Seeds solitary, suspended. — Leaves all opposite.

1. **Clematis.** Climbing by the leafstalks, or erect herbs.

Tribe II. ANEMONEÆÆ. Sepals imbricated in the bud. Petals none or very small and stamen-like. Achenia numerous or several. Seed solitary. — Stem-leaves often opposite or whorled, forming an involucre.

* Seed suspended. Sepals 4–20.

2. **Anemone.** Achenia numerous, in a head, pointed or tailed, not ribbed nor inflated. Involucre leaf-like and remote from the flower.

3. **Hepatica.** Achenia several, not ribbed. Involucre close to the flower, of 3 simple leaves, and resembling a calyx.

4. **Thalictrum.** Achenia 4–10, ribbed, grooved, or inflated. Involucre none, or leaf-like, and remote from the flowers.

** Seed erect. Sepals 3–5, caducous.

5. **Trautvetteria.** Achenia inflated and 4-angled. Involucre none.

Tribe III. RANUNCULÆÆ. Sepals imbricated in the bud. Petals evident, often with a scale or pore inside. Achenia numerous. Seed solitary.

6. **Ranunculus.** Sepals not appendaged. Achenia in a head. Seed erect.

7. **Myosurus.** Sepals spurred at the base. Achenia in a long spike. Seed suspended.

Tribe IV. HELLEBORINEÆÆ. Sepals imbricated in the bud, deciduous, rarely persistent, petal-like. Petals (*nectaries* of the early botanists) tubular, irregular, or 2-lipped, often none. Pods (follicles) few, rarely single, few-several-seeded. — Leaves all alternate.

* Flower regular. Pods several-seeded. Herbs.

8. **Isopyrum.** Petals none (in our species). Pods few. Leaves compound.

9. **Caltha.** Petals none. Pods several. Leaves kidney-shaped, undivided.

10. **Trollius.** Petals many, minute and stamen-like, hollowed near the base. Pods 8-15, sessile. Leaves palmately divided.
11. **Coptis.** Petals 5-6, small, hollowed at the apex. Pods 3-7, long-stalked. Sepals deciduous. Leaves trifoliolate.
12. **Helleborus.** Petals 8-10, small, tubular, 2-lipped. Pods several, sessile. Sepals 5, persistent, turning green with age.
13. **Aquilegia.** Petals 5, spur-shaped, longer than the 5 deciduous sepals. Pods 5.
* * Flower unsymmetrical and irregular. Pods several-seeded.
14. **Delphinium.** Upper sepal spurred. Petals 4, of two forms; the upper pair with long spurs, enclosed in the spur of the calyx.
15. **Aconitum.** Upper sepal hooded, covering the two long-clawed small petals.
* * * Flower symmetrical. Pods ripening only one seed. Shrubby.
16. **Zanthorhiza.** Petals 5, small, 2-lobed, with claws. Stamens 5-10. Flowers in drooping compound racemes.
- Tribe V. CIMICIFUGEÆ.** Sepals imbricated, falling off as the flower opens. Petals small and flat, or none. Pistils 1-several. Fruit a 2-several-seeded pod or berry. All the leaves alternate.
17. **Hydrastis.** Flower solitary. Pistils several in a head, becoming berries in fruit, 2-seeded. Leaves simple, lobed. Petals none.
18. **Actæa.** Flowers in a single short raceme. Pistil single, forming a many-seeded berry. Leaves 2-3-ternately compound. Petals manifest, but small.
19. **Cimicifuga.** Flowers in long spiked racemes. Pistils 1-8, in fruit forming dry and several-seeded pods. Leaves 2-3-ternately compound.

1. CLÉMATIS, L. VIRGIN'S-BOWER.

Sepals 4, or rarely more, colored, the valvate margins turned inwards in the bud. Petals none, or small. Achenia numerous in a head, bearing the persistent styles as naked, hairy, or plumose tails.—Perennial herbs or vines, mostly a little woody, and climbing by the bending or clasping of the leafstalks, rarely low and erect. Leaves opposite. (*Κλημαρίς*, a name of Dioscorides for a climbing plant with long and lithe branches.)

§ 1. **ATRÁGENE, L.** *Some of the outer filaments enlarged and gradually passing into small spatulate petals: peduncles bearing single large flowers: the thin sepals widely spreading.*

1. **C. verticillàris, DC.** Woody-stemmed climber, almost glabrous; leaves trifoliolate, with slender common and partial petioles; leaflets ovate or slightly heart-shaped, pointed, entire, or on sterile stems 1-3-toothed or lobed; flower bluish-purple, 2' - 3' across; tails of the fruit plumose. (*Atragene Americana, Sims.*)—Rocky places in mountainous districts, Maine and Western New England to Virginia, Wisconsin, and northwestward: rare. May.—A pair of leaves with a peduncle between them, developed in spring from each of the opposite buds, gives the appearance of a whorl, whence the specific name.

§ 2. **CLEMATIS** proper. *Petals entirely wanting.*

* *Peduncles bearing single large nodding flowers: calyx leathery: anthers linear.*

← *Stem low, erect and mostly simple: calyx silky outside, greenish.*

2. **C. ochroleuca, Ait.** Leaves simple and entire, ovate or sometimes 3-lobed, almost sessile, silky beneath; tails of the fruit very plumose.—Copses, Long Island, Staten Island (*Dr. Allen*), Pennsylvania, and Virginia: rare. May.

— — *Stems climbing: leaves pinnate: calyx (and foliage) glabrous or puberulent.*

3. **C. Viórna**, L. (LEATHER-FLOWER.) Calyx ovate and at length bell-shaped; the purplish *sepals very thick and leathery, tipped with short recurved points*; the long tails of the fruit *very plumose*; leaflets 3–7, ovate or oblong, sometimes slightly cordate, 2–3-lobed or entire; uppermost leaves often simple. — Rich soil, Pennsylvania, Ohio, and southward. May–Aug.

4. **C. Pitcheri**, Torr. & Gray. Calyx bell-shaped; the dull purplish *sepals with narrow and slightly margined recurved points*; tails of the fruit *filiform and barely pubescent*; leaflets 3–9, ovate or somewhat cordate, entire or 3-lobed, much reticulated; uppermost leaves often simple. — Illinois on the Mississippi, and southward. June.

5. **C. cylindrica**, Sims. Calyx cylindraceous below, the upper half of the bluish-purple *sepals dilated* and widely spreading, with *broad and wavy thin margins*; tails of the fruit *silky*; leaflets 5–9, thin, varying from oblong-ovate to lanceolate, entire or 3–5-parted. — Virginia near Norfolk, and southward. May–Aug.

* * *Flowers in paniced clusters, polygamo-dicæious: sepals thin: anthers oblong.*

6. **C. Virginiana**, L. (COMMON VIRGIN'S-BOWER.) Smooth; leaves bearing 3 ovate acute leaflets, which are cut or lobed, and somewhat heart-shaped at the base; tails of the fruit plumose. — River-banks, &c., common; climbing over shrubs. July, August. — The axillary peduncles bear clusters of numerous white flowers (sepals obovate, spreading); the fertile succeeded in autumn by the conspicuous feathery tails of the fruit.

2. ANEMONE, L. ANÉMONE. WIND-FLOWER.

Sepals few or many, petal-like. Petals none, or in No. 1 resembling abortive stamens. Achenia pointed or tailed, flattened, not ribbed. Seed suspended. — Perennial herbs with radical leaves; those of the stem 2 or 3 together, opposite or whorled, and forming an involucre remote from the flower. (Name from *ἀνεμος*, the wind, because the flower was thought to open only when the wind blows.)

§ 1. **PULSATÍLLA**, Tourn. *Carpels numerous in a head, with long and hairy styles which in fruit form feathery tails, as in Clematis: flower large, usually with some glandular bodies like abortive stamens answering to petals, but minute or indistinct.*

1. **A. patens**, L., var. **Nuttalliana**. (PASQUE-FLOWER.) Villous with long silky hairs; flower erect, developed before the leaves; which are ternately divided, the lateral divisions 2-parted, the middle one stalked and 3-parted, the segments deeply once or twice cleft into narrowly linear and acute lobes; lobes of the involucre like those of the leaves, at the base all united into a shallow cup; sepals 5–7, purplish or whitish (1'–1½'' long), spreading when in full anthesis. (A. Nuttalliana, DC. Pulsatilla Nuttalliana, ed. 2. P. patens, var. Wolfgangiana, Trautv.) — Prairies, Illinois (Bebb), Wisconsin (Lapham), thence northward and westward. March–April. — A span high. Tail of carpels 2' long. (Eu. Siberia.)

§ 2. *Carpels very numerous in a dense head, tipped with short and nearly naked styles, thickly clothed with very long and matted wool when ripe.*

* *Low or slender plants, somewhat pubescent, always simple-stemmed, with a mostly sessile 2-3-leaved involucre far below the flower.*

2. **A. Caroliniana**, Walt. (CAROLINA ANEMONE.) Stem 3'-6' high from a round tuber; root-leaves once or twice 3-parted or cleft; involucre 3-parted, its wedge-shaped divisions 3-cleft; sepals 10-20, oblong-linear, purple or whitish; head of fruit oblong. — Illinois (*O. Everett, J. W. Powell, M. S. Bebb, E. Hall, T. J. Hale, &c.*) and southward. May. Apparently passes into the South American *A. decapetala*.

3. **A. parviflora**, Michx. (SMALL-FLOWERED A.) Stem 3'-12' high from a slender rootstock; root-leaves 3-parted, their broadly wedge-shaped divisions crenate-incised or lobed; involucre 2-3-leaved; sepals 5 or 6, oval, white; head of fruit globular. — Lake Superior, northward and westward. May, June.

* * *Taller, commonly branching above or producing two or more peduncles: sepals 5-8, silky or downy beneath (4''-6'' long), oval or oblong.*

4. **A. multifida**, DC. (MANY-CLEFT A.) Silky-hairy (6'-12' high); principal involucre 2-3-leaved, bearing one naked and one or two 2-leaved peduncles; leaves of the involucre short-petioled, similar to the root-leaves, twice or thrice 3-parted and cleft, their divisions linear; sepals 5-8, obtuse, red, sometimes greenish-yellow or whitish; head of fruit spherical or oval. — Rocks, Western Vermont and Northern New York, Lake Superior, &c. rare. June.

5. **A. cylindrica**, Gray. (LONG-FRUITED A.) Slender (2° high), clothed with silky hairs; flowers 2-6, on very long and upright naked peduncles; leaves of the involucre long-petioled, twice or thrice as many as the flower-stalks, 3-divided; their divisions wedge-shaped, the lateral 2-parted, the middle one 3-cleft; lobes cut and toothed at the apex; sepals 5, rather obtuse, greenish-white; head of fruit cylindrical (1' long). — Sandy or dry woods, Massachusetts and Rhode Island to Illinois and northwestward. May. — Peduncles 7'-12' long, all appearing together from the same involucre, and naked throughout, or sometimes part of them with involucels, as in the next.

6. **A. Virginiana**, L. (VIRGINIAN A.) Hairy; principal involucre 3-leaved; the leaves long-petioled, 3-parted; their divisions ovate-lanceolate, pointed, cut-serrate, the lateral 2-parted, the middle 3-cleft; peduncles elongated, the earliest naked, the others with a 2-leaved involucel at the middle; sepals 5, acute, greenish (in one variety white and obtuse); head of fruit oval or oblong. — Woods and meadows; common. June-August. — Plant 2°-3° high; the upright peduncles 6'-12' long. In this and the next species the first flowerstalk is leafless; but from the same involucre soon proceed one or two lateral ones, which are 2-leaved at the middle; these partial involucre in turn giving rise to similar peduncles, thus producing a succession of flowers through the summer.

§ 3. *Carpels fewer, the achænia and the short slender styles merely pubescent.*

7. **A. Pennsylvanica**, L. (PENNSYLVANIAN A.) Hairy, rather low; involucre sessile; the primary ones 3-leaved, bearing a naked peduncle, and soon a pair of branches or peduncles with a 2-leaved involucre at the middle, which branch similarly in turn; their leaves broadly wedge-shaped, 3-cleft, cut and

toothed; radical leaves 5-7-parted or cleft; sepals 5, obovate, white (6"-9" long); head of fruit spherical; the carpels flat, orbicular. — W. New England to Illinois and northwestward. June-Aug.

8. **A. nemorosa**, L. (WIND-FLOWER. WOOD A.) Low, smoothish; stem perfectly simple, from a filiform rootstock, slender, leafless, except the involucre of 3 long-petioled trifoliolate leaves, their leaflets wedge-shaped or oblong, and toothed or cut, or the lateral ones (var. *QUINQUEFOLIA*) 2-parted; a similar radical leaf in sterile plants solitary from the rootstock; peduncle not longer than the involucre: sepals 4-7, oval, white, sometimes tinged with purple outside; carpels only 15-20, oblong, with a hooked beak. — Margin of woods. April, May. — A delicate vernal species; the flower 1' broad. (Eu.)

3. HEPÁTICA, Dill. LIVER-LEAF. HEPATICA.

Involucre simple and 3-leaved, very close to the flower, so as to resemble a calyx; otherwise as in *Anemone* (of which this genus should strictly be viewed as only a section). — Leaves all radical, heart-shaped and 3-lobed, thickish and persistent through the winter, the new ones appearing later than the flowers, which are single, on hairy scapes. (Name from a fancied resemblance to the liver in the shape of the leaves.)

1. **H. triloba**, Chaix. (ROUND-LOBED HEPATICA.) Leaves with 3 ovate obtuse or rounded lobes; those of the involucre also obtuse. — Woods; common eastward; flowering soon after the snow leaves the ground in spring. Sepals 6-9, blue, purplish, or nearly white. Achenia several, in a small loose head, ovate-oblong, pointed, hairy. (Eu.)

2. **H. acutiloba**, DC. (SHARP-LOBED HEPATICA.) Leaves with 3 ovate and pointed lobes, or sometimes 5-lobed; those of the involucre acute or acutish. — Woods, Vermont and New York to Wisconsin. Sepals 7-12, pale purple, pink, or nearly white. Perhaps runs into the other.

4. THALÍCTRUM, Tourn. MEADOW-RUE.

Sepals 4 or more, petal-like or greenish. Petals none. Achenia 4-15, grooved or ribbed, or else inflated. Seed suspended. — Perennials, with 2-3-ternately compound leaves, the divisions and the leaflets stalked. Flowers in corymbs or panicles, often polygamous or dioecious. (Derivation obscure.)

§ 1. **SYNDÉSMON**, Hoffm. *Between Thalicttrum and Anemone, having all its stem-leaves in the form of an involucre at the top, and the stamens shorter than the 5-10 white and conspicuous sepals; but the stigma depressed-truncate, and the ovoid sessile carpels terete, many-angled, with deep intermediate grooves: flowers perfect.*

1. **T. anemonoides**, Michx. (RUE-ANEMONE.) Glabrous; stem and slender petiole of radical leaf (a span high) rising from a cluster of thickened tuberous roots; the latter 2-3-ternately compound; leaflets roundish, somewhat 3-lobed at the end, cordate at the base, long-petiolulate, those of the 2-3-leaved 1-2-ternate involucre similar; flowers several in an umbel; sepals oval ($\frac{1}{2}$ ' long, rarely pinkish) not early deciduous. — Woods, common, flowering in early

spring, along with and considerably resembling *Anemone nemorosa*. Rarely the sepals are 3-lobed like the leaflets.

§ 2. *Leaves alternate along the stem: no involucre: roots fibrous: flowers comparatively small and numerous, paniced: sepals 4 or 5, usually falling early.*

* *Flowers diœcious or sometimes polygamous, in ample panicles: filaments slender: stigmas elongated, linear or subulate, mostly unilateral: achenia sessile or short-stipitate, ovoid, pointed, strongly several-angled and grooved.*

2. **T. dioicum**, L. (EARLY MEADOW-RUE.) Smooth and pale or glaucous, 1°–2° high; *leaves all with general petioles; leaflets drooping*, rounded and 3–7-lobed; flowers purplish and greenish; the yellowish anthers linear, mucronate, drooping on *fine capillary filaments*.—Rocky woods, &c.; common. April, May.

3. **T. purpurascens**, L. (PURPLISH M.) Taller (2°–4° high, the stem usually purplish); *stem-leaves sessile* (without general petiole) or nearly so; leaflets roundish or oblong and more or less 3-lobed, thickish, pale and usually minutely pubescent beneath, the margin mostly revolute and the veining conspicuous; panicles compound; *flowers* (sepals, filaments, &c.) *greenish and purplish*; *anthers* linear or oblong-linear, mucronulate, *drooping on capillary filaments* which are manifestly broadened at the summit. (T. *Virginianum elatius*, &c., *Moris*. T. *rugosum*, *Ait.*? T. *pubescens*, *Pursh*. T. *revolutum*, *DC.*)—Dry uplands and rocky hills, S. New England to Michigan, Illinois and southward. May, June.—Sometimes nearly glabrous throughout, often minutely pubescent, and in

Var. **ceriferum**, C. F. Austin, mss., with the lower surface of the leaves, sepals, and mostly the fruit thickly beset with waxy atoms. Plant often growing with the other, and exhaling a peculiar odor.

4. **T. Cornuti**, L. (TALL M.) Smooth or obscurely pubescent, 4°–8° high; *stem-leaves sessile*; leaflets nearly as in the last, but usually thinner and less revolute and veiny and the lobes more acute; panicles very compound; *flowers white*, the fertile ones with some stamens; *anthers not drooping*, small, oblong, blunt, the *white filaments* decidedly thickened upwards. (T. *rugosum*, *Pursh.*, *DC.* T: *corynellum*, *DC.*)—Wet meadows and along rivulets, common, especially eastward. July–Sept.

* * *Flowers all perfect, corymbed; the filaments strongly club-shaped or inflated under the small and short anther: stigma short and unilateral: achenia long-stipitate.*

5. **T. clavatum**, DC. Size and appearance of No. 2, but leaves only twice ternate; flowers white and fewer; achenia 5–10, flat, somewhat crescent-shaped, tapering into the slender stipe.—Mountains of S. Virginia and southward. June.

5. TRAUTVETTERIA, Fischer & Meyer. FALSE BUGBANE.

Sepals 3–5, usually 4, concave, petal-like, very caducous. Petals none. Achenia numerous, in a head, membranaceous, compressed-4-angled and inflated. Seed erect.—A perennial herb, with palmately-lobed leaves, all alternate, and corymbose white flowers. (Dedicated to *Prof. Trautvetter*, a Russian botanist.)

1. **T. palmata**, Fischer & Meyer. (*Cimicifuga palmata*, Michx.) Woods, along streams, Virginia and Kentucky along the mountains: also sparingly in Ohio and Illinois. July, Aug. — Root-leaves large, 5-9-lobed; the lobes toothed and cut. Stems 2°-3° high.

6. RANUNCULUS, L. CROWFOOT. BUTTERCUP.

Sepals 5. Petals 5, flat, with a little pit or scale at the base inside. Achenia numerous, in a head, mostly flattened, pointed; the seed erect. — Annuals or perennials: stem-leaves alternate. Flowers solitary or somewhat corymbed, yellow, rarely white. (Sepals and petals rarely only 3, the latter often more than 5. Stamens occasionally few in number.) — (A Latin name for a little frog; applied by Pliny to these plants, the aquatic species growing where frogs abound.)

§ 1. **BATRACHIUM**, DC. — *Petals with a spot or naked pit at the base, white, or only the claw yellow: achenia marginless, transversely wrinkled: aquatic perennials, with the immersed foliage repeatedly dissected (mostly by threes) into capillary divisions: peduncles 1-flowered.*

1. **R. divaricatus**, Schrank. (STIFF WATER-CROWFOOT.) *Leaves all under water and sessile or nearly so, the divisions and subdivisions short, spreading in one roundish plane, rigid, keeping their form without collapsing when withdrawn from the water. (R. circinatus, Sibthorp.)* — Ponds and slow streams, northward and westward, much rarer than the next. June - Aug. (Eu.)

2. **R. aquatilis**, L., var. **trichophyllus**, Chaix. (COMMON WHITE WATER-CROWFOOT.) *Leaves all under water and mostly petioled, their capillary divisions and subdivisions rather long and soft, usually collapsing more or less when withdrawn from the water.* — Common, especially in slow-flowing waters. June - Aug. (Eu.)

Var. **heterophyllus**, DC. (FLOATING W.) *Uppermost leaves floating, rounded and 3-5-lobed, the lobes wedge-shaped. (R. aquatilis, Bigel, ed. 3.)* — Roxbury and Newton, near Boston, *Bigelow*; but not met with for many years: was possibly introduced from Europe, where this form is common.

§ 2. **RANUNCULUS** proper. *Petals with a little scale at the base (yellow).*
* *Achenia smooth.*

— *Aquatic, perennial: immersed leaves filiformly dissected, as in the preceding.*

3. **R. multifidus**, Pursh. (YELLOW WATER-CROWFOOT.) *Stems floating or immersed, with the leaves all repeatedly 3-forked into long filiform divisions, or sometimes creeping in the mud, the emersed leaves with shorter and linear or wedge-shaped divisions, or else kidney-shaped and sparingly lobed or toothed; flower deep bright yellow, $\frac{1}{2}$ '-1' in diameter; petals 5-8, much larger than the calyx; carpels in a round head, pointed with a straight beak. (R. lacustris, Beck & Tracy, and R. Purshii, Richards, both in the year 1823. R. Gmelini, DC. (1818) is an older name, belonging to a small northern form of the species; but R. multifidus, Pursh, is the oldest, 1814, and apparently free for use.) Eastern New England to Illinois and northward. May - July. — Out of water it is often pubescent, especially in*

Var. terréstris, which differs from the ordinary emersed forms by the stems ascending from the base and paniculately several-flowered at the summit, where the leaves are reduced to oblong or linear bracts; no immersed dissected leaves. — Ann Arbor, Michigan, on muddy banks, *Miss Clark*.

← ← *Terrestrial but growing in very wet places, glabrous or nearly so: root perennial: leaves all entire or barely toothed, all or else all but the lowest lanceolate or linear; carpels forming a globular head.* (SPEARWORT.)

4. **R. alismæfólius**, Geyer. (WATER PLANTAIN SPEARWORT.) Stems hollow, ascending (1° – 2° high), often rooting from the lower joints; leaves lanceolate or the lowest oblong, mostly denticulate ($3'$ – $5'$ long), contracted into a margined petiole which expands into a membranaceous clasping base; petals 5–7, bright yellow, much longer than the calyx ($3''$ – $4''$ long); *carpels flattened, large, pointed with a long and straight narrow subulate beak.* — Common, especially northward. June–Aug. — Intermediate in appearance between *R. Flammula* and *Lingua*, and has been confounded with both, but most resembles the latter.

5. **R. Flámmula**, L. (SMALLER SPEARWORT.) Stem reclining or ascending, rooting below; leaves lanceolate or linear, or the lowest oblong-lanceolate, entire or nearly so, mostly petioled ($1'$ – $2'$ long); petals 5–7, much longer than the calyx, bright yellow; *carpels flattish but turgid, mucronate with a short abrupt point.* — Shore of L. Ontario and northward: rare, and only a small form (var. *INTERMEDIUS*) met with in this country, a span high, with flowers $3''$ in diameter, passing into

Var. réptans. (CREEPING S.) Small, slender, the *filiform creeping stems rooting at all the joints* ($3'$ – $6'$ long; leaves linear, spatulate, or oblong ($\frac{1}{2}'$ – $1'$ long). — Gravelly or sandy shores and inundated banks; very common northward. June–Sept. (Eu.)

6. **R. oblongifólius**, Ell. Stem erect or ascending, often pubescent below, slender (1° high), *diffusely branched above and many-flowered; leaves serrate or denticulate; the lower long-petioled, ovate or oblong* ($\frac{1}{2}'$ – $1\frac{1}{2}'$ long); the uppermost linear; flowers $3''$ – $5''$ broad; *petals 5, twice the length of the calyx, bright yellow; stamens numerous; carpels minute, almost globular, tipped with a very small sessile stigma.* (*R. pusillus*, var. *Torr. & Gr. Fl. R. Texensis, Engelm.*) Wet prairies, Salem, Illinois, *Bebb*, and in S. States. June.

7. **R. pusillus**, Poir. Stem ascending, weak, loosely branching ($6'$ – $18'$ long); *leaves entire or obscurely denticulate; the lowest round-ovate or heart-shaped* ($\frac{1}{2}'$ long), long-petioled, the upper oblong or lanceolate ($1'$ – $1\frac{1}{2}'$ long); *flowers very small; petals 1–5, yellowish, scarcely exceeding the calyx and the 3–10 stamens; carpels very turgid, tipped with a minute sessile stigma.* — Wet places, S. New York and southward along the coast. June–Aug.

← ← ← *Terrestrial, with annual root, spreading by runners, glabrous: leaves all rounded and undivided but coarsely crenate; carpels in fruit forming an oblong head.*

8. **R. Cymbalària**, Pursh. (SEA-SIDE CROWFOOT.) Flowering stems leafless ($3'$ – $6'$ high), 1–7-flowered; leaves clustered at the root and on the joints of the long rooting runners, roundish-heart-shaped or kidney-shaped,

rather fleshy, long-petioled; petals 5-8; carpels striate on the sides. — Sandy shores, from New Jersey northward, and along the Great Lakes to Illinois and westward: also at salt springs. June-Aug.

+ + + + *Terrestrial, but often in wet places: root perennial: some or all of the leaves cleft or divided.*

++ *Root-leaves not divided to the very base.*

9. **R. rhomboideus**, Goldie. *Dwarf, hairy; root-leaves roundish or rhombic-ovate, rarely subcordate, toothed or crenate; lowest stem-leaves similar or 3-5-lobed; the upper 3-5-parted, almost sessile, the lobes linear; carpels orbicular with a minute beak, in a spherical head; petals large, exceeding the calyx.* (Also *R. brevicaulis* & *ovàlis*, Hook.) — Prairies, Michigan to Illinois and northward. April, May. — Stems 3'-6' high, sometimes not longer than the root-leaves. Flower deep yellow, as large as in No. 14.

10. **R. abortivus**, L. (SMALL-FLOWERED C.) *Glabrous and very smooth; primary root-leaves round heart-shaped or kidney-form, barely crenate, the succeeding ones often 3-lobed or 3-parted; those of the stem and branches 3-5-parted or divided, subsessile; their divisions oblong or narrowly wedge-form, mostly toothed; carpels in a globular head, mucronate with a minute curved beak; petals shorter than the reflexed calyx.* — Shady hillsides and along brooks, common. April-June. — Stem erect, 6'-2° high, at length branched above, the pale yellow flowers very small in proportion.

Var. **micranthus**. *Pubescent; root-leaves seldom at all heart-shaped, some of them 3-parted or 3-divided; divisions of the upper stem-leaves more linear and entire; peduncles more slender.* (*R. micranthus*, Nutt.) — Massachusetts (near Boston, C. J. Sprague), Michigan, Illinois, and westward.

11. **R. sceleratus**, L. (CURSED C.) *Smooth and glabrous; root-leaves 3-lobed, rounded; lower stem-leaves 3-parted, the lobes obtusely cut and toothed, the uppermost almost sessile, with the lobes oblong-linear and nearly entire; carpels barely mucronulate, very numerous, in oblong or cylindrical heads; petals scarcely exceeding the calyx.* — Wet ditches: appearing as if introduced. June-Aug. — Stem thick and hollow, 1° high. Leaves thickish. Juice acrid and blistering. Flowers small, pale yellow. (Eu.)

12. **R. recurvatus**, Poir. (HOOKED C.) *Hirsute; leaves of the root and stem nearly alike, long-petioled, deeply 3-cleft, large; the lobes broadly wedge-shaped, 2-3-cleft, cut and toothed towards the apex; carpels in a globular head, flat and margined, conspicuously beaked by the long and recurved hooked styles; petals shorter than the reflexed calyx, pale.* — Woods, common. May, June. — Stem 1°-2° high.

++ ++ *All the leaves ternately divided to the very base, or compound, and the divisions cleft or cut: achenia flat.*

a. *Head of carpels oblong: petals pale, not exceeding the calyx.*

13. **R. Pennsylvanicus**, L. (BRISTLY C.) *Hirsute with rough spreading bristly hairs; stem stout, erect; divisions of the leaves stalked, somewhat ovate, unequally 3-cleft, sharply cut and toothed, acute; carpels pointed with a sharp straight beak.* — Wet places, common. June-Aug. — A coarse plant, 2°-3° high, with inconspicuous flowers.

b. *Head of carpels globular; petals bright yellow, much larger than the calyx.*

14. **R. fascicularis**, Muhl. (EARLY C.) Low, pubescent with close-pressed silky hairs; root a cluster of thickened fleshy fibres; radical leaves appearing pinnate, the long-stalked terminal division remote from the sessile lateral ones, itself 3-5-divided or parted and 3-5-cleft, the lobes oblong or linear; stems ascending; petals spatulate-oblong, twice the length of the spreading calyx; carpels scarcely margined, tipped with a slender straight or rather curved beak. — Rocky hills. April, May. — Plant 5'-9' high; the bright yellow flower 1' broad: petals rather distant, the base scarcely broader than the scale, often 6 or 7.

15. **R. repens**, L. (CREEPING C.) Low, hairy or nearly glabrous; stems ascending, and some of them forming long runners; leaves 3-divided; the divisions all stalked (or at least the terminal one), broadly wedge-shaped or ovate, unequally 3-cleft or parted and variously cut; peduncles furrowed; petals obovate, much larger than the spreading calyx; carpels strongly margined, pointed by a stout straightish beak. — Moist or shady places, wet meadows, &c., May-Aug. — Extremely variable in size and foliage, commencing to flower by upright stems in spring before the long runners are formed. Flowers as large as those of No. 14, or often larger. (Eu.)

16. **R. bulbosus**, L. (BULBOUS C. BUTTERCUPS.) Hairy; stem erect from a bulb-like base; radical leaves 3-divided; the lateral divisions sessile, the terminal stalked and 3-parted, all wedge-shaped, cleft and toothed; peduncles furrowed; petals round, wedge-shaped at the base, much longer than the reflexed calyx; carpels tipped with a very short beak. — Fields; very abundant only in E. New England; rare in the interior. May-July. — A foot high. Leaves appearing as if pinnate. Petals often 6 or 7, deep glossy yellow, the corolla more than an inch broad. (Nat. from Eu.)

17. **R. acris**, L. (TALL C. or BUTTERCUPS.) Hairy; stem erect (2°-3° high); leaves 3-divided; the divisions all sessile and 3-cleft or parted, their segments cut into lanceolate or linear crowded lobes; peduncles not furrowed; petals obovate, much longer than the spreading calyx. — Fields; common eastward. June-Aug. — Plant twice the height of the preceding; the flower nearly as large, but not so deep yellow. — The *Buttercups* are avoided by cattle, on account of their very acrid or even blistering juice, which, however, being volatile, is dissipated in drying, when these plants are cut with hay. (Nat. from Eu.)

* * *Achenia beset with rough points or small prickles: annuals.*

18. **R. muricatus**, L. Nearly glabrous; lower leaves roundish or reniform, 3-lobed, coarsely crenate; the upper 3-cleft, wedge-form at the base; petals longer than the calyx; carpels flat, spiny-tuberculate on the sides, strongly beaked, surrounded with a wide and sharp smooth margin. — Eastern Virginia and southward. (Nat. from Eu.)

19. **R. parviflorus**, L. Hairy, slender, and diffuse; lower leaves roundish-cordate, 3-cleft, coarsely toothed or cut; the upper 3-5-parted; petals not longer than the calyx; carpels minutely hispid and rough, beaked, narrowly margined. — Norfolk, Virginia, and southward. (Nat. from Eu.)

7. **MYOSÛRUS**, Dill MOUSE-TAIL.

Sepals 5, spurred at the base. Petals 5, small and narrow, raised on a slender claw, at the summit of which is a nectariferous hollow. Stamens 5-20. Achenia numerous, somewhat 3-sided, crowded on a very long and slender spike-like receptacle (whence the name, from $\mu\hat{\upsilon}\varsigma$, a mouse, and $\omicron\upsilon\rho\acute{\alpha}$, a tail), the seed suspended. — Little annuals, with tufted narrowly linear-spatulate root-leaves, and naked 1-flowered scapes. Flowers small, greenish.

1. **M. mínimus**, L. Carpels blunt. — Alluvial ground, Illinois and Kentucky, thence south and west; apparently indigenous. (Eu.)

8. **ISOPÝRUM**, L. (ENÉMIÓN, Raf.)

Sepals 5, petal-like, deciduous. Petals 5, minute, wanting in the American species. Stamens 10-40. Pistils 3-6 or more, pointed with the styles. Pods ovate or oblong, 2-several-seeded. — Slender smooth herbs, with 2-3-ternately compound leaves; the leaflets 2-3-lobed. Flowers axillary and terminal, white. (Name from $\acute{\iota}\sigma\omicron\varsigma$, equal, and $\pi\upsilon\rho\acute{\omicron}\varsigma$, wheat; of no obvious application.)

1. **I. biternátum**, Torr. & Gray. Petals none; pistils 3-6 (commonly 4), divaricate in fruit, 2-3-seeded; seeds even. ♀ — Moist shady places, Ohio to Illinois, Kentucky, and westward. May. — Fibres of the root thickened here and there into little tubers. Aspect and size of the plant much as in *Thalictrum anemonoides*.

9. **CALTHA**, L. MARSH MARIGOLD.

Sepals 5-9, petal-like. Petals none. Pistils 5-10, with scarcely any styles. Pods (follicles) compressed, spreading, many-seeded. Glabrous perennials, with round and heart-shaped or kidney-form, large, undivided leaves. (Name from $\kappa\acute{\alpha}\lambda\alpha\theta\omicron\varsigma$, a goblet, in allusion to the golden flower-cup or calyx.)

1. **C. palústris**, L. (MARSH MARIGOLD.) Stem hollow, furrowed; leaves round or kidney-shaped, either crenate or nearly entire; sepals broadly oval (bright yellow). — Swamps and wet meadows, common northward. April, May. — This well-known plant is used as a pot-herb in spring, when coming into flower, under the name of COWSLIPS; but the Cowslip is a totally different plant, namely, a species of Primrose. The *Caltha* should bear with us, as in England, the popular name of *Marsh Marigold*. (Eu.)

10. **TRÓLLIUS**, L. GLOBE-FLOWER.

Sepals 5-15, petal-like. Petals numerous, small, 1-lipped, the concavity near the base. Stamens and pistils numerous. Pods 9 or more, sessile, many-seeded. — Smooth perennials with palmately parted and cut leaves, like *Ranunculus*, and large solitary terminal flowers. (Name thought to be derived from the old German word *troll*, a globe, or something round.)

1. **T. láxus**, Salisb. (SPREADING GLOBE-FLOWER.) Sepals 5-6, spreading; petals 15-25, inconspicuous, much shorter than the stamens. — Deep swamps, New Hampshire to Delaware and Michigan. May. — Flowers twice the size of the common Buttercup; the sepals spreading, so that the

name is not appropriate, as it is to the *European Globe-flower* of the gardens, nor is the blossom showy, being pale greenish-yellow, or nearly white.

11. CÓPTIS, Salisb. GOLDTHREAD.

Sepals 5-7, petal-like, deciduous. Petals 5-7, small, club-shaped, hollow at the apex. Stamens 15-25. Pistils 3-7, on slender stalks. Pods divergent, membranaceous, pointed with the style, 4-8-seeded.—Low smooth perennials, with ternately divided root-leaves, and small white flowers on scapes. (Name from *κόπτω*, to cut, alluding to the divided leaves.)

1. **C. trifolia, Salisb.** (THREE-LEAVED GOLDTHREAD.) Leaflets 3, obovate-wedge-form, sharply toothed, obscurely 3-lobed; scape 1-flowered.—Bogs, abundant northward; extending south to Maryland along the mountains. May.—Root of long, bright yellow, bitter fibres. Leaves evergreen, shining. Scape naked, slender, 3'-5' high. (Eu.)

12. HELLÉBORUS, L. HELLEBORE.

Sepals 5, petal-like or greenish, persistent. Petals 8-10, very small, tubular, 2-lipped. Pistils 3-10, sessile, forming coriaceous many-seeded pods.—Perennial herbs of the Old World, with ample palmate or pedate leaves, and large, solitary, nodding, early vernal flowers. (Name from *ἔλαιον*, to injure, and *βόσκη*, food, from their well-known poisonous properties.)

1. **H. víridis, L.** (GREEN HELLEBORE.) Root-leaves glabrous, pedate, calyx spreading, greenish.—Near Brooklyn and Jamaica, Long Island, and Bucks Co., Penn., *Martindale*. (Adv. from Eu.)

13. AQUILÈGIA, Tourn. COLUMBINE.

Sepals 5, regular, colored like the petals. Petals 5, all alike, with a short spreading lip, produced backwards into large hollow spurs, much longer than the calyx. Pistils 5, with slender styles. Pods erect, many-seeded.—Perennials, with 2-3-ternately compound leaves, the leaflets lobed. Flowers large and showy, terminating the branches. (Name from *aquila*, an eagle, from some fancied resemblance of the spurs to talons.)

1. **A. Canadénsis, L.** (WILD COLUMBINE.) Spurs nearly straight; stamens and styles longer than the ovate sepals.—Rocks, common. April-June.—Flowers 2' long, scarlet, yellow inside (or rarely all over), nodding, so that the spurs turn upward, but the stalk becomes upright in fruit.—More graceful than the

A. vulgáris, L., the common GARDEN COLUMBINE, of Europe, with hooked spurs, which is beginning to escape from cultivation in some places.

14. DELPHÍNIUM, Tourn. LARKSPUR.

Sepals 5, irregular, petal-like; the upper one prolonged into a spur at the base. Petals 4, irregular, the upper pair continued backwards into long spurs which are enclosed in the spur of the calyx, the lower pair with short claws:

rarely only 2 united into one. Pistils 1-5, forming many-seeded pods in fruit. — Leaves palmately divided or cut. Flowers in terminal racemes. (Name from *Delphin*, in allusion to the shape of the flower, which is sometimes not unlike the classical figures of the dolphin.)

* *Perennials, indigenous; pistils 3-5.*

1. **D. exaltatum**, Ait. (TALL LARKSPUR.) Leaves deeply 3-5-cleft; the divisions narrow wedge-form, diverging, 3-cleft at the apex, acute; *racemes wand-like*, paniced, *many-flowered*; spur straight; *Pods 3, erect*. — Rich soil, Penn. to Michigan and southward. July. — Stem 2°-5° high. Flowers purplish-blue, downy.

2. **D. tricorne**, Michx. (DWARF L.) Leaves deeply 5-parted, their divisions unequally 3-5-cleft; the lobes linear, acutish; *raceme few-flowered, loose*; spur straightish, ascending; *Pods strongly diverging*. — W. Penn. to Illinois and southward. April, May. — Root a tuberous cluster. Stem simple, 6'-12' high. Flowers bright blue, sometimes white.

3. **D. azureum**, Michx. (AZURE L.) Leaves deeply 3-5-parted, the divisions 2-3 times cleft; the lobes all narrowly linear; *raceme strict*; spur ascending, usually curved upwards; *Pods erect*. — Wisconsin, Illinois, and southward. May, June. — Stem 1°-2° high, slender, often softly pubescent. Flowers sky-blue or whitish.

** *Annual, introduced: pistil single.*

4. **D. CONSÓLIDA**, L. (FIELD L.) Leaves dissected into narrow linear lobes; *racemes rather few-flowered, loose*; pedicels shorter than the bracts; *petals all combined into one body*. — Pennsylvania and Virginia, escaped from grain-fields or gardens; and sparingly along roadsides farther north. (Nat. from Eu.)

15. ACONITUM, Tourn. ACONITE. MONKSHOOD. WOLFSBANE.

Sepals 5, petal-like, very irregular; the upper one (helmet) hooded or helmet-shaped, larger than the others. Petals 2 (the 3 lower wanting entirely, or very minute rudiments among the stamens), consisting of small spur-shaped bodies raised on long claws and concealed under the helmet. Pistils 3-5. Pods several-seeded. Seed-coat usually wrinkled or scaly. — Perennials, with palmately cleft or dissected leaves, and showy flowers in racemes or panicles. (The ancient Greek and Latin name, said to be derived from *Acone*, in Bithynia.)

1. **A. uncinatum**, L. (WILD MONKSHOOD.) Glabrous; *stem slender, erect*, but weak and disposed to climb; *leaves deeply 3-5-lobed*, petioled; the lobes ovate-lanceolate, coarsely toothed; *flowers blue*; *helmet erect, obtusely conical*, compressed, slightly pointed or beaked in front. — Rich shady soil along streams, S. W. New York, and southward along the mountains. June-Aug.

2. **A. reclinatum**, Gray. (TRAILING WOLFSBANE.) Glabrous; stems trailing (3°-8° long); *leaves deeply 3-7-cleft*, petioled, the lower orbicular in outline; the divisions wedge-form, incised, often 2-3-lobed; *flowers white*, in very loose panicles; *helmet soon horizontal, elongated-conical*, with a straight beak in front. — Cheat Mountain, Virginia, and southward in the Alleghanies. Aug. — Lower leaves 5'-6' wide. Flowers 9'' long, nearly glabrous.

16. ZANTHORHIZA, Marshall. SHRUB YELLOW-ROOT.

Sepals 5, regular, spreading, deciduous. Petals 5, much smaller than the sepals, concave and obscurely 2-lobed, raised on a claw. Stamens 5 to 10. Pistils 5-15, bearing 2 pendulous ovules. Pods 1-seeded, oblong, the short style becoming dorsal in its growth. — A low shrubby plant; the bark and the long roots deep yellow and bitter. Flowers polygamous, brown purple, in compound drooping racemes, appearing, along with the 1-2-pinnate leaves, from large terminal buds in early spring. (Name compounded of *ξανθός*, *yellow*, and *ρίζα*, *root*.)

1. *Z. apiifolia*, L'Her. — Shady banks of streams, Sherburne, New York, Dr. Douglass, and from the mountains of Pennsylvania southward. — Stems clustered, 1°-2° high. Leaflets cleft and toothed. — The rootstocks of this, and also of the next plant, were used as a yellow dye by the aborigines.

17. HYDRÁSTIS, L. ORANGE-ROOT. YELLOW PUCCOON.

Sepals 3, petal-like, falling away when the flower opens. Petals none. Pistils 12 or more in a head, 2-ovuled: stigma flat, 2-lipped. Ovaries becoming a head of crimson 1-2-seeded berries in fruit. — A low perennial herb, sending up in early spring, from a thick and knotted yellow rootstock, a single radical leaf, and a simple hairy stem, which is 2-leaved near the summit, and terminated by a single greenish-white flower. (Name perhaps from *ἕδωρ*, *water*, and *δράω*, *to act*, alluding to the active properties of the juice.)

1. *H. Canadensis*, L. — Rich woods, New York to Wisconsin and southward: rare. — Leaves rounded, heart-shaped at the base, 5-7-lobed, doubly serrate, veiny, when full grown in summer 4'-9' wide.

18. ACTÆA, L. BANE BERRY.

Sepals 4 or 5, falling off when the flower expands. Petals 4-10, small, flat, spatulate, on slender claws. Stamens numerous, with slender white filaments. Pistil single: stigma sessile, depressed, 2-lobed. Fruit a many-seeded berry. Seeds smooth, flattened, and packed horizontally in 2 rows. — Perennials, with ample 2-3-ternately compound leaves, the ovate leaflets sharply cleft and toothed, and a short and thick terminal raceme of white flowers. (Name from *ἀκτῆ*, the *Elder*, from some resemblance in the leaves.)

1. *A. spicata*, L., var. *rubra*, Michx. (RED BANE BERRY.) *Raceme ovate*; petals rhombic-spatulate, much shorter than the stamens; *pedicels slender*; *berries cherry-red*, oval. (*A. brachypétala*, DC.) — Rich woods, common, especially northward. April, May. — Plant 2° high. (Eu.)

2. *A. alba*, Bigel. (WHITE BANE BERRY.) Taller and rather smoother than the preceding; *raceme oblong*; *petals slender*, mostly truncate at the end, appearing to be transformed stamens; *pedicels thickened in fruit*, as large as the peduncle and red, the globular-oval *berries white*. (*A. spicata*, var. *alba*, Michx., and ed. 2. *A. pachypoda*, Ell.) — Rich woods, flowering a week or two later than the other, and more common westward and southward. — White berries rarely occur with slender pedicels, also red berries with thick pedicels: but these are perhaps the result of crossing.

19. CIMICÍFUGA, L. BUGBANE.

Sepals 4 or 5, falling off soon after the flower expands. Petals, or rather transformed stamens, 1-8, small, on claws, 2-horned at the apex. Stamens as in *Actæa*. Pistils 1-8, forming dry dehiscent pods in fruit. — Perennials, with 2-3-ternately-divided leaves, the leaflets cut-serrate, and white flowers in elongated wand-like racemes. (Name from *cimex*, a bug, and *fugo*, to drive away; the Siberian species being used as a bugbane.)

§ 1. MACRÔTYS, Raf. *Pistil solitary, sometimes 2-3: seeds smooth, flattened and packed horizontally in the pod in two rows, as in Actæa: stigma broad and flat.*

1. *C. racemosa*, Ell. (BLACK SNAKEROOT.) Racemes very long; pods ovoid, sessile. — Rich woods, Maine and Vermont to Wisconsin, and southward. July. — Stem 3°-8° high, from a thick knotted rootstock; the racemes in fruit becoming 1°-3° long.

§ 2. CIMICIFUGA, L. *Pistils 3-8: seeds flattened laterally, covered with chaffy scales, and occupying one row in the membranaceous pods: style awl-shaped: stigma minute.*

2. *C. Americana*, Michx. (AMERICAN BUGBANE.) Racemes slender, paniced; ovaries mostly 5, glabrous; pods stalked, flattened, veiny, 6-8-seeded. — Mountains of Southern Pennsylvania and southward throughout the Alleghanies. Aug.-Sept. — Plant 2°-4° high, more slender than the preceding.

ADONIS AUTUMNALIS, L., the PHEASANT'S EYE of Europe, has been found growing spontaneously in Western New York, and in Kentucky.

NIGÉLLA DAMASCÈNA, L., the FENNEL-FLOWER, which offers a remarkable exception, in having the pistils partly united into a compound ovary, so as to form a several-celled pod, grows nearly spontaneously around gardens.

PÆONIA, the PÆONY, of which *P. OFFICINALIS* is familiar in gardens, forms a sixth tribe of this order, distinguished by a leafy persistent calyx, and a fleshy disk surrounding the base of the follicular pistils.

ORDER 2. MAGNOLIÁCEÆ. (MAGNOLIA FAMILY.)

Trees or shrubs, with the leaf-buds covered by membranous stipules, poly-petalous, hypogynous, polyandrous, polygynous; the calyx and corolla colored alike, in three or more rows of three, and imbricated (rarely convolute) in the bud. — Sepals and petals deciduous. Anthers adnate. Pistils many, mostly packed together and covering the prolonged receptacle, cohering with each other, and in fruit forming a sort of fleshy or dry cone. Seeds 1 or 2 in each carpel, anatropous: albumen fleshy: embryo minute. — Leaves alternate, not toothed, marked with minute transparent dots, feather-veined. Flowers single, large. Bark aromatic and bitter. — There are only two Northern genera, *Magnolia* and *Liriodendron*.

1. MAGNŌLIA, L. MAGNOLIA.

Sepals 3. Petals 6-9. Stamens imbricated, with very short filaments, and long anthers opening inwards. Pistils aggregated on the long receptacle and coherent in a mass, together forming a fleshy and rather woody cone-like red fruit; each carpel at maturity opening on the back, from which the 1 or 2 berry-like seeds hang by an extensile thread composed of unrolled spiral vessels. Inner seed-coat bony. — Buds conical, the coverings formed of the successive pairs of stipules, each pair enveloping the leaf next above, which is folded lengthwise, and applied straight against the side of the next stipular sheath, and so on. (Named after *Magnol*, Professor of Botany at Montpellier in the 17th century.)

* *Leaves all scattered along the branches: leaf-buds silky.*

1. **M. glauca**, L. (SMALL OR LAUREL MAGNOLIA. SWEET BAY.) *Leaves* oblong or oval, obtuse, white beneath; *flower* globular, white, 2' long, very fragrant; petals broad; cone of fruit small, oblong. — Swamps, from near Cape Ann and New York southward, near the coast; in Pennsylvania as far west as Cumberland Co. June-Aug. — Shrub 4°-20° high, with thickish leaves, which farther south are evergreen, and sometimes oblong-lanceolate.

2. **M. acuminata**, L. (CUCUMBER-TREE.) *Leaves* oblong, pointed, green and a little pubescent beneath; *flower* oblong bell-shaped, glaucous-green tinged with yellow, 2' long; cone of fruit small, cylindrical. — Rich woods, W. New York to Ohio and southward. May, June. — Tree 60-90 feet high. Leaves thin, 5'-10' long. Fruit 2'-3' long, when young slightly resembling a small cucumber, whence the common name.

3. **M. macrophylla**, Michx. (GREAT-LEAVED MAGNOLIA.) *Leaves* obovate-oblong, cordate at the narrowed base, pubescent and white beneath; *flower* open bell-shaped, white, with a purple spot at the base, petals ovate, 6' long; cone of fruit ovoid. — Rockcastle and Kentucky Rivers, S. E. Kentucky and southward. Occasionally planted farther north. May, June. — Tree 20°-40° high. Leaves 2½°-3½° long.

** *Leaves crowded on the summit of the flowering branches in an umbrella-like circle: leaf-buds glabrous: flowers white, slightly scented.*

4. **M. Umbrella**, Lam. (UMBRELLA-TREE.) *Leaves* obovate-lanceolate, pointed at both ends, soon glabrous, petals obovate-oblong, 4'-5' long. (*M. tripétala*, L.) — York and Lancaster counties, Penn. (*Prof. Porter*), to Virginia and Kentucky along the Alleghanies. May. — A small tree. Leaves 1°-3° long. Fruit rose-color, 4'-5' long, ovoid-oblong.

5. **M. Fraseri**, Walt. (EAR-LEAVED UMBRELLA-TREE.) *Leaves* oblong-obovate or spatulate, auriculate at the base, glabrous; petals obovate-spatulate, with narrow claws, 4' long. (*M. auriculata*, Lam.) — Virginia and Kentucky along the Alleghanies, and southward. April, May. — Tree 30°-50° high. Leaves 8'-12' long. Flower more graceful and cone of fruit smaller than in the preceding.

M. cordata, Michx., the YELLOW CUCUMBER-TREE, of Georgia, and the

M. grandiflora, L., the GREAT LAUREL MAGNOLIA, of the Southern States (a noble tree, remarkable for its deliciously fragrant great flowers, and

thick evergreen leaves, which are shining and deep green above and rusty-colored beneath), are the only remaining North American species. The former is hardy as far north as Cambridge. One tree of the latter bears the winter and blossoms near Philadelphia. The Umbrella-tree attains only a small size in New England, where *M. macrophylla* is precarious.

2. LIRIODÉNDRON, L. TULIP-TREE.

Sepals 3, reflexed. Petals 6, in two rows, making a bell-shaped corolla. Anthers linear, opening outwards. Pistils flat and scale-form, long and narrow, imbricating and cohering together in an elongated cone, dry, separating from each other and from the long and slender axis in fruit, and falling away whole, like a samara or key, indehiscent, 1-2-seeded in the small cavity at the base. Buds flat, sheathed by the successive pairs of flat and broad stipules joined at their edges, the folded leaves bent down on the petiole so that their apex points to the base of the bud. (Name from *λίριον*, *lily* or *tulip*, and *δένδρον*, *tree*.)

1. **L. Tulipifera**, L. — Rich soil, S. New England to Michigan, Illinois, and southward. May, June. — A most beautiful tree, sometimes 140° high and 8°-9° in diameter in the Western States, where it is wrongly called POPLAR. Leaves very smooth, with 2 lateral lobes near the base, and 2 at the apex, which appears as if cut off abruptly by a broad shallow notch. Petals 2' long, greenish-yellow marked with orange. Cone of fruit 3' long.

ORDER 3. ANONACEÆ. (CUSTARD-APPLE FAMILY.)

Trees or shrubs, with naked buds and no stipules, a calyx of 3 sepals, and a corolla of 6 petals in two rows, valvate in the bud, hypogynous, polyandrous. — Petals thickish. Anthers adnate, opening outwards: filaments very short. Pistils several or many, separate or cohering in a mass, fleshy or pulpy in fruit. Seeds anatropous, large, with a crustaceous seed-coat, and a minute embryo at the base of the *ruminated* albumen. — Leaves alternate, entire, feather-veined. Flowers axillary, solitary. Bark, &c. acrid-aromatic or fetid. — A tropical family, except one genus in the United States, viz.:

1. ASÍMINA, Adans. NORTH AMERICAN PAPAW.

Petals 6, increasing after the bud opens; the outer set larger than the inner. Stamens numerous in a globular mass. Pistils few, ripening 1-4 large and oblong pulpy several-seeded fruits. Seeds horizontal, flat, enclosed in a fleshy aril. — Shrubs or small trees, with unpleasant odor when bruised; the lurid flowers solitary from the axils of last year's leaves. (Name from *Asiminier*, of the French colonists.)

1. **A. triloba**, Dunal. (COMMON PAPAW.) Leaves thin, obovate-lanceolate, pointed; petals dull-purple, veiny, round-ovate, the outer ones 3-4 times as long as the calyx. — Banks of streams in rich soil, W. New York and Penn. to Illinois and southward. April, May. — Tree 10°-20° high; the

young shoots and expanding leaves clothed with a rusty down, soon glabrous. Flowers appearing with the leaves, $1\frac{1}{2}'$ wide. Fruits 3' - 4' long, yellowish, sweet and edible in autumn.

A. *PARVIFLORA*, a smaller-flowered and small-fruited low species, probably does not grow so far north as Virginia.

ORDER 4. MENISPERMACEÆ. (MOONSEED FAMILY.)

Woody climbers, with palmate or peltate alternate leaves, no stipules; the sepals and petals similar, in three or more rows, imbricated in the bud; hypogynous, diœcious, 3 - 6-gynous; fruit a 1-seeded drupe, with a large or long curved embryo in scanty albumen. — Flowers small. Stamens several. Ovaries nearly straight, with the stigma at the apex, but often incurved in fruiting, so that the seed and embryo are bent into a crescent or ring. Properties bitter-tonic and narcotic. — Chiefly a tropical family; there are only three species, belonging to as many genera, in the United States.

1. **Cocculus.** Stamens, petals, and sepals each 6. Anthers 4-celled.

2. **Menispermum.** Stamens 12-24, slender. Petals 6-8. Anthers 4-celled.

3. **Calycoocarpum.** Stamens in the sterile flowers 12; in the fertile flowers 6, abortive. Petals none. Anthers 2-celled.

1. CŒCCULUS, DC. COCCULUS.

Sepals, petals, and stamens 6, the two latter short. Anthers 4-celled. Pistils 3-6 in the fertile flowers: style pointed. Drupe and seed as in Moonseed. Cotyledons narrowly linear and flat. — Flowers in axillary racemes or panicles. (An old name, from *coccum*, a berry.)

1. **C. Carolinus**, DC. Minutely pubescent; leaves downy beneath, ovate or cordate, entire or sinuate-lobed, variable in shape; flowers greenish; the petals in the sterile ones auriculate-inflexed below around the filaments; drupe red (as large as a small pea). — River-banks, S. Illinois, Virginia, and southward. July, Aug.

2. MENISPERMUM, L. MOONSEED.

Sepals 4-8. Petals 6-8, short. Stamens 12-20 in the sterile flowers, as long as the sepals: anthers 4-celled. Pistils 2-4 in the fertile flowers, raised on a short common receptacle: stigma broad and flat. Drupe globular, the mark of the stigma near the base, the ovary in its growth after flowering being strongly incurved, so that the (wrinkled and grooved) laterally flattened stone (putamen) takes the form of a large crescent or a ring. The slender embryo therefore is horseshoe-shaped: cotyledons filiform. — Flowers white, in axillary panicles. (Name from *μήνη*, moon, and *σπέρμα*, seed.)

1. **M. Canadense**, L. (CANADIAN MOONSEED.) Leaves peltate near the edge, 3-7-angled or lobed. — Banks of streams; common. June, July. Drupes black with a bloom, ripe in September, looking like frost grapes.

3. CALYCOCARPUM, Nutt. CUPSEED.

Sepals 6. Petals none. Stamens 12 in the sterile flowers, short: anthers 2-celled. Pistils 3, spindle-shaped, tipped with a radiate many-cleft stigma. Drupe not incurved; but the thin crustaceous putamen hollowed out like a cup on one side. Embryo foliaceous, heart-shaped. — Flowers greenish-white, in long racemose panicles. (Name composed of *κάλυξ*, a cup, and *καρπός*, fruit, from the shape of the shell.)

1. **C. Lyoni**, Nutt. (*Menispermum Lyoni*, *Pursh.*) — Rich soil, S. Kentucky and southward. May. — Stems climbing to the tops of trees. Leaves large, thin, deeply 3–5-lobed, cordate at the base; the lobes acuminate. Drupe an inch long, globular, greenish; the shell crested-toothed on the edge of the cavity.

ORDER 5. BERBERIDACEÆ. (BARBERRY FAMILY.)

Shrubs or herbs, with the sepals and petals both imbricated in the bud in two or more rows of 2–4 each; the hypogynous stamens as many as the petals and opposite to them; anthers opening by 2 valves or lids hinged at the top. (Podophyllum is an exception, and Jeffersonia as respects the sepals in one row.) *Pistil single.* Filaments short. Style short or none. Fruit a berry or a pod. Seeds few or several, anatropous, with albumen. Embryo small, except in Berberis. Leaves alternate.

* Petals and stamens 6. Fruit few-seeded.

1. **Berberis**. Shrubs, with yellow flowers and wood; a pair of glandular spots on the base of each petal. Fruit a berry.
2. **Caulophyllum**. Herb, with greenish flowers: petals thick, much shorter than the sepals. Ovary soon bursting; the two seeds left naked.
3. **Diphyleia**. Herb with white flowers; petals much longer than sepals. Berry 2–4-seeded.

** Petals 6–9 Stamens 8–18. Fruit many-seeded. Herbs.

4. **Jeffersonia**. Petals and stamens usually 8: anthers opening by uplifted valves. Pod opening by a lid.
5. **Podophyllum**. Petals 6–9. Stamens 12–18: anthers not opening by uplifted valves. Fruit a large berry.

1. BÉRBERIS, L. BARBERRY.

Sepals 6, roundish, with 2–6 bractlets outside. Petals 6, obovate, concave, with two glandular spots inside above the short claw. Stamens 6. Stigma circular, depressed. Fruit a 1–few-seeded berry. Seeds erect, with a crustaceous integument. — Shrubs, with yellow wood and inner bark, yellow flowers in drooping racemes, sour berries, and 1–9-foliolate leaves. Stamens irritable. (Derived from *Berbérys*, the Arabic name of the fruit.)

1. **B. vulgàris**, L. (COMMON BARBERRY.) Leaves scattered on the fresh shoots of the season, mostly reduced to sharp triple or branched spines; from which the next season proceed rosettes or fascicles of obovate-oblong closely bristly-toothed leaves (the short petiole jointed!), and drooping *many-flowered racemes*; petals entire; *berries oblong, scarlet.* — Thickets and waste grounds in

E. New England, where it has become thoroughly wild : elsewhere occasionally spontaneous. May, June. (Nat. from Eu.)

2. **B. Canadensis**, Pursh. (AMERICAN BARBERRY.) Leaves repandly toothed, the teeth less bristly-pointed; *racemes few-flowered*; petals notched at the apex; *berries oval* (otherwise as in No. 1). — Alleghanies of Virginia and southward : *not* in Canada. June. — Shrub 1°–3° high.

2. CAULOPHYLLUM, Michx. BLUE COHOSH.

Sepals 6, with 3 small bractlets at the base, ovate-oblong. Petals 6 thick and gland-like somewhat kidney-shaped or hooded bodies, with short claws, much smaller than the sepals, one at the base of each of them. Stamens 6: anthers oblong. Pistil gibbous: style short: stigma minute and unilateral: ovary bursting soon after flowering by the pressure of the 2 erect, enlarging seeds, and withering away; the spherical seeds naked on their thick seed-stalks, looking like drupes; the fleshy integument turning blue: albumen of the texture of horn. — A perennial glabrous herb, with matted knotty rootstocks, sending up in early spring a simple and naked stem, terminated by a small raceme or panicle of yellowish-green flowers, and a little below bearing a large triterately compound leaf, without any common petiole (whence the name, from *καυλός*, *stem*, and *φύλλον*, *leaf*; the stem seeming to form a stalk for the great leaf). Leaflets obovate wedge-form, 2–3-lobed.

1. **C. thalictroides**, Michx. (Also called PAPPOUSE-ROOT.) *Leontice thalictroides*, L. — Deep rich woods; common westward. April, May. — Stems 1°–2½° high. Flowers appearing while the leaf is yet small. A smaller biterminate leaf often at the base of the panicle. Whole plant glaucous when young, also the seeds, which are as large as peas.

3. DIPHYLLÈIA, Michx. UMBRELLA-LEAF.

Sepals 6, fugacious. Petals 6, oval, flat, larger than the sepals. Stamens 6: anthers oblong. Ovary oblong: style hardly any: stigma depressed. Ovules 5 or 6, attached to one side of the cell below the middle. Berry few-seeded. Seeds oblong, with no aril. — A perennial glabrous herb, with thick horizontal rootstocks, sending up each year either a huge centrally peltate and cut-lobed, rounded, umbrella-like, radical leaf, on a stout stalk, or a flowering stem bearing two similar (but smaller and more 2-cleft) alternate leaves which are peltate near one margin, and terminated by a cyme of white flowers. (Name composed of *δίς*, *twice*, and *φύλλον*, *leaf*.)

1. **D. cymosa**, Michx. Wet or springy places, mountains of Virginia and southward. May. — Root-leaves 1°–2° in diameter, 2-cleft, each division 5–7-lobed; lobes toothed. Berries blue.

4. JEFFERSONIA, Barton. TWIN-LEAF.

Sepals 4, fugacious. Petals 8, oblong, flat. Stamens 8: anthers oblong-linear, on slender filaments. Ovary ovoid, soon gibbous, pointed: stigma 2-lobed. Pod pear-shaped, opening half-way round horizontally, the upper part

making a lid. Seeds many in several rows on the lateral placenta, with a fleshy lacerate aril on one side. — A perennial glabrous herb, with matted fibrous roots, long-petioled root-leaves, parted into 2 half-ovate leaflets, and simple naked 1-flowered scapes. (Named in honor of *Thomas Jefferson*.)

1. **J. diphýlla**, Pers. — Woods, W. New York to Wisconsin and southward. April, May. — Low. Flower white, 1' broad: the parts rarely in threes or fives. — Called *Rheumatism-root* in some places.

5. **PODOPHYLLUM**, L. MAY-APPLE. MANDRAKE.

Flower-bud with 3 green bractlets, which early fall away. Sepals 6, fugacious. Petals 6 or 9, obovate. Stamens as many as the petals in the Himalayan species, twice as many in ours: anthers linear-oblong, not opening by uplifted valves. Ovary ovoid: stigma sessile, large, thick, and undulate. Fruit a large fleshy berry. Seeds covering the very large lateral placenta, in many rows, each seed enclosed in a pulpy aril, all forming a mass which fills the cavity of the fruit. — Perennial herbs, with creeping rootstocks and thick fibrous roots. Stems 2-leaved, 1-flowered. (Name from *πούς*, a foot, and *φύλλον*, a leaf, from a fancied resemblance of the 5-7-parted leaf to the foot of some web-footed animal.)

1. **P. peltátum**, L. Stamens 12-18; leaves 5-9-parted; the lobes oblong, rather wedge-shaped, somewhat lobed and toothed at the apex. — Rich woods, common. May. — Flowerless stems terminated by a large round 7-9-lobed leaf, peltate in the middle like an umbrella. Flowering stems bearing two one-sided leaves, with the stalk fixed near their inner edge; the nodding white flower from the fork nearly 2' broad. Fruit ovoid, 1'-2' long, ripe in July, sweet and slightly acid, edible. The leaves and roots are drastic and poisonous! — Found in Ohio, by *W. C. Hampton*, with two carpels!

ORDER 6. **NYMPHÆACEÆ**. (WATER-LILY FAMILY.)

Aquatic perennial herbs, with horizontal rootstocks and peltate or sometimes only cordate leaves floating or emersed; the ovules borne on the sides or back (or when solitary hanging from the summit) of the cells, not on the ventral suture; the embryo enclosed in a little bag at the end of the albumen next the hilum, except in Nelumbium, which has no albumen. Radicle hardly any: cotyledons thick and fleshy, enclosing a well-developed plumule. — Flowers axillary, solitary. Leaves rolled inwards in vernalion. Rootstocks very obscurely exogenous in structure. — Comprises a few genera, which differ so much in the flower and fruit, that, for the sake of convenient definition, we have formerly treated as separate orders the following suborders:

SUBORDER I. **CABÓMBEÆ**. (WATER-SHIELD FAMILY.)

Sepals and petals each 3 or sometimes 4, hypogynous and persistent. Stamens definite (6-18). Pistils 2-18, free and distinct, coriaceous and

indehiscent, 1-3-seeded on the dorsal suture. — Stems slender, leafy, coated with mucilage. Flowers small.

1. **Brasenia.** Stamens 12-18: filaments slender. Leaves all peltate.

SUBORDER II. **NELUMBONÆÆ.** (NELUMBO FAMILY.)

Sepals and petals numerous in several rows, passing gradually into each other, and with the indefinitely numerous stamens hypogynous and deciduous. Pistils several, 1-ovuled, separately immersed in the obconical receptacle, which is much enlarged and broadly top-shaped at maturity, the imbedded nut-like fruits resembling small acorns. Embryo large; no albumen. — Petioles and peduncles all from the tuberous rootstock, the centrally peltate leaves and the flowers large.

2. **Nelumbium.** Character of the Suborder.

SUBORDER III. **NYMPHÆACEÆ** PROPER. (WATER-LILY F.)

Sepals 4-6, and petals numerous in many rows, persistent or decaying away, either hypogynous or variously adnate to the surface of the compound 8-30-celled ovary, which is formed by the union of as many carpels; the numerous ovules inserted over the whole inner face of the cells, except at the ventral suture. Stigmas radiate as in Poppy. Fruit baccate, with a firm rind. Petioles and peduncles from a thick rootstock.

3. **Nymphæa.** Petals adnate to the ovary, large; the stamens on its summit.

4. **Nuphar.** Petals, very small and stamen-like, and stamens inserted under the ovary.

1. **BRASÈNIA,** Schreber. WATER-SHIELD.

Sepals 3 or 4. Petals 3-4, linear, sessile. Stamens 12-18: filaments filiform: anthers innate. Pistils 4-18, forming little club-shaped indehiscent pods: stigma linear. Seeds 1-2, pendulous on the dorsal suture! — Rootstock creeping. Leaves alternate, long-petioled, centrally peltate, oval, floating on the water. Flowers axillary, small, dull-purple. (Name of uncertain origin.)

1. **B. peltata,** Pursh. (*Hydropeltis purpurea, Michx.*) — Ponds and slow streams. June-Aug. — Leaves entire, 2' - 3' across. (Also a native of Puget Sound, Japan, Australia, and Eastern India!)

CABÓMBA, the other genus of the group, occurs from N. Carolina southward.

2. **NELÚMBIUM,** Juss. NELUMBO. SACRED BEAN.

The only genus of the suborder. (*Nelumbo* is the Ceylonese name of the East India species, the pink-flowered *N. speciosum.*)

1. **N. luteum,** Willd. (YELLOW NELUMBO, or WATER CHINQUEPIN.) Corolla pale yellow: anthers tipped with a slender hooked appendage. — Waters of the Western and Southern States; rare in the Middle States: introduced into the Delaware below Philadelphia. Near Woodstown and Swedesboro, New Jersey. Big Sodus Bay, L. Ontario, and in the Connecticut near Lyme; perhaps introduced there by the aborigines. June-Aug. — Leaves usually raised high out of the water, circular in outline, with the centre de-

pressed or cupped, 1° – 2° in diameter. Flower 5'–10' broad. Tubers farinaceous and edible. Seeds also eatable. Embryo like that of *Nymphæa* on a large scale. Cotyledons thick and fleshy, enclosing a plumule of 1 or 2 well-formed young leaves, enclosed in a delicate stipule-like sheath.

3. NYMPHÆA, Tourn. WATER-NYMPH. WATER-LILY.

Sepals 4, green outside, nearly free. Petals numerous, in many rows, the innermost gradually passing into stamens, imbricately inserted all over the surface of the ovary. Stamens indefinite, inserted on the ovary, the outer with dilated filaments. Ovary 18–30-celled, the concave summit tipped with a globular projection at the centre, around which are the radiate stigmas; these project at the margin, and are extended into linear and incurved sterile appendages. Fruit depressed-globular, covered with the bases of the decayed petals, maturing under water. Seeds enveloped by a sac-like aril.—Flowers white, rose-color, or blue, very showy. (Dedicated by the Greeks to the Water-Nymphs.)

1. *N. odorata*, Ait. (SWEET-SCENTED WATER-LILY.) Leaves orbicular, cordate-cleft at the base to the petiole (5'–9' wide), the margin entire; stipules broadly triangular or almost kidney-shaped, notched at the apex, appressed to the rootstock; flower white, very sweet scented (often as much as $5\frac{1}{2}'$ in diameter when fully expanded, opening early in the morning, closing in the afternoon); petals obtuse; aril much longer than the distinctly stipitate oblong seeds (these about $1\frac{1}{2}''$ long; anthers blunt).—Ponds and still or slow-flowing water: common eastward and southward. June–Sept.—Varies with pinkish-tinged and rarely with bright pink-red flowers (especially at Barnstable, Mass.), the leaves often crimson underneath, — and in size by gradations into

Var. *minor*, Sims (*N. minor*, DC., &c.), with leaves only 2'–5' and flowers 2'–3' broad.—Shallow water, in cold bogs and in sandy soil.

2. *N. tuberosa*, Paine, Cat. Pl. Oneida, 1865. (TUBER-BEARING W.) Leaves reniform-orbicular, mostly larger (8'–15' wide) and more prominently ribbed than the last, green both sides; rootstock bearing numerous spontaneously detaching often compound tubers; flower scentless (or with a slight odor as of apples, white, never pinkish, $4\frac{1}{2}'$ –9' in diameter, the petals, as in *N. alba*, proportionally broader and blunter than in No. 1); the fruit more depressed, and with fewer but much larger (i. e. twice as broad) globular-ovoid seeds which when mature are barely enclosed by the aril, and not stipitate. (*N. alba*, Nutt. Gen. *N. reniformis*, DC.? scarcely of Walter, which is very obscure.)—Lakes, slow rivers, &c., W. New York (from Oneida Lake, Paine) and near Meadville, Penn. (*W. L. Chaffin*) to Michigan, Illinois, and probably in the Southern States. July–Sept.

4. NÛPHAR, Smith. YELLOW POND-LILY. SPATTER-DOCK.

Sepals 5, 6, or sometimes more, colored, or partly green outside, roundish. Petals numerous, small and thickish, stamen-like or scale-like, inserted with the very numerous and short stamens on the receptacle under the ovary, not

surpassing the disk-like 8-24-rayed sessile stigma. Fruit ovoid, naked, usually ripening above water. Aril none.—Leaves with a deep sinus at the base. Flowers yellow or sometimes tinged with purple, produced all summer. (*Noûφap* of Dioscorides, from the Egyptian name.)—Our various forms seem to include only two species.

1. **N. ádvena**, Ait. (COMMON Y.) *Sepals 6, unequal; petals shorter than the stamens* and resembling them, thick and fleshy, truncate; stigma 12-24-rayed; ovary and fruit not contracted into a narrow neck under the stigma; thin submersed leaves seldom appearing; floating or emersed and erect leaves thick, varying from roundish to ovate or almost oblong in outline, the sinus open, or (var. *VARIÉGATUM*, Engelm., flower often partly purplish) closed or narrow.—Very common, in still or stagnant water.

2. **N. lûteum**, Smith. (SMALLER Y.) *Sepals 5, nearly equal; petals longer and dilated upwards*; stigma 12-16-rayed; fruit globular, with a short narrow neck; earlier and submersed leaves very thin and delicate, roundish, the floating ones oval and usually with a narrow or closed sinus.—The only specimen seen like the European (expanded flower fully 2' across) is from "Mannyoung, 7 miles from Philadelphia," in herb. *Collins*, now *Durand*. (Eu.)

Var. **pûmilum**. (SMALL Y.) Flower $1\frac{1}{2}$ ' - 1" across when outspread; leaves 1' - 5' long. (*N. pumilum*, *Hoppe*. *N. Kalmiana*, *Pursh*.)—Ponds, N. England to Penn. and northward. (Eu.)

N. POLYSÉPALUM, Engelm., with very large flowers and numerous sepals, occurs far west.

N. SAGITTIFÓLIA, Pursh, of N. Carolina and southward, has narrow and long leaves. Both perhaps run into No. 1.

ORDER 7. SARRACENIACEÆ. (PITCHER-PLANTS.)

Polyandrous and hypogynous bog-plants, with hollow pitcher-form or trumpet-shaped leaves,—comprising one plant in the mountains of Guiana, another (*Darlingtonia*, *Torr.*) in California, and the following genus in the Atlantic United States.

1. SARRACÉNIA, Tourn. SIDE-SADDLE FLOWER.

Sepals 5, with 3 bractlets at the base, colored, persistent. Petals 5, oblong or obovate, incurved, deciduous. Stamens numerous, hypogynous. Ovary compound, 5-celled, globose, crowned with a short style, which is expanded at the summit into a very broad and petal-like, 5-angled, 5-rayed, umbrella-shaped body; the 5 delicate rays terminating under the angles in as many little hooked stigmas. Capsule with a granular surface, 5-celled, with many-seeded placentæ in the axis, 5-valved. Seeds anatropous, with a small embryo at the base of fleshy albumen.—Perennials, yellowish-green and purplish; the hollow leaves all radical, with a wing on one side, and a rounded arching hood at the apex. Scape naked, 1-flowered: flower nodding. (Named by Tournefort in honor of *Dr. Sarrazin* of Quebec, who first sent our Northern species, and a botanical account of it, to Europe.)

1. **S. purpurea**, L. (SIDE-SADDLE FLOWER. PITCHER-PLANT. HUNTS-MAN'S CUP.) *Leaves pitcher-shaped*, ascending, curved, broadly winged; the hood erect, open, round heart-shaped; *flower deep purple*; the fiddle-shaped petals arched over the (greenish-yellow) style. — Varies rarely with greenish-yellow flowers, and without purple veins in the foliage. (*S. heterophylla*, *Eaton*.) — Peat-bogs; common from N. England to Minnesota, N. Illinois, and southward east of the Alleghanies. June. — The curious leaves are usually half filled with water and drowned insects: the inner face of the hood is clothed with stiff bristles pointing downward. Flower globose, nodding on a scape a foot high: it is difficult to fancy any resemblance between its shape and a side-saddle, but it is not very unlike a pillion.

2. **S. flava**, L. (TRUMPETS.) *Leaves long* (1° – 3°) *and trumpet-shaped*, erect, with an open mouth, the erect hood rounded, narrow at the base; wing almost none; *flower yellow*, the petals becoming long and drooping. — Bogs, Virginia and southward. April.

ORDER 8. PAPAVERACEÆ. (POPPY FAMILY.)

Herbs with milky or colored juice, regular flowers with the parts in twos or fours, fugacious sepals, polyandrous, hypogynous, the ovary 1-celled with 2 or more parietal placentæ. — Sepals 2, rarely 3, falling when the flower expands. Petals 4–12, spreading, imbricated and often crumpled in the bud, early deciduous. Stamens rarely as few as 16, distinct. Fruit a dry 1-celled pod (in the Poppy imperfectly many-celled, in *Glaucium* 2-celled). Seeds numerous, anatropous, often crested, with a minute embryo at the base of fleshy and oily albumen. — Leaves alternate, without stipules. Peduncles mostly 1-flowered. Juice narcotic or acrid.

* Ovary incompletely several-celled by the projecting placentæ.

1. **Papaver**. Stigmas united into a radiate crown on the summit of the ovary. Pod opening by chinks or pores. Petals crumpled in the nodding flower-bud.

** Ovary strictly 1-celled. Pod opening by valves, leaving the 2–6 filiform parietal placentæ as a framework.

— Petals twice as many as the sepals, usually more or less crumpled in the flower-bud.

2. **Argemone**. Stigmas (sessile) and placentæ 4–6. Pod and leaves prickly.

3. **Stylophorum**. Stigmas and placentæ 3–4. Style distinct. Pod bristly.

4. **Chelidonium**. Stigmas and placentæ 2. Pod linear, smooth.

— Petals 4–6 times as many as the 2 sepals, not crumpled in the erect flower-bud.

5. **Sanguinaria**. Stigma 2-grooved. Placentæ 2. Petals white.

*** Ovary and elongated pod 2-celled by a spongy partition.

6. **Glaucium**. Stigma 2-lobed or 2-horned. Petals 4.

1. PAPAVER, L. POPPY.

Sepals mostly 2. Petals mostly 4. Stigmas united in a flat 4–20-rayed crown, resting on the summit of the ovary and capsule; the latter short and turgid, with 4–20 many-seeded placentæ projecting like imperfect partitions, opening by as many pores or chinks under the edge of the stigma. — Herbs

with a white juice; the flower-buds nodding. (Derivation obscure.)—Three annual species of the Old World are sparingly adventive; viz.:

1. **P. SOMNIFERUM**, L. (COMMON POPPY.) *Smooth*, glaucous; leaves clasping, wavy, incised and toothed; *pod globose*; corolla mostly white or purple.—Near dwellings in some places. (Adv. from Eu.)

2. **P. DUBIUM**, L. (SMOOTH-FRUITED CORN-POPPY.) Pinnatifid leaves and the long stalks *bristly*; *Pods club-shaped, smooth*; corolla light scarlet.—Cult. grounds, Westchester, Penn. and southward: rare. (Adv. from Eu.)

3. **P. ARGEMONE**, L. (ROUGH-FRUITED C.) Smaller, with finer-cut leaves and paler flowers than the last; *Pods club-shaped and bristly*.—Waste grounds, near Philadelphia, *Dr. Dieffenbaugh*. (Adv. from Eu.)

2. ARGEMONE, L. PRICKLY POPPY.

Sepals 2 or 3, often prickly. Petals 4–6. Style almost none: stigmas 3–6, radiate. Pod oblong, prickly, opening by 3–6 valves at the top. Seeds crested.—Annuals or biennials, with prickly bristles and yellow juice. Leaves sessile, sinuate-lobed, and with prickly teeth, often blotched with white. Flower-buds erect, short-peduncled. (Name from *ἀργέμα*, a disease of the eye, for which the juice was a supposed remedy.)

1. **A. MEXICANA**, L. (MEXICAN P.) Flowers yellow, rarely white.—Waste places, southward. July–Oct. (Adv. from trop. Amer.)

3. STYLÓPHORUM, Nutt. CELANDINE POPPY.

Sepals 2, hairy. Petals 4. Style distinct, columnar: stigma 2–4-lobed. Pods bristly, 2–4-valved to the base. Seeds conspicuously crested.—Perennial low herbs, with stems naked below and oppositely 2-leaved, or sometimes 1–3-leaved, and umbellately 1–few-flowered at the summit; the flower-buds and the pods nodding. Leaves pinnately parted or divided. Juice yellow. (Name from *στυλος*, *style*, and *φέρω*, *I bear*, indicating one of the distinctive characters.)

1. **S. diphyllum**, Nutt. Leaves pale or glaucous beneath, smoothish, deeply pinnatifid into 5 or 7 oblong sinuate-lobed divisions, and the root-leaves often with a pair of smaller and distinct leaflets; peduncles equalling the petioles; flower deep yellow (2' broad); stigmas 3 or 4; pod oval.—Damp woods, W. Penn. to Wisconsin, and southward. May.—Foliage and flower resembling Celandine.

4. CHELIDONIUM, L. CELANDINE.

Sepals 2. Petals 4. Stamens 16–24. Style nearly none: stigma 2-lobed. Pod linear, slender, smooth, 2-valved, the valves opening from the bottom upwards. Seeds crested.—Perennial herb with brittle stems, saffron-colored acrid juice, pinnately divided or 2-pinnatifid and toothed or cut leaves, and small yellow flowers in a pedunculate umbel; the buds nodding. (Name from *χελιδών*, the *Swallow*, because, according to Dioscorides, it begins to flower at the time the swallows appear.)

1. **C. MAJUS**, L. (CELANDINE.) Waste grounds near dwellings. May-Aug. (Adv. from Eu.)

5. **SANGUINÀRIA**, Dill. BLOOD-ROOT.

Sepals 2. Petals 8-12, spatulate-oblong, the inner narrower. Stamens about 24. Style short; stigma 2-grooved. Pod oblong, turgid, 1-celled, 2-valved. Seeds with a large crest.—A low perennial, with thick prostrate rootstocks, surcharged with red-orange acrid juice, sending up in earliest spring a rounded palmate-lobed leaf, and a 1-flowered naked scape. Flower white, handsome, the bud erect, the petals not crumpled. (Name from the color of the juice.)

1. **S. Canadensis**, L. — Open rich woods; common. April, May.

6. **GLAÛCIUM**, Tourn. HORN-POPPY.

Sepals 2. Petals 4. Style none: stigma 2-lobed or 2-horned. Pod very long and linear, completely 2-celled by a spongy false partition; seeds crestless. — Annuals or biennials, with saffron-colored juice, clasping leaves, and solitary yellow flowers. (The Greek name, *γλαύκιον*, from the glaucous foliage.)

1. **G. LUTEUM**, Scop. Lower leaves pinnatifid; upper ones sinuate-lobed and toothed, cordate-clasping; pods rough, (6' - 10' long). — Waste places S. E. New England, Maryland, and Virginia; not common. (Adv. from Eu.)

ORDER 9. **FUMARIACEÆ**. (FUMITORY FAMILY.)

Delicate smooth herbs, with watery juice, compound dissected leaves, irregular flowers, with 4 somewhat united petals, 6 diadelphous stamens, and 2-merous pods and seeds like those of the Poppy Family. — Sepals 2, small and scale-like. Corolla flattened, closed; the 4 petals in two pairs; the outer with spreading tips, and one or both of them spurred or saccate at the base; the inner pair narrower, and with their callous crested tips united over the stigma. Stamens in two sets of 3 each, placed opposite the larger petals, hypogynous; their filaments often united; the middle anther of each set 2-celled, the lateral ones 1-celled. Pod 1-celled, either 1-seeded and indehiscent, or several-seeded with 2 parietal placentæ. — Leaves usually alternate, without stipules. (Slightly bitter, innocent plants.)

* Pod slender: the 2 valves separating from the persistent filiform placentæ.

1. **Adlumia**. Corolla heart-shaped, persistent; petals all united. Seeds crestless.

2. **Dicentra**. Corolla heart-shaped or 2-spurred at the base. Seeds crested.

3. **Corydalis**. Corolla 1-spurred at the base. Seeds crested or arilled.

** Pod fleshy, indehiscent, globular, 1-seeded.

4. **Fumaria**. Corolla 1-spurred at the base. Seed crestless.

1. **ADLÛMIA**, Raf. CLIMBING FUMITORY.

Petals all permanently united in a cordate-ovate corolla, becoming spongy-cellular and persistent, enclosing the small, few-seeded pod. Seeds not crested.

Stigma 2-crested. Filaments monadelphous below in a tube which is adherent to the corolla, diadelphous at the summit. — A climbing biennial, with thrice-pinnate leaves, cut-lobed delicate leaflets, and ample panicles of drooping white or purplish flowers. (Dedicated by Rafinesque to *Major Adlum*.)

1. **A. cirrhosa**, Raf. — Wet woods; common westward. June–Oct. — A handsome vine, with delicate foliage, climbing by the slender young leaf-stalks over high bushes; cultivated for festoons and bowers in shaded places.

2. DICÉNTRA, Bork. DUTCHMAN'S BREECHES.

Petals slightly cohering into a heart-shaped or 2-spurred corolla, either deciduous or withering-persistent. Stigma 2-crested and sometimes 2-horned. Filaments slightly united in two sets. Pod 10–20-seeded. Seeds crested. — Low, stemless perennials (as to our wild species) with ternately compound and dissected leaves, and racemose nodding flowers. Pedicels 2-bracted. (Name from *dis*, twice, and *κέντρον*, a spur; — accidentally printed DICLÝTRA in the first instance, which by an erroneous conjecture was afterwards changed into DIÉLYTRA.)

1. **D. Cucullària**, DC. (DUTCHMAN'S BREECHES.) Scape and slender-petioled leaves from a sort of *granulate bulb*; lobes of the leaves linear; *raceme simple*, few-flowered; *corolla with 2 divergent spurs* longer than the pedicel; *crest of the inner petals minute*. — Rich woods, especially westward. — A very delicate plant, sending up in early spring, from the cluster of grain-like tubers crowded together in the form of a scaly bulb, the finely cut leaves and the slender scape, bearing 4–10 pretty, but odd, white flowers tipped with cream-color.

2. **D. Canadénsis**, DC. (SQUIREL CORN.) Subterranean shoots bearing scattered *grain-like tubers* (resembling peas or grains of Indian corn, yellow); leaves and raceme as in No. 1; *corolla merely heart-shaped*, the spurs very short and rounded; *crest of the inner petals conspicuous, projecting*. — Rich woods, Maine to Wisconsin and Kentucky, especially northward. April, May. — Flowers greenish-white tinged with rose, with the fragrance of Hyacinths.

3. **D. eximia**, DC. Subterranean shoots *scaly*; divisions and lobes of the leaves broadly oblong; *raceme compound, clustered*; *corolla oblong, 2-saccate* at the base; *crest of the inner petals projecting*. — Rocks, W. New York, rare (*Thomas, Sartwell*), and Alleghanies of Virginia. May–Aug. — Coarser-leaved than the others; scapes 6'–10' high.

3. CORÝDALIS, Vent. CORYDALIS.

Corolla 1-spurred at the base (on the upper side), deciduous. Style persistent. Pod many-seeded. Seeds crested or arilled. Flowers in racemes. Our species are biennial, leafy-stemmed, and pale or glaucous. (The ancient Greek name for the Fumitory.)

1. **C. glauca**, Pursh. (PALE CORYDALIS.) *Stem upright*; racemes panicled; spur of the *purplish and yellow-tipped corolla* very short and rounded; *pods erect*, slender, elongated. — Rocky places: common: 6'–2° high. May–Aug.

2. **C. flavula**, Raf. *Corolla pale-yellow, 3''–4'' long*; *spur very short*; *tips of the outer petals pointed, wing-crested* on the back, longer than the inner; *seeds*

acutely margined, rugose-reticulated; aril loose: otherwise as in the next. — Pennsylvania to Wisconsin and southward.

3. **C. aurea**, Willd. (GOLDEN C.) *Stems low or decumbent*; racemes simple; *corolla golden-yellow*, $\frac{1}{2}$ ' long; slightly decurved *spur somewhat shorter than the pedicel*, not half the length of the rest of the flower; *tips of outer petals blunt, crestless and naked* on the back, little longer than the inner; pods usually pendent; *seeds smooth* and even, or sometimes very minutely reticulated, *turgid, marginless*, partly covered by the scale-shaped aril. — Rocky places, Vermont to Penn., Wisconsin, and northward. April–July. — Var. **MICRANTHA**, Engelm., is a state with minute *spurless* flowers (probably fertilized in the bud), and ascending pods, on very short pedicels. — W. Illinois and St. Louis, *Riehl*.

(**C. MONTANA**, Engelm., or perhaps rather **C. AUREA**, var. **OCCIDENTALIS**, Engelm., Missouri to Texas and westward, differs from the Eastern **C. aurea** in the often ascending spur nearly equalling the rest of the corolla and longer than the pedicel, erect or ascending pods, and seeds lenticular with acutish margin. — **C. CURVISLIQUA**, Engelm., of Southwest, differs from this in longer 4-angular pods ascending on very short pedicels, the acute-margined seed *muricate!* — **C. CRYSTALLINA**, Engelm., of Southwest, differs from this in the very broad but short wing on tip of outer petals, short few-seeded pod covered with crystalline vesicles, and sharper-margined tubercular-reticulated seeds.)

4. **FUMARIA**, L. FUMITORY.

Corolla 1-spurred at the base. Style deciduous. Fruit indehiscent, small, globular, 1-seeded. Seeds crestless. — Branched and leafy-stemmed annuals, with finely dissected compound leaves, and small flowers in dense racemes or spikes. (Name from *fumus*, smoke.)

1. **F. OFFICINALIS**, L. (COMMON FUMITORY.) Sepals ovate-lanceolate, acute, sharply toothed, narrower and shorter than the corolla (which is flesh-color tipped with crimson); fruit slightly notched. — Waste places, about dwellings. (Adv. from Eu.)

ORDER 10. **CRUCIFERÆ**. (MUSTARD FAMILY.)

Herbs, with a pungent watery juice and cruciform tetradynamous flowers: fruit a silique or silicle. — Sepals 4, deciduous. Petals 4, hypogynous, regular, placed opposite each other in pairs, their spreading limbs forming a cross. Stamens 6, two of them inserted lower down and shorter (rarely only 4 or 2). Pod 2-celled by a thin partition stretched between the two marginal placentæ, from which when ripe the valves separate, either much longer than broad (a *silique*), or short (a *silicle* or *pouch*), sometimes indehiscent and nut-like (*nucumentaceous*), or separating across into 1-seeded joints (*lomenteaceous*). Seeds campylotropous, without albumen, filled by the large embryo, which is curved or folded in various ways: i. e. the *cotyledons accumbent*, viz. their margins on one side applied to the radicle, so that the cross-section of the seed appears thus \propto ;

or else *incumbent*, viz. the back of one cotyledon applied to the radicle, thus \ominus . In these cases the cotyledons are plane; but they may be folded upon themselves and round the radicle, as in Mustard, where they are conduplicate, thus \odot . In *Leavenworthia* alone the whole embryo is straight. — Leaves alternate, no stipules. Flowers in terminal racemes or corymbs: pedicels rarely bracted. — A large and very natural family, of pungent or acrid, but not poisonous plants. (Characters of genera taken from the pods and seeds; the flowers being nearly alike in all.)

I. SILIQUOSÆ. Pod long, a *siliqua*, opening by valves.

Tribe I. ARABIDÆÆ. Pod elongated (except in *Nasturtium*). Seeds orbicular or oval, more or less flattened. Cotyledons accumbent, plane.

* Seeds small and turgid, seldom half the breadth of the turgid pod.

1. ***Nasturtium*.** Pod terete, linear, oblong or even globular. Flowers white or yellow.
* * Seeds flat or flattish, scarcely narrower than the partition, except in some of No. 5.
2. ***Leavenworthia*.** Pod oblong, flat; the valves nerveless. Seeds winged: embryo straight! Flowers white or purplish, with a yellowish base. Leaves all radical.
3. ***Dentaria*.** Pod flat, lanceolate; the valves nerveless, opening elastically from the base. Seeds wingless, on broad seedstalks. Flowers white or purple. Stem naked below.
4. ***Cardamine*.** Pod flat, linear or lanceolate; the valves nerveless, opening elastically. Seeds wingless, on slender stalks. Flowers white or purple. Stem leafy, at least below.
5. ***Arabis*.** Pod linear, elongated, flat or flattish, rarely almost terete; the valves commonly with a prominent midrib, or veiny, not opening elastically. Flowers white, whitish, or purple. Stems leafy, at least below.
6. ***Barbarea*.** Pod linear, more or less 4-sided, the rigid valves being keeled by a prominent midrib. Seeds wingless. Flowers yellow.

Tribe II. SISYMBRIÆÆ. Pod elongated. Seeds thickish, mostly oblong. Cotyledons incumbent, narrow, plane.

7. ***Erysimum*.** Pod sharply 4-angled, linear. Flowers yellow.
8. ***Sisymbrium*.** Pods terete, 4-6-angled, or flattish. Flowers white or yellow.

Tribe III. BRASSICÆÆ. Pod linear or oblong. Seeds globular. Cotyledons incumbent and conduplicate, folded round the radicle.

9. ***Brassica* (including *Sinapis*).** Pod beaked or pointed beyond the end of the valves, or tipped with a rigid style, nearly terete, or 4-sided. Flowers yellow or whitish.

II. SILICULOSÆ. Pod short, a *silicle* or pouch, opening by valves.

Tribe IV. ALYSSINÆÆ. Pod oval or oblong, flattened parallel to the broad partition, or globular. Cotyledons accumbent, plane

1. ***Nasturtium*.** Pod terete or globular with many small seeds.
10. ***Draba*.** Pod flat, oval, oblong, or even linear, many-seeded.
11. ***Alyssum*.** Pod flat, orbicular, 2-4-seeded.
12. ***Vesicaria*.** Pod globular-inflated, 4-several-seeded.

Tribe V. CAMELINÆÆ. Pod turgid or somewhat flattened parallel with the broad partition. Cotyledons incumbent, narrow.

13. ***Camellina*.** Pod pear-shaped, many-seeded: style slender. Flowers yellow.
14. ***Subularia*.** Pod globular, few-seeded: style none. Flowers white.

Tribe VI. LEPIDINÆÆ and THLASPIDÆÆ. Pod short; the boat-shaped valves conduplicate or much flattened contrary to the narrow partition. Flowers white.

15. ***Capsella*.** Pod many-seeded, obcordate-triangular, wingless. Cotyledons incumbent.
16. ***Thlaspi*.** Pod several-seeded, obovate or obcordate, winged. Cotyledons accumbent.
17. ***Lepidium*.** Pod 2-seeded, flat, scale-shaped. Cotyledons incumbent or accumbent.
18. ***Senebiera*.** Pod 2-seeded, didymous; the valves rugose, separating at maturity from the little partition as two closed one-seeded nutlets. Cotyledons incumbent, narrow.

III. LOMENTACEÆ. Pod articulated, separating across into joints.

Tribe VII. CAKILINEÆ. Cotyledons plane and accumbent, as in Tribe 1.

19. **Cakile.** Pod short, 2-jointed: the joints 1-celled and 1-seeded.

Tribe VIII. RAPHANÆÆ. Cotyledons conduplicate and incumbent, as in Tribe 3.

20. **Raphanus.** Pod elongated, several-seeded, transversely intercepted.

1. NASTÚRTIUM, R. Br. WATER-CRESS.

Pod a short silique or a silicle, varying from oblong-linear to globular, terete or nearly so. Seeds small, turgid, marginless, in 2 irregular rows in each cell (except in No. 2). Cotyledons accumbent. — Aquatic or marsh plants, with yellow or white flowers, and commonly pinnate or pinnatifid leaves, usually glabrous. (Name from *Nasus tortus*, a convulsed nose, alluding to the effect of its pungent qualities.)

§ 1. *Petals white, twice the length of the calyx: pods linear: leaves pinnate.*

1. **N. OFFICINALE**, R. Br. (TRUE WATER-CRESS.) Stems spreading and rooting; leaflets 3–11, roundish or oblong, nearly entire; pods (6"–8" long) ascending on slender widely spreading pedicels. ♀ — Brooks and ditches: escaped from cultivation. (Nat. from Eu.)

§ 2. *Petals yellow or yellowish, seldom much exceeding the calyx: pods linear, oblong, or even ovoid or globular: leaves mostly pinnatifid.*

* *Perennial from creeping or subterranean shoots: flowers rather large, bright yellow.*

2. **N. SYLVÉSTRE**, R. Br. (YELLOW CRESS.) Stems ascending; leaves pinnately parted, the divisions toothed or cut, lanceolate or linear; pods ($\frac{1}{2}$ ' long) on slender pedicels, linear and narrow, bringing the seeds into one row; style very short. — Wet meadows, Massachusetts to Virginia: rare. (Nat. from Eu.)

3. **N. sinuatum**, Nutt. Stems low, diffuse; leaves pinnately cleft, the short lobes nearly entire, linear-oblong; pods linear-oblong (4"–6" long), on slender pedicels; style slender. — Banks of the Mississippi and westward. June.

* * *Annual or biennial, rarely perennial? with simple fibrous roots: flowers small or minute, greenish or yellowish: leaves somewhat lyrate.*

4. **N. sessiliflorum**, Nutt. Stems erect, rather simple; leaves obtusely incised or toothed, obovate or oblong; flowers minute, nearly sessile; pods elongated-oblong (5"–6" long), thick; style very short. — W. Illinois to Tennessee and southward. April–June.

5. **N. obtusum**, Nutt. Stems much branched, diffusely spreading; leaves pinnately parted or divided, the divisions roundish and obtusely toothed or repand; flowers minute, short-pedicelled; pods longer than the pedicels, varying from linear-oblong to short-oval; style short. — With No. 3 and 4.

7. **N. palústre**, DC. (MARSH CRESS.) Stem erect; leaves pinnately cleft or parted, or the upper lacinate; the lobes oblong, cut-toothed; pedicels about as long as the small flowers and mostly longer than the oblong, ellipsoid, or ovoid pods; style short. — Wet places or in shallow water; common. June–Sept. — Flowers only 1"–1½" long. Stems 1°–3° high. — The typical form with oblong pods is rare (W. New York, *Dr. Sartwell*). Short pods and hirsute stems and leaves are common. Var. **HISPIDUM** (*N. hispidum*, DC.) is a form, with ovoid or globular pods. (Eu.)

§ 3. *Petals white, much longer than the calyx: pods ovoid or globular: leaves undivided, or the lower ones pinnatifid: root perennial.* (Armoracia.)

7. **N. lacustre**, Gray, Gen. Ill. 1, p. 132. (LAKE CRESS.) Aquatic; immersed leaves 1-3-pinnately dissected into numerous capillary divisions; emerged leaves oblong, entire, serrate, or pinnatifid; pedicels widely spreading; *pods ovoid, one-celled, a little longer than the style.* (N. natans, var. Americanum, Gray. Armoracia Americana, Arn.)—Lakes and rivers, N. E. New York to Illinois and southwestward. July - Aug.—Near N. amphibium.

8. **N. ARMORACIA**, Fries. (HORSE RADISH.) Root-leaves very large, oblong, crenate, rarely pinnatifid; those of the stem lanceolate; fruiting pedicels ascending; *pods globular* (seldom formed); *style very short.* (Cochlearia Armoracia, L.)—Roots large and long;—a well-known condiment. Escaped from cultivation into moist ground. (Adv. from Eu.)

2. LEAVENWORTHIA, Torr. LEAVENWORTHIA.

Pod broadly linear or oblong, flat; the valves nerveless, but minutely reticulate-veined. Seeds in a single row in each cell, flat, surrounded by a wing. Embryo straight! or the short radicle only slightly bent in the direction which if continued would make the orbicular cotyledons accumbent.—Little biennial or hyemal annuals, glabrous and stemless, with lyrate root-leaves and short one-few-flowered scapes. (Named in honor of the late M. C. Leavenworth.)

1. **L. Michauxii**, Torr.—On flat rocks and barrens, S. E. Kentucky and southwestward: rare. March - May.—Scapes 2'-4' high. Petals purple, rose color, or nearly white, with a golden yellow or yellowish base, or rarely yellow throughout? (L. aurea, Torr.), cuneate-obcordate or emarginate; the flowers rather large for the size of the plant.

3. DENTARIA, L. TOOTHWORT. PEPPER-ROOT.

Pod lanceolate, flat, as in Cardamine, but broader. Seedstalks broad and flat. Cotyledons petioled, their margins somewhat infolding each other.—Perennials, with long, horizontal, fleshy, sometimes interrupted, scaly or toothed rootstocks, of a pleasant pungent taste; the simple stems leafless below, bearing 2 or 3 petioled compound leaves about the middle, and terminated by a single corymb or short raceme of large white or purple flowers. (Name from dens, a tooth.)

1. **D. diphylla**, L. *Rootstock long and continuous, toothed; stem-leaves 2, similar to the radical ones, close together, of 3 rhombic-ovate coarsely toothed leaflets; petals white.*—Rich woods, Maine to Kentucky. May.—Rootstocks 5'-10' long, crisp, tasting like Water-Cress.

2. **D. máxima**, Nutt. *Rootstock interrupted, forming a string of thick toothed tubers; stem-leaves (2-7) mostly 3 and alternate; leaflets 3, ovate, obtuse, coarsely toothed and incised, often 2-3-cleft; petals pink.* (D. laciniata, var. δ , Torr. & Gr.)—Northern New York (Watertown, Crave; Utica, Paine) to Penn.: rare. May.—Seldom taller, but the leaves often smaller, than in No. 1.

3. **D. heterophylla**, Nutt. *Rootstock a chain of 2 or 3 narrow-oblong and obscurely toothed tubers; stem-leaves 2 or 3, alternate (rarely opposite), divided into*

3 lanceolate or linear-oblong somewhat toothed (sometimes incised or 2-cleft) leaflets; root-leaves of 3 rounded or cuneate-ovate incised leaflets; petals purple. — Penn. (near Philadelphia, *Dr. Diffenbaugh*) and Washington (*Bebb*) to Kentucky and southward. April, May.

4. **D. laciniata**, Muhl. Rootstock as in the last; stem-leaves 3 in a whorl, 3-parted; the leaflets linear or lanceolate, irregularly cut or cleft into prominent narrow teeth, the lateral ones deeply 2-lobed; root-leaves similarly dissected; petals pale purple or nearly white. — Rich soil along streams, W. New England to Wisconsin and Kentucky. April, May.

5. **D. multifida**, Muhl. More slender and delicate than the last; rootstock similar; stem-leaves 3 or sometimes 2, alternate, opposite, or whorled, finely 2-3-ternately divided, or the subdivisions parted, into very narrowly linear lobes; root-leaves similarly 3-4-ternately divided; flowers (4-7) white. — Southern Kentucky (doubtless) and southward. — This appears to pass into No. 4; and all these species except the first probably run together.

4. CARDÁMINE, L. BITTER CRESS.

Pod linear, flattened, usually opening elastically from the base; the valves nerveless and veinless, or nearly so. Seeds in a single row in each cell, wingless; their stalks slender. Cotyledons accumbent. — Flowers white or purple. (A Greek name, in Dioscorides, for some Cress, from the cordial or cardiacal qualities.) — Runs into *Dentaria* on the one hand, into *Arabis* on the other.

* Root perennial: leaves simple or 3-foliolate.

1. **C. rhomboidea**, DC. (SPRING CRESS.) Stems upright from a tuberiferous base, simple; root-leaves round and rather heart-shaped; lower stem-leaves ovate or rhombic-oblong, somewhat petioled, the upper almost lanceolate, all somewhat angled or sparingly toothed; pods linear-lanceolate, pointed with a slender style tipped with a conspicuous stigma; seeds round-oval. — Wet meadows and springs; common. Flowers large, white. April-June.

Var. **purpurea**, Torr. Lower (4'-6' high), and slightly pubescent; leaves rounder; flowers rose-purple, appearing earlier. — Along streams in rich soil, W. New York and mountains of Penn. to Wisconsin and northward.

2. **C. rotundifolia**, Michx. (MOUNTAIN WATER-CRESS.) Stems branching, weak or decumbent, making long runners; root fibrous; leaves all much alike, roundish, somewhat angled, often heart-shaped at the base, petioled, the lowest frequently 3-lobed or of 3 leaflets; pods linear-awl-shaped, pointed with the style; stigma minute; seeds oval-oblong. — Cool shaded springs, Pennsylvania and southward along the mountains. May, June. — Leaves with just the taste of the true Water-Cress. Flowers white, smaller than in No. 1.

3. **C. bellidifolia**, L. Dwarf (2'-3' high), tufted; leaves ovate, entire, or sometimes 3-lobed (4" long), on long petioles; pods upright, linear; style nearly none. — Alpine summits of the White Mountains and Katahdin, Maine (*J. W. Chickering*). July. — Flowers 1-5, white. Pods 1' long, turgid, the convex valves 1-nerved. (Eu.)

* * Root perennial: leaves pinnate: flowers showy.

4. **C. pratensis**, L. (CUCKOO FLOWER.) Stem ascending from a short

rootstock, simple; leaflets 7-13, those of the lower leaves rounded and stalked; of the upper ones oblong or linear, entire, or slightly angled-toothed; petals (white or rose-color) thrice the length of the calyx. — Wet places and bogs, Vermont to New Jersey, Wisconsin, and northward: rare. May. (Eu.)

* * * *Root mostly biennial or annual: leaves pinnate: flowers small, white.*

5. **C. hirsuta**, L. (SMALL BITTER CRESS.) Beset with scattered hairs, or glabrous; stems (3'-12' high) erect or ascending from the spreading cluster of root-leaves; their leaflets rounded, those of the upper leaves oblong or linear and often confluent, all either toothed, angled, or entire; pods linear, slender, erect or ascending in line with the pedicel; style very short or almost none. (C. *Pennsylvanica*, *Muhl.*; usually taller and more leafy-stemmed than the true European *C. hirsuta*, the stamens always 6, and the pods less erect.) — Wet places: common. May-July. (Eu.)

Var. **sylvatica** (C. *sylvatica*, *Link.*, C. *Virginica*, *Michx.*) grows in drier places, is more slender, and has pods more erect than their ascending or spreading pedicels, the style evident. — Commoner southward. (Eu.)

5. **ÁRABIS**, L. ROCK CRESS.

Pod linear, flattened; the valves plane or convex, more or less 1-nerved in the middle, or longitudinally veiny. Seeds usually margined or winged. Cotyledons accumbent. — Leaves undivided. Flowers white or purple. (Name from the country, *Arabia*. See *Linn. Phil. Bot.* § 235.)

§ 1. *Seeds in one row in each cell, being nearly as broad as the partition.*

* *Low, chiefly biennials, diffuse or spreading from the base.*

1. **A. Ludoviciana**, Meyer. Nearly glabrous, often annual; leaves all pinnately parted into oblong or linear few-toothed or entire divisions, those of the lower leaves numerous; flowers small, white; pods rather broadly linear and spreading, flat; seeds wing-margined. (Cardamine *Ludoviciana*, *Hook.*) — Open grounds, Virginia? to Illinois and southward.

2. **A. lyrata**, L. *Root biennial*; plant mostly glabrous, except the *lyrate-pinnatifid* root-leaves, stem-leaves scattered, spatulate or linear with a tapering base, sparingly toothed or entire; petals white, much longer than the yellowish calyx; pods long and slender, flat, ascending or spreading; the seeds marginless. — On rocks, New England to Kentucky along the mountains, Minnesota and northward. April-July. — Radicle sometimes oblique, or even dorsal.

A. **Thaliana**, L., resembles the last, but the root-leaves are hardly if at all lyrate, the stem more strict, flowers smaller, and the cotyledons uniformly incumbent; so it is referred to *Sisymbrium*, p. 70.

3. **A. petraea**, Lam. *Root perennial, multicapital*; leaves sparingly pinnatifid-toothed or incised, sometimes entire; petals rose-color or nearly white; pods shorter and less flat than in *A. lyrata*: otherwise similar. — Rocks, L. Superior? Willoughby Mountain, Vermont, *H. Mann*.

4. **A. dentata**, Torr. & Gray. Roughish-pubescent, slender (1°-2° high); leaves oblong, very obtuse, unequally and sharply toothed; those of the stem numerous, half-clasping and eared at the base, of the root broader and tapering into a short petiole; petals (whitish) scarcely exceeding the calyx,

Pods widely spreading, very slender, short-stalked; style scarcely any; seeds marginless. — New York and Illinois to Virginia and Kentucky. May, June.

* * *Erect and simple leafy-stemmed biennials, with white or whitish flowers, narrow but flattened ascending or erect pods, and wingless seeds.*

5. **A. patens**, Sulliv. Downy with spreading hairs, erect (1° – 2° high); stem-leaves oblong-ovate, acutish, coarsely toothed or the uppermost entire, partly clasping by the heart-shaped base; petals (bright white, $4''$ long) twice the length of the calyx; *pedicels slender, spreading; pods spreading or ascending, tipped with a distinct style.* — Central Ohio (rocky banks of the Scioto, *Sullivant*), Pennsylvania (Huntingdon Co. to the Schuylkill, *Porter*); also in E. Tennessee. April, May.

6. **A. hirsuta**, Scop. Rough-hairy, sometimes smoothish, strictly erect (1° – 2° high); stem-leaves oblong or lanceolate, entire or toothed, partly clasping by a somewhat arrow-shaped or heart-shaped base; petals (greenish-white) small, but longer than the calyx; *pedicels and pods strictly upright; style scarcely any.* — Rocks, common, especially northward. May, June. (Eu.)

* * * *Erect and simple leafy-stemmed biennials (1° – 3° high), with small whitish flowers, recurved-spreading or pendulous flat pods ($3'$ – $4'$ long), and broadly winged seeds, their stalks adherent to the partition.*

7. **A. lævigata**, DC. Smooth and glaucous, upright; *stem-leaves partly clasping* by the arrow-shaped base, lanceolate or linear, sparingly cut-toothed or entire; petals scarcely longer than the calyx; *Pods long and narrow, recurved-spreading* on ascending or merely spreading pedicels. (This is also *A. heterophylla*, *Nutt.*) — Rocky places, Maine to Wisconsin and Kentucky. May.

8. **A. Canadensis**, L. (SICKLE-POD.) Stem upright, smooth above; *stem-leaves pubescent, pointed at both ends*, oblong-lanceolate, sessile, the lower toothed; petals twice the length of the calyx, oblong-linear; *Pods very flat, scythe-shaped, hanging* on rough-hairy pedicels ($2''$ wide). — (*A. falcata*, *Michx.*) Woods and ravines; not rare, especially westward. June–Aug.

* * * * *Tall and leafy-stemmed biennials or perennials, with rather showy flowers, the pink-purple petals long-clawed; anthers sagittate and when old arcuate-recurved; the widely spreading and rather short pods nearly truncate; seeds marginless.* (IODANTHUS, *Torr. & Gray.*) — Transition to *Thelypodium* and *Streptanthus*.

9. **A. hesperidoides**. Glabrous (1° – 3° high), often branched above; root-leaves round or heart-shaped, on slender petioles; stem-leaves ovate-oblong and ovate-lanceolate ($2'$ – $6'$ long), membranaceous, veiny, sharply and often doubly toothed, tapering to each end, the lower into a winged petiole, sometimes bearing a pair or two of small lateral lobes; pods on short diverging pedicels, pointed by a short style. (*Hesperis pinnatifida*, *Michx.* *Iodanthus hesperidoides*, *Torr. & Gray.*) — Alluvial river-banks, Ohio, Kentucky and southwestward. May, June.

§ 2. **TURRITIS**, Dill. *Seeds not so broad as the partition, occupying two more or less distinct rows in each cell, at least when young: strict and very leafy-stemmed biennials; the cauline leaves partly clasping by a sagittate base.* (Very gla-

brous in the following species, except the base of the stem and the lower or tuft of radical leaves, these mostly hirsute.)

10. **A. perfoliata**, Lam. (TOWER MUSTARD.) Tall (2° – 4° high); glaucous; stem-leaves oblong or ovate-lanceolate, entire, half-clasping by the sagittate base; petals yellowish-white, little longer than the calyx; pods very narrow (3' long) and pedicels strictly erect. (*Turritis glabra*, L.)—Rocks and fields, scarce and perhaps introduced southward; more common northward. (Eu.)

11. **A. Drummóndii**, Gray. Scarcely glaucous, 1° – 2° high; stem-leaves lanceolate or oblong-linear and sagittate (1'–2' long) with narrow auricles, or the lowest spatulate; petals white or rose-color, fully twice the length of the calyx; pedicels and flat pods loosely erect, or ascending, or even spreading; seeds wing-margined, when mature little narrower than the partition. (*Turritis stricta*, Graham.)—Rocky places, from the St. Lawrence in Canada East, to Lewiston (Clinton), Lake Superior, and northwestward; also "Chenango Co. New York," Northern Illinois, Vasey. — Pods $2\frac{1}{2}'$ – $3\frac{1}{2}'$ long, or in var. (*T. brachycarpa*, Torr. & Gray) only 1'–2' long.

6. BARBARÈA, R. Br. WINTER CRESS.

Pod linear, terete or somewhat 4-sided; the valves being keeled by a mid-nerve. Seeds in a single row in each cell, marginless. Cotyledons accumbent. — Mostly biennials resembling *Nasturtium*; flowers yellow. (Anciently called The Herb of St. Barbara.)

1. **B. vulgaris**, R. Br. (COMMON WINTER CRESS. YELLOW ROCKET.) Smooth; lower leaves lyrate, the terminal division round and usually large, the lateral 1–4 pairs or rarely wanting; upper leaves obovate, cut-toothed, or pin-natifid at the base; pods erect or slightly spreading; or in var. *stricta*, appressed; in var. *arcuata*, ascending on spreading pedicels. — Low grounds and roadsides: apparently introduced, but indigenous from L. Superior northward and westward. (Eu.)

2. **B. præcox**, R. Br. (EARLY WINTER C.), with 5–8 pairs of lateral lobes to the leaves, and longer pods on very thick pedicels, — yet probably only a variety of the other, — somewhat cultivated from New York southward as a winter salad, under the name of SCURVY-GRASS, — is beginning to run wild. (Eu.)

7. ERÝSIMUM, L. TREACLE MUSTARD.

Pod linear, 4-sided; the valves keeled with a strong midrib. Seeds in a single row in each cell, oblong, marginless. Cotyledons (often obliquely) incumbent. Calyx erect. — Chiefly biennials, with yellow flowers; the leaves not clasping. (Name from *ἐρύω*, to draw blisters.)

1. **E. cheiranthoides**, L. (WORM-SEED MUSTARD.) Minutely roughish, branching, slender; leaves lanceolate, scarcely toothed; flowers small; pods small and short (7"–12" long), very obtusely angled, ascending on slender divergent pedicels. — Banks of streams, New York, Penn., Illinois, and northward. July. (Eu.)

2. **E. ásperum**, DC., var. **Arkansánum**, Nutt. (WESTERN WALL-FLOWER.) Minutely roughish-hoary; stem simple; leaves lanceolate, some-

what toothed; *Pods nearly erect on very short pedicels, elongated (3'–4' long), exactly 4-sided; stigma 2-lobed.* — Ohio (on limestone cliffs) to Illinois, and southwestward. June, July. — Plant stout, 1°–2° high; the crowded bright orange-yellow flowers as large as those of the Wall-flower. Petals orbicular, on very slender claws.

8. SISÝMBRIUM, L. HEDGE MUSTARD.

Pod terete, flattish, or 4–6-sided; the valves 1–3-nerved. Seeds oblong, marginless. Cotyledons incumbent. Calyx open. — Flowers small, white or yellow. (An ancient Greek name for some plant of this family.) Ours are annuals or biennials.

1. **S. OFFICINALE**, Scop. (HEDGE MUSTARD.) *Leaves runcinate; flowers very small, pale yellow; pods awl-shaped, close pressed to the stem, scarcely stalked.* — Waste places. May–Sept. — An unsightly branched weed, 2°–3° high. (Nat. from Eu.)

2. **S. THALIANA**, Gaud. (MOUSE-EAR CRESS.) *Leaves obovate or oblong, entire or barely toothed; flowers white; pods linear, somewhat 4-sided, longer than the slender spreading pedicels.* (*Arabis Thaliana*, L.; the plant resembles *A. lyrata*.) — Old fields and rocks, Massachusetts to Kentucky. April, May. — A span high, slender, branched, hairy at the base. (Nat. from Eu.)

3. **S. CANÉSCENS**, Nutt. (TANSY MUSTARD.) *Leaves 2-pinnatifid, often hoary or downy, the divisions small and toothed; flowers yellowish, very small; pods in long racemes, oblong-club-shaped or oblong-linear, shorter than their mostly horizontal pedicels; seeds 2-ranked in each cell.* — Penn. and New York (Lucifer Falls, Tompkins Co., *J. W. Chickering*) to Lake Superior, thence southward and westward. June–Aug.

S. SOPHIA, L., with slender linear pods nearly erect on ascending pedicels, and one-ranked seeds is nat. from Eu. in Canada East. — **S. INCISUM**, Engelm., differing only in the shorter pods widely spreading on horizontal pedicels, is wild beyond the Mississippi.

9. BRÁSSICA (Brassica and Sinàpis), Tourn.

Pod linear or oblong, nearly terete or 4-sided, with a stout 1-seeded beak or a rigid style; valves 1–5-nerved. Seeds globose, one-rowed. Cotyledons incumbent, folded around the radicle. — Annuals or biennials, with yellow flowers. Lower leaves mostly lyrate, incised, or pinnatifid. (The Latin name of Cabbage. *Sinapis* is the Greek *Σίναπι*, which is said to come from the Celtic *nap*, a turnip.)

1. **B. SINAPSTRUM**, Boissier (or **SINAPIS ARVÉNSIS**, L., the English CHARLOCK), with knotty pods, fully one third occupied by a stout 2-edged beak (which is either empty or one-seeded), the upper leaves barely toothed, is a noxious weed in grain-fields, from Pennsylvania and New York westward. (Adv. from Eu.)

2. **B.** (or **SINÀPIS**) **ÁLBA**. (WHITE MUSTARD.) Pods bristly, ascending on spreading pedicels, more than half its length occupied by the sword-shaped one-seeded beak; leaves all pinnatifid; seeds pale. (Cult. and adv. from Eu.)

3. **B.** (or **SINÀPIS**) **NÌGRA.** (BLACK MUSTARD.) Pods smooth ($\frac{1}{2}$ long), 4-cornered (the valves only 1-nerved), erect on appressed pedicels forming a slender raceme, tipped with a stout persistent style; seeds dark brown, smaller and more pungent than in the last; lower leaves with a large terminal lobe and a few small lateral ones.—Fields and waste places, or cultivated. (Adv. from Eu.)

B. CAMPÈSTRIS, L., in the form of the **RUTABAGA** and the **TURNIP**, sometimes persists a year or two in neglected grounds.

10. **DRÀBA, L.** WHITLOW-GRASS.

Pouch oval, oblong, or even linear, flat; the valves plane or slightly convex; the partition broad. Seeds several or numerous, in 2 rows in each cell, marginless. Cotyledons accumbent. Calyx equal. Filaments not toothed.—Low herbs with entire or toothed leaves, and white or yellow flowers; the pubescence often stellate. (Name from $\delta\rho\acute{\alpha}\beta\eta$, *acid*, in allusion to the pungency of the leaves.)

§ 1. **DRABA, DC.** *Petals not notched nor cleft.*

* *Perennial or biennial, leafy-stemmed: flowers white: pods twisted when ripe.*

1. **D. ramosissima, Desv.** *Diffusely much branched* and forming many radical tufts, perennial (5'–8' high), pubescent; leaves laciniate-toothed, linear-lanceolate, the lower oblanceolate; racemes corymbosely-branched; pods hairy, oval-oblong or lanceolate (2''–5'' long), on slender spreading pedicels, tipped with a long style.—Cliffs, Harper's Ferry, Natural Bridge, &c., Virginia to Kentucky River, and southward. April, May.

2. **D. arábisans, Michx.** Slightly pubescent, the perennial root bearing rather numerous radical tufts; flowering stems (6'–10' high) *erect and mostly simple*; leaves oblong-lanceolate, linear, or the lower spatulate, *sparingly toothed*; racemes short, usually simple; pods glabrous, oblong-lanceolate (5'–6' long), *acute*, on rather short and spreading pedicels, *pointed with a short but distinct style*.—Rocky banks, N. Vermont and New York towards the St. Lawrence, also Akron, Ohio (*Clinton*), and shores of L. Huron and L. Superior. May, June.—Petals rather large. Too near some forms of the next.

3. **D. incana, L.** *Hoary-pubescent*, biennial or somewhat perennial, the radical tuft seldom branching; leaves shorter, raceme more strict, *petals smaller*, and *Pods shorter and blunter* than in the last, *often pubescent*, on short erect pedicels; style very short or none.—Dry rocks, Willoughby Mountain, Vermont, *Tuckerman, H. Mann.*; also high northward. (Eu.)

* * *Annual or biennial: leafy stems short: flowers white, or in No. 5 yellow: style none. (Leaves oblong or obovate, hairy, sessile.)*

4. **D. brachycarpa, Nutt.** Low (2'–4' high), minutely pubescent; stems leafy to the base of the dense at length elongated raceme; leaves narrowly oblong or the lowest ovate (2''–4'' long), few toothed or entire; flowers small; pods smooth, narrowly oblong, *acutish* (2'' long), *about the length of the ascending or spreading pedicels*.—Dry hills, Illinois, Kentucky, and southward. April.—Petals sometimes minute, sometimes none.

5. **D. nemorosa**, L. Leaves oblong or somewhat lanceolate, more or less toothed; *racemes elongated* (4' - 8' long in fruit); petals emarginate, small; *Pods elliptical-oblong, half the length of the horizontal or widely-spreading pedicels*, pubescent (*D. nemoralis*, Ehrh.), or smooth (*D. lutea*, DC.). — Fort Gratiot, Michigan, and northward. (Eu.)

6. **D. cuneifolia**, Nutt. Leaves obovate, wedge-shaped, or the lowest spatulate, toothed; *raceme somewhat elongated in fruit* (1' - 3'), at length equaling the naked peduncle; petals emarginate, much longer than the calyx; *Pods oblong-linear, minutely hairy, longer than the horizontal pedicels*. — Grassy places, Illinois, Kentucky, and southward. March, April.

7. **D. Caroliniana**, Walt. Small (1' - 5' high); leaves obovate, mostly entire; peduncles scape-like; petals usually twice the length of the calyx; *raceme short or corymbose in fruit* ($\frac{1}{2}$ ' - 1' long); *Pods broadly linear, smooth, much longer than the ascending pedicels*. — Sandy and waste fields, Rhode Island to Wisconsin, and southward. March - May. — Petals often wanting in the later racemes, especially in the

Var. **micrantha** (*D. micrantha*, Nutt.), with minutely rough-hairy pods. With the other, westward, *Bebb*, &c.

§ 2. ERÓPHILA, DC. *Petals 2-cleft. (Annual or biennial: flowers white.)*

8. **D. vérna**, L. (WHITLOW-GRASS.) Small (scapes 1' - 3' high); leaves all radical, oblong or lanceolate; racemes elongated in fruit; pods varying from round-oval to oblong-lanceolate, smooth, shorter than the pedicels. — Sandy waste places and roadsides. April, May. — Not found north of Lower Canada: perhaps introduced. (Eu.)

11. ALÝSSUM, Tourn. ALYSSUM.

Like Vesicaria but with a flat pouch: only one or two seeds in a cell: flowers yellow or white. Filaments often toothed. (Greek name of a plant reputed to check the hiccup, as the etymology denotes.) They are plants of the Old World, two adventive species deserving a mere mention, and one indigenous, rare and doubtful.

1. **A. MARÍTIMUM**, L. (SWEET ALYSSUM), with green or slightly hoary linear leaves, honey-scented small white flowers, and 2-seeded pods, commonly cult., begins to be spontaneous southward. (Adv. from Eu.)

2. **A. CALYCÍNUM**, L., a dwarf hoary annual, with linear-spatulate leaves, pale yellow or whitish petals little exceeding the persistent calyx, and orbicular sharp-margined 4-seeded pod, the style minute, occurs in grass-land at Amherst, Mass., *Tuckerman*. (Adv. from Eu.)

3. **A. Lescúrii**. (*Vesicaria? Lescurii*, ed. 2.) Somewhat pubescent, but green; stems diffusely ascending from a biennial root: *leaves* oblong or oval, sparingly toothed, those of the stem *half-clasping by a sagittate base*; racemes elongated, many-flowered; pedicels ascending; filaments inflated at the base; *style half the length of the hispid orbicular or broadly oval flat pod; seeds wing-margined, 1-4 in each cell, on free stalks*. — Hills near Nashville, Tennessee, *Leo Lesquereux*. April, May. — Petals golden yellow, rather large. Ambiguous between this genus and the next.

12. VESICÀRIA, Tourn. BLADDER-POD.

Pouch globular or inflated, with a broad mostly orbicular partition; the hemispherical or convex thin valves nerveless. Seeds few or several, flat. Cotyledons accumbent. Filaments toothless. — Low herbs, pubescent or hoary with stellate hairs. Flowers mostly yellow. (Name from *vesica*, a bladder.)

1. *V. Shórtii*, Torr. & Gray. Minutely hoary all over; stems spreading or decumbent from an annual or biennial root; leaves oblong or lanceolate with a tapering base, repand-toothed or nearly entire; raceme at length elongated, with filiform diverging pedicels; petals light yellow; style filiform, much longer than the small globose about 4-seeded pod; seeds marginless. — Rocky banks of Elkhorn Creek near Lexington, Kentucky (*Short*), and Kentucky River near Frankfort, *Lesquereux*. May, June.

13. CAMÉLINA, Crantz. FALSE FLAX.

Pouch obovoid or pear-shaped, pointed, turgid, flattish parallel to the broad partition: valves 1-nerved. Seeds numerous oblong. Cotyledons incumbent. Style slender. Flowers small, yellow. (Name from *χαμαί*, *dwarf*, and *λίνον*, *flax*. It has been fancied to be a sort of degenerate flax.)

1. *C. SATIVA*, Crantz. Annual; leaves lanceolate and arrow-shaped; pods margined, large. A weed in flax-fields, &c. (Adv. from Eu.)

14. SUBULÀRIA, L. AWLWORT.

Pouch ovoid or globular, with a broad partition; the turgid valves 1-nerved. Seeds several. Cotyledons long and narrow, incumbently folded transversely, i. e. the cleft extending to the radicular side of the curvature. Style none. — A dwarf stemless perennial, aquatic; the tufted leaves awl-shaped (whence the name). Scape naked, few-flowered, 1'–3' high. Flowers minute, white.

1. *S. aquática*, L. — Margin of lakes in Maine, *Nuttall*, &c. Echo Lake, Franconia, New Hampshire, *Tuckerman*. June, July. (Eu.)

15. CAPSÉLLA, Vent. SHEPHERD'S PURSE.

Pouch obcordate-triangular, flattened contrary to the narrow partition; the valves boat-shaped, wingless. Seeds numerous. Cotyledons incumbent. — Annuals: flowers small, white. (Name a diminutive of *capsula*, a pod.)

1. *C. BURSA-PASTÓRIS*, Mœnch. Root-leaves clustered, pinnatifid or toothed; stem-leaves arrow-shaped, sessile. — Waste places; the commonest of weeds. April–Sept. (Nat. from Eu.)

16. THLÁSPI, Tourn. PENNYCRESS.

Pouch orbicular, obovate, or obcordate, flattened contrary to the narrow partition, the midrib or keel of the boat-shaped valves extended into a wing. Seeds 2–8 in each cell. Cotyledons accumbent. Petals equal. — Low plants, with root-leaves undivided, stem-leaves arrow-shaped and clasping, and small white or purplish flowers. (Ancient Greek name, from *θλάω*, to *crush*, of a Cress the seeds of which were bruised and used like Mustard.)

1. **T. ARVENSE**, L. (FIELD P. or MITHRIDATE MUSTARD.) A smooth annual, with broadly winged pod $\frac{1}{2}$ ' in diameter, several-seeded, deeply notched at top; style minute. — Waste places, shore of Lake Huron and in Lower Canada; also Virginia. (Nat. from Eu.)

17. LEPIDIUM, L. PEPPERWORT. PEPPERGRASS.

Pouch roundish, much flattened contrary to the narrow partition; the valves boat-shaped and keeled. Seeds solitary in each cell, pendulous. Cotyledons incumbent, or in No. 1 accumbent! Flowers small, white or greenish. (Name from *λεπίδιον*, a little scale, alluding to the small flat pods.) — Ours are annuals or biennials, except the last.

* *Leaves all with a tapering base; the upper linear or lanceolate and entire, the lower and often the middle ones incised or pinnatifid: pods orbicular or oval, with a small notch at the top: the style minute or none: stamens only 2.*

1. **L. Virginicum**, L. (WILD PEPPERGRASS.) *Cotyledons accumbent* and seed minutely margined; *pod marginless* or obscurely margined at the top; petals present, except in some of the later flowers. — June–Sept. A common roadside weed, which has immigrated from farther South.

2. **L. intermedium**, Gray. *Cotyledons incumbent* as in the following; *pod minutely wing-margined at the top*; petals sometimes conspicuous, rarely wanting; otherwise nearly as in No. 1. — Dry places, from Northern Michigan and Illinois northward and westward.

3. **L. RUDERÀLE**, L. More diffuse, the smaller and oval *pods and the seeds marginless*; *petals always wanting*. — Roadsides, near Boston, Philadelphia, &c.; not common. (Adv. from Eu.)

* * *Stem-leaves with a sagittate partly clasping base, rather crowded.*

4. **L. CAMPÈSTRE**, L. Minutely soft downy; leaves arrow-shaped, somewhat toothed; *pods ovate, winged, rough*, the style longer than the narrow notch. — Old fields, Mass. and New York to Virginia: rare. (Nat. from Eu.)

5. **L. DRÀBA**, L. Perennial, *obscurely hoary*; leaves oval or oblong, the upper with broad clasping auricles; flowers corymbose; *pods heart-shaped, wingless, thickish, entire, tipped with a conspicuous style*. — Astoria, near New York, D. C. Eaton. (Adv. from Eu.)

18. SENEBIÈRA, DC. WART-CRESS. SWINE-CRESS.

Pouch flattened contrary to the narrow partition; the two cells indchiscent, but falling away at maturity from the partition as closed nutlets, strongly wrinkled or tuberculate, 1-seeded. Cotyledons narrow and incumbently folded transversely. Low and diffuse or prostrate annuals or biennials, with minute whitish flowers. Stamens often only 2. (Dedicated to *J. Senbier*, a distinguished vegetable physiologist.)

1. **S. didyma**, Pers. Leaves 1–2-pinnately parted; *pods notched at the apex, rough-wrinkled*. (*S. pinnatifida*, DC. · *Lepidium didymum*, L.) — Waste places, at ports, Philadelphia to Virginia, &c.: an immigrant from farther South.

2. **S. CORONÓPUS**, DC. Leaves less divided, with narrower lobes; *Pods not notched at the apex, tubercled.* Virginia, *Pursh.* Newport, Rhode Island, *Robbins, &c.* (Adv. from Eu.)

19. CAKÍLE, TOURN. SEA-ROCKET.

Pod short, 2-jointed across, fleshy, the upper joint separating at maturity; each indehiscent, 1-celled and 1-seeded, or the lower sometimes seedless. Seed erect in the upper, suspended in the lower joint. Cotyledons obliquely accumbent. — Seaside fleshy annuals. Flowers purplish. (An old Arabic name.)

1. **C. Americana**, Nutt. (AMERICAN SEA-ROCKET.) Leaves obovate, sinuate and toothed; lower joint of the fruit obovoid, emarginate; the upper ovate, flattish at the apex. — Coast of the Northern States and of the Great Lakes. July–Sept. — Joints nearly even and fleshy when fresh; the upper one 4-angled and appearing more beaked when dry.

20. RÁPHANUS, L. RADISH.

Pods linear or oblong, tapering upwards, 2-jointed; the lower joint often seedless and stalk-like; the upper necklace-form by constriction between the seeds, with no proper partition. Style long. Seeds spherical as in Cabbage, &c. — Annuals or biennials. (The ancient Greek name from $\rho\acute{\alpha}$, *quickly*, and $\phi\alpha\acute{\iota}\nu\omega$, *to appear*, alluding to the rapid germination.)

1. **R. RAPHANSTRUM**, L. (WILD RADISH. JOINTED CHARLOCK.) Pods necklace-form, long-beaked; leaves lyre-shaped, rough; petals yellow, turning whitish or purplish, veiny. — A troublesome weed in fields, E. New England to Pennsylvania. (Adv. from Eu.)

R. SATIVUS, L., GARDEN RADISH, with pink-purple or whitish flowers, and thick knobby and pointed pods, with irregular fleshy partitions between the seeds, occasionally becomes spontaneous for a year or two.

ORDER 11. CAPPARIDACEÆ. (CAPER FAMILY.)

Herbs (when in northern regions), *with cruciform flowers, but 6 or more not tetradynamous stamens, a 1-celled pod with 2 parietal placentæ, and kidney-shaped seeds.* — Pod as in Cruciferae, but with no partition: seeds similar, but the embryo coiled rather than folded. — Leaves alternate, mostly palmately compound. — Often with the acrid or pungent qualities of Cruciferae (as in *capers*, the flower-buds of *Capparis spinosa*); also commonly bitter and nauseous. Represented within our limits only by

1. POLANÍSIA, Raf. POLANISIA.

Sepals 4. Petals 4, with claws, notched at the apex. Stamens 8–32, unequal. Receptacle not elongated, bearing a gland behind the base of the ovary. Pod linear or oblong, veiny, turgid, many-seeded. — Fetid annuals, with glandular or clammy hairs. Flowers in leafy racemes. (Name from $\pi\omicron\lambda\upsilon\varsigma$, *many*, and $\acute{\alpha}\nu\iota\sigma\omicron\varsigma$, *unequal*, points in which the genus differs in its stamens from Cleome.)

1. **P. graveolens**, Raf. Leaves with 3 oblong leaflets; stamens about 11, scarcely exceeding the petals: style short; pod slightly stipitate. — Gravelly shores, from Connecticut (near Hartford) and W. Vermont to Wisconsin and Kentucky. June — Aug. — Flowers small: calyx and filaments purplish: petals yellowish-white.

ORDER 12. RESEDÀCÆ. (MIGNONETTE FAMILY.)

*Herbs, with unsymmetrical 4-7-merous small flowers, a fleshy one-sided hypogynous disk between the petals and the (3-40) stamens, bearing the latter. Calyx not closed in the bud. Pod 3-6-lobed, 3-6-horned, 1-celled with 3-6 parietal placentæ, opening at the top before the seeds (which are as in Order 11) are full grown. — Leaves alternate, with only glands for stipules. Flowers in terminal spikes or racemes. — A small and unimportant family, of the Old World, represented by the Mignonette (*Reseda odorata*) and the Dyer's Weed.*

1. RESÈDA, L. MIGNONETTE. DYER'S ROCKET.

Petals 4-7, cleft, unequal. Stamens 12-40, on one side of the flower. (Name from *resedo*, to calm, in allusion to supposed sedative properties.)

1. **R. LUTÈOLA**, L. (DYER'S WEED or WELD.) Leaves lanceolate; calyx 4-parted; petals 4, greenish-yellow; the upper one 3-5-cleft, the two lateral 3-cleft, the lower one linear and entire; pods depressed. — Roadsides, New York, &c. — Plant 2° high. Used for dyeing yellow. (Adv. from Eu.)

ORDER 13. VIOLÀCÆ. (VIOLET FAMILY.)

Herbs, with a somewhat irregular 1-spurred corolla of 5 petals, 5 hypogynous stamens with adnate introrse anthers conniving over the pistil, and a 1-celled 3-valved pod with 3 parietal placentæ. — Sepals 5, persistent. Petals imbricated in the bud. Stamens with their short and broad filaments continued beyond the anther-cells, and often coherent with each other. Style usually club-shaped, with the simple stigma turned to one side. Valves of the capsule bearing the several-seeded placentæ on their middle: after opening, each valve as it dries folds together lengthwise firmly, projecting the seeds. Seeds anatropous, rather large, with a hard seed-coat, and a large and straight embryo nearly as long as the albumen: cotyledons flat. — Leaves alternate, with stipules. Flowers axillary, nodding. (Roots slightly acrid or emetic.) — Two genera in the Northern United States.

1. SÒLEA, Ging., DC. GREEN VIOLET.

Sepals not prolonged at the base. Petals nearly equal in length, but the lower one larger and gibbous or saccate at the base, more notched than the others at the apex. Stamens completely united into a sheath enclos-

ing the ovary, and bearing a broad gland on the lower side. Style hooked at the summit. — A homely perennial herb, with stems leafy to the top, and 1–3 small greenish-white flowers in the axils, on short recurved pedicels. (Named in honor of *W. Sole*, author of an essay on the British Mints.)

1. **S. cóncolor**, Ging. (*Viola concolor*, *Pursh*, &c.) — Woods, New York to Illinois and southward. June. — Plant 1°–2° high. Leaves oblong, pointed at both ends, entire. Pod 1' long.

2. VIOLA, L. VIOLET. HEART'S-EASE.

Sepals extended into ears at the base. Petals somewhat unequal, the lower one spurred at the base. Stamens closely surrounding the ovary, often slightly cohering with each other; the two lower ones bearing spurs which project into the spur of the corolla. Besides these conspicuous blossoms, which appear in spring, others are produced later (especially in the stemless species), on shorter peduncles or on runners, usually concealed under the leaves; these never open nor develop petals, but are fertilized in the young bud, producing pods which are far more fruitful than the ordinary blossoms. (The ancient Latin name of the genus.)

§ 1. **STEMLESS VIOLETS**: *the leaves and scapes all from a subterranean rootstock, flowering in early spring, and bearing fruitful apetalous flowers all summer.*

* *Flowers yellow: rootstock creeping and producing summer runners.*

1. **V. rotundifolia**, Michx. (ROUND-LEAVED VIOLET.) Leaves round-ovate, heart-shaped, slightly crenate; lateral petals bearded and marked with brown lines; spur very short. — Cold woods, Maine to Michigan, and south along the Alleghanies. — Smoothish: leaves 1' broad at flowering, increasing to 3' or 4' in the summer, then lying flat on the ground, shining above.

* * *Flowers white (small, short-spurred); lower petals striped with lilac veins: rootstock creeping and producing summer runners or subterranean filiform branches. (No. 2 and No. 4, however different, seem to be connected by No. 3.)*

2. **V. lanceolata**, L. (LANCE-LEAVED VIOLET.) Smooth; leaves lanceolate, erect, blunt, tapering into a long-margined petiole, almost entire; petals beardless. — Damp soil, Maine to Illinois, Kentucky, and southward; common eastward.

3. **V. primulæfolia**, L. (PRIMROSE-LEAVED V.) Smooth or a little pubescent; leaves oblong or ovate, abrupt or somewhat heart-shaped at the base; petals often acute, the lateral ones usually sparingly bearded. (*V. acuta*, *Bigelow*.) — Damp soil; with No. 2.

4. **V. blanda**, Willd. (SWEET WHITE V.) Leaves round-heart-shaped or kidney-form, minutely pubescent; petals mostly beardless. — Damp places, everywhere, Maine to Wisconsin and Kentucky. — Flowers faintly sweet-scented.

* * * *Flowers purplish or violet: rootstocks slender and creeping, often producing numerous summer runners.*

V. ODORATA, L. (SWEET OR ENGLISH VIOLET), cultivated in gardens, from Europe, belongs to this section, and is becoming sparingly spontaneous in some places.

5. **V. palústris**, L. (MARSH V.) Smooth; leaves round-heart-shaped and kidney-form, slightly crenate; flowers (small) pale lilac with purple streaks, nearly beardless; *spur very short* and obtuse.—Alpine summits of the White Mountains, New Hampshire, and high northward. June. (Eu.)

6. **V. Selkirkii**, Pursh., Goldie., 1822. (GREAT-SPURRED V.) Small and delicate; the filiform rootstock fibrose-rooted, no runners above ground; smooth, except the round-heart-shaped crenate leaves, which are minutely hairy on the upper surface and have a deep narrowed sinus; *spur very large*, thickened at the end, almost as long as the beardless pale violet petals. (*V. umbrosa*, Fries, 1828. *V. Kamtschatica*, Gingins, 1826.)—Damp and shady soil, W. Massachusetts to Chatauque Co., N. Y. (*Clinton*), L. Superior (*Robbins*), and northward: rare.—Scapes and petioles 1'–2', the leaf $\frac{1}{2}$ '–1 $\frac{1}{4}$ ' long, thin; the spur 3'' long. (Eu.)

* * * * *Flowers violet or purple (or rarely almost white): rootstocks fleshy and thickened or tuberous, mostly erect or ascending, producing neither runners nor runner-like subterranean branches.*

7. **V. cucullàta**, Ait. (COMMON BLUE V.) Rootstocks thickly dentate with fleshy teeth, branching and forming compact masses; *leaves all long-petioled and upright, heart-shaped with a broad sinus*, varying to kidney-shaped and dilated-triangular, smooth, or more or less pubescent, the sides at the base rolled inwards when young, obtusely serrate; *lateral and often the lower petals bearded*; spur short and thick; stigma slightly beaked or beakless.—Low grounds, common everywhere.—Very variable in size, shape of leaves and sepals, and in the color and size of the flowers, which are deep or pale violet-blue or purple, sometimes nearly white, or variegated with white. Scapes 3'–10' high. Passes by intermediate forms of all sorts into

Var. **palmàta**. (HAND-LEAF V.) *Leaves variously 3–7-cleft* or parted, or the earlier ones entire on the same individual. (*V. palmàta*, L.)—Common, especially southward.

Var. **cordàta**. Leaves chiefly round-heart-shaped and *prostrate*, sometimes villous, sometimes nearly glabrous, small. (*V. villòsa* and *V. cordata*, Walt. *V. soròria*, Willd.)—Common southward; a variety growing in drier soil or more exposed situations.

8. **V. sagittàta**, Ait. (ARROW-LEAVED V.) Smoothish or hairy; *leaves on short and margined*, or the later often on long and naked *petioles*, varying from *oblong-heart-shaped to halberd-shaped, arrow-shaped, oblong-lanceolate or ovate*, denticulate, sometimes cut-toothed near the base, the lateral or occasionally all the (pretty large purple-blue) petals bearded; spur short and thick; stigma beaked. (*V. ovàta*, Nutt., and *V. emarginàta*, Le Conte, are states of this variable species.)—Dry or moist open places, New England to Illinois and southward.—Rootstock nearly as in the preceding, into which some forms seem to pass.

9. **V. delphinifòlia**, Nutt. (LARKSPUR V.) *Leaves all palmately or pedately 5–7-parted*, divisions 2–3-cleft; lobes linear; *lateral petals bearded*; stigma short-beaked; otherwise like the next.—Rich prairies, Illinois and westward.

10. **V. pedàta**, L. (BIRD-FOOT V.) Nearly smooth; rootstock short and very thick, erect, not scaly; *leaves all 3–5-divided*, or the earliest only parted, the lateral divisions 2–3-parted, all linear or narrowly spatulate, sometimes 2–

3-toothed or cut at the apex; *petals beardless*; stigma nearly beakless. — Sandy or gravelly soil, New England to Illinois and southward. — Flower large, 1' broad, pale or deep lilac-purple or blue.

Var. **bicolor**. A very handsome variety, with the two upper petals deep violet, and as it were velvety, like a pansy, occurs sparingly from Massachusetts to Maryland, &c.

§ 2. LEAFY-STEMMED VIOLETS: *all but the last perennial from short rootstocks.*

* *Leaf-bearing from base to summit, usually branching and flowering all summer: stipules entire or barely toothed, not foliaceous.*

11. **V. canina**, L., var. **sylvéstris**, Regel. (DOG V.) Low (3'–8' high); stems ascending, mostly simple, from the base at length producing creeping branches; leaves heart-shaped, or the lowest kidney-form, crenate, the uppermost slightly pointed; stipules lanceolate, fringe-toothed; *spur cylindrical, half the length of the light violet petals*, the lateral ones slightly bearded; stigma beaked. (*V. sylvestris*, Lam. *V. Muhlenbergii*, Torr., and former ed.) — Damp or wet shady places: common. May–July. (Eu.)

12. **V. rostrata**, Pursh. (LONG-SPURRED V.) Stems ascending (3'–6' high); leaves roundish-heart-shaped, serrate, the upper acute; stipules lanceolate, fringed-toothed, large; *spur slender* ($\frac{1}{2}$ ' long), *longer than the pale violet beardless petals*; style straight and slender; stigma terminal, beakless. — Shaded hillsides, Maine to Ohio and Kentucky, and southwards in the Alleghanies: rather rare. June, July.

13. **V. striata**, Ait. (PALE V.) Stems angular, ascending (6'–10' high); leaves heart-shaped, finely serrate, often acute; stipules oblong-lanceolate, large, strongly fringed-toothed; *spur thickish, much shorter than the cream-colored or white petals*, the lateral ones bearded, the lower striped with purplish lines; stigma beaked. — Low grounds; common, especially westward. April–Oct.

14. **V. Canadensis**, L. (CANADA V.) Upright (1°–2° high); leaves heart-shaped, pointed, serrate; *stipules ovate-lanceolate, entire*; petals white or whitish inside, the upper ones mostly tinged with violet beneath, the lateral bearded; *spur very short*; stigma beakless. — Rich woods; common northward and along the Alleghanies. May–Aug.

* * *Simple stems erect, naked below, 2–4-leaved above: stipules nearly entire: flowers yellow, in spring and early summer: stigma bearded on each side.*

15. **V. pubescens**, Ait. (DOWNY YELLOW V.) Softly pubescent (6'–12' high); *leaves very broadly heart-shaped*, toothed, somewhat pointed; stipules ovate or ovate-lanceolate, large; spur extremely short; lower petals veined with purple. — Woods; common.

Var. **eriocarpa**, Nutt. More pubescent, stout, 1°–2° high; pods woolly. (*V. eriocarpa*, Schwein.) — Common westward.

Var. **scabriuscula**, Torr. & Gray. Smaller and greener, slightly pubescent; stems often decumbent (4'–10' high). — Rhode Island to Kentucky.

16. **V. hastata**, Michx. (HALBERD-LEAVED V.) Nearly glabrous, slender (4'–10' high); *stem-leaves halberd-shaped* or oblong-heart-shaped, slightly serrate, acute; stipules ovate, small; spur very short. — Woods, N. Ohio (near Painesville, *Miss Shattuck*), mountains of Penn., and southward: rare.

* * * *Leaf-bearing throughout from an annual, biennial, or sometimes short-lived perennial root; the stipules large, leaf-like and lyrate-pinnatifid.*

17. **V. TRICOLOR**, L. (PANSY. HEART'S-EASE.) Stem angled and branched; leaves roundish, or the upper oval and the lowest heart-shaped, crenate or entire; petals variable in color or variegated (yellow, whitish, violet-blue and purple); — in var. **ARVENSIS** shorter or little longer than the calyx. — Dry or sandy soil, New York to Illinois and southward: the variety seeming like a native plant. April–Sept. (Nat. from Eu.)

ORDER 14. CISTACEÆ. (ROCK-ROSE FAMILY.)

Low shrubs or herbs, with regular flowers, distinct and hypogynous mostly indefinite stamens, a persistent calyx, a 1-celled 3–5-valved pod with as many parietal placentæ borne on the middle of the valves, and orthotropous albuminous seeds. — Sepals 5; the two external often small, like bracts, or sometimes wanting; the three others a little twisted in the bud. Petals 3 or 5, convolute in the opposite direction from the calyx in the bud. Anthers short, innate, on slender filaments. Style single or none. Ovules few or many, on slender stalks, with the orifice at their apex. Embryo long and slender, straightish or curved, in mealy albumen: cotyledons narrow. — Leaves simple and mostly entire, the lower usually opposite, and the upper alternate. (Inert plants. A small family: mostly of the Mediterranean region.)

1. **Helianthemum**. Petals 5, crumpled in the bud, fugacious. Stamens and ovules numerous in the petal-bearing flowers.
2. **Hudsonia**. Petals 5, fugacious. Stamens 9–30. Style long and slender. Pod strictly 1-celled, 2–6-seeded.
3. **Lechea**. Petals 3, persistent. Stamens 3–12. Style none. Pod partly 3-celled, the imperfect partitions bearing broad 2-seeded placentæ.

1. HELIANTHEMUM, Tourn. ROCK-ROSE.

Petals 5, crumpled in the bud, fugacious. Style short or none in our species: stigma 3 lobed. Capsule strictly 1-celled. Embryo curved in the form of a hook or ring. — Flowers in most N. American species of two sorts, viz., 1. the *primary* or earlier ones, with large petals, indefinitely numerous stamens, and many-seeded pods; 2. *secondary*, or later ones, which are much smaller and in clusters, with small petals or none, 3–10 stamens, and much smaller 3–few-seeded pods. The yellow flowers open only once, in sunshine, and cast their petals by the next day. (Name from *ἥλιος*, the sun, and *ἄνθεμον*, flower.)

1. **H. Canadense**, Michx. (FROST-WEED.) *Petal-bearing flowers solitary; the small secondary flowers clustered in the axils of the leaves, nearly sessile; calyx of the large flowers hairy-pubescent; of the small ones hoary, like the stem and lower side of the lanceolate-oblong leaves.* — A variety is more hoary, and with a stronger tendency to multiply the minute clustered flowers. — Sandy or gravelly dry soil, Maine to Wisconsin and southward. June–Aug. — Stems at first simple. Corolla of the large flowers 1' wide, producing pods 3' long:

Pods of the smaller flowers not larger than a pin's head. — Late in autumn, crystals of ice shoot from the cracked bark at the root, whence the popular name.

2. *H. corymbosum*, Michx. *Flowers all clustered at the summit of the stem or branches, the petal-bearing ones at length on slender stalks: calyx woolly.* — Pine barrens, New Jersey and southward along the coast.

2. HUDSONIA, L. HUDSONIA.

Petals 5, fugacious (lasting but a day), much larger than the calyx. Stamens 9–30. Style long and slender: stigma minute. Pod oblong, enclosed in the calyx, strictly 1-celled, with 1 or 2 seeds attached near the base of each nerve-like placenta. Embryo coiled into the form of a closed hook. — Bushy heath-like little shrubs (seldom a foot high), covered all over with the small awl-shaped or scale-like persistent downy leaves, producing numerous (small but showy) bright yellow flowers crowded along the upper part of the branches. (Named in honor of *Wm. Hudson*, an early English botanist.)

1. *H. ericoides*, L. Downy but greenish; leaves slender, awl-shaped, loose; flowers on slender naked stalks. — Dry sandy soil near the coast, N. Maine to Virginia. May.

2. *H. tomentosa*, Nutt. Hoary with down; leaves oval or narrowly oblong, short, close-pressed and imbricated; flowers sessile (sandy coasts from Maine to Maryland), — or short-peduncled, the leaves also narrower: Maine (at Harrison, *J. Blake*) and along the shores of the Great Lakes to Minnesota. May, June.

3. LÉCHEA, L. PINWEED.

Petals 3, narrow, flat in the bud: not longer than the calyx, withering-persistent. Stamens 3–12. Style scarcely any: stigmas 3, plumose. Pod globular, partly 3-celled; the 3 broad and thin placentæ borne on imperfect partitions, each bearing 2 seeds on the face towards the valve: in our species, the placentæ curve backwards and partly enclose the seeds. Embryo straightish. — Homely perennial herbs, with very small greenish or purplish flowers, in summer. (Named in honor of *John Leche*, a Swedish botanist.)

1. *L. major*, Michx. *Hairy; stem upright (1°–2° high, stout), simple, producing slender prostrate branches from the base; leaves elliptical, mucronate-pointed, alternate and opposite or sometimes whorled; flowers densely crowded in paniced clusters; pedicels shorter than the very small globose-triangular pod; sepals narrower than its valves.* — Sterile grounds: common, especially southward.

2. *L. thymifolia*, Pursh. *Hoary with appressed hairs, especially the decumbent stout leafy shoots from the base; flowering stems ascending, loosely branched, with the leaves linear or oblanceolate; those of the shoots elliptical, whorled, crowded; flowers scattered in small and loose clusters; pedicels as long as the globose pods.* — Sandy coast, Maine to New Jersey and southward. — Scarcely a foot high, tufted, rigid; the pods larger than in No. 1.

3. *L. Novæ-Cæsariæ*, C. F. Austin, ined. Intermediate in appearance between No. 1 and the taller forms of No. 4; leaves of the former, but

smaller ($\frac{1}{2}$ long) *elliptical* or *linear-oblong*, often opposite or whorled; *flowers in narrow rather close panicles*; pedicels longer than the oval pods; the two outer or *bract-like sepals very slender, mostly longer than the others*. — Open dry grounds, N. New Jersey and adjacent part of New York, *C. F. Austin*.

4. **L. minor**, Lam. *Minutely hairy*; stems slender, upright or diffuse; leafy shoots densely tufted at the base; *leaves linear*; *flowers loosely racemed* on the slender branchlets; pedicels mostly longer than the oval pods. — Dry open soil: common. June–Sept. — Plant 5'–15' high, slender, running into numberless variations according to the soil, season, and exposure. Pods smaller than in No. 2.

ORDER 15. DROSERACEÆ. (SUNDEW FAMILY.)

Bog-herbs, mostly glandular-haired, with regular hypogynous flowers, pentamerous and withering-persistent calyx, corolla, and stamens, the anthers fixed by their middle and turned outwards, and a 1-celled pod with twice as many styles or stigmas as there are parietal placentæ. — Calyx imbricated. Petals convolute. Seeds numerous, anatropous, with a short and minute embryo at the base of the albumen. — Leaves circinate in the bud, i. e. rolled up from the apex to the base as in Ferns. (A small family, of no known qualities, except a slight bitterness, &c.; the Sundews impart a purple stain to paper.) Only one genus within our limits, viz.

1. DRÓSERÁ, L. SUNDEW.

Stamens 5. Styles 3, or sometimes 5, deeply 2-parted so that they are taken for 6 or 10, slender, stigmatose above on the inner face. Pod 3- (rarely 5-) valved; the valves bearing the numerous seeds on their middle for the whole length. — Low perennials or biennials; the leaves clothed with reddish gland-bearing bristles, in our species all in a tuft at the base; the naked scape bearing the flowers in a 1-sided raceme-like inflorescence, which nods at the undeveloped apex, so that the fresh-blown flower (which opens only in sunshine) is always highest. (The glands of the leaves exude drops of a clear glutinous fluid, glittering like dew-drops, whence the name, from *δροσερός*, *dewy*.)

1. **D. rotundifolia**, L. (ROUND-LEAVED SUNDEW.) *Leaves orbicular, abruptly narrowed into the spreading hairy petioles*; seeds spindle-shaped, the coat loose and chaff-like; flowers white, the parts sometimes in sixes. — Peat-bogs, common, especially northward. July–Aug. (Eu.)

2. **D. longifolia**, L. *Leaves spatulate-oblong, tapering into the long rather erect naked petioles*; seeds oblong, with a rough close coat; flowers white. (*D. intermedia*, Hayne.) — Bogs; less common. June–Aug. — Plant raised on its prolonged caudex when growing in water. (Eu.)

3. **D. linearis**, Goldie. (SLENDER SUNDEW.) *Leaves linear, obtuse, the blade (2'–3' long, scarcely 2" wide) on naked erect petioles* about the same length; seeds oblong, with a smooth and perfectly close coat; flowers white. — Shore of Lake Superior. July.

4. **D. filifórmis**, Raf. (THREAD-LEAVED SUNDEW.) *Leaves very long and filiform*, erect, with no distinction between blade and stalk; seeds spindle-shaped; flowers numerous, purple rose-color ($\frac{1}{2}$ broad). — Wet sand, near the coast, Plymouth, Massachusetts to New Jersey, and southward. Aug. — Scapes 6'–12' high, and the singular leaves nearly as long, from a bulb-like base or corm.

DIONÆA MUSCÍPULA, Ellis, the VENUS'S FLY-TRAP, — so noted for the extraordinary irritability of its leaves, closing quickly at the touch, — is a native of the sandy savannas of the eastern part of North Carolina. It differs in several respects from the character of the order given above; the stamens being 15, the styles united into one, and the seeds all at the base of the pod.

ORDER 16. **HYPERICACEÆ.** (ST. JOHN'S-WORT FAMILY.)

Herbs or shrubs, with opposite entire dotted leaves and no stipules, regular hypogynous flowers, the petals mostly oblique and convolute in the bud, and many or few stamens commonly collected in 3 or more clusters or bundles. Pod 1-celled with 2–5 parietal placentæ, and as many styles, or 3–7-celled by the union of the placentæ in the centre: dehiscence mostly septical. — Sepals 4 or 5, imbricated in the bud, herbaceous, persistent. Petals 4 or 5, mostly deciduous. Styles persistent, at first sometimes united. Seeds numerous, small, anatropous, with no albumen. Embryo cylindrical. — Plants with a resinous juice (acrid and balsamic), dotted with pellucid or dark glands, usually smooth. Leaves mostly sessile. Flowers solitary or cymose.

1. **Ascyrum**. Sepals 4, very unequal. Petals 4, oblique, convolute, yellow.
2. **Hypericum**. Sepals 5. Petals 5, oblique, convolute, yellow.
3. **Elodes**. Sepals 5. Petals 5, equal-sided, imbricated, purplish. Glands alternating with the stamen-clusters.

1. **ÁSCYRUM**, L. ST. PETER'S-WORT.

Sepals 4; the 2 outer very broad and leaf-like; the inner much smaller. Petals 4, oblique, very deciduous, convolute in the bud. Stamens numerous; the filaments distinct and scarcely in clusters. Pod strictly 1-celled; 2–4-valved. — Low, rather shrubby, smooth plants, with pale black-dotted leaves, and nearly solitary light yellow flowers. (An ancient Greek name of some plant, from *a*, *without*, and *σκόπος*, *roughness*.)

1. **A. stáns**, Michx. (ST. PETER'S-WORT.) Stem rather simple, 2-edged, 1°–2° high, stout; *leaves oval or oblong, somewhat clasping, thickish; petals obovate*; styles 3 or 4. — Pine barrens, Long Island to Penn. and southward. July, Aug. — Flowers showy, almost sessile: outer sepals round-heart-shaped.

2. **A. Crux-Ándreæ**, L. (ST. ANDREW'S CROSS.) Low, much branched and decumbent; *leaves narrowly obovate-oblong, contracted at the base, thin; petals linear-oblong*; styles 2, very short; pod flat. — Pine barrens, New Jersey to Illinois, and southward. July–Sept. — Petals scarcely exceeding the outer sepals, approaching each other in pairs over them, in the form of a St. Andrew's cross.

2. *HYPÉRICUM*, L. ST. JOHN'S-WORT.

Sepals 5, somewhat equal. Petals 5, oblique, convolute in the bud. Stamens commonly united or clustered in 3–5 parcels: no interposed glands. Pod 1-celled or 3–5-celled. Seeds usually cylindrical.—Herbs or shrubs, with cymose yellow flowers. (An ancient Greek name, of obscure meaning.)

§ 1. *Stamens very numerous, 5-adelphous: pod 5–7-celled, with the placentæ turned far back into the cells: perennial: flowers very large: styles united.*

1. *H. pyramidatum*, Ait. (GREAT ST. JOHN'S-WORT.) Branches 2–4-angled; leaves ovate-oblong, partly clasping; petals narrowly obovate, not deciduous until after they wither; stigmas capitate.—Banks of rivers: rare. New England and Penn. to Wisconsin and Illinois. July.—Plant 3°–5° high. Leaves 2'–3' long. Petals 1' long. Pod $\frac{3}{4}$ ' long, conical.

§ 2. *Stamens very numerous, obscurely if at all clustered: styles 3 (No. 2 excepted), more or less united into one and the sepals foliaceous, except in No. 9.*

* *Bushy shrubs, 1°–6° high, leafy to the top: pod 3–5-celled.*

2. *H. Kalmianum*, L. (KALM'S ST. JOHN'S-WORT.) Branches 4-angled: branchlets 2-edged; leaves crowded, *glaucous*, oblanceolate (1'–2' long); flowers few in a cluster (1' wide); *pods ovate, 5-celled.*—Wet rocks, Niagara Falls and Northern lakes. Aug.

3. *H. prolificum*, L. (SHRUBBY ST. JOHN'S-WORT.) Branchlets 2-edged; leaves lanceolate-oblong, mostly obtuse, narrowed at the base; flowers numerous, in single or compound clusters; *pods oblong, 3-celled.*—New Jersey to Michigan, Illinois, and southward. July–Sept.—Varies greatly in size, &c.

Var. *densiflorum*. Exceedingly branched above, 1°–6° high, the branches slender and crowded with smaller leaves; flowers smaller ($\frac{1}{2}$ '– $\frac{3}{4}$ ' in diameter) and more numerous, in crowded compound cymes. (*H. densiflorum*, & *H. galioides*, *Pursh.*)—Pine barrens of New Jersey to glades of Kentucky, and southward.

** *Perennial herbs or in No. 4 and 5 a little woody at the base: pod one-celled with 3 parietal placentæ, or incompletely 3–4-celled.*

4. *H. adpressum*, Barton. Stem simple, herbaceous, from a slightly woody creeping base (1°–2° high), obscurely 4-angled below and 2-edged above; *leaves ascending, lanceolate* or linear-oblong, often acute, thin; cyme leafy at the base, few-flowered; *sepals linear-lanceolate; pods ovoid-oblong, partly 3–4-celled.*—Moist places, Rhode Island (*Olney*), New Jersey, Pennsylvania, and southwestward. July–Aug.—Leaves $1\frac{1}{2}$ ' long. Petals bright yellow, 3''–5'' long.

5. *H. dolabriforme*, Vent. Stems branched from the decumbent base, woody below (6'–20' high), terete; *leaves linear-lanceolate*, widely spreading, veinless; cyme leafy, few-flowered; *sepals oblong or ovate-lanceolate*, about the length of the very oblique petals (5''–6'' long); *pods ovate-conical, pointed, strictly 1-celled*, the walls very thick and hard. (*H. procumbens*, *Michx.*)—Dry hills and rocks, barrens of Kentucky and westward. June–Aug.

6. *H. nudiflorum*, Michx. Stems branched, woody at the base, sharply 4-angled or almost winged above (2°–4° high); *leaves oblong or oval-lanceolate*, obtuse, obscurely veined, pale (2'–2 $\frac{1}{2}$ ' long); cyme compound, many-flowered,

naked; *sepals oblong*; *Pods ovate-conical, pointed, almost 3-celled*; seeds slender cylindrical, minutely pitted. — Low grounds, Pennsylvania, Virginia, and southward. July.

7. **H. sphærocárpón**, Michx. Stems mostly simple, herbaceous, with a somewhat woody base, angled with 4 very narrow salient lines (1° – 2° high); *leaves oblong-linear*, greener above and narrower than in the preceding; the naked cyme similar; *sepals ovate*; *Pods depressed-globular or ovoid-conical, strictly 1-celled*; seeds oblong, rough-pitted. — Rocky banks of the Ohio and its tributaries, S. W. Ohio to Illinois and southward. July–Sept. — Flowers small.

8. **H. ellipticum**, Hook. Stem simple, herbaceous (1° high), obscurely 4-angled; *leaves spreading, elliptical-oblong*, obtuse, thin; cyme nearly naked, rather few-flowered; *sepals oblong*; *Pods ovoid, very obtuse*, purple, 1-celled. — Wet places, New England and Pennsylvania to Lake Superior and northward. July, Aug. — Petals light yellow, 3'' long.

9. **H. angulòsum**, Michx. Stem slender, strict, simple, sharply 4-angled, herbaceous (1° – 2° high); *leaves ascending, opaque, ovate or oblong-lanceolate, acute* ($\frac{1}{2}$ '–1' long), closely sessile by a broad base; cyme compound, naked, the scattered flowers racemose on its ascending branches; *sepals herbaceous, erect*, enclosing the ovoid 1-celled pod; *styles 3, separate*. — Wet pine barrens of New Jersey and southward. July–Sept. — Petals copper-yellow, 4''–5'' long, furnished with a tooth on one side.

§ 3. *Stamens very numerous, in 3 or 5 clusters: styles 3 separate and usually diverging: pod 3-celled: calyx erect: petals and anthers with black dots.*

10. **H. perforátum**, L. (COMMON ST. JOHN'S-WORT.) Stem much branched and corymbed, somewhat 2-edged (producing runners from the base); *leaves elliptical-oblong or linear-oblong, with pellucid dots*; petals (deep yellow) twice the length of the *lanceolate acute sepals*; flowers numerous, in open leafy cymes. — Fields, &c. June–Sept. — Too well known as a pernicious weed, which it is difficult to extirpate. Juice very acrid. (Nat. from Eu.)

11. **H. corymbòsum**, Muhl. Conspicuously marked with both black and pellucid dots: stem terete, sparingly branched; *leaves oblong*, the base either obtuse or somewhat clasping; *flowers crowded* (small); *petals pale yellow*, much longer than the *oblong sepals*, styles not longer than the pod. — Damp places; common. July–Sept. — Leaves larger and flowers much smaller than in the last: petals 2''–3'' long, marked with black lines as well as dots. — Too near *H. maculatum*, Walt., of the South, which has more clasping leaves and very long and slender styles.

12. **H. gravèolens**, Buckley. Like the last, but with larger leaves and fewer much larger bright yellow flowers, lanceolate acute sepals, and long erect styles; common in the mountains of N. Carolina, doubtless also in S. Virginia.

§ 4. *Stamens 5–12, distinct or in 3 clusters: pod (brown purple) 1-celled, with 3 strictly parietal placenta: styles short, distinct; petals oblong or linear: sepals narrow, erect: slender annuals, with 4-angular branches; flowering all summer.*

13. **H. mütillum**, L. Stem flaccid, widely branching (6'–10' high); *leaves ovate or oblong, obtuse, partly clasping, 5-nerved*; cymes leafy; *Pods ovate-conical, rather longer than the calyx*. (*H. parviflorum*, Muhl.) — Low grounds, everywhere. — Flowers 2'' broad.

Var. **gymnanthum** (*H. gymnanthum*, *Engelm. & Gray*), is a form, or perhaps species, with strict stem and branches, or often unbranched, more clasping heart-shaped stem-leaves, and a naked cyme, the floral leaves being reduced to small awl-shaped bracts; so that in aspect it approaches the next. — Newcastle Co., Delaware, *Canby*, and Illinois, *E. Hall*, thence southward.

14. **H. Canadense**, L. Stem strict (6'–15' high), with the branches erect; *leaves linear*, 3-nerved at the base, obtuse; cymes naked; *Pods conical-oblong*, usually much longer than the calyx. — Wet, sandy soil: common. June–Oct. — Flowers deep yellow, 2"–3" broad when expanded.

Var. **major** is a large form, 1°–2° high, with lanceolate leaves 1½' long, 3" wide, the upper acute. — L. Superior, *Robbins*; S. New York and southward.

15. **H. Drummóndii**, Torr. & Gray. Stem and the mostly alternate bushy branches rigid, erect (10'–18' high); *leaves linear-subulate*, nearly erect, 1-nerved (3"–9" long); *flowers scattered* along the upper part of the leafy branches, *short-pedicelled*; *Pods ovoid*, not longer than the calyx. (*Sarothra Drummóndii*, *Grev. & Hook.*) — W. Illinois and southward, in dry soil.

16. **H. Saróthra**, Michx. (ORANGE-GRASS. PINE-WEED.) Stem and bushy branches thread-like, wiry (4'–9' high); *leaves minute awl-shaped scales*, appressed; *flowers minute*, mostly sessile and scattered along the erect branches; *Pods ovate-lanceolate*, acute, much longer than the calyx. (*Sarothra gentianoides*, L.) — Sandy fields: common. June–Oct.

3. ELÓDES, Adans. MARSH ST. JOHN'S-WORT.

Sepals 5, equal, erect. Petals 5, equal-sided, oblong, naked, imbricated in the bud. Stamens 9 (rarely more), united in 3 sets; the sets separated by as many large orange-colored glands. Pod 3-celled, oblong: styles distinct. — Perennial herbs, in marshes or shallow water, with small close clusters of flesh-colored flowers in the axils of the leaves and at the summit of the stem. (Name *ἑλώδης*, growing in marshes, accidentally changed to *ELODĒA* by Jussieu, who was followed by Pursh, &c.)

1. **E. Virgínica**, Nutt. *Leaves closely sessile or clasping by a broad base*, oblong or ovate, very obtuse; filaments united below the middle. (*Hypericum Virgínicum*, L.) — Common in swamps. July, Aug.

2. **E. petioláta**, Pursh. *Leaves tapering into a short petiole*, oblong: filaments united beyond the middle. — From New Jersey south and westward.

ORDER 17. ELATINACEÆ. (WATER-WORT FAMILY.)

Little marsh annuals, with membranaceous stipules between the opposite dotless leaves, minute axillary flowers like Chickweeds, but the pod 2–5-celled, and the seeds as in St. John's-wort. The principal genus is

1. ELÁTINE, L. WATER-WORT.

Sepals 2–4, persistent. Petals 2–4, hypogynous. Stamens as many, rarely twice as many, as the petals. Styles, or sessile capitate stigmas, 2–4. Pod 2–4-celled, several-many-seeded, 2–4-valved; the partitions left attached to the

axis, or evanescent. Seeds cylindrical, straightish or curved. (A Greek name for some obscure herb.)

1. **E. Americana**, Arnott. Dwarf (1' high), creeping, rooting in the mud, tufted; leaves obovate; flowers sessile; sepals, petals, stamens, and stigmas 2, rarely 3; seeds 5 or 6 in each cell, rising from the base. (Peplis Americana, Pursh. Crypta minima, Nutt.)—Margin of ponds, &c., N. Hampshire, to Illinois, Virginia, and southwestward. Pod very thin and delicate; the seeds large in proportion, straightish.

ORDER 18. CARYOPHYLLACEÆ. (PINK FAMILY.)

Herbs, with opposite entire leaves, symmetrical 4-5-merous flowers, with or without petals; the distinct stamens no more than twice the number of the sepals, either hypogynous or perigynous; styles 2-5 (or rarely united into one); seeds attached to the base or the central column of the 1-celled (rarely 3-5-celled) pod, with a slender embryo coiled or curved around the outside of mealy albumen, in Dianthus nearly straight.—Bland herbs; the stems usually swollen at the joints; uppermost leaves rarely alternate. Leaves often united at the base. Calyx persistent. Styles stigmatic along the inside. Seeds amphitropous or campylotropous.

Tribe I. SILENEÆ. Sepals united into a tube or cup. Petals and stamens borne on the stipe or stalk of the ovary, the former with slender claws, to the base of which the corresponding filaments often adhere, included in the calyx-tube, mostly convolute in the bud. Seeds numerous.—Stipules none. Flowers often large and showy.

* Calyx with scaly bractlets or small leaves at the base. Seeds flattened on the back, attached by their face: embryo nearly straight.

1. **Dianthus.** Calyx terete, mostly cylindrical. Styles 2.

** Calyx naked. Seeds globular or kidney-shaped: embryo curved or coiled.

2. **Saponaria.** Calyx terete. Styles 2.

3. **Vaccaria.** Calyx 5-angled and in fruit 5-winged. Styles 2.

4. **Silene.** Calyx 5-toothed. Styles 3.

5. **Lychnis.** Calyx 5-toothed or 6-lobed. Styles 5, rarely 4.

Tribe II. ALSINEÆ. Sepals separate to the base or nearly so, imbricated in the bud. Petals when present without claws, mostly imbricated in the bud, and with the stamens inserted at the base of the sessile ovary, or into a little disk which often coheres with the base of the calyx. Pod splitting into valves or teeth, several - many-seeded. Stamens opposite the sepals, when not more numerous than they.—Low herbs. Stipules none.

* Styles opposite the sepals, or, when fewer, opposite those which are exterior in the bud.

6. **Arenaria.** Petals entire. Styles usually 3. Pod short, splitting into 3 or 6 valves.

7. **Stellaria.** Petals 2-cleft or none. Styles usually 3. Pod short, splitting to the base.

8. **Holosteum.** Petals denticulate or notched at the end. Styles usually 3. Pod opening at the apex by 6 teeth. Seeds fixed by their face.

9. **Cerastium.** Petals notched at the end or 2-cleft. Styles 5 or 4 (as many as the petals). Pod usually elongated, opening at the apex by 10 or 8 teeth. Seeds fixed edgewise.

** Styles alternate with the sepals: stamens as many as they, sometimes twice as many.

10. **Sagina.** Petals 4 or 5, undivided, or none. Styles 4 or 5. Pod 4-5-valved.

Tribe III. ILLECEBREÆ. Sepals separate or more or less united below. Petals without long claws, or minute, or often none, inserted under the sessile ovary or on the calyx. Pod 1-celled and splitting into valves, or a one-seeded utricle.—Leaves with dry,

scale-like stipules (except in *Scleranthus*), the uppermost sometimes alternate. Flowers mostly small.

* Pod (capsule) many-seeded. Styles 3-5. Petals usually conspicuous.

11. **Spergularia.** Styles 3-5. Leaves opposite.
12. **Spergula.** Styles 5. Valves of the pod opposite the sepals. Leaves whorled.
* * Pod (utricle) 1-seeded. Styles 2, often united. Petals none or minute.
13. **Anychia.** Stamens on the base of the 5-parted awnless calyx. Style hardly any.
14. **Paronychia.** Stamens on the base of the 5-parted calyx; the sepals bristle-pointed. Style 1, two-cleft at the top.
15. **Scleranthus.** Stamens borne on the throat of the indurated 5-cleft and pointless calyx. Styles 2. Stipules none.

Tribe IV. MOLLUGINEÆ. Stamens alternate with the sepals when of the same number, when fewer alternate with the cells of the 3-celled ovary. Partitions of the pod persistent on the valves. Leaves not truly opposite, otherwise as in Tribe II.

16. **Mollugo.** Petals none. Stamens 3-5. Stigmas 3. Pod many-seeded.

1. DIÁNTHUS, L. PINK. CARNATION.

Calyx cylindrical, nerved or striate, 5-toothed, subtended by 2 or more imbricated bractlets. Stamens 10. Styles 2. Pod 1-celled, 4-valved at the apex. Seeds flattish on the back: embryo scarcely curved.—Ornamental plants, of well-known aspect and value in cultivation. (Name from *Διός*, of *Jupiter*, and *ἄθος*, *flower*, i. e. Jove's own flower.) Two insignificant annual species are rarely spontaneous.

1. **D. ARMÉRIA, L.** (DEPTFORD PINK.) Flowers in close clusters; bractlets of the calyx and *bracts lance-awl-form*, herbaceous, downy, as long as the tube; leaves linear, *hairy*; petals small, rose-color with white dots, crenate.—Fields, &c., Virginia to E. Massachusetts. July.—(Adv. from Eu.)

2. **D. PRÓLIFER, L.** (PROLIFEROUS PINK.) *Smooth*, slender; flowers clustered; *bractlets ovate, dry*, concealing the calyx; leaves few, narrow, linear, erect; petals small, pink.—Near Philadelphia, *C. E. Smith.* (Adv. from Eu.)

2. SAPONÀRIA, L. SOAPWORT.

Calyx tubular, terete, nerveless, 5-toothed, naked at the base. Stamens 10. Styles 2. Pod short-stalked, 1-celled, or partly 2-celled at the base, 4-toothed at the apex.—Flowers clustered. (Name from *sapo*, soap, the mucilaginous juice forming a lather with water.)

1. **S. OFFICINÀLIS, L.** (COMMON SOAPWORT. BOUNCING BET.) Clusters corymbed; petals crowned with an appendage at the top of the claw; leaves oval-lanceolate.—Roadsides, &c. July-Sept.—A stout perennial, with large rose-colored flowers, commonly double. (Adv. from Eu.)

3. VACCÀRIA, Medik. COW-HERB.

Calyx naked at the base, ovoid-pyramidal, 5-angled, 5-toothed, enlarged and wing-angled in fruit. Petals not crowned. Stamens 10. Styles 2. Pod incompletely 4-celled at the base.—A smooth annual, with pale red flowers in corymbed cymes, and ovate-lanceolate leaves. (Name from *vacca*, a cow.)

1. **V. VULGÀRIS, Host.** (*Saponaria Vaccaria, L.*)—Escaped from gardens and becoming spontaneous in some places. (Adv. from Eu.)

4. SILÈNE, L. CATCHFLY. CAMPION.

Calyx 5-toothed, 10-many-nerved, naked at the base. Stamens 10. Styles 3, rarely 4. Pod 1-celled, sometimes 3-celled at least at the base, opening by 3 or 6 teeth at the apex. — Flowers solitary or in cymes. Petals mostly crowned with a scale at the base of the blade. (Name from *σίαλον*, *saliva*, from the viscid exudation on the stems and calyx of many species. The English name *Catchfly* alludes to the same peculiarity.)

* *Calyx bladderly-inflated: perennial: flowers paniced, white, in summer.*

1. **S. stellata**, Ait. (STARRY CAMPION.) *Leaves in whorls of 4, ovate-lanceolate, taper-pointed; calyx bell-shaped; petals cut into a fringe, crownless.* (*Cucùbalus stellatus*, L.) — Wooded banks, Rhode Island to Wisconsin, and southward. — Stem 3° high, with a large and open pyramidal panicle. Corolla $\frac{3}{4}$ ' broad.

2. **S. nivea**, DC. *Leaves opposite, lanceolate or oblong, taper-pointed; calyx oblong; petals wedge-form, 2-cleft, minutely crowned.* — Columbia, Pennsylvania, to Ohio and Illinois: rare. — Stem 1°–2° high, almost smooth. Flowers few, larger than in the last.

3. **S. INFLATA**, Smith. (BLADDER CAMPION.) *Glaucous; leaves opposite, ovate-lanceolate; calyx globular, much inflated, elegantly veined; petals 2-cleft, nearly crownless.* — Fields and roadsides, E. New England to Penn. — A foot high. Flowers loosely cymose. (Nat. from Eu.)

* * *Calyx elongated or club-shaped, not inflated except by the enlarging pod: flowers cymose or clustered: perennial, pubescent with viscid hairs, especially the calyx: petals crowned, red or rose-color.*

4. **S. Pennsylvànica**, Michx. (WILD PINK.) *Stems low (4'–8' high); root-leaves narrowly spatulate, nearly glabrous, tapering into hairy petioles; stem-leaves (2 or 3 pairs) lanceolate; flowers clustered, short-stalked; calyx club-shaped; petals wedge-form, slightly notched and eroded, pink.* — Gravelly places, E. New England to Penn., Kentucky, and southward. April–June.

5. **S. Virginica**, L. (FIRE PINK. CATCHFLY.) *Stems slender (1°–2° high); leaves thin, spatulate, or the upper oblong-lanceolate; flowers few and loosely cymose, peduncled; calyx oblong-cylindrical, soon obconical; petals oblong, 2-cleft, deep crimson; the limb 1' long.* — Open woods, W. New York (*Dr. Sartwell*) to Illinois and southward. June–Aug.

6. **S. règia**, Sims. (ROYAL CATCHFLY.) *Stem roughish, erect (3°–4° high); leaves thickish, ovate-lanceolate, acute; flowers numerous, short-stalked, in clusters, forming a strict panicle; calyx ovoid-club-shaped in fruit; petals spatulate-lanceolate, mostly undivided, deep scarlet.* — Prairies, Ohio to Illinois, and southward. July.

7. **S. rotundifolia**, Nutt. (ROUND-LEAVED CATCHFLY.) *Viscid-hairy; stems weak, branched, decumbent (2° long); leaves thin, round, abruptly pointed, the lower obovate; flowers few, loosely cymose, stalked; calyx elongated; petals 2-cleft and cut-toothed, deep scarlet.* — Shaded banks of the Ohio, and in Kentucky. June–Aug. — Leaves and flowers large. — The last three probably run together.

* * * *Calyx not inflated, except by the enlarging pod: annuals.*

+ *Glabrous, a portion of each joint of the stem glutinous: flowers pink.*

8. **S. ARMERIA**, L. (SWEET-WILLIAM CATCHFLY.) Glaucescent; leaves ovate-lanceolate; flowers in flat cymes, open in sunshine; calyx club-shaped; petals notched, crowned with awl-shaped scales. — Escaped from gardens: rare. (Adv. from Eu.)

9. **S. antirrhina**, L. (SLEEPY C.) Stem slender (8'–30' high); leaves lanceolate or linear; flowers small, paniculate; calyx ovoid; petals obcordate, crowned, opening transiently in sunshine. — Dry soil: common in waste places. June–Sept.

++ *Viscid-pubescent: flowers white or nearly so, opening at night, sweet-scented.*

10. **S. NOCTURNA**, L. (NIGHT C.) Leaves short, the lower spatulate, the upper linear; flowers small, alternate in a 1-sided spike; petals 2-parted. — Introduced sparingly in Pennsylvania, according to Schweinitz. (Adv. from Eu.)

11. **S. NOCTIFLORA**, L. (NIGHT-FLOWERING C.) Viscid-hairy, tall (1°–3° high); lower leaves large and spatulate; the upper lanceolate; flowers few, peduncled; calyx-tube elongated (over 1' long), soon ovoid, with awl-shaped teeth; petals rather large, 2-parted, crowned. — Cultivated grounds. (Nat. from Eu.)

* * * *Dwarf, tufted, smooth, perennial: flowering shoots 1-flowered.*

12. **S. ACAULIS**, L. (MOSS CAMPION.) Tufted like a moss (1'–2' high), leaves linear, crowded; flowers almost sessile, or rarely on a naked peduncle; petals purple or rarely white, notched or entire, crowned. — Alpine summits of the White Mountains, New Hampshire. July. (Eu.)

5. LÝCHNIS, TOURN. LYCHNIS. COCKLE.

Styles 5, rarely 4; and pod opening by as many or twice as many teeth: otherwise nearly as in *Silene*. Calyx in one species with leaf-like lobes. (Ancient Greek name for a scarlet or flame-colored species, from *λύχνος*, a light or lamp.)

1. **L. VESPERTINA**, Sibth. (EVENING L.) Biennial, usually dioecious, viscid pubescent, in foliage, &c., like *Silene noctiflora*; but 5 styles; calyx much shorter, the fertile enlarging and broadly ovoid in fruit, with lance-linear teeth; flowers white or pinkish, opening at evening. (*L. dioica*, *L.* in part.) — Cult. or waste grounds: scarce. (Adv. from Eu.)

2. **L. GITHAGO**, Lam. (CORN COCKLE.) Annual, clothed with long soft appressed hairs; flowers long-peduncled; calyx-lobes similar to the long and linear leaves, surpassing the broad and crownless purple-red petals, falling off in fruit. (*Agrostémna Githago*, *L.*, & ed. 2.) — A weed in wheat-fields, too common, the black seeds of Cockle being injurious to the appearance of the flour. (Adv. from Eu.)

6. ARENÀRIA, L. SANDWORT.

Sepals 5. Petals 5, entire, sometimes barely notched, rarely wanting. Stamens 10. Styles 3, rarely more or fewer, opposite as many sepals. Pod short, splitting into as many or twice as many valves as there are styles, few-many-

seeded. — Low, usually tufted herbs, with sessile exstipulate leaves and small white flowers. (Name from *arena*, sand, in which many of the species grow.) — The following sections are by many botanists taken for genera, as they were in the former edition.

§ 1. **ARENARIA** proper. *Pod splitting wholly or part-way down into 3 or at length into 6 valves: seeds many, naked at the hilum.*

1. **A. SERPYLLIFOLIA**, L. (THYME-LEAVED SANDWORT.) Diffusely branched, roughish (2'–6' high); leaves ovate, acute, small; cymes leafy; sepals lanceolate, pointed, 3–5-nerved, about as long as the petals and the 6-toothed pod. — A low annual, in sandy waste places. June–Aug. (Nat. from Eu.)

§ 2. **ALSINE**, (Tourn.) Wahl. *Pod splitting to the base into 3 entire valves: seeds many, usually rough, naked at the hilum: flowers solitary and terminal or cymose: root in our species perennial.*

* *Leaves small, rigid, awl-shaped or bristle-shaped.*

2. **A. squarrosa**, Michx. (PINE-BARREN S.) Densely tufted from a deep perpendicular root; leaves closely imbricated, but spreading, awl-shaped, short, channelled; branches naked and minutely glandular above, several-flowered; sepals obtuse, ovate, shorter than the pod. (Alsine, ed. 2.) — In pure sand, S. New York, New Jersey, and southward along the coast. May–July.

3. **A. stricta**, Michx. Erect, or usually diffusely spreading from a small root, smooth; leaves slender, between awl-shaped and bristle-form, with many others clustered in the axils; cyme diffuse, naked, many-flowered; sepals pointed, 3-ribbed, ovate, as long as the pod. (Alsine Michauxii, Fenzl.) — Rocks and dry wooded banks, Vermont to Wisconsin and Kentucky. July. — The specific name is a bad one, as there is nothing strict about the plant.

* * *Leaves soft and herbaceous, filiform-linear: petals retuse or notched.*

4. **A. pátula**, Michx. Diffusely branched from the slender root; stems filiform (6'–10' long); branches of the cyme diverging; peduncles long; sepals lanceolate, acuminate, 3–5-nerved. (Alsine, ed. 2.) — Cliffs of Kentucky River, mountains of Western Virginia, and southward.

5. **A. Grœnlândica**, Spreng. (MOUNTAIN S.) Densely tufted from slender roots, smooth; flowering stems filiform, erect (2'–4' high), few-flowered; sepals oblong, obtuse, nerveless. (Stellaria Grœnlandica, Retz. Alsine, ed. 2.) — Summit of the Shawangunk, Catskill, and Adirondack Mountains, New York, of all the higher mountains of New England, and northward; alpine or subalpine. At Bath, Maine, on river-banks near the sea. June–Aug. — Leaves and peduncles 3"–6" long; flowers large in proportion.

A. **GLABA**, Michx., of the mountain-tops in Carolina, may occur on those of Virginia, and is perhaps a large form of the above.

§ 3. **MœHRÍNGIA**, L. *Parts of the flower sometimes in fours: pod as in § 1, but the young ovary 3-celled: seeds rather few, smooth and with a thickish appendage (strophiole) at the hilum: perennials, with flaccid broadish leaves.*

6. **A. lateriflora**, L. Sparingly branched, erect, minutely pubescent; leaves oval or oblong obtuse ($\frac{3}{4}$ '–1' long); peduncles 2- (rarely 3–4-) flowered,

soon becoming lateral; sepals oblong, obtuse. — Gravelly shores, &c., Rhode Island to Pennsylvania, Wisconsin, and northward. May, June. (Eu.)

§ 4. **AMMADÈNIA**, Gmelin, (Honkénya, Ehrh.) Pod fleshy, splitting into as many valves as there are styles (3, rarely 4 or 5); the ovary more or less 3- (5-) celled: seeds few, smooth, short-beaked at the naked hilum: disk under the ovary more prominent than usual, glandular, 10-lobed; flowers almost sessile in the axils of fleshy leaves, sometimes dioecious or polygamous: root perennial.

7. **A. peploides**, L. Stems (simple or forking from long rootstocks, 6'–10' high) and ovate partly-clasping leaves (8"–10" long) very fleshy. (Honkénya peploides, Ehrh., ed. 2.) — Sands of the sea-shore, New Jersey to Maine and northward. June. (Eu.)

7. STELLÀRIA, L. CHICKWEED. STARWORT.

Sepals 4–5. Petals 4–5, deeply 2-cleft, sometimes none. Stamens 8, 10, or fewer. Styles 3, rarely 4 or 5, opposite as many sepals. Pod ovoid, 1-celled, opening by twice as many valves as there are styles, several-many-seeded. Seeds naked. — Flowers (white) solitary or cymose, terminal, or appearing lateral by the prolongation of the stem from the upper axils. (Name from *stella*, a star, in allusion to the star-shaped flowers.)

* Stems spreading, flaccid, marked longitudinally with one or two pubescent lines: leaves ovate or oblong, $\frac{1}{2}$ '– $2\frac{1}{2}$ ' long.

1. **S. MÈDIA**, Smith. (COMMON CHICKWEED.) Annual* or nearly so; lower leaves on hairy petioles; petals shorter than the calyx, 2-parted; stamens 3–10. — Everywhere in damp grounds. (Nat. from Eu.)

2. **S. pùbera**, Michx. (GREAT CHICKWEED.) Root perennial; leaves all sessile; petals longer than the calyx, deeply 2-cleft; stamens 10. — Shaded rocks, Pennsylvania to Kentucky and southward. May.

** Stems erect or spreading: wholly glabrous perennials, with sessile and narrow or small leaves: stamens usually 10, perigynous.

+ Scaly-bracted: petals 2-parted, equalling or surpassing the calyx.

3. **S. longifolia**, Muhl. (LONG-LEAVED STITCHWORT.) Stem erect, weak, often with rough angles (8'–18' high); leaves linear, acutish at both ends, spreading; cymes naked and at length lateral, peduncled, many-flowered, the slender pedicels spreading; petals 2-parted, longer than the calyx; seeds smooth. — Grassy places: common, especially northward. June, July. (Eu.)

4. **S. longipes**, Goldie. (LONG-STALKED S.) Shining or somewhat glaucous, very smooth; leaves ascending, lanceolate or linear-lanceolate, acute, broadest at the base, rather rigid; cyme terminal, few-flowered, the long pedicels strictly erect; petals longer than the calyx; seeds smooth. — Maine to Wisconsin: rare, common farther north. (Eu.)

5. **S. uliginosa**, Murr. (SWAMP S.) Stems weak, decumbent or diffuse, at length prolonged, leaving the naked and usually sessile cymes lateral; leaves lanceolate or oblong, veiny; petals and ripe pods as long as the calyx; seeds roughened. (S. aquatica, Pollich.) — Swamps and rills, Pennsylvania (Darlington, &c.), New Hampshire (Blake), and northward in British America. (Eu.)

← ← *Leafy-bracted, the flowers terminal or in the forks of the stem or of leafy branches; bracts foliaceous: petals 2-parted, small or often none: styles 3-4: pod longer than the calyx.*

6. **S. crassifolia**, Ehrhart. Stems diffuse or erect, flaccid; leaves rather fleshy, varying from linear-lanceolate to oblong; petals longer than the calyx, or wanting; seeds rugose-roughened. — (An apetalous 4-6-androus state is *Sagina fontinalis*, *Short & Peter*.) Springy places, E. Kentucky (*Short*), Ringwood, Illinois (*Vasey*), and northward. April-June. (Eu.)

7. **S. borealis**, Bigelow. (NORTHERN S.) Stems erect or spreading, flaccid, many times forked, at length resolved into a leafy cyme; leaves varying from broadly lanceolate to ovate-oblong; petals 2-5, shorter than the calyx, or oftener none; sepals acute; styles usually 4; seeds smooth. — Shaded or wet places, Rhode Island to Wisconsin and northward. June-Aug. — Var. **ALPÉSTRIS** (*S. alpestris*, *Fries*, *S. Fenzlii*, *Regel*) has the later flowers more cymose, and their bracts small and partly scarious, also the seeds obscurely reticulated or roughish. — Lake Superior, *Dr. Robbins*. (Eu.)

8. **S. humifusa**, Rottbœll. Spreading or creeping; stems or branches (2' high) 1-3-flowered; leaves fleshy, ovate or oblong (2''-3'' long); petals a little longer than the calyx; seeds smooth. — Northern border of Maine on the St. John's (*G. L. Goodale*), and high northward. June. (Eu.)

8. HOLÓSTEUM, L. JAGGED CHICKWEED.

Sepals 5. Petals 5, usually jagged or denticulate at the point. Stamens 3-5, rarely 10. Styles mostly 3. Pod ovoid, 1-celled, many-seeded, opening at the top by 6 teeth. Seeds rough, flattened on the back, attached by the inner face. — Annuals or biennials, with several (white) flowers in an umbel, borne on a long terminal peduncle. (Name composed of *όλος*, *all*, and *όστέον*, *bone*, by antiphrasis, these plants being soft and tender.)

1. **H. UMBELLATUM**, L. Leaves oblong; peduncle and upper part of the stem glandular-pubescent; pedicels reflexed after flowering. — Hills around Lancaster, Pennsylvania, abundant, *Prof. Porter*. (Nat. from Eu.)

9. CERÁSTIUM, L. MOUSE-EAR CHICKWEED.

Sepals 5, rarely 4. Petals as many, 2-lobed or cleft, rarely entire. Stamens twice as many, or fewer. Styles equal in number to the sepals and opposite them. Pod 1-celled, usually elongated, membranaceous, opening at the apex by twice as many teeth as there were styles, many-seeded. Seeds rough. — Flowers white, in terminal cymes. (Name from *κέρας*, a *horn*, alluding to the shape of the pods in many species.)

§ 1. *Petals 2-cleft or obcordate: parts of the flower in fives: pods (except in No. 5) longer than the calyx, and usually more or less curved.*

1. **C. VULGATUM**, L. (MOUSE-EAR CHICKWEED.) Annual, hairy and rather clammy, nearly erect (4'-9' high); leaves ovate or obovate; bracts herbaceous; flowers (small) in close clusters at first; pedicels even in fruit not longer than the acute sepals; petals shorter than the calyx. — Grassy places, eastward and southward: not common. May-July. (The names of this and the next were

transposed by Linnæus, and by continental botanists ever since.) — Stamens often 5. — The var. ? SEMIDECÁNDRUM, which has more *lengthened fruit-bearing pedicels*, is here hardly met with. (Nat. from Eu.)

2. **C. VISCOSUM**, L. (LARGER M.) Perennial; stems clammy-hairy, spreading (6'–15' long); *leaves oblong*; upper bracts scarious-margined; flowers at first clustered, the fruiting *pedicels* longer, the earlier ones mostly much *longer than the obtuse sepals*; petals equalling the calyx. — Fields and copses: common, perhaps indigenous to the country. May–July. (Nat. from Eu.)

3. **C. NUTANS**, Raf. Annual, very clammy-pubescent; stems erect, slender, grooved, diffusely branched (6'–20' high); cyme loose and open, *many-flowered*; *leaves oblong-lanceolate*, acute, the lowest spatulate; peduncles mostly elongated; petals longer than the calyx; *Pods nodding on the stalks, curved upwards, thrice the length of the calyx.* — Moist places, Vermont to Minnesota and southward. May–July. — Var. BRACHYPODUM, Engelm., W. Illinois and southwestward, has *pedicels shorter than the pods.*

4. **C. OBLONGIFOLIUM**, Torr. Perennial; stems ascending, villous (6'–12' high), *many-flowered*; *leaves oblong-lanceolate* and ovate; peduncles clammy-hairy; petals (2-lobed) and ripe *Pods about twice the length of the calyx.* — Rocky places, New York to N. Virginia and Illinois: rare. May–July. — Stouter and larger flowered than the following species.

5. **C. ARVENSE**, L. (FIELD CHICKWEED.) Perennial; stems ascending or erect, tufted, downy, slender (4'–8' high), naked and *few-several-flowered* at the summit; *leaves linear*; petals obcordate, more than twice the length of the calyx; *Pods scarcely longer than the calyx.* Dry or rocky places, New England to Wisconsin and northward. May–July. (Eu.)

§ 2. **MËNCHINA**, Ehrhart. *Petals entire or merely retuse: parts of the flower commonly in fours: pod ovate, not longer than the calyx.*

6. **C. QUATERNELLUM**, Fenzl. Smooth and glaucous annual; stem simple, erect (2'–4' high), 1–2-flowered; leaves lanceolate, acute; petals not exceeding the calyx; stamens 4. (*Sagina erecta*, L. *Mœnchia quaternella*, Ehrh.) — Near Baltimore, in dry ground. (Adv. from Eu.)

10. SAGINA, L. PEARLWORT.

Sepals 4 or 5. Petals 4 or 5, undivided, or often none. Stamens as many as the sepals, rarely twice their number. Styles as many as the sepals and alternate with them. Pod many-seeded, 4–5-valved to the base; valves opposite the sepals. — Little, matted herbs, with thread-like or awl-shaped leaves, no stipules, and small flowers terminating the stems or branches; in summer. (Name from *sagina*, fattening; of dubious application.)

* *Parts of the flower in fours, rarely with some few in fives.*

1. **S. PROCUMBENS**, L. Perennial, depressed or spreading on the ground, glabrous; leaves linear-thread-shaped; apex of the peduncle often hooked soon after flowering; *petals shorter than the broadly ovate obtuse sepals*, sometimes none. — Springy places and damp rocks, coast of Maine to Pennsylvania. (Eu.)

2. **S. APÉTALA**, L. Annual, erect, with more slender leaves, narrower sepals, and *petals none or obsolete.* — Dry soil, New York and Pennsylvania to Illinois: scarce, seemingly native? (Eu.)

* * *Parts of the flower in fives, the stamens not rarely 10.*

3. **S. subulata**, Wimmer. Perennial (or apparently annual), ascending; the peduncles and calyx with the margins of the upper leaves at first glandular-pubescent; leaves short, often bristly-tipped, not fascicled in the axils; peduncles slender; petals equalling or shorter than the calyx (*S. Elliottii*, Fenzl.), — or in var. *SMITHII*, a slender form, seemingly annual, *apetalous*, at least in the later flowers. — Near Philadelphia, in waste ground, and sandy fields, &c., Somers' Point, New Jersey, *C. E. Smith*. — This form has the aspect of No. 2. Seeds minutely roughened. (Perhaps nat. from Eu.)

4. **S. nodosa**, Fenzl. Perennial, tufted, glabrous, or glandular above; stems ascending (3'–5' high; lower leaves thread-form, the upper short and awl-shaped, with minute ones fascicled in their axils so that the branchlets appear knotty; petals much longer than the calyx. (*Spergula nodosa*, L.) — Wet sandy soil, along the coast of Maine, New Hampshire, also Lake Superior, and northward. (Eu.)

11. SPERGULARIA, Pers. SAND-SPURRY.

Sepals 5. Petals 5, entire. Stamens 2–10. Styles and valves of the many-seeded pod 3, very rarely 5, when the valves alternate with the sepals! Embryo not coiled into a complete ring. — Low herbs, mostly on or near the sea-coast, with filiform or linear somewhat fleshy opposite leaves, and smaller ones often clustered in the axils: stipules scaly-membranaceous: flowering all summer. (Name altered from *Spergula*.) — Genus also known under the name of *LEPIGONUM*, Fries. Our species, which perhaps run together, are here arranged in view of Kindberg's monograph, but with some reduction. They are all annuals and subperennials.

1. **S. rubra**, Presl., var. **campéstris**. Nearly glabrous, except the summit of the prostrate slender stems, peduncles, and sepals, which are usually glandular-pubescent; leaves filiform; stipules triangular-awl-shaped; pods and pink-red corolla small (2''), hardly equalling or exceeding the calyx; seeds rough with projecting points, semi-obovate or gibbous-wedge-shaped, wingless. — Sandy or gravelly dry soil, New England to Virginia along and near the coast, but rarely maritime. (Eu.)

2. **S. salina**, Presl. Larger and more decidedly fleshy than the preceding, with ovate stipules, and peduncles rarely longer than the pod, which is longer than the calyx (3'' long); petals pale; seeds obovate-rounded and roughened with points, wingless or narrow-winged. — Brackish sands, &c., coast of New England to Virginia and southward. (Eu.)

3. **S. média**, Presl. Distinguished from the last mainly by the smooth seeds, either winged or wingless; peduncles equalling or exceeding the pod, which is 2'' or 3'' long, and a little exceeds the calyx. (*Lepigonum medium*, Fries. *L. leiospermum*, Kindberg.) — Salt marshes and sands, with the last. (Eu.)

Var. **macrocarpa**. (*S. macrocarpa*, Presl. *Lepigonum marinum*, Fries.) Root more woody and perennial; pod 3½''–5'' long, surpassing the calyx; seeds also larger, rounded, broadly winged, or a few wingless. — Sea-beaches, rare northward. (Eu.)

12. SPÉRGULA, L. SPURREY.

Stamens 5 or 10. Styles 5. The 5 valves of the pod opposite the sepals. Embryo spirally annular. Leaves in whorls. Otherwise as in Spergularia. (Name from *spargo*, to scatter, from the seeds.)

1. **S. ARVÉNSIS**, L. (CORN SPURREY.) Annual; leaves numerous in the whorls, thread-shaped (1' - 2' long); stipules minute; flowers white, in a stalked paniced cyme; seeds rough. — Grain-fields. (Adv. from Eu.)

13. ANÝCHIA, Michx. FORKED CHICKWEED.

Sepals 5, scarcely concave, indistinctly mucronate on the back, greenish. Petals none. Stamens 2-3, rarely 5. Stigmas 2, sessile. Utricle 1-seeded, larger than the calyx. Radicle turned downwards. — Small, many times forked annuals, with small stipules, and minute flowers in the forks, produced all summer. (Same derivation as the next genus.)

1. **A. dichótoma**, Michx. Erect or spreading; leaves varying from lanceolate to elliptical, somewhat petioled. Varies much; in woods or rich soil being very smooth, erect (6' - 10' high) and capillary, with long joints, the leaves broader and thinner (5'' - 10'' long), and the flowers more stalked (*A. capillacea*, Nutt. and *Queria Canadensis* L.); in sterile or parched soil it is pubescent, low and spreading, short-jointed, narrower-leaved, and the flowers nearly sessile and more clustered (*A. dichotoma*, DC.). Common throughout.

14. PARONÝCHIA, Tourn. WHITLOW-WORT.

Sepals 5, linear or oblong concave, awned at the apex. Petals bristle-form, or minute teeth, or none. Stamens 5. Style 2-cleft at the apex. Utricle 1-seeded, enclosed in the calyx. Radicle ascending. — Tufted herbs, with dry and silvery stipules, and clustered flowers. (Greek name for a *whitlow*, and for a plant thought to cure it.)

1. **P. argyrócoma**, Nutt. Forming broad tufts, branched, *spreading*; leaves linear; flowers densely clustered, surrounded by conspicuous large silvery bracts; calyx hairy, short-awned; petals mere teeth between the stamens. ¶ — Slides in the Notch of the White Mountains, New Hampshire, and bare summits above. Alleghany Mountains from Virginia southward. July.

2. **P. dichótoma**, Nutt. Smooth, tufted; stems (6' - 12' high) *ascending* from a rather woody base; leaves and bracts *awl-shaped*; cymes open, many-times forked; sepals short-pointed; minute bristles in place of petals. ¶ — Rocks, Harper's Ferry, Virginia, and southwestward. July - Sept.

15. SCLERÁNTHUS, L. KNAWEL.

Sepals 5, united below in an indurated cup, enclosing the 1-seeded utricule. Petals none. Stamens 10 or 5. Styles 2, distinct. — Homely little weeds, with awl-shaped leaves, obscure greenish clustered flowers, and no stipules. (Name from *σκληρός*, *hard*, and *ἄνθος*, *flower*, from the hardened calyx-tube.)

1. **S. ANNUS**, L. Much branched, spreading (3' - 5' high); flowers sessile in the forks; calyx-lobes scarcely margined. — Waste places. (Nat. from Eu.)

16. MOLLÛGO, L. INDIAN-CHICKWEED.

Sepals 5, white inside. Petals none. Stamens hypogynous, 5 and alternate with the sepals, or 3 and alternate with the 3 cells of the ovary. Stigmas 3. Pod 3-celled, 3-valved, loculicidal, the partitions breaking away from the many-seeded axis. — Low homely annuals, much branched; the stipules obsolete. (An old Latin name for some soft plant.)

1. **M. verticillata**, L. (CARPET-WEED.) Prostrate, forming patches; leaves spatulate, clustered in whorls at the joints, where the 1-flowered pedicels form a sort of sessile umbel; stamens usually 3. — Sandy river-banks, and cultivated grounds. June – Sept. (An immigrant from farther south.)

ORDER 19. PORTULACACEÆ. (PURSLANE FAMILY.)

Herbs, with succulent leaves, and regular but unsymmetrical flowers; viz., sepals usually fewer than the petals; the stamens opposite the petals when of the same number, but often indefinite: otherwise nearly as Chickweeds. — Sepals 2, rarely 3 or 5. Petals 5, or sometimes none. Stamens mostly 5 – 20. Styles 3 – 8, united below, or distinct, stigmatic along the inside. Pod 1 – 5-celled, with few or many campylotropous seeds rising on stalks from the base, or from a central placenta. Embryo curved around mealy albumen. — Insipid and innocent herbs, with entire leaves. Corolla opening only in sunshine, mostly ephemeral, then shrivelling.

* ANOMALOUS PORTULACACEÆ. Sepals 5, bearing the stamens.

1. **Sesuvium**. Petals none. Stamens 5 – 60. Pod 3 – 5-celled, opening across by a lid.

** TRUE PORTULACACEÆ. Sepals (2) fewer than the petals: pod 1-celled.

2. **Portulaca**. Stamens 7 – 20, on the partly adherent calyx. Pod opening by a lid.

3. **Talinum**. Stamens more numerous than the petals, hypogynous. Pod many-seeded.

4. **Claytonia**. Stamens as many as the hypogynous petals, and attached to their base. Pod 3 – 6-seeded.

1. SESÛVIUM, L. SEA PURSLANE.

Calyx 5-parted, purplish inside, persistent, free. Petals none. Stamens 5 – 60, inserted on the calyx. Styles 3 – 5, separate. Pod 3 – 5-celled, many-seeded, circumscissile, the upper part falling off as a lid. — Prostrate maritime herbs, with succulent stems, opposite leaves, and axillary or terminal flowers. (An unexplained name.)

1. **S. Portulacastrum**, L. Root perennial; leaves lanceolate-oblong, flattish; flowers sessile or short-peduncled; stamens many. — Coast of New Jersey and southward. July – Sept.

2. PORTULÀCA, Tourn. PURSLANE.

Calyx 2-cleft; the tube cohering with the ovary below. Petals 5, rarely 6, inserted on the calyx with the 7 – 20 stamens, fugacious. Style mostly 3 – 8-parted. Pod 1-celled, globular, many-seeded, opening transversely, the upper part (with the upper part of the calyx) separating as a lid. — Fleshy annuals, with mostly scattered leaves. (An old Latin name, of unknown meaning.)

1. *P. OLERÀCEA*, L. (COMMON PURSLANE.) Prostrate, very smooth; leaves obovate or wedge-form; flowers sessile (opening only in sunny mornings); sepals keeled; petals pale yellow; stamens 7-12; style deeply 5-6-parted; flower-bud flat and acute. — Cultivated and waste grounds: common. (Nat. from Eu.)

P. RETÙSA, Engelm., closely resembling the common Purslane, is indigenous west of the Mississippi.

P. GRANDIFLÒRA, with terete leaves and showy flowers, cult. for ornament, begins to be spontaneous around gardens.

3. TALÌNUM, Adans. TALINUM.

Sepals 2, distinct and free, deciduous. Petals 5, ephemeral. Stamens 10-30. Style 3-lobed at the apex. Pod 3-celled at the base when young, 3-valved, with many seeds on a globular stalked placenta. (Derivation obscure.)

1. *T. teretifòlium*, Pursh. Leafy stems low, tuberous at the base; leaves linear, cylindrical; peduncle long (3'-6') and naked, bearing an open cyme of pink flowers ($\frac{2}{3}$ ' broad); stamens 15-20. ♀ — Serpentine rocks, Westchester, Penn., Falls of St. Croix River, Wisconsin, and southward. June-Aug.

4. CLAYTÒNIA, L. SPRING-BEAUTY.

Sepals 2, ovate, free, persistent. Stamens 5, adhering to the short claws of the petals. Style 3-cleft at the apex. Pod 1-celled, 3-valved, 3-6-seeded. — Our two species are perennials, sending up simple stems in early spring from a small deep tuber, bearing a pair of opposite leaves, and a loose raceme of pretty flowers. Corolla rose-color with deeper veins, opening for more than one day! (Named in honor of *Dr. John Clayton*, one of our earliest botanists, who contributed to Gronovius the materials for the *Flora Virginica*.)

1. *C. Virgínica*, L. Leaves linear-lanceolate, elongated (3'-6' long). — Moist open woods: common, especially westward and southward.

2. *C. Caroliniàna*, Michx. Leaves spatulate-oblong or oval-lanceolate (1'-2' long). — W. New Hampshire, to Wisconsin, and southward along the Alleghanies.

ORDER 20. MALVÀCEÆ. (MALLOW FAMILY.)

Herbs or shrubs, with alternate stipulate leaves and regular flowers, the calyx valvate and the corolla convolute in the bud, numerous stamens monadelphous in a column, which is united at the base with the short claws of the petals, 1-celled anthers, and kidney-shaped seeds. — Sepals 5, united at the base, persistent, often involucellate with a whorl of bractlets, forming a sort of exterior calyx. Petals 5. Anthers kidney-shaped, opening along the top. Pistils several, with the ovaries united in a ring, or forming a several-celled pod. Seeds with little albumen: embryo curved, the leafy cotyledons variously doubled up. — Mucilaginous, innocent plants, with tough bark, and palmately-veined leaves. Flowerstalks with a joint, axillary.

Tribe I. MALVEÆ. Columns of stamens anther-bearing at the top. Ovaries and pods (carpels) 5-20 or more, closely united in a ring around a central axis, from which they separate after ripening.

* Stigmas occupying the inner face of the styles : carpels 1-seeded, falling away separately.

1. **Althæa.** Involucel of 6 to 9 bractlets.
2. **Malva.** Involucel of 3 bractlets. Petals obovate. Carpels rounded, beakless.
3. **Callirrhoe.** Involucel of 3 bractlets or none. Petals truncate. Carpels beaked.
4. **Napæa.** Involucel none. Flowers diœcious. Stamens few.

** Stigmas terminal, capitate : carpels 1 - few-seeded, usually dehiscent.

- 5 **Malvastrum.** Involucel of 3 bractlets or none. Seeds solitary in the cells, ascending.
6. **Sida.** Involucel none. Seed solitary in the cells, pendulous.
7. **Abutilon.** Involucel none. Seeds 3 or more in each cell.
8. **Modiola.** Involucel of 3 bractlets. Seeds 2 in each cell, and with a transverse partition between them.

Tribe II. HIBISCEÆ. Column of stamens anther-bearing for a considerable part of its length, naked and 5-toothed at the very apex. Pod mostly 5-celled, loculicidal, leaving scarcely any axis in the centre after opening.

9. **Kosteletzkya.** Involucel of several bractlets. Pod 5-celled, 5-seeded.
10. **Hibiscus.** Involucel of many bractlets. Pod 5-celled, many-seeded.

1. ALTHÆA, L. MARSH-MALLOW.

Calyx surrounded by a 6-9-cleft involucel. Otherwise as in *Malva*. (Name from *ἄλθω*, to cure, in allusion to its healing properties.)

1. **A. OFFICINALIS, L.** (COMMON MARSH-MALLOW.) Stem erect; leaves ovate or slightly heart-shaped, toothed, sometimes 3-lobed, velvety-downy: peduncles axillary, many flowered. — Salt marshes, coast of New England and New York. Aug., Sept. — Flowers pale rose-color. Perennial root thick, abounding in mucilage, the basis of the *Pâtes de Guimauve*. (Nat. from Eu.)

2. MÁLVA, L. MALLOW.

Calyx with a 3-leaved involucel at the base, like an outer calyx. Petals obovate. Styles numerous, stigmatic down the inner side. Fruit depressed, separating at maturity into as many 1-seeded and indehiscent round kidney-shaped blunt carpels as there are styles. Radicle pointing downwards. (An old Latin name, from *μαλᾶχη*, soft, alluding to the emollient leaves.)

1. **M. ROTUNDIFÓLIA, L.** (COMMON MALLOW.) Stems procumbent from a deep biennial root; leaves round-heart-shaped, on very long petioles, crenate, obscurely-lobed; petals twice the length of the calyx, whitish; carpels pubescent, even. — Waysides and cultivated grounds: common. (Nat. from Eu.)

2. **M. SYLVÉSTRIS, L.** (HIGH M.) Biennial; stem erect, branched (2°-3° high); leaves sharply 5-7-lobed; petals thrice the length of the calyx, large, purple and rose-color; carpels wrinkled-veiny. — Waysides. (Adv. from Eu.)

3. **M. CRÍSPA, L.** (CURLLED M.) A tall, erect annual, with round and angled toothed and crisped leaves, and small sessile flowers crowded in the axils, — sparingly escaped from old gardens. (Adv. from Eu.)

4. **M. MOSCHÁTA, L.** (MUSK M.) A low perennial; with the stem-leaves 5-parted, and the divisions once or twice parted or cleft into linear lobes, faintly musky-scented, the flowers rose-color or white (1½' in diameter) on short pe-

duncles crowded on the stem and branches, the fruit downy: has escaped from gardens to waysides. (Adv. from Eu.)

5. **M. ÁLCEA**, L., with the *stem-leaves only once 5-parted or cleft*, the lobes incised, large flowers like No. 4, but the fruit smooth, and bractlets of the involucl ovate: has escaped from gardens in Chester Co., Penn. (Adv. from Eu.)

3. CALLÍRRHOË, Nutt. CALLIRRHOË.

Calyx either naked or with a 3-leaved involucl at its base. Petals wedge-shaped and truncate (usually red-purple). Styles, &c. as in *Malva*. Carpels 10-20, straightish, with a short empty beak, separated within from the 1-seeded cell by a narrow projection, indehiscent or partly 2-valved. Radicle pointing downwards. — Flowers perfect.

1. **C. triangulata**, Gray. Hairy-pubescent; stems nearly erect (2° high) from a tuberous root; leaves triangular or halberd-shaped, or the lower rather heart-shaped, coarsely crenate; the upper incised or 3-5-cleft; flowers paniced, short-pedicelled (purple); *involucl as long as the calyx; carpels short-pointed, crestless*. (*Malva triangulata*, *Leavenworth*. M. Houghtonii, *Torr. & Gray*.) — Dry prairies, Wisconsin, Illinois, and southward. July.

2. **C. alcæoides**, Gray. Strigose-pubescent; stems slender (1° high) from a perennial root; lower leaves triangular-heart-shaped, incised; the upper 5-7-parted, lacinate, the uppermost divided into linear segments; flowers corymbose, on slender peduncles (rose-color or white); *involucl none; carpels obtusely beaked, crested and strongly wrinkled on the back*. (*Sida alcæoides*, *Michx.*) — Barren oak-lands, S. Kentucky and Tennessee.

4. NAPÆA, Clayt. GLADE MALLOW.

Calyx naked at the base, 5-toothed. Petals entire. *Flowers diœcious*; the staminate flowers destitute of pistils, with 15-20 anthers; the fertile with a short column of filaments but usually no anthers. Styles 8-10, stigmatic along the inside. Fruit depressed-globular, separating when ripe into as many kidney-shaped 1-seeded beakless and scarcely dehiscent carpels as there are styles. Radicle pointing downwards. — A tall and roughish perennial herb, with very large 9-11-parted lower leaves, the pointed lobes pinnatifid-cut and toothed, and small white flowers in paniced clustered corymbs. (Named by Clayton from *νᾶπη*, a wooded valley or glade, or, poetically, the nymph of the groves.)

1. **N. diœica**, L. (*Sida diœica*, *Cav.*) — Limestone valleys, Penn. and southward to the Valley of Virginia, west to Ohio and Illinois: rare. July.

5. MALVÁSTRUM, Gray. FALSE MALLOW.

Calyx with an involucl of 2 or 3 bractlets, or none. Petals notched at the end or entire. Styles 5 or more: stigmas capitate. Carpels as in *Malva*, or else as in *Sida*, but the solitary kidney-shaped seed ascending and the radicle pointing downwards, as in the former. — Flowers perfect. (Name altered from *Malva*.)

1. **M. angústum**, Gray. Slightly hairy; stem erect (6'-9' high) from an annual root; leaves lance-oblong or linear, with scattered fine callous teeth; flowers in the upper axils, on peduncles shorter than the broadly ovate-triangular sepals; bractlets and stipules setaceous; petals yellow, scarcely exceeding the calyx; carpels 5, kidney-shaped, with smooth sides, at length 2-valved. — Rock Island in the Mississippi, Illinois, *Engelmann, Parry*. Aug.

2. **M. coccíneum**, Gray (*Sida coccinea*, Pursh), a low and hoary perennial, with 5-parted or pedate leaves, and short spikes or racemes of showy pink-red flowers, the petals very much longer than the calyx; the carpels 10 or more, reticulated on the sides and indehiscent: abounds on the plains from Iowa and Minnesota westward.

6. SIDA, L. SIDA.

Calyx naked at the base, 5-cleft. Petals entire, usually oblique. Styles 5 or more, tipped with capitate stigmas: the ripe fruit separating into as many 1-seeded carpels, which are closed, or commonly 2-valved at the top, and tardily separate from the axis. Seed pendulous. Embryo abruptly bent; the radicle pointing upwards. — Flowers perfect. (A name used by Theophrastus.)

1. **S. Napæa**, Cav. A smooth, tall (4°-10° high) perennial; leaves 5-cleft, the lobes oblong and pointed, toothed; flowers (white) umbellate-corymbed, large; carpels 10, pointed. (*Napæa lævis* & *hermaphrodita*, L.) — Rocky river-banks, Penn., York Co., &c., *Porter*. Kanawha Co., Virginia, *Rev. J. M. Brown*. (Cultivated in old gardens.)

2. **S. Ellióttii**, Torr. & Gray. A smooth, erect perennial (1°-4° high); leaves linear, serrate, short-petioled; peduncles axillary, 1-flowered, short; flowers (yellow) rather large; carpels 9-10, slightly and abruptly pointed, forming a depressed fruit. — Sandy soil, S. Virginia and southward. May - Aug.

3. **S. SPINOSA**, L. Annual weed, minutely and softly pubescent, low (10'-20' high), much branched; leaves ovate-lanceolate or oblong, serrate, rather long-petioled; peduncles axillary, 1-flowered, shorter than the petiole; flowers (yellow) small; carpels 5, combined into an ovate fruit, each splitting at the top into 2 beaks. — A little tubercle at the base of the leaves on the stronger plants gives the specific name, but it cannot be called a spine. — Waste places: common southward. (Nat. from Trop. Amer. or Afr.)

7. ABÛTILON, Tourn. INDIAN MALLOW.

Carpels 2-9-seeded, at length 2-valved. Radicle ascending or pointing inwards. Otherwise as in *Sida*. (Name of unknown origin.)

1. **A. AVICÉNNÆ**, Gærtn. (VELVET-LEAF.) Tall annual (4° high); leaves roundish-heart-shaped, taper-pointed, velvety; peduncles shorter than the leaf-stalks; corolla yellow; pods 12-15, hairy, beaked. — Waste places, escaped from gardens. (Adv. from India.)

8. MODIOLA, Moench. MODIOLA.

Calyx with a 3-leaved involucl. Petals obovate. Stamens 10-20. Stigmas capitate. Carpels 14-20, kidney-shaped, pointed, and at length 2-valved at the

top; the cavity divided into two by a cross partition, with a single seed in each cell. — Humble, procumbent or creeping annuals or biennials, with cut leaves and small purplish flowers solitary in the axils. (Name from *modiolus*, the broad and depressed fruit resembling in shape the Roman measure of that name.)

1. **M. multifida**, Mœnch. Hairy; leaves 3–5-cleft and incised; stamens 15–20; fruit hispid at the top. — Low grounds, Virginia and southward.

9. KOSTELÉTZKYA, Presl. KOSTELETKYA.

Pod depressed, with a single seed in each cell. Otherwise as *Hibiscus*. (Named after *V. F. Kosteletzky*, a Bohemian botanist.)

1. **K. Virginica**, Presl. Roughish-hairy perennial (2°–4° high); leaves halberd-shaped and heart-shaped; the lower 3-lobed. (*Hibiscus Virginicus*, L.) — Marshes on the coast, New York to Virginia and southward. Aug. — Corolla 2' wide, rose-color. Column slender.

10. HIBÍSCUS, L. ROSE-MALLOW.

Calyx involuclate at the base by a row of numerous bractlets, 5-cleft. Column of stamens long, bearing anthers for much of its length. Styles united: stigmas 5, capitate. Fruit a 5-celled loculicidal pod. Seeds several or many in each cell. — Herbs or shrubs, usually with large and showy flowers. (An old Greek and Latin name of unknown meaning.)

* *Indigenous, tall perennials (4°–8°), flowering late in summer.*

1. **H. Moscheutos**, L. (SWAMP ROSE-MALLOW.) *Leaves ovate*, pointed, toothed, the lower 3-lobed, whitened underneath with a fine soft down, glabrous or slightly downy beneath; the 1-flowered peduncles sometimes united at the base with the petioles; *calyx not inflated*; *pod and seeds smooth* or nearly so. — Brackish marshes along the coast, sometimes extending up rivers far beyond the influence of salt water (as above Harrisburg, Penn.): also Onondaga Lake, N. Y., and westward, usually within the influence of salt springs. — Corolla 5'–6' in diameter, light rose-color or white, with or without a crimson eye.

2. **H. grandiflorus**, Michx. *Leaves soft-downy both sides, the lower broadly ovate and heart-shaped*; *pod very hirsute*; seeds smooth; — otherwise resembling the last. — Illinois (Lawrence Co., *Fritchey*) and southward.

3. **H. militaris**, Cav. (HALBERD-LEAVED R.) *Smooth throughout*; lower leaves ovate-heart-shaped, toothed, 3-lobed; *upper leaves halberd-form*, the short lateral lobes spreading at the base, the middle one prolonged and taper-pointed; peduncles slender; *fruiting calyx inflated*; *seeds hairy*. — River-banks, Penn. to Illinois and southward. — Corolla 4'–5' in diameter, flesh-color.

* * *Escaped from gardens or grounds.*

4. **H. TRIÏNUM**, L. (BLADDER KETMIA.) A low, rather *hairy annual*; upper leaves 3-parted, with lanceolate divisions, the middle one much the longest; *fruiting calyx inflated, membranaceous, 5-winged*; corolla sulphur-yellow with a blackish eye, ephemeral; hence the name *Flower-of-an-hour*. (Adv. from Eu.)

5. **H. SYRIACUS**, L. (SHRUBBY ALTHÆA of the gardeners.) *Tall shrub*, smooth; leaves wedge-ovate, pointed, cut-toothed or lobed; corolla usually rose-color. — Roadsides and copses, Pennsylvania, &c. Sept. (Adv. from Eu.)

ORDER 21. TILIACEÆ. (LINDEN FAMILY.)

Trees (rarely herbs), with the mucilaginous properties, fibrous bark, valvate calyx, &c., of the Mallow Family; but the sepals deciduous, petals imbricated in the bud, the stamens usually polyadelphous, and the anthers 2-celled;—represented in Northern regions only by the genus,

1. TÍLIA, L. LINDEN. BASSWOOD.

Sepals 5. Petals 5, spatulate-oblong. Stamens numerous: filaments cohering in 5 clusters with each other (in European species), or with the base of a spatulate petal-like body placed opposite each of the real petals. Pistil with a 5-celled ovary and 2 half-anatropous ovules in each, a single style, and a 5-toothed stigma. Fruit a sort of woody globular nut, becoming 1-celled and 1-2-seeded. Embryo with a taper radicle, and a pair of leaf-like somewhat heart-shaped and lobed cotyledons, which are a little folded.—Fine trees, with soft and white wood, more or less heart-shaped and serrate leaves (oblique and often truncate at the base), deciduous stipules, and small cymes of flowers, hanging on an axillary peduncle which is united to a leaf-like bract. Flowers cream-color, honey-bearing, fragrant. (The classical Latin name of the genus.)

1. **T. Americana**, L. (BASSWOOD.) Leaves green and glabrous or nearly so, thickish.—Rich woods, May, June.—This familiar tree is rarely called *Lime-tree*, oftener *White-wood*, commonly *Basswood*; the name (now obsolete in England) alluding to the use of the inner bark for mats and cordage.

Var. **pubescens**. Leaves softly pubescent underneath, often thin. (*T. pubescens*, *Ait.*)—Common from Maryland southward and westward.

2. **T. heterophylla**, Vent. (WHITE BASSWOOD.) Leaves larger, smooth and bright green above, silvery-whitened with a fine down underneath. (*T. alba*, *Michx.*)—Mountains of Penn. to Kentucky and southward.

T. EUROPEÆ, the EUROPEAN LINDEN, which is planted in and near our cities as an ornamental tree, is at once distinguished from any native species by the absence of the petal-like scales among the stamens. This tree (the *Lin*) gave the family name to *Linnaeus*.

ORDER 22. CAMELLIACEÆ.* (CAMELLIA FAMILY.)

Trees or shrubs, with alternate simple feather-veined leaves, and no stipules, the regular flowers hypogynous and polyandrous, the sepals and petals both imbricated in aestivation, the stamens more or less united at the base with each other (monadelphous or 3-5-adelphous) and with the base of the petals.—Anthers 2-celled, introrse. Fruit a woody 3-5-celled loculicidal pod. Seeds few, with little or no albumen. Embryo large, with broad cotyledons.—A family with showy flowers, the types of which are the well-known CAMELLIA and the more important TEA PLANT,—represented in this country by the two following genera.

* Name of same date as TERNSTROMIACEÆ, and preferable.

1. **STUÁRTIA**, Catesby. STUARTIA.

Sepals 5, rarely 6, ovate or lanceolate. Petals 5, rarely 6, obovate, crenulate. Stamens monadelphous below. Pod 5-celled. Seeds 1 or 2 in each cell, crustaceous, anatropous, ascending. Embryo straight, nearly as long as the albumen: radicle longer than the cotyledons. — Shrubs with membranaceous deciduous oblong-ovate serrulate leaves, soft-downy beneath, and large short-peduncled flowers solitary in their axils. (Named for *John Stuart, Lord Bute*.)

1. **S. Virginica**, Cav. Petals 5 white (1' long); sepals ovate; style 1; stigma 5-toothed; pod globular, blunt; seeds not margined. (S. Malachodéndron, L.) — Woods, Virginia and southward.

S. PENTÁGYNA, L'Her., with cream-colored flowers, 5 styles, and an angled and pointed pod, may be found in the Alleghanies of Southern Virginia.

2. **GORDÓNIA**, Ellis. LOBLOLLY BAY.

Sepals 5, rounded, concave. Petals 5, obovate. Stamens 5-adelphous, one cluster adhering to the base of each petal. Style 1. Pod ovoid, 5-valved; the valves separating from the persistent axis; cells 2–8-seeded. Seeds pendulous. Embryo straightish, with a short radicle, and thin longitudinally plaited cotyledons. — Shrubs or small trees, with large and showy white flowers on axillary peduncles. (Dedicated by Dr. Garden to his "old master, *Dr. James Gordon of Aberdeen*," and by Ellis to a London nurseryman of the same name.)

1. **G. Lasiánthus**, L. (LOBLOLLY BAY.) Leaves coriaceous and persistent, lanceolate-oblong, narrowed at the base, minutely serrate, smooth and shining; pod pointed; seeds winged above. Swamps near the coast. Virginia and southward. May–July. — Petals 1½' long.

ORDER 23. **LINÀCÆÆ**. (FLAX FAMILY.)

Herbs (rarely shrubs) with the regular and symmetrical hypogynous flowers 4–6-merous throughout, strongly imbricated calyx and convolute petals, the 5 stamens monadelphous at the base, and an 8–10-seeded pod, having twice as many cells as there are styles. Represented by the genus,

1. **LÍNUM**, L. FLAX.

Sepals (persistent), petals, stamens, and styles 5, regularly alternate with each other. Pod of 5 united carpels (into which it splits in dehiscence) and 5-celled, with 2 seeds hanging from the summit of each; but each cell is partly or completely divided into two by a false partition which projects from the back of the carpel, thus becoming 10-celled. Seeds anatropous, mucilaginous, flattened, containing a large embryo with plano-convex cotyledons. — Herbs, with a tough fibrous bark, simple and sessile entire leaves (alternate or often opposite), without stipules, but often with glands in their place, and with corymbose or panicled flowers. Corolla usually ephemeral. (The classical name of the Flax.)

* *Indigenous species, glabrous, 1°–3° high, with yellow flowers, in summer.*

1. **L. Virginianum**, L. Stem erect from the base and with the corymbose spreading or recurving branches *terete and even*; leaves oblong or lanceolate,

or the lower spatulate and often opposite; flowers scattered, small (barely 3" long); sepals ovate, pointed, smooth-edged or nearly so, equalling the depressed 10-celled pod; styles distinct. — Dry woods: common. — Root apparently annual; but the plant propagated by suckers from the base of the stem.

2. **L. striatum**, Walt. Stems gregarious, erect or ascending from a creeping or decumbent base, slightly viscid, and with the mostly racemose short branches striate with about 4 sharp wing-like angles decurrent from the leaves; these broader than in the last, and mostly oblong, usually with all the lower ones opposite; flowers more crowded; sepals scarcely equalling the very small brownish pod: otherwise nearly as in No. 1. (*L. oppositifolium*, *Engelm.*) — Wet or boggy grounds, New England to Virginia and southward. — Generally confounded with *L. Virginianum* (figured for it in *Reichenb. Ic. Exot.*), but well distinguished by Walter, except that the stem-leaves are commonly opposite up to the first branch: here described from the indications given by *C. F. Austin*.

3. **L. sulcatum**, Riddell, 1836. Stem strictly erect from an annual root, and with the upright or ascending branches striate-angled or grooved; leaves linear, acute, or the upper subulate, rather rigid; a pair of dark glands in place of stipules: sepals ovate-lanceolate and sharp-pointed, strongly 3-nerved and (like the bracts) with rough-bristly-glandular margins, scarcely longer than the ovoid-globose incompletely 10-celled pod; styles united almost to the middle. (*L. Boottii*, *Planchon*, 1848.) — Dry soils, Rhode Island to Illinois and south-westward. — Flowers and pods twice as large as in the preceding. (*L. rigidum*, *Pursh.*, of the Western plains, probably in Minnesota, is dwarf, glaucous, and has the styles united almost to the top.)

* * *Escaping from cultivation, blue-flowered, annual.*

4. **L. USITATISSIMUM**, L. (COMMON FLAX), is occasionally spontaneous in fields.

ORDER 24. GERANIACEÆ. (GERANIUM FAMILY.)

Plants (chiefly herbs) with perfect and generally symmetrical hypogynous flowers; the stamens, counting sterile filaments, as many or commonly twice as many, and the lobes or cells (1-few-ovuled) of the ovary as many, as the sepals, an axis of the dry fruit persisting. — Seeds without albumen, except in *Oxalis*. The flower of *Impatiens* is partly, and that of *Tropæolum* still more unsymmetrical. Herbage often strong-scented, but never punctate with pellucid dots. — As a whole the order, here recombined as it was founded by *Jussieu*, is hard to define. Of late it has generally been broken into several small orders: the principal ones here stand as suborders, with only one or two genera to each. — *TROPÆOLUM*, the GARDEN NASTURTIUM, occupies a position between the first and the second suborder.

SUBORDER I. GERANIEÆ. (GERANIUM FAMILY PROPER.)

Flowers 5-merous and symmetrical; the persistent sepals imbricated and the petals usually convolute in the bud: 5 glands of the receptacle

alternate with the petals. Stamens somewhat monadelphous. Ovary deeply 5-lobed, the 5 two-ovuled carpels and the lower part of the long styles adnate to a long and beak-like prolongation of the receptacle, from which, when ripe and dry, the small and membranaceous pods are torn off, and carried away by the styles,— which, as they separate from the beak from the base upwards, are elastically recurved or revolute, the solitary seed falling out. Embryo filling the seed (no albumen); cotyledons folded together and bent down on the short radicle. — Strong-scented herbs (or the Pelargoniums, which have somewhat irregular flowers, shrubby plants), with opposite or alternate stipulate leaves, and astringent roots.

1. **Geranium.** Stamens with anthers 10, rarely 5. The recurving bases of the styles or tails of the carpels in fruit naked inside.
2. **Erodium.** Stamens with anthers only 5. Tails of the carpels in fruit bearded inside, often spirally twisted.

SUBORDER II. **LIMNANTHEÆ.** (LIMNANTHES FAMILY.)

Flowers 3-merous or 5-merous, completely symmetrical and regular; the persistent sepals ovate in the bud: glands of the receptacle alternate with the petals. Stamens distinct. No beak or prolongation of the receptacle beyond the ovary, which consists of 3 or 5 almost distinct, at length fleshy and separating, indehiscent, 1-seeded carpels, united by a common style. Seeds without albumen: cotyledons very thick and fleshy, the short radicle included by their heart-shaped bases. — Tender low annuals, with pinnate alternate leaves and no stipules. — Consists of the pretty-flowered Californian LIMNANTHES, and of

3. **Flerkea.** Sepals, minute petals, and lobes of the ovary 3: stamens 6.

SUBORDER III. **BALSAMINEÆ.** (BALSAM FAMILY.)

Flowers mostly unsymmetrical, 5-merous as to the stamens and pistil; the sepals and petals irregular, usually unsymmetrical and of fewer pieces, imbricated in the bud, all petaloid and deciduous, the larger piece with an ample sac or spur: no glands: filaments distinct, short. Fruit a fleshy 5-celled pod or berry: no albumen: the straight embryo with thick cotyledons and a short radicle. — Tender herbs, the succulent stems gorged with a bland watery juice; the leaves simple, mostly alternate, without stipules.

4. **Impatiens.** Inner or lateral petals unequally 2-lobed. Pod bursting elastically into 5 valves, several-seeded.

SUBORDER IV. **OXALIDEÆ.** (SORREL FAMILY.)

Flowers 5-merous, regular and symmetrical, decandrous; the persistent sepals imbricated and the petals convolute in the bud: no glands alternate with the latter. Stamens often monadelphous at the base. Fruit a

5-celled loculicidal pod or a berry. Seeds 2 or few in each cell, with a straight embryo in a little fleshy albumen. — Leaves compound: juice sour.

5. **Oxalis.** Styles 5, separate. Pod oblong: the valves not falling away. Leaflets usually obcordate.

1. GERANIUM, L. CRANESBILL.

Stamens 10 (sometimes only 5 in No. 2), all with perfect anthers, the 5 longer with glands at their base (alternate with the petals). Styles smooth inside in fruit when they separate from the axis. — Stems forking. Peduncles 1-3-flowered. (An old Greek name, from *γέρανος*, a crane; the long fruit-bearing beak thought to resemble the bill of that bird.)

* *Rootstock perennial.*

1. **G. maculatum**, L. (WILD CRANESBILL.) Stem erect, hairy; leaves about 5-parted, the wedge-shaped divisions lobed and cut at the end; sepals slender-pointed; petals entire, light purple, bearded on the claw ($\frac{1}{2}$ ' long). — Open woods and fields. April-July. — Leaves somewhat blotched with whitish as they grow old.

** *Root biennial or annual: flowers small.*

2. **G. Carolinianum**, L. (CAROLINA C.) Stems at first erect, diffusely branched from the base, hairy; leaves about 5-parted, the divisions cleft and cut into numerous oblong-linear lobes; peduncles and pedicels short; sepals *awn-pointed*, as long as the emarginate (pale rose-color) petals; carpels hairy; seeds *ovoid-oblong, very minutely reticulated*. — Barren soil and waste places: common. May-Aug. — Depauperate forms, except by the seeds, are hardly distinguishable from

3. **G. dissectum**, L. (CUT-LEAVED C.) More slender and spreading, with narrower lobes to the leaves, and smaller red-purple petals notched at the end; seeds *short-ovoid or globular, finely and strongly reticulated*. — Waste grounds, scarce. (Nat. from Eu.)

4. **G. columbinum**, L. (LONG-STALKED C.) Minutely hairy, with very slender decumbent stems; leaves 5-7-parted and cut into narrow linear lobes; peduncles and pedicels *filiiform and elongated*; sepals awned, about equaling the purple petals, enlarging after flowering; carpels *glabrous*; seeds nearly as in No. 3. — Along the Susquehanna, Lancaster Co., &c., Prof. Porter. Alexandria, Virginia, A. H. Curtiss. June, July. (Nat. from Eu.)

5. **G. pusillum**, L. (SMALL-FLOWERED C.) Stems procumbent, slender, minutely pubescent; leaves *rounded kidney-form*, 5-7-parted, the divisions wedge-shaped, mostly 3-lobed; sepals *awnless*, about as long as the (purplish) petals; seeds *smooth*. — Waste places, New York and Massachusetts: rare. (Nat. from Eu.)

6. **G. Robertianum**, L. (HERB ROBERT.) Sparsely hairy, diffuse; strong-scented, leaves *3-divided or pedately 5-divided, the divisions twice pinnatifid*; sepals awned, shorter than the (red-purple) petals; pods *wrinkled*; seeds *smooth*. — Moist woods and shaded ravines: common northward. June-Oct. (Eu.)

2. ERÒDIUM, L'Her. STORKSBILL.

The 5 shorter stamens sterile or wanting. Styles in fruit twisting spirally, bearded inside. Otherwise as *Geranium*. (Name from *ἔρωδιός*, a heron.)

1. *E. cicutarium*, L'Her. Annual, hairy; stems low, spreading; leaves pinnate; the leaflets sessile, 1-2-pinnatifid; peduncles several-flowered. — New York, Pennsylvania, &c.: scarce. (Adv. from Eu.)

3. FLÛRKEA, Willd. FALSE MERMAID.

Sepals 3. Petals 3, shorter than the calyx, oblong. Stamens 6. Ovaries 3, opposite the sepals, united only at the base; the style rising in the centre: stigmas 3. Fruit of 3 (or 1-2) roughish fleshy achenia. Seed anatropous, erect, filled by the large embryo with its hemispherical fleshy cotyledons. — A small and inconspicuous annual, with minute solitary flowers on axillary peduncles. (Named after *Flörke*, a German botanist.)

1. *F. proserpinacoides*, Willd. — Marshes and river-banks, W. New England to Wisconsin and Kentucky. April-June. — Leaflets 3-5, lanceolate, sometimes 2-3-cleft. Taste slightly pungent.

4. IMPÀTIENS, L. BALSAM. JEWEL-WEED.

Calyx and corolla colored alike and not clearly distinguishable. Sepals apparently only 4; the anterior one, which is notched at the apex, probably consisting of two combined; the posterior one (appearing anterior as the flower hangs on its stalk) largest, and forming a spurred sac. Petals 2, unequal-sided and 2-lobed (each consisting of a pair united). Stamens 5, short: filaments appendaged with a scale on the inner side, the 5 scales connivent and united over the stigma: anthers opening on the inner face. Ovary 5-celled: stigma sessile. Pod with evanescent partitions, and a thick axis bearing the several anatropous seeds, 5-valved, the valves coiling elastically and projecting the seeds in bursting. Embryo straight: albumen none. — Leaves simple, alternate, without stipules, in our species ovate or oval, coarsely toothed, petioled. Flowers axillary or paniced, often of two sorts, viz., — the larger ones, as described above, which seldom ripen seeds; — and very small ones, which are fertilized early in the bud; their floral envelopes never expand, nor grow to their full size, but are forced off by the growing pod and carried upwards on its apex. (Name from the sudden bursting of the pods when touched, whence also the popular appellation, *Touch-me-not*, or *Snap-weed*.)

1. *I. pallida*, Nutt. (PALE TOUCH-ME-NOT.) Flowers pale-yellow, sparingly dotted with brownish-red; sac dilated and very obtuse, broader than long, tipped with a short incurved spur. — Moist shady places and along rills, in rich soil; most common northward. July-Sept. — Larger and greener than the next, with larger flowers: a spotless variety in N. Vermont and New Hampshire, *Miss Lumbard*.

2. *I. fúlva*, Nutt. (SPOTTED TOUCH-ME-NOT.) Flowers orange-color, thickly spotted with reddish-brown; sac longer than broad, acutely conical, tapering into a strongly inflexed spur. — Rills and shady moist places: common, especially

southward. June–Sept.—Plant 2°–4° high: the flowers loosely paniced at the ends of the branches, hanging gracefully on their slender nodding stalks, the open mouth of the cornucopiæ-shaped sepal upward.—A variety is not rare with *spotless flowers*, which differs from the *I. Noli-tangere* of Europe in the more inflexed spur and smaller petals. Spur rarely wanting.

5. ÓXALIS, L. WOOD-SORREL.

Sepals 5, persistent. Petals 5, their bases sometimes united, withering after expansion. Stamens 10, usually monadelphous at the base, alternately shorter. Styles 5, distinct. Pod oblong, membranaceous, 5-celled, more or less 5-lobed, each cell opening on the back; the valves persistent, being fixed to the central column or axis by the partitions. Seeds 2 or more in each cell, pendulous from the axis, anatropous, their outer coat loose and separating. Embryo large and straight in fleshy albumen: cotyledons flat.—Herbs, with sour watery juice, alternate or radical leaves, mostly of 3 obcordate leaflets, which close and droop at nightfall. Several species produce (like *Impatiens*) small peculiar flowers, which are precociously fertilized in the bud and are particularly fruitful; and the ordinary flowers are often dimorphous or even trimorphous in the relative length of the stamens and styles. (Name from *ὄξύς*, *sour*.)

* *Stemless perennials: leaves and scapes a rootstock or bulb: leaflets broadly obcordate: flowers nearly 1' broad; cells of the pod few-seeded.*

1. *O. Acetosélla*, L. (COMMON WOOD-SORREL.) *Rootstock creeping and scaly-toothed; scape 1-flowered (2'–5' high); petals white with reddish veins, often notched.—Deep cold woods, Massachusetts to Penn., L. Superior and northward: also southward in the Alleghanies. June. (Eu.)*

2. *O. violácea*, L. (VIOLET W.) *Nearly smooth; bulb scaly; scapes umbellately several-flowered (5'–9' high), longer than the leaves; petals violet.—Rocky places; most common southward. May, June.*

* * *Stems leafy, branching: peduncles axillary: flowers small: cells several-seeded.*

3. *O. stricta*, L. (YELLOW W.) *Annual or perennial by running subterranean shoots; stems at first erect; stipules absent; peduncles 2–6-flowered, longer than the leaves; petals yellow; pods elongated, erect in fruit.—Copses and cultivated grounds: common. May–Sept.—Varies greatly in appearance and in the size of its flowers.*

O. corniculáta, L., if in this district, and truly distinct from *O. stricta*, should be known by the stipules at the base of the petioles. (Eu.)

ORDER 25. RUTACEÆ. (RUE FAMILY.)

Plants with simple or compound leaves, dotted with pellucid glands, abounding with a pungent or bitter-aromatic acrid volatile oil, producing hypogynous almost always regular 3–5-merous flowers, the stamens as many or twice as many as the sepals (rarely more numerous); the 2–5 pistils separate or combined into a compound ovary of as many cells, raised on a prolongation of the receptacle (gynophore) or glandular disk.—Embryo large, curved or

straight, usually in fleshy albumen. Styles commonly united or cohering, even when the ovaries are distinct. Fruit usually capsular. Leaves alternate or opposite. Stipules none. — A large family, chiefly of the Old World and the Southern hemisphere; the PROPER RUTACEÆ, represented in gardens by the Rue (*Ruta graveolens*, L.) and Fraxinella (*Dictamnus Fraxinella*, L.) chiefly herbs, but the rest are shrubs or trees. — The AURANTIÆ or Orange Family, recently appended to this order, has baccate fruit, seeds without albumen, and stamens sometimes almost indefinitely numerous. — Our two indigenous genera are

1. **Zanthoxylum.** Flowers dioecious: ovaries 3–5, separate, forming fleshy pods.

2. **Ptelea.** Flowers polygamous: ovary 2-celled, forming a samara, like that of Elm.

1. ZANTHÓXYLUM, Colden. PRICKLY ASH.

Flowers dioecious. Sepals 4 or 5, obsolete in one species. Petals 4 or 5, imbricated in the bud. Stamens 4 or 5 in the sterile flowers, alternate with the petals. Pistils 2–5, separate, but their styles conniving or slightly united. Pods thick and fleshy, 2-valved, 1–2-seeded. Seed-coat crustaceous, black, smooth and shining. Embryo straight, with broad cotyledons. — Shrubs or trees, with mostly pinnate leaves, the stems and often the leafstalks prickly. Flowers small, greenish or whitish. (Name from *ξανθός*, *yellow*, and *ξύλον*, *wood*: therefore more properly spelled with an initial X.)

1. **Z. Americanum**, Mill. (NORTHERN PRICKLY ASH. TOOTHACHE-TREE.) Leaves and flowers in axillary clusters; leaflets 4–5 pairs and an odd one, ovate-oblong, downy when young; calyx none; petals 5; pistils 3–5, with slender styles; pods short-stalked. — Rocky woods and river-banks: common northward. April, May. — A prickly shrub, with yellowish-green flowers appearing before the leaves. Bark, leaves, and pods very pungent and aromatic.

2. **Z. Carolinianum**, Lam. (SOUTHERN P.) *Glabrous*; leaflets 3–5 pairs and an odd one, ovate or ovate-lanceolate, oblique, *shining above*; flowers in a terminal cyme, appearing after the leaves; sepals and petals 5; pistils 3, with short styles; pods sessile. — Sandy coast of Virginia, and southward. June. — A small tree with very sharp prickles.

2. PTÉLEA, L. SHRUBBY TREFOIL. HOP-TREE.

Flowers polygamous. Sepals 3–5. Petals 3–5, imbricated in the bud. Stamens as many. Ovary 2-celled: style short: stigmas 2. Fruit a 2-celled and 2-seeded samara, winged all round, nearly orbicular. — Shrubs, with 3-foliate leaves, and greenish-white small flowers in compound terminal cymes. (The Greek name of the Elm, here applied to a genus with similar fruit.)

1. **P. trifoliata**, L. Leaflets ovate, pointed, downy when young. — Rocky places, Penn. to Wisconsin and southward. June. — A tall shrub. Fruit bitter, used as a substitute for hops. Odor of the flowers disagreeable.

AILÁNTHUS GLANDULÓSUS, Desf., called TREE OF HEAVEN, — but whose blossoms, especially the staminate ones, are redolent of anything but “airs from

heaven," — is much planted as a shade-tree, especially in towns, and is inclining to spread from seed in Pennsylvania. It belongs to the order SIMARUBACEÆ, which differs from Rutaceæ in the absence of dots in the leaves. The tree is known by its very long pinnate leaves of many leaflets, and small polygamous greenish flowers in panicles, the female producing 2-5 thin, linear-oblong, veiny samaras. (Adv. from China.)

ORDER 26. ANACARDIACEÆ. (CASHEW FAMILY.)

Trees or shrubs, with a resinous or milky acrid juice, dotless alternate leaves, and small, often polygamous, regular, pentandrous flowers, with a 1-celled and 1-ovuled ovary, but with 3 styles or stigmas. — Petals imbricated in the bud. Seed without albumen, borne on a curved stalk that rises from the base of the cell. Stipules none. Juice or exhalations often poisonous. — Represented here only by the genus

1. RHŪS, L. SUMACH.

Sepals 5. Petals 5. Stamens 5, inserted under the edge or between the lobes of a flattened disk in the bottom of the calyx. Fruit small and indehiscent, a sort of dry drupe. — Leaves (simple in *R. Cōtinus*, the *Smoke-Plant* of gardens) usually compound. Flowers greenish-white or yellowish. (The old Greek and Latin name of the genus.)

§ 1. SŪMAC, DC. *Flowers polygamous, in a terminal thyrsoid panicle: fruit globular, clothed with acid crimson hairs; the stone smooth: leaves odd-pinnate. (Not poisonous.)*

1. **R. typhina**, L. (STAGHORN SUMACH.) *Branches and stalks densely velvety-hairy; leaflets 11-31, pale beneath, oblong-lanceolate, pointed, serrate, rarely laciniate.* — Hillsides. June. — Shrub or tree 10°-30° high, with orange-colored wood.

2. **R. glabra**, L. (SMOOTH S.) *Smooth, somewhat glaucous; leaflets 11-31, whitened beneath, lanceolate-oblong, pointed, serrate.* — Rocky or barren soil. June, July. — Shrub 2°-12° high.

3. **R. copallina**, L. (DWARF S.) *Branches and stalks downy; petioles wing-margined between the 9-21 oblong or ovate-lanceolate (often entire) leaflets, which are oblique or unequal at the base, smooth and shining above.* — Rocky hills. July. — Shrub 1°-7° high, with running roots.

§ 2. TOXICODÉNDRON, Tourn. *Flowers polygamous, in loose and slender axillary panicles: fruit globular, glabrous, whitish or dun-colored; the stone striate: leaves odd-pinnate or 3-foliolate, thin. (Poisonous to the touch.)*

4. **R. venenata**, DC. (POISON S. OR DOGWOOD.) *Smooth, or nearly so; leaflets 7-13, obovate-oblong, entire.* (*R. Vénix*, L.) — Swamps. June. — Shrub 6°-18° high. The most poisonous species: also called *Poison Elder*.

5. **R. Toxicodéndron**, L. (POISON IVY. POISON OAK.) *Climbing by rootlets over rocks, &c., or ascending trees; leaflets 3, rhombic-ovate, mostly pointed, and rather downy beneath, variously notched, sinuate, or cut-lobed, — or else entire, then it is *R. radicans*, L.* — Thickets, low grounds, &c. June.

§ 3. **LOBADIUM**, Raf. *Flowers polygamo-dioecious, in clustered scaly-bracted spikes like catkins, preceding the leaves: disk 5-parted, large: fruit as in § 1, but flattish: leaves 3-foliolate. (Not poisonous.)*

6. **R. aromática**, Ait. (FRAGRANT S.) Leaves pubescent when young, thickish when old; leaflets 3, rhombic-ovate, unequally cut-toothed, the middle one wedge-shaped at the base; flowers pale yellow.—Dry rocky soil, from Vermont westward and southward. April, May.—A straggling bush; the crushed leaves sweet-scented.

ORDER 27. VITACEÆ. (VINE FAMILY.)

Shrubs with watery juice, usually climbing by tendrils, with small regular flowers, a minute or truncated calyx, its limb mostly obsolete, and the stamens as many as the valvate petals and opposite them! Berry 2-celled, usually 4-seeded.—Petals 4–5, very deciduous, hypogynous or perigynous. Filaments slender: anthers introrse. Pistil with a short style or none, and a slightly 2-lobed stigma: ovary 2-celled, with 2 erect anatropous ovules from the base of each cell. Seeds bony, with a minute embryo at the base of the hard albumen, which is grooved on one side.—Stipules deciduous. Leaves palmately veined or compound: tendrils and flower-clusters opposite the leaves. Flowers small, greenish. (Young shoots, foliage, &c., acid.)—Consists of Vitis and one or two nearly allied genera.

1. VITIS, Tourn. GRAPE.

Calyx very short, usually with a nearly entire border or none at all, filled with an adnate fleshy disk which bears the petals and stamens.—Flowers in a compound thyrus; pedicels mostly umbellate-clustered. (The classical Latin name.)

§ 1. **VITIS** proper. *Petals 5, cohering at the top, separating at the base, and so the corolla usually falls off without expanding: 5 thick glands or lobes of the disk alternating with the stamens: flowers polygamous or dioecious in all the American species, exhaling a fragrance like that of Mignonette: leaves simple, rounded and heart-shaped, often variously and variably lobed.*

* *Leaves woolly beneath, when lobed having obtuse or rounded sinuses.*

1. **V. Labrusca**, L. (NORTHERN FOX-GRAPE.) *Branchlets and young leaves very woolly; leaves continuing rusty-woolly beneath; fertile panicles compact; berries large.—Moist thickets: common. June. Fruit ripe in Sept. or Oct., dark purple or amber-color, with a tough musky pulp. Improved by cultivation, it has given rise to the Isabella Grape, &c.*

2. **V. æstivalis**, Michx. (SUMMER GRAPE.) *Young leaves downy with loose cobwebby hairs beneath, smoothish when old, green above; fertile panicles compound, long and slender: berries small, black with a bloom.—Thickets: common. May, June.—Berries pleasant, ripe in Oct.*

* * *Leaves smooth or nearly so and bright green both sides, commonly pubescent on the veins beneath, either incisely lobed or undivided.*

3. **V. cordifolia**, Michx. (WINTER OR FROST GRAPE.) *Leaves thin,*

not shining, heart-shaped, acuminate, sharply and coarsely toothed, often obscurely 3-lobed; *panicles compound, large and loose; berries small, blue or black with a bloom, very acerb, ripening after frosts.* — Var. *RIPARIA*, has the leaves broader and cut-lobed. (*V. riparia, Michx.*) — Thickets and river-banks: common. May, June. — Flowers very sweet-scented.

4. *V. vulpina*, L. (MUSCADINE OR SOUTHERN FOX-GRAPE.) *Leaves shining both sides, small, rounded with a heart-shaped base, very coarsely toothed with broad and bluntish teeth, seldom lobed; panicles small, densely flowered; berries large* ($\frac{1}{2}$ ' - $\frac{3}{4}$ ' in diameter), *musky, purplish without a bloom, with a thick and tough skin, ripe early in autumn.* — River-banks, Maryland to Kentucky and southward. May. — Bark of stem close, not separating in strips as in the other species. Branchlets minutely warty. This is the original of the *Catawba* and the *Scuppernong Grape*, &c.

§ 2. *CISSUS*, L. *Petals* (5 in our species) *expanding before or when they fall: disk thick and broad, usually 4-5-lobed: flowers commonly perfect: tendrils fewer.*

5. *V. indivisa*, Willd. Nearly glabrous; *leaves heart-shaped or truncate at the base, coarsely and sharply toothed, acuminate, not lobed; panicle small and loose; style slender; berries of the size of a pea, 1-3-seeded.* — River-banks, West Virginia, Ohio, and southward. June.

6. *V. bipinnata*, Torr. & Gray. Nearly glabrous, bushy and rather upright; *leaves twice pinnate or ternate, the leaflets cut-toothed; flowers cymose; calyx 5-toothed; disk very thick, adherent to the ovary; berries black, obovate.* — Rich soils, Virginia, Kentucky, and southward.

2. AMPELÓPSIS, Michx. VIRGINIAN CREEPER.

Calyx slightly 5-toothed. Petals concave, thick, expanding before they fall. Disk none. — Leaves digitate, with 5 oblong-lanceolate sparingly serrate leaflets. Flower-clusters cymose. Tendrils fixing themselves to trunks or walls by dilated sucker-like disks at their tips. (Name from *ἄμπελος, a vine, and ὄψις, appearance.*)

1. *A. quinquefolia*, Michx. — A common woody vine, in low or rich grounds, climbing extensively, sometimes by rootlets as well as by its disk-bearing tendrils, blossoming in July, ripening its small blackish berries in October. Also called *American Ivy*, and still less appropriately, *Woodbine*. Leaves turning bright crimson in autumn.

ORDER 28. RHAMNACEÆ. (BUCKTHORN FAMILY.)

Shrubs or small trees, with simple leaves, small and regular flowers (sometimes apetalous), with the 4 or 5 perigynous stamens as many as the valvate sepals and alternate with them, accordingly opposite the petals! Drupe or pod with only one erect seed in each cell, not arilled. — Petals folded inwards in the bud, hooded or concave, inserted along with the stamens into the edge of the fleshy disk which lines the short tube of the calyx and sometimes unites it to the lower part of the 2-5-celled ovary.

Ovules solitary, anatropous. Stigmas 2-5. Embryo large, with broad cotyledons, in sparing fleshy albumen. — Flowers often polygamous, sometimes diœcious. Leaves mostly alternate: stipules small or obsolete. Branches often thorny. (Slightly bitter and astringent: the fruit often mucilaginous, commonly rather nauseous or drastic.)

* Calyx and disk free from the ovary.

1. **Berchemia**. Petals sessile, entire, as long as the calyx. Drupe with thin flesh and a 2-celled bony putamen.
2. **Rhamnus**. Petals small, short-clawed, notched, or none. Drupe berry-like, with the 2-4 separate seed-like nutlets concave on the back: cotyledons leaf-like, revolute.
3. **Frangula**. Petals, &c. as in No. 2. Seed-like nutlets convex on the back: cotyledons flat, fleshy.

** Calyx with the disk adherent to the base of the ovary.

4. **Ceanothus**. Petals long-clawed, hooded. Fruit dry, at length dehiscent.

1. BERCHÈMIA, Necker. SUPPLE-JACK.

Calyx with a very short and roundish tube; its lobes equalling the 5 oblong sessile acute petals, longer than the stamens. Disk very thick and flat, filling the calyx-tube and covering the ovary. Drupe oblong, with thin flesh and a bony 2-celled putamen. — Woody high-climbing twiners, with the pinnate veins of the leaves straight and parallel, the small greenish-white flowers in small panicles. (Name unexplained, probably personal.)

1. **B. volùbilis**, DC. Glabrous; leaves oblong-ovate, acute, scarcely serrulate; style short. — Damp soils, Virginia, and southward. June. — Ascending tall trees. Stems tough and very lithe, whence the popular name.

2. RHÁMNUS, Tourn. BUCKTHORN.

Calyx 4-5-cleft; the tube campanulate, lined with the disk. Petals small, short-clawed, notched at the end, wrapped around the short stamens, or sometimes none. Ovary free, 2-4-celled. Drupe berry-like (black), containing 2-4 separate seed-like nutlets, of cartilaginous texture, which are grooved on the back, as is the contained seed. Cotyledons foliaceous, the margins revolute. — Shrubs or small trees, with loosely pinnately veined leaves, and greenish polygamous or diœcious flowers, in axillary clusters. (The ancient Greek name, from the numerous branchlets.)

* Lobes of the calyx, petals, and stamens 4.

1. **R. cathárticus**, L. (COMMON BUCKTHORN.) *Leaves ovate*, minutely serrate; *fruit 3-4-seeded*; branchlets thorny. — Cultivated for hedges; sparingly naturalized eastward. May, June. (Nat. from Eu.)

2. **R. lanceolátus**, Pursh. *Leaves oblong-lanceolate* and acute, or on flowering shoots oblong and obtuse, finely serrulate, smooth or minutely downy beneath; petals deeply notched; *fruit 2-seeded*. Hills and river-banks, Penn. (Mercersburg, *Prof. Green*) to Illinois, and westward. May. — Shrub tall, not thorny; the yellowish-green flowers of two forms on distinct plants, both perfect: one with the short pedicels clustered in the axils and with a short included style; the other with the pedicels oftener solitary, the style longer and exerted.

* * *Lobes of the calyx and stamens 5 : petals wanting.*

3. **R. alnifolius**, L'Her. A low shrub, leaves oval, acute, serrate, nearly straight-veined : fruit 3-seeded. — Swamps, Maine to Penn., Illinois, and northward. June.

3. **FRÁNGULA**, Tourn. ALDER-BUCKTHORN.

Seeds not grooved or concave (but convex) on the back. Cotyledons plane, thick. Flowers perfect; the lobes of the calyx, petals, and stamens almost always 5. Leaves with nearly straight parallel veins. Otherwise as in Rhamnus. (Name from *frango*, to break, in allusion to the brittleness of the stems.)

1. **F. Caroliniána**, Gray. Thornless shrub or small tree; leaves (3'–5' long) oblong, obscurely serrulate, nearly glabrous, deciduous; flowers in one form umbelled, in another solitary in the axils, short-peduncled; drupe globose, 3-seeded. — Secaucus swamp, New Jersey, *Dr. Post, W. H. Leggett*, and riverbanks. Virginia to Kentucky and southward. June.

4. **CEANÒTHUS**, L. NEW JERSEY TEA. RED-ROOT.

Calyx 5-lobed; the lobes colored and incurved; the lower part with the thick disk cohering with the ovary, the upper separating across in fruit. Petals hooded, spreading, on slender claws longer than the calyx. Filaments also elongated. Fruit 3-lobed, dry and splitting into its 3 carpels when ripe. Seed as in Frangula. — Shrubby plants; the flowers in little umbel-like clusters, which are crowded in dense panicles or corymbs at the summit of naked flower-branches: calyx and pedicels colored like the petals. (An obscure name in Theophrastus, probably misspelled.)

1. **C. Americanus**, L. (NEW JERSEY TEA.) Leaves ovate or oblong-ovate, 3-ribbed, serrate, downy beneath, often heart-shaped at the base: common peduncles elongated. — Dry woodlands. July. — Stems 1°–3° high from a dark red root: branches downy. Flowers in pretty white clusters. — The leaves were used for tea during the American Revolution; and the manufacture has been recently revived in Pennsylvania.

2. **C. ovalis**, Bigelow. Leaves narrowly oval or elliptical-lanceolate, finely glandular-serrate, glabrous or nearly so, as well as the short common peduncles. — Dry rocks, W. Vermont to Wisconsin, and westward. May. — The white flowers larger than in No. 1, more corymbed: leaves narrower, smooth, mostly acute at both ends.

ORDER 29. **CELASTRACEÆ**. (STAFF-TREE FAMILY.)

Shrubs with simple leaves, and small regular flowers, the sepals and the petals both imbricated in the bud, the 4 or 5 perigynous stamens as many as the petals and alternate with them, inserted on a disk which fills the bottom of the calyx and sometimes covers the ovary. Seeds arilled. — Ovules one or few (erect or pendulous) in each cell, anatropous: styles united into one. Fruit 2–5-celled, free from the calyx. Embryo large, in fleshy albumen: cotyledons broad and thin. Stipules minute and fugacious. Pedicels jointed. — Represented by two genera.

1. **CELÁSTRUS**, L. STAFF-TREE. SHRUBBY BITTER-SWEET.

Flowers polygamo-dicecious. Petals (crenulate) and stamens 5, inserted on the margin of a cup-shaped disk which lines the base of the calyx. Pod globose (orange-color and berry-like), 3-celled, 3-valved, loculicidal. Seeds 1 or 2 in each cell, erect, enclosed in a pulpy scarlet aril. — Leaves alternate. Flowers small, greenish, in raceme-like clusters terminating the branches. (An ancient Greek name for some evergreen, which our plant is not.)

1. **C. scándens**, L. (WAX-WORK. CLIMBING BITTER-SWEET.) Twinning shrub; leaves ovate-oblong, finely serrate, pointed. — Along streams and thickets. June. — The opening orange-colored pods, displaying the scarlet covering of the seeds, are very ornamental in autumn.

2. **EUÒNYMUS**, TOURN. SPINDLE-TREE.

Flowers perfect. Sepals 4 or 5, united at the base, forming a short and flat calyx. Petals 4–5, rounded, spreading. Stamens very short, inserted on the edge or face of a broad and flat 4–5-angled disk, which coheres with the calyx and is stretched over the ovary, adhering to it more or less. Style short or none. Pod 3–5-lobed, 3–5-valved, loculicidal. Seeds 1–4 in each cell, enclosed in a red aril. — Shrubs, with 4-sided branchlets, opposite serrate leaves, and loose cymes of small flowers on axillary peduncles. (Derivation from εὖ, *good*, and ὄνομα, *name*, because it has the bad reputation of poisoning cattle. *Tourn.*)

1. **E. atropurpùreus**, Jacq. (BURNING-BUSH. WAAHOO.) Shrub tall (6°–14° high) and upright; leaves *petioled*, oval-oblong, pointed; parts of the (dark-purple) flower commonly in fours; pods *smooth, deeply lobed*. — New York to Wisconsin and southward: also cultivated. June. — Ornamental in autumn, by its copious crimson fruit, drooping on long peduncles.

2. **E. Americanus**, L. (STRAWBERRY BUSH.) Shrub low, upright or straggling (2°–5° high); leaves *almost sessile, thickish*, bright green, varying from ovate to oblong-lanceolate, acute or pointed; parts of the greenish-purple flowers mostly in fives; pods *rough-warty, depressed*, crimson when ripe; the aril scarlet. — Wooded river-banks, Western New York to Illinois and southward. June.

Var. **obovátus**, Torr. & Gray. Trailing, with rooting branches; flowering stems 1°–2° high; leaves thin and dull, obovate or oblong. (*E. obovatus*, *Nutt.*) — Low or wet places: the commoner form.

ORDER 30. **SAPINDÀCEÆ**. (SOAPBERRY FAMILY.)

Trees, shrubs, with simple or compound leaves, mostly unsymmetrical and often irregular flowers; the 4–5 sepals and petals both imbricated in æstivation; the 5–10 stamens inserted on a fleshy (perigynous or hypogynous) disk; a 2–3-celled and lobed ovary, with 1–2 (or rarely more) ovules in each cell; and the embryo (except Staphylea) curved or convolute, without albumen. — A large and diverse order, the true Sapindaceæ principally tropical, here represented only by the Buckeyes.

SUBORDER I. **STAPHYLEACEÆ.** BLADDER-NUT FAMILY.

Flowers (perfect) regular : stamens as many as the petals. Ovules 1-8 in each cell. Seeds bony, with a straight embryo in scanty albumen. — Shrubs with opposite pinnately compound leaves, both stipulate and stipellate.

1. **Staphylea.** Lobes of the colored calyx and petals 5, erect. Stamens 5. Fruit a 3-celled bladderly-inflated pod.

SUBORDER II. **SAPINDACEÆ** proper (including HIPPOCASTANEÆ.)

Flowers (often polygamous) mostly unsymmetrical and irregular; the stamens commonly more numerous than the petals or sepals, but rarely twice as many. Ovules 1 or 2 in each cell. Albumen none. Embryo curved or convolute, rarely straight : cotyledons thick and fleshy. — Leaves alternate or sometimes opposite, destitute of stipules, mostly compound.

2. **Æsculus.** Calyx 5-lobed. Petals 4 or 5. Stamens commonly 7. Fruit a leathery pod. Leaves opposite, digitate.

SUBORDER III. **ACERINEÆ.** MAPLE FAMILY.

Flowers (polygamous or diœcious) small, regular, but usually unsymmetrical. Petals often wanting. Ovary 2-lobed and 2-celled, with a pair of ovules in each cell. Winged fruits 1-seeded. Albumen none. Embryo coiled or folded; the cotyledons long and thin. — Leaves opposite, simple or compound.

3. **Acer.** Flowers polygamous. Leaves simple, or rarely digitately compound.

4. **Negundo.** Flowers diœcious. Leaves pinnate, with 3-5 leaflets.

1. **STAPHYLËA, L.** BLADDER-NUT.

Calyx deeply 5-parted, the lobes erect, whitish. Petals 5, erect, spatulate, inserted on the margin of the thick perigynous disk which lines the base of the calyx. Stamens 5, alternate with the petals. Pistil of 3 several-ovuled carpels, united in the axis, their long styles lightly cohering. Pod large, membranaceous, inflated, 3-lobed, 3-celled, at length bursting at the summit; the cells containing 1-4 bony anatropous seeds. Aril none. Embryo large and straight, in scanty albumen; cotyledons broad and thin. — Upright shrubs, with opposite pinnate leaves of 3 or 5 serrate leaflets, and white flowers in drooping raceme-like clusters, terminating the branchlets. Stipules and stipels deciduous. (Name from *σταφύλη*, a cluster.)

1. **S. trifolia, L.** (AMERICAN BLADDER-NUT.) Leaflets 3, ovate, pointed. — Thickets, in moist soil. May. — Shrub 10° high, with greenish striped branches.

2. **ÆSCULUS, L.** HORSE-CHESTNUT. BUCKEYE.

Calyx tubular, 5-lobed, often rather oblique or gibbous at the base. Petals 4, sometimes 5, more or less unequal, with claws, nearly hypogynous. Stamens

7 (rarely 6 or 8) : filaments long and slender, often unequal. Style 1 : ovary 3-celled, with 2 ovules in each, only one of which, or one in each cell, forms a seed. Seed very large, with a thick and shining coat, and a large and round pale scar, without albumen. Cotyledons very thick and fleshy, their contiguous faces more or less united, remaining under ground in germination : plumule 2-leaved : radicle curved. — Trees or shrubs. Leaves opposite, digitate : leaflets serrate, straight-veined, like a Chestnut-leaf. Flowers in a terminal thyrsus or dense panicle, often polygamous, the greater portion with imperfect pistils and sterile. Pedicels jointed. Seeds farinaceous, but imbued with a bitter and narcotic principle. (The ancient name of some Oak or other mast-bearing tree.)

§ 1. *ÆSCULUS* proper. *Fruit covered with prickles when young.*

1. *Æ. HIPPOCÀSTANUM*, L. (COMMON HORSE-CHESTNUT.) Corolla spreading, white spotted with purple and yellow, of 5 petals ; stamens declined ; leaflets 7. — Commonly planted. (Adv. from Asia via Eu.)

2. *Æ. glàbra*, Willd. (FETID or OHIO BUCKEYE.) Stamens curved, longer than the pale yellow corolla of 4 upright petals ; leaflets 5. — River-banks, W. Penn. and Virginia to Michigan and Kentucky. June. — A large tree ; the bark exhaling an unpleasant odor, as in the rest of the genus. Flowers small, not showy.

§ 2. *PÀVIA*, Boerhaave. *Fruit smooth : petals 4, conniving ; the 2 upper smaller and longer than the others, with a small and rounded blade on a very long claw.*

3. *Æ. flàva*, Ait. (SWEET BUCKEYE.) *Stamens included in the yellow corolla ; calyx oblong-campanulate ; leaflets 5, sometimes 7, glabrous, or often minutely downy underneath.* — Rich woods, Virginia to Ohio, Indiana, and southward. May. A large tree or a shrub.

Var. *purpuràscens*. Flowers (both calyx and corolla) tinged with flesh-color or dull purple ; leaflets commonly downy beneath. (*Æ. discolor*, Pursh.) — From West Virginia southward and westward.

4. *Æ. Pàvia*, L. (RED BUCKEYE.) Stamens not longer than the corolla, which is bright red, as well as the *tubular calyx* ; leaflets glabrous or soft-downy beneath. — Fertile valleys, Virginia, Kentucky, and southward. May. — A shrub or small tree.

3. *ÀCER*, Tourn. MAPLE.

Flowers polygamo-diœcious. Calyx colored, 5- (rarely 4-12-) lobed or parted. Petals either none, or as many as the lobes of the calyx, equal, with short claws if any, inserted on the margin of the lobed disk, which is either perigynous or hypogynous. Stamens 4-12. Ovary 2-celled, with » pair of ovules in each : styles 2, long and slender, united only below, stigmatic down the inside. From the back of each ovary grows a wing, converting the fruit into two 1-seeded, at length separable samaras or keys. Seed without albumen. Embryo variously coiled or folded, with large and thin cotyledons. — Trees, or sometimes shrubs, with opposite palmately-lobed leaves, and small flowers. Pedicels not jointed. (The classical name, from the Celtic *ac*, hard.)

* *Flowers in terminal racemes, greenish, appearing after the leaves: stamens 6-8.*

1. **A. Pennsylvanicum**, L. (STRIPED MAPLE.) Leaves 3-lobed at the apex, finely and sharply doubly serrate; the short lobes taper-pointed and also serrate; *racemes drooping, loose; petals obovate*; fruit with large diverging wings. (*A. striatum*, Lam.)—Rich woods, Maine to Wisconsin, and northward along the Alleghanies to Virginia and Kentucky. June.—A small and slender tree, with light-green bark striped with dark lines, and greenish flowers and fruit. Also called *Striped Dogwood* and *Moose-Wood*.

2. **A. spicatum**, Lam. (MOUNTAIN M.) Leaves downy underneath, 3- (or slightly 5-) lobed, coarsely serrate, the lobes taper-pointed; *racemes upright, dense, somewhat compound; petals linear-spatulate*; fruit with small very divergent wings. (*A. montanum*, Ait.)—Moist woods, with the same range as No. 1. June.—A tall shrub, forming clumps.

** *Flowers umbellate-corymbed, greenish-yellow, appearing with the leaves.*

3. **A. saccharinum**, Wang. (SUGAR OR ROCK M.) Leaves 3-5-lobed, with rounded sinuses and pointed sparingly sinuate-toothed lobes, either heart-shaped or nearly truncate at the base, whitish and smooth or a little downy on the veins beneath; flowers from terminal leaf-bearing and lateral leafless buds, drooping on very slender hairy pedicels; calyx hairy at the apex; petals none; wings of the fruit broad, slightly diverging.—Rich woods, especially northward and along the mountains southward. April, May.—A large and handsome tree.

Var. **nigrum**. (BLACK SUGAR-M.) Leaves scarcely paler beneath, but often minutely downy, the lobes wider, the sinus at the base often closed. (*A. nigrum*, Michx.)—With the ordinary form.

*** *Flowers in umbel-like clusters arising from separate lateral buds, and much preceding the leaves: stamens 3-6.*

4. **A. dasycarpum**, Ehrhart. (WHITE OR SILVER M.) Leaves very deeply 5-lobed with the sinuses rather acute, silvery-white (and when young downy) underneath, the divisions narrow, cut-lobed and toothed; flowers (greenish-yellow) on short pedicels; *petals none; fruit woolly when young*, with large divergent wings.—River-banks; most common southward and westward. March-April.—A fine ornamental tree.

5. **A. rubrum**, L. (RED OR SWAMP M.) Leaves 3-5 lobed, with acute sinuses, whitish underneath; the lobes irregularly serrate and notched, acute, the middle one usually longest; *petals linear-oblong*; flowers (scarlet, crimson, or sometimes yellowish) on very short pedicels; but the *smooth fruit* on prolonged drooping pedicels.—Swamps and wet woods. April.—A small tree, with reddish twigs; the leaves varying greatly in shape, turning bright crimson in early autumn.

4. NEGÚNDO, Mœnch. ASH-LEAVED MAPLE. BOX-ELDER.

Flowers diœcious. Calyx minute, 4-5-cleft. Petals none. Stamens 4-5. Disk none.—Sterile flowers in clusters on capillary pedicels, the fertile in drooping racemes, from lateral buds. Leaves pinnate, with 3 or 5 leaflets. (Name unmeaning.)

1. **N. aceroides**, Mœnch. (*Acer Negundo*, L.) Leaflets smoothish when old, very veiny, ovate, pointed, toothed; fruit smooth, with large rather incurved wings.—River-banks. Pennsylvania to Wisconsin, and southward. April.—A small but handsome tree, with light-green twigs, and very delicate drooping clusters of small greenish flowers, rather earlier than the leaves.

ORDER 31. POLYGALACEÆ. (MILKWORT FAMILY.)

Plants with irregular hypogynous flowers, 4–8 diadelphous or monadelphous stamens, their 1-celled anthers opening at the top by a pore or chink; the fruit a 2-celled and 2-seeded pod.—Represented in this country only by the genus

1. POLÝGALA, Tourn. MILKWORT.

Flower very irregular. Calyx persistent, of 5 sepals, of which 3 (the upper and the 2 lower) are small and often greenish, while the two lateral or inner (called *wings*) are much larger, and colored like the petals. Petals 3, hypogynous, connected with each other and with the stamen-tube, the middle (lower) one keel-shaped and often crested on the back. Stamens 6 or 8: their filaments united below into a split sheath, or into 2 sets, cohering more or less with the petals, free above: anthers 1-celled, often cup-shaped, opening by a hole or broad chink at the apex. Ovary 2-celled, with a single anatropous ovule pendulous in each cell: style prolonged and curved: stigma various. Fruit a small, loculicidal 2-seeded pod, usually rounded and notched at the apex, much flattened contrary to the very narrow partition. Seeds with a caruncle, or variously shaped appendage, at the hilum. Embryo large, straight, with flat and broad cotyledons, surrounded by a sparing albumen.—Bitter plants (low herbs in temperate regions), with simple entire often dotted leaves, and no stipules: sometimes (as in the last two species) bearing concealed flowers next the ground, which are fertilized in the closed bud. (An old name composed of *πολύς*, *much*, and *γάλα*, *milk*, from a fancied property of its increasing this secretion.)

§ 1. *Biennials or annuals, with alternate leaves, and yellow flowers, which are disposed to turn greenish in drying: crest of the keel (lower petal) small: flowering all summer.*

1. **P. lutea**, L. Low; flowers (bright orange-yellow) in solitary ovate or oblong heads ($\frac{3}{4}$ ' thick,) terminating the stem or simple branches; leaves (1'–2' long) obovate or spatulate; lobes of the *caruncle* nearly as long as the seed.—Sandy swamps, New Jersey and southward, near the coast.

2. **P. ramosa**, Ell. Flowers (citron-yellow) in numerous short and dense spike-like racemes collected in a flat-topped compound cyme; leaves oblong-linear, the lowest spatulate or obovate; seeds ovoid; minutely hairy, twice the length of the *caruncle*. (*P. cymosa*, Poir., not of Walt. *P. corymbosa*, Nutt.)—Damp pine-barrens, Delaware and southward. (The allied *P. cymosa*, Walt., which is *P. graminifolia*, Poir., *P. attenuata*, Nutt. and *P. acutifolia*, Torr. & Gray,—known by its simpler cymes, stem naked above, narrower leaves, and globular seeds with no caruncle,—may occur in S. Virginia.)

§ 2. *Annuals, with all the leaves alternate: flowers in spikes, heads, or racemes terminating the stem or branches, purple or rose-color, in summer: none subterranean.*

* *Corolla conspicuously crested on the keel: the claws of the true petals united into a long and slender cleft tube much surpassing the wings.*

3. **P. incarnata**, L. Glauous; stem slender, sparingly branched; leaves minute and linear-awl-shaped; spike cylindrical; flowers flesh-color; caruncle longer than the narrow stalk of the hairy seed. — Dry soil, Penn. to Wisconsin and southward; rather rare.

* * *Corolla minutely or inconspicuously crested; the true petals not longer but mostly shorter than the wings: seed pear-shaped.*

4. **P. sanguinea**, L. Stem sparingly branched above, leafy to the top; leaves oblong-linear; heads globular, at length oblong, very dense (4"–5" thick), bright red-purple (rarely paler or even white); pedicels scarcely any; wings broadly ovate, closely sessile, longer than the pod; the 2-parted caruncle almost equalling the seed. — Sandy and moist ground: common.

5. **P. fastigiata**, Nutt. Stem slender, at length corymbosely branched; leaves narrowly linear, small; spikes short and dense (3" in diameter); the small rose-purple flowers on pedicels of about the length of the pod; wings obovate or oval-oblong, narrowed at the base, scarcely exceeding the pod; bracts deciduous with the flowers or fruits; caruncle as long as and nearly enveloping the stalk-like base of the minutely hairy seed. (*P. sanguinea*, Torr. & Gr., excl. syn.; not of Nutt., nor L.) — Pine barrens of New Jersey (Nuttall) and Delaware to Kentucky and southward.

6. **P. Nuttallii**, Torr. & Gr. (*Fl.* 1, p. 670, excl. syn. & descr.) Resembles the last, but usually lower; spikes more cylindrical; flowers duller or greenish purple, on shorter pedicels; the awl-shaped scaly bracts persistent on the axis after the flowers or fruits fall; seed very hairy, the caruncle smaller. (*P. sanguinea*, Nutt., not of L. *P. Mariana*, &c., *Pluk. t.* 437. *P. ambigua*, Torr. & Gr. *Fl.*, not of Nutt.) — Dry sandy soil, coast of Massachusetts to Kentucky and southward. — Spike sometimes rather loose.

7. **P. Curtissii**, n. sp. Slender (9' high); leaves, &c., as in the two preceding; flowers rose-purple, in elongated and loose racemes; the ascending pedicels and the narrow oblong erect wings fully twice the length of the pod; bracts persistent, those of the lower and remoter flowers foliaceous; caruncle small, on one side of the stalk-like base of the very hairy seed, which is conspicuously apiculate at the broader end. — Near Alexandria, Virginia, *A. H. Curtiss*. — Most related to *P. Chapmanii* of Florida.

§ 3. *Annuals with at least the lower stem-leaves whorled in fours, sometimes in fives: spikes terminating the stem and branches; fl. summer and autumn.*

* *Spikes short and thick (4"–9" in diameter; the axis rough with the squarrose bracts persisting after the fall of the (middle-sized) rose or greenish purple flowers: crest of the keel small.*

8. **P. cruciata**, L. Stems (3'–10' high) almost winged at the angles, with spreading opposite branches; leaves nearly all in fours, linear and somewhat spatulate or oblanceolate; spikes sessile or nearly so, wings broadly deltoid-ovate, slightly heart-shaped, tapering to a bristly point, or rarely pointless; caruncle

nearly as long as the seed. — Margin of swamps, Maine to Virginia and southward near the coast, and along the Great Lakes.

9. **P. brevifolia**, Nutt. Rather slender, branched above; leaves scattered on the branches, narrower; *spikes peduncled; wings lanceolate-ovate, pointless or barely mucronate.* — Margin of sandy bogs, Rhode Island (*Olney*), New Jersey, and southward. Too near the last.

* * *Spikes slender (about 2" thick), the bracts falling with the flowers, which are small, greenish-white or barely tinged with purple, the crest of the keel larger.*

10. **P. verticillata**, L. Slender (6'–10' high), much branched; *stem-leaves all whorled, those of the branches scattered, linear, acute; spikes peduncled, dense, acute; wings round, clawed; the 2-lobed caruncle half the length of the seed.* — Dry soil: common.

11. **P. ambigua**, Nutt. Very slender, loosely branched; *lowest stem-leaves in fours, the rest scattered; spikes long-peduncled, more slender, the flowers often purplish and scattered; wings oval; caruncle shorter; otherwise nearly as in No. 10,* — of which it is probably a mere variety. — Dry soil, from New York and Pennsylvania southward.

§ 4. *Perennial, with alternate leaves throughout, and white flowers in a solitary close spike: no subterranean blossoms.*

12. **P. Sénega**, L. (SENECA SNAKEROOT.) Stems several from thick and hard knotty rootstocks, simple (6'–12' high); leaves lanceolate or oblong-lanceolate, with rough margins; flowers almost sessile; wings round-obovate, concave; crest short; caruncle nearly as long as the seed. — Rocky soil, W. New England to Wisconsin and southward. May, June.

Var. **latifolia**, Torr. & Gray. Taller, sometimes branched; leaves ovate or ovate-lanceolate, 2'–4' long, tapering to each end. — Kentucky, *Short*.

§ 5. *Biennials and perennials, with showy, rose-purple, conspicuously crested flowers; also bearing colorless and inconspicuous more fertile ones, with imperfect corollas, fertilized in the bud, on subterranean branches.*

13. **P. polygama**, Walt. Stems numerous from the biennial root, mostly simple, ascending, very leafy (6' to 9' high); *leaves oblanceolate or oblong, alternate; terminal raceme many-flowered, the broadly obovate wings longer than the keel; stamens 8; radical flowers racemed on short runners beneath the ground; lobes of the caruncle 2, scale-like, shorter than the seed.* — Dry sandy soil; common eastward. July.

14. **P. paucifolia**, Willd. *Perennial; flowering stems short (3'–4' high), rising from long and slender prostrate or subterranean shoots, which also bear concealed fertile flowers; lower leaves small and scale-like, scattered: the upper leaves ovate, petioled, crowded at the summit of the stem; flowers 1–3, large, peduncled; wings obovate, rather shorter than the conspicuously fringe-crested keel; stamens 6; caruncle of 2 or 3 awl-shaped lobes longer than the seed.* — Woods, in light soil, especially northward, extending southward along the Alleghanies. May. — A delicate plant, with large and very handsome flowers, 9" long, rose-purple, or rarely pure white. Sometimes called *Flowering Wintergreen*, but more appropriately FRINGED POLYGALA.

ORDER 32. LEGUMINOSÆ. (PULSE FAMILY.)

Plants with papilionaceous or sometimes regular flowers, 10 (rarely 5 and sometimes many) monadelphous, diadelphous, or rarely distinct stamens, and a single simple free pistil, becoming a legume in fruit. Seeds mostly without albumen. Leaves alternate, with stipules, usually compound. One of the sepals inferior (i. e. next the bract); one of the petals superior (i. e. next the axis of the inflorescence).—A very large order (nearly free from noxious qualities), of which the principal representatives in northern temperate regions belong to the first of the three suborders it comprises.

SUBORDER I. PAPILIONACEÆ. PROPER PULSE FAMILY.

Calyx of 5 sepals, more or less united, often unequally so. Corolla perigynous (inserted into the base of the calyx), of 5 irregular petals (or very rarely fewer), imbricated in the bud, more or less distinctly *papilionaceous*, i. e. with the upper or odd petal, called the *vexillum* or *standard*, larger than the others and enclosing them in the bud, usually turned backward or spreading; the two lateral ones, called the *wings*, oblique and exterior to the two lower petals, which last are connivent and commonly more or less coherent by their anterior edges, forming a body named the *carina* or *keel*, from its resemblance to the keel or prow of a boat, and which usually encloses the stamens and pistil. Stamens 10, very rarely 5, inserted with the corolla, monadelphous, diadelphous (mostly with 9 united in one set in a tube which is cleft on the upper side, i. e. next the standard, and the tenth or upper one separate), or occasionally distinct. Ovary 1-celled, sometimes 2-celled by an intrusion of one of the sutures, or transversely 2-many-celled by cross-division into joints: style simple: ovules amphitropous, rarely anatropous. Cotyledons large, thick or thickish: radicle incurved.—Leaves simple or simply compound, the earliest ones in germination usually opposite, the rest alternate: leaflets almost always quite entire. Flowers perfect, solitary and axillary, or in spikes, racemes, or panicles.

A. Stamens monadelphous or diadelphous.

Tribe I. GENISTEÆ. Shrubs or herbs, never climbing, with simple or palmately compound leaves, and peduncles terminal or opposite the leaves. Stamens monadelphous: anthers of two forms. Pod continuous.

1. **Lupinus.** Calyx deeply 2-lipped. Keel scythe-shaped, pointed. Pod flat.
2. **Crotalaria.** Calyx 5-lobed. Keel scythe-shaped, pointed. Pod inflated. Leaves simple.
3. **Gentia.** Keel straight, deflexed. Pod usually flat. Leaves simple.

Tribe II. TRIFOLIEÆ. Herbs, not climbing, with 3 (rarely 5-7-) foliolate leaves, the veinlets of the leaflets often running into minute teeth, and the stipules united with the base of the petiole. Peduncles axillary. Stamens diadelphous: anthers uniform. Pod small and 1-few-seeded, or coiled.

4. **Trifolium.** Flowers capitate. Pods membranaceous, 1-6-seeded. Petals adherent to the stamen-tube.
5. **Melilotus.** Flowers racemed. Pods coriaceous, wrinkled, 1-2-seeded.
6. **Medicago.** Flowers racemed or spiked. Pods curved or coiled, 1-few-seeded.

Tribe III. GALEGEÆ. Herbs or woody plants (twining only in No. 12), not tendril-bearing, with pinnate or sometimes palmately compound leaves. Stamens diadelphous or rarely monadelphous: anthers uniform (or the 5 alternate ones sometimes smaller in No. 7). Pod continuous, at least not transversely jointed.

- * **PSORALIEÆ.** Glandular-dotted shrubs or herbs. Ovary 1-3 ovuled; the legume small and indehiscent, 1-seeded, rarely 2-seeded.
- 7. **Psoralea.** Corolla truly papilionaceous. Stamens 10, more or less diadelphous, half of the anthers often smaller or less perfect. Leaves 3-5-foliolate.
- 8. **Dalea.** Corolla imperfectly papilionaceous. Stamens 9 or 10, monadelphous; the cleft tube of filaments bearing 4 of the petals about its middle. Leaves pinnate.
- 9. **Petalostemon.** Corolla scarcely at all papilionaceous. Stamens 5, monadelphous; the cleft tube of filaments bearing 4 of the petals on its summit. Leaves pinnate.
- 10. **Amorpha.** Corolla of only one petal! Stamens 10, monadelphous at the base. Leaves pinnate.
- ** **TEPHROSIEÆ.** Trees, shrubs, or herbs. Ovary several-many-ovuled: pod flattened, several-seeded, 2-valved. Leaves pinnate. Standard large and broad.
- 11. **Robinia.** Wings of the corolla free from the keel. Pod flat, thin, margined on one edge. Trees or shrubs: leaflets stipellate.
- 12. **Wistaria.** Wings free from the keel. Pod tumid, marginless. Woody twiners: leaflets obscurely stipellate.
- 13. **Tephrosia.** Wings cohering with the keel. Pod flat, marginless. Herbs: leaflets not stipellate.
- *** **ASTRAGALEÆ.** Chiefly herbs, with pinnate leaves and axillary racemes or spikes. Pod mostly turgid or inflated. Style beardless. Standard narrow, erect.
- 14. **Astragalus.** Anthers distinctly 2-celled. Keel not tipped with a point or sharp appendage. Pod turgid, with one or both the sutures turned in, sometimes dividing the cell lengthwise into two.
- 15. **Oxytropis.** Keel of the corolla tipped with a point: otherwise as Astragalus.
- 16. **Glycyrrhiza.** Anthers with the two cells confluent. Pod short, few-seeded, nearly indehiscent, glandular-prickly.

Tribe IV. HEDYSAREÆ. Herbs, with pinnate or pinnately 1-3-foliolate leaves, not tendril-bearing. Anthers uniform, except in No. 21. Pod (loment) transversely 2-several-jointed, the reticulated 1-seeded joints remaining closed, or sometimes reduced to one such joint.

* Leaves pinnate, with several leaflets, not stipellate.

- 17. **Æschynomene.** Stamens equally diadelphous (5 & 5). Calyx 2-lipped. Pod several-jointed: joints square.
- 18. **Hedysarum.** Stamens unequally diadelphous (9 & 1). Calyx 5-cleft. Pod several-jointed: joints roundish.
- ** Leaves pinnately 3-foliolate, rarely 1-foliolate.
- 19. **Desmodium.** Stamens diadelphous (9 & 1) or monadelphous below. Calyx 2-lipped. Pod several-jointed. Flowers all of one sort and complete. Leaflets stipellate.
- 20. **Lespedeza.** Stamens diadelphous (9 & 1): anthers uniform. Pod 1-2-jointed. Flowers often of 2 sorts, the more fertile ones apetalous. Leaflets not stipellate.
- 21. **Stylosanthes.** Stamens monadelphous: anthers of 2 sorts. Pod 1-2-jointed. Calyx deciduous, the tube narrow and stalk-like. Leaflets not stipellate.

Tribe V. VICIEÆ. Herbs, with abruptly pinnate leaves, usually climbing, the common leafstalk continued into a tendril or bristle. Peduncles axillary. Stamens chiefly diadelphous (9 & 1). Pod continuous, 2-valved, few-several-seeded. Cotyledons very thick, remaining under ground in germination.

- 22. **Vicia.** Wings adherent to the keel. Style filiform, bearded with a tuft or ring of hairs at the apex.
- 23. **Lathyrus.** Wings nearly free. Style somewhat dilated and flattened upwards, bearded down the inner face.

Tribe VI. PHASEOLEÆ. Twining or sometimes only trailing plants (herbs in temperate regions), with pinnately 3- (rarely 1- or 5-7-) foliolate leaves, commonly stipellate. Peduncles or flowers axillary; the pedicels usually clustered on the thickened nodes of the raceme. Anthers uniform. Pod continuous, not jointed, nor more than 1-celled, except by cellular matter sometimes deposited between the seeds, 2-valved. Cotyledons thick, and rising above ground little changed in germination, or sometimes becoming foliaceous.

* Leaves pinnate.

12. **Wistaria.** Woody twiner: leaflets 9-13. Keel barely incurved, obtuse.

24. **Aplous.** Herbaceous twiner: leaflets 5-7. Keel slender and much incurved or coiled.

** Leaves 3-foliolate. Ovules and seeds several. Flowers not yellow.

25. **Phaseolus.** Keel strongly incurved or coiled: standard recurved-spreading. Style bearded lengthwise.

26. **Centrosema.** Calyx short, 5-cleft. Standard with a spur at the base: keel broad, merely incurved. Style minutely bearded next the stigma.

27. **Clitorea.** Calyx tubular, 5-lobed. Standard erect, spurless: keel scythe-shaped. Style bearded down the inner face.

28. **Amphicarpea.** Calyx tubular, 4-5-toothed. Standard erect: keel almost straight. Style beardless. Some nearly apetalous fertile flowers next the ground.

29. **Galactia.** Calyx 4-cleft, the upper lobe broadest and entire. Style beardless. Bract and bractlets minute, mostly deciduous.

*** Leaves 1-3-foliolate. Ovules and seeds only one or two. Flowers yellow.

30. **Rhynchosia.** Keel scythe-shaped. Calyx 4-5-parted. Pod short.

B. Stamens all separate.

Tribe VII. SOPHOREÆ and PODALYRIÆ. Stamens 10, distinct; the corolla being truly papilionaceous.

31. **Baptisia.** Calyx 4-5-lobed. Pod inflated. Herbs: leaves palmately 3-foliolate or simple.

32. **Cladrastis.** Calyx 5-toothed. Pod very flat. Tree, with pinnate leaves.

SUBORDER II. CÆSALPINIÆ. BRASILETTO FAMILY.

Corolla imperfectly or not at all papilionaceous, sometimes nearly regular, imbricated in the bud, the upper or odd petal inside and enclosed by the others. Stamens 10 or fewer, commonly distinct, inserted on the calyx. Seeds anatropous, often with albumen. Embryo straight.

* Flowers imperfectly papilionaceous, perfect.

33. **Cereis.** Calyx campanulate, 5-toothed. Pod flat, wing-margined. Leaves simple.

** Flowers not at all papilionaceous, perfect.

34. **Cassia.** Calyx of 5 nearly distinct sepals. Leaves simply and abruptly pinnate.

*** Flowers not at all papilionaceous, polygamous or diœcious.

35. **Gymnocladus.** Tree: leaves all doubly pinnate. Calyx-tube elongated, at its summit bearing 5 petals resembling the calyx-lobes. Stamens 10.

36. **Gleditschia.** Trees thorny: leaves simply and doubly pinnate. Calyx-tube short; its lobes, petals, and stamens 3-5.

SUBORDER III. MIMOSÆ. MIMOSA FAMILY.

Flower regular. Corolla valvate in æstivation, often united into a 4-5-lobed cup, hypogynous, as are the (often very numerous) exerted stamens. Embryo straight. Leaves twice pinnate.

37. **Desmanthus.** Petals distinct. Stamens 5 or 10. Pod smooth.

38. **Schrankia.** Petals united below into a cup. Stamens 8 or 10. Pod covered with small prickles or rough projections.

1. LUPINUS, Tourn. LUPINE.

Calyx very deeply 2-lipped. Sides of the standard reflexed: keel scythe-shaped, pointed. Sheath of the monadelphous stamens entire: anthers alternately oblong and roundish. Pod oblong, flattened, often knotty by constrictions between the seeds. Cotyledons thick and fleshy. Herbs, with palmately 1-15-foliolate leaves, stipules adnate to base of the petiole, and showy flowers in terminal racemes or spikes. (Name from *Lupus*, a wolf, because these plants were thought to devour the fertility of the soil.)

1. *L. perennis*, L. (WILD LUPINE.) Perennial, somewhat hairy; stem erect (1° - 2°); leaflets 7-11, oblanceolate; flowers in a long raceme; pods very hairy. — Sandy soil: common. May, June. — Flowers showy, purplish-blue, rarely pale.

2. CROTALÀRIA, L. RATTLE-BOX.

Calyx 5-cleft, scarcely 2-lipped. Standard large, heart-shaped: keel scythe-shaped. Sheath of the monadelphous stamens cleft on the upper side: 5 of the anthers smaller and roundish. Pod inflated, oblong, many-seeded. — Herbs with simple leaves. Flowers yellow. (Name from *κρόταλον*, a rattle; the loose seeds rattling in the coriaceous inflated pods.)

1. *C. sagittalis*, L. Annual, hairy (3'-6' high); leaves oval or oblong-lanceolate, scarcely petioled; stipules united and decurrent on the stem, so as to be inversely arrow-shaped; peduncles few-flowered; corolla not longer than the calyx. — Sandy soil, Mass. to Virginia near the coast, Illinois and southward. July.

3. GENÍSTA, L. WOAD-WAXEN. WHIN.

Calyx 2-lipped. Standard oblong-oval, spreading: keel oblong, straight, deflexed. Stamens monadelphous, the sheath entire; 5 alternate anthers shorter. Pod mostly flat and several-seeded. — Shrubby plants, with simple leaves, and yellow flowers. (Name from the Celtic *gen*, a bush.)

1. *G. tinctoria*, L. (DYER'S GREEN-WEED.) Low, not thorny, with striate-angled erect branches; leaves lanceolate; flowers in spiked racemes. — Eastern New York and Massachusetts, especially Essex Co., where it is thoroughly established on sterile hills. June. (Adv. from Eu.)

4. TRIFOLIUM, L. CLOVER. TREFOIL.

Calyx persistent, 5-cleft, the teeth bristle-form. Corolla mostly withering or persistent; the claws of all the petals, or of all except the oblong or ovate standard, more or less united below with the stamen-tube: keel short and obtuse. Tenth stamen more or less separate. Pods small and membranous, often included in the calyx, 1-6-seeded, indehiscent, or opening by one of the sutures. — Tufted or diffuse herbs. Leaves mostly palmately, sometimes pinnately 3-foliolate: leaflets often toothed. Stipules united with the petiole. Flowers in heads or spikes. (Name from *tres*, three, and *folium*, a leaf.)

* *Flowers sessile in dense heads: corolla purple or purplish, withering away after flowering, tubular below; the petals more or less coherent with each other.*

+ *Calyx-teeth silky-plumose, longer than the whitish corolla: root annual.*

1. **T. ARVENSE**, L. (RABBIT-FOOT OR STONE CLOVER.) Silky, branching (5'-10' high); leaflets oblanceolate; heads becoming very soft-silky and grayish, oblong or cylindrical. — Old fields, &c. (Nat. from Eu.)

++ *Calyx scarcely hairy except a bearded ring in the throat, shorter than the rose-purple elongated-tubular corolla. (Short-lived perennials: flowers sweet-scented.)*

2. **T. PRATENSE**, L. (RED C.) Stems ascending, somewhat hairy; leaflets oval or obovate, often notched at the end and marked on the upper side with a pale spot; stipules broad, bristle-pointed; heads ovate, sessile. — Fields and meadows; largely cultivated. (Adv. from Eu.)

3. **T. MEDIUM**, L. (ZIGZAG C.) Stems zigzag, smoothish; leaflets oblong, entire, and spotless; heads mostly stalked; flowers deeper purple, otherwise too like the last. — Dry hills, E. Massachusetts. (Adv. from Eu.)

* * *Flowers pedicelled in umbel-like round heads on a naked peduncle, their short pedicels reflexed when old: corolla white or rose-color, withering-persistent and turning brownish in fading; the tubular portion short.*

4. **T. REFLEXUM**, L. (BUFFALO C.) Annual or biennial; stems ascending, downy; leaflets obovate-oblong, finely toothed; stipules thin, ovate; calyx-teeth hairy; pods 3-5-seeded. — Western New York (rare) to Illinois and southward. — Heads and flowers larger than in No. 2: standard rose-red; wings and keel whitish.

5. **T. STOLONIFERUM**, Muhl. (RUNNING BUFFALO-C.) Smooth, perennial; stems with long runners from the base; leaflets broadly obovate or obcordate, minutely toothed; heads loose; pods 2-seeded. — Open woodlands and prairies, Ohio to Illinois, Kentucky, and westward. — Flowers white, tinged with purple. Probably a variety of the last.

6. **T. REPENS**, L. (WHITE C.) Smooth, perennial; the slender stems spreading and creeping; leaflets inversely heart-shaped or merely notched, obscurely toothed; stipules scale-like, narrow; petioles and especially the peduncles very long; heads small and loose; calyx much shorter than the white corolla; pods about 4-seeded. — Fields and copses, everywhere. Here probably introduced, but indigenous northward.

7. **T. CAROLINIANUM**, Michx. (CAROLINA C.) Somewhat pubescent small perennial, procumbent, in tufts; leaflets wedge-obovate and slightly notched; stipules ovate, foliaceous; heads small on slender peduncles; calyx-teeth lanceolate nearly equalling the purplish corolla; standard pointed; pods 4-seeded. — Nat. from Southern States in waste grounds below Philadelphia (C. E. Smith); probably wild in S. Virginia. May.

* * * *Flowers short-pedicelled in close heads, reflexed when old: corolla yellow, persistent, turning dry and chestnut-brown with age, the standard becoming hood-shaped: annuals, fl. in summer.*

7. **T. AGRARIUM**, L. (YELLOW OR HOP-C.) Smoothish, somewhat upright (6'-12' high); leaflets obovate-oblong, all three from the same point (palmate) and nearly sessile; stipules narrow, cohering with the petiole for more than half its length. — Sandy fields, Massachusetts to Virginia. (Nat. from Eu.)

8. **T. PROCUMBENS**, L. (LOW HOP-C.) Stems spreading or ascending, pubescent (3'–6' high); *leaflets wedge-obovate*, notched at the end; *the lateral at a small distance from the other* (pinnately 3-foliolate); *stipules ovate, short*.—Sandy fields and roadsides, New England to Virginia. Also var. *MINUS* (T. minus, *Relhan*), with smaller heads, the standard not much striate with age. With the other; also Kentucky, in cultivated grounds. (Nat. from Eu.)

5. MELILÏTUS, Tourn. MELILOT. SWEET CLOVER.

Flowers much as in Clover, but in spiked racemes, small: corolla deciduous, free from the stamen-tube. Pod ovoid, coriaceous, wrinkled, longer than the calyx, scarcely dehiscent, 1–2-seeded.—Annual or biennial herbs, fragrant in drying, with pinnately 3-foliolate leaves; leaflets toothed. (Name from μέλι, *honey*, and Λωρός, some leguminous plant.)

1. **M. OFFICINALIS**, Willd. (YELLOW MELILOT.) Upright (2°–4° high); leaflets obovate-oblong, obtuse; *corolla yellow*; the petals nearly of equal length.—Waste or cultivated grounds. (Adv. from Eu.)

2. **M. ALBA**, Lam. (WHITE M.) Leaflets truncate; *corolla white*; the standard longer than the other petals. (M. leucántha, *Koch*.)—In similar places to the last, and much like it. (Adv. from Eu.)

6. MËDICÀGO, L. MËDICK.

Flowers nearly as in Melilotus. Pod 1–several-seeded, scythe-shaped, incurved, or variously coiled.—Leaves pinnately 3-foliolate. Stipules often cut. (Μηδική, the name of Lucerne, because it came to the Greeks from Media.)

1. **M. SATIVA**, L. (LUCERNE.) Upright, smooth, perennial; leaflets obovate-oblong, toothed; *flowers (purple) racemed*; pods spirally twisted.—Cultivated for green fodder, rarely spontaneous. (Adv. from Eu.)

2. **M. LUPULÏNA**, L. (BLACK MËDICK. NONESUCH.) Procumbent, pubescent, annual; leaflets wedge-obovate, toothed at the apex; *flowers in short spikes (yellow)*; pods kidney-form, 1-seeded.—Waste places. (Adv. from Eu.)

3. **M. MACULÀTA**, Willd. (SPOTTED MËDICK.) Spreading or procumbent annual, somewhat pubescent; leaflets obcordate, with a purple spot, minutely toothed; *peduncles 3–5-flowered*; flowers yellow; pods compactly spiral, of 2 or 3 turns, compressed, *furrowed on the thick edge*, and fringed with a double row of curved prickles.—Introduced with wool into waste grounds in some places. (Adv. from Eu.)

4. **M. DENTICULÀTA**, Willd. Nearly glabrous; pods loosely spiral, deeply reticulated, and with a thin keeled edge: otherwise like the last; in similar places, eastward. (Adv. from Eu.)

7. PSORÀLEA, L. PSORALEA.

Calyx 5-cleft, persistent, the lower lobe longest. Stamens diadelphous or sometimes monadelphous. Pod seldom longer than the calyx, thick, often wrinkled, indehiscent, 1-seeded.—Perennial herbs, usually sprinkled all over or roughened (especially the calyx, pods, &c.) with glandular dots or points.

Leaves mostly 3-5-foliolate. Flowers spiked or racemed, white or mostly blue-purple. Root sometimes tuberous and farinaceous. (Name, ψωραλέος, *scurfy*, from the glands or dots.)

* *Leaves pinnately 3-foliolate.*

1. **P. Onóbrychis**, Nutt. Nearly smooth and free from glands, erect (3°-5° high); leaflets lanceolate-ovate, taper-pointed (3' long); stipules and bracts awl-shaped; racemes elongated; peduncle shorter than the leaves; pods roughened and wrinkled. — River-banks, Ohio to Ill. and southward. July.

2. **P. stipulata**, Torr. & Gray. Nearly smooth and glandless; stems diffuse; leaflets ovate-elliptical, reticulated; stipules ovate; flowers in heads on rather short peduncles; bracts broadly ovate, sharp-pointed. — Rocks, Falls of the Ohio, Kentucky. June, July.

3. **P. melilotoides**, Michx. Somewhat pubescent, more or less glandular; stems erect (1°-2° high), slender; leaflets lanceolate or narrowly oblong; spikes oblong, long-peduncled; stipules awl-shaped; bracts ovate or lanceolate, taper-pointed; pods strongly wrinkled transversely. (Also *P. eglandulosa*, Ell.) — Dry soil, Ohio to Illinois, Virginia, and southward. June.

** *Leaves palmately 3-5-foliolate.*

4. **P. floribunda**, Nutt. Slender, erect, much branched and bushy (2°-4° high), minutely hoary-pubescent when young; leaflets varying from linear to obovate-oblong ($\frac{1}{2}$ ' - $1\frac{1}{2}$ ' long), glandular-dotted; racemes paniced; lobes of the calyx and bracts ovate, acute; pod glandular. — Prairies of Illinois and southwestward. June-Sept. — Flowers 2" or 3" long.

5. **P. argophylla**, Pursh. Silvery silky-white all over, erect, divergently branched (1°-3° high); leaflets elliptical-lanceolate; spikes interrupted; lobes of the calyx and bracts lanceolate. — High plains, N. Wisconsin, and westward. June. — Flowers 4" - 5" long.

6. **P. esculenta**, Pursh. Roughish hairy all over; stem stout (5' - 15' high) and erect from a tuberous or turnip-shaped farinaceous root; leaflets 5, obovate- or lanceolate-oblong; spikes oblong, dense, long-peduncled; lobes of the calyx and bracts lanceolate, nearly equalling the corolla ($\frac{1}{2}$ ' long). — High plains, N. W. Wisconsin, *Mr. Spears, T. J. Hale, &c.*, and westward. June. THE POMME BLANCHE, OR POMME DE PRAIRIE, of the Voyageurs.

8. DALEA, L. DALEA.

Calyx 5-cleft or toothed. Corolla imperfectly papilionaceous: petals all on claws: the standard heart-shaped, inserted in the bottom of the calyx: the keel and wings borne on the middle of the monadelphous sheath of filaments, which is cleft down one side. Stamens 10, rarely 9. Pod membranaceous, 1-seeded, indehiscent, enclosed in the persistent calyx. — Mostly herbs, more or less glandular-dotted with minute stipules; the small flowers in terminal spikes or heads. (Named for *Thomas Dale*, an English botanist.)

1. **D. alopecuroides**, Willd. Erect annual (1°-2° high), glabrous, except the dense and cylindrical silky-villous spike; leaves pinnate, of many linear-oblong leaflets; corolla whitish. — Alluvial soil, Illinois and southward. July. (Numerous species occur farther southwest.)

9. PETALOSTĒMON, Michx. PRAIRIE CLOVER.

Calyx 5-toothed. Corolla indistinctly papilionaceous: petals all on thread-shaped claws, 4 of them nearly similar and spreading, borne on the top of the monadelphous and cleft sheath of filaments, alternate with the 5 anthers; the fifth (standard) inserted in the bottom of the calyx, heart-shaped or oblong. Pod membranaceous, enclosed in the calyx, indehiscent, 1-seeded. — Chiefly perennial herbs, upright, glandular-dotted, with crowded odd-pinnate leaves, minute stipules, and small flowers in very dense terminal and peduncled heads or spikes. (Name combined of the two Greek words for petal and stamen, alluding to the peculiar union of these organs in this genus.)

1. **P. violáceus**, Michx. Smoothish: leaflets 5, narrowly linear; heads globose-ovate, or oblong-cylindrical when old; bracts pointed, not longer than the silky-hoary calyx; *corolla rose-purple*. — Dry prairies, Michigan to Minnesota and southward. July.

2. **P. candidus**, Michx. Smooth; leaflets 7-9, lanceolate or linear-oblong; heads oblong, when old cylindrical; bracts awned, longer than the nearly glabrous calyx; *corolla white*. — With No. 1. July.

3. **P. villósus**, Nutt. *Soft-downy or silky* all over; leaflets 13-17, linear or oblong, small (4" - 5" long); *spikes cylindrical* (1' - 5' long), short-peduncled, soft-villous; *corolla rose-color*. — N. Wisconsin (Lake Pepin, &c. *T. J. Hale*) and westward.

10. AMÓRPHA, L. FALSE INDIGO.

Calyx inversely conical, 5-toothed, persistent. Standard (the other petals entirely wanting!) wrapped around the stamens and style. Stamens 10, monadelphous at the very base, otherwise distinct. Pod oblong, longer than the calyx, 1-2-seeded, roughened, tardily dehiscent. — Shrubs, with odd-pinnate leaves; the leaflets marked with minute dots, usually stipellate. Flowers violet or purple; crowded in clustered terminal spikes. (Name, *ἄμορφη*, *wanting form*, from the absence of four of the petals.)

1. **A. fruticósa**, L. (FALSE INDIGO.) *Rather pubescent or smoothish*; leaflets 8-12 pairs, oval, scattered; pods 2-seeded. — River-banks S. Penn. to Wisconsin and southward. June. — A tall shrub: very variable.

2. **A. canéscens**, Nutt. (LEAD-PLANT.) *Low* (1°-3° high), *whitened with hoary down*; leaflets 15-25 pairs, elliptical, crowded, small, smoothish above with age; pods 1-seeded. — Prairies and crevices of rocks, Michigan to Wisconsin and southwestward. July. — Supposed to indicate lead-ore.

11. ROBINIA, L. LOCUST-TREE.

Calyx short, 5-toothed, slightly 2-lipped. Standard large and rounded, turned back, scarcely longer than the wings and keel. Stamens diadelphous. Pod linear, flat, several-seeded, margined on the seed-bearing edge, at length 2-valved. — Trees or shrubs, often with prickly spines for stipules. Leaves odd-pinnate, the ovate or oblong leaflets stipellate. Flowers showy, in hanging axillary racemes. Base of the leaf-stalks covering the buds of the next year. (Named in honor of *John Robin*, herbalist to Henry IV. of France, and his son *Vespasian Robin*, who first cultivated the Locust-tree in Europe.)

1. **R. Pseudacacia**, L. (COMMON LOCUST OR FALSE ACACIA.) Branches naked; *racemes slender, loose*; flowers white, fragrant; pod smooth. — S. Pennsylvania to S. Illinois and southward. Commonly cultivated as an ornamental tree, and for its valuable timber: naturalized in many places. June.

2. **R. viscosa**, Vent. (CLAMMY L.) *Branchlets and leafstalks clammy*; flowers crowded in oblong racemes, tinged with rose-color, nearly inodorous; pod glandular-hispid. — Virginia and southward. Cultivated, like the last, a smaller tree. June.

3. **R. hispida**, L. (BRISTLY L. OR ROSE ACACIA.) *Branchlets and stalks bristly*; flowers large and deep rose-color, inodorous; pods glandular-hispid. — Varies with less bristly or nearly naked branchlets; also with smaller flowers, &c. — Mountains of Virginia and southward: commonly cultivated. May, June. — Shrub 3°–8° high.

12. WISTARIA, Nutt. WISTARIA.

Calyx campanulate, somewhat 2-lipped; upper lip of 2 short teeth, the lower of 3 longer ones. Standard roundish, large, turned back, with 2 callosities at its base: keel scythe-shaped: wings doubly auricled at the base. Stamens diadelphous. Pod elongated, thickish, knobby, stipitate, many-seeded, at length 2-valved. Seeds large. — Woody twiners, climbing high, with minute stipules, pinnate leaves of 9–13-ovate-lanceolate leaflets, with or without minute stipels, and dense racemes of large and showy lilac-purple flowers. (Dedicated to the late *Professor Wistar*, of Philadelphia.)

1. **W. frutescens**, DC. Downy or smoothish when old; wings of the corolla with one short auricle and an awl-shaped one as long as the claw. (*W. speciosa*, Nutt.) — Alluvial grounds, W. Virginia to Illinois and southward. May. — Sometimes cultivated for ornament, as is the still handsomer Chinese species.

13. TEPHRÛSIA, Pers. HOARY PEA.

Calyx about equally 5-cleft. Standard roundish, usually silky outside, turned back, scarcely longer than the coherent wings and keel. Stamens monadelphous or diadelphous. Pod linear, flat, several-seeded, 2-valved. — Hoary perennial herbs, with odd-pinnate leaves, and white or purplish racemed flowers. Leaflets mucronate, veiny. (Name from *τεφρός*, ash-colored or hoary.)

1. **T. Virginiana**, Pers. (GOAT'S RUE. CATGUT.) *Silky-villous* with whitish hairs when young; stem erect and simple (1°–2° high), leafy to the top; leaflets 17–29, linear-oblong; flowers large and numerous, clustered in a terminal oblong dense raceme or panicle, yellowish-white marked with purple. — Dry sandy soil. June, July. — Roots long and slender, very tough.

2. **T. spicata**, Torr. & Gray. *Villous with rusty hairs*; stems branched below, straggling or ascending (2° long), few-leaved; leaflets 9–15, obovate or oblong-wedge-shaped, often notched; flowers few, in a loose interrupted very long-peduncled spike, reddish. — Dry soil, Delaware and southward. July.

3. **T. hispídula**, Pursh. Hairy with some long and rusty or only minute and appressed pubescence; stems slender (9'–24' long), divergently branched,

straggling; leaflets 5-15, oblong, varying to obovate-wedge-shaped and oblanceolate; *peduncles longer than the leaves, 2-4-flowered*; flowers reddish-purple. — Dry sandy soil, Virginia and southward.

14. ASTRÁGALUS, L. MILK-VETCH.

Calyx 5-toothed. Corolla usually long and narrow: standard narrow, equaling or exceeding the wings and blunt keel, its sides reflexed or spreading. Stamens diadelphous. Stigma minute, terminal. Pod several-many-seeded, various, mostly turgid, one or both sutures usually projecting into the cell, either slightly or so as to divide the cavity lengthwise into two. Seed-stalks slender. Chiefly herbs (ours perennials), with odd-pinnate leaves and spiked or racemed flowers. (The ancient Greek name of a leguminous plant, as also of the ankle-bone; but the connection between the two is past all guess.)

§ 1. *Pod very thick and juicy when fresh, not stalked in the calyx, 2-celled, indehiscent, or tardily separable into 2 closed portions: stems low, decumbent or ascending: leaflets numerous.*

1. **A. caryocarpus**, Ker. (GROUND PLUM.) Pale and minutely appressed-pubescent; leaflets narrowly oblong; flowers in a short spike-like raceme: *corolla violet-purple; fruit glabrous, ovate-globular, more or less pointed, about ¾' in diameter, very thick-walled, cellular or corky when dry.* Upper Mississippi River, thence westward and southward. May.

2. **A. Mexicánus**, A. DC. Smoother, or pubescent with looser hairs, larger; leaflets roundish, obovate, or oblong; flowers larger (10''-12'' long); calyx softly hairy; *corolla cream-color, bluish only at the tip; fruit globular, very obtuse and pointless, 1' or more in diameter: otherwise like the last: the unripe fruits of both resemble green plums, — whence the popular name, — and are eaten, raw or cooked, by travellers.* (A. trichocalyx, Nutt.) — Prairies and open plains, from Illinois opposite St. Louis westward and southward.

3. **A. Platténsis**, Nutt. Loosely villous; stipules conspicuous; leaflets oblong, often glabrous above; flowers crowded in a short spike or oblong head, cream-color often tinged or tipped with purple; *fruit ovate, pointed, and with the calyx villous, — in var. TENNESSEÉNSIS (A. Tennesseeensis, Gray, in Chaym. S. Fl.) oblong and slightly curved, much less fleshy and thick than in the preceding.* — Gravelly or sandy banks of Illinois River (Vasey, Slosson, Bebb), thence southward and westward. May.

§ 2. *Pod dry and dehiscent, more or less thin-walled, turgid, not stalked in the calyx, completely 2-celled only in No. 4.*

4. **A. Canadénsis**, L. Tall and erect (1°-4° high), somewhat pubescent; leaflets 21-27, oblong; *flowers greenish cream-color, very numerous, in long and close spikes; pods ovoid-oblong, coriaceous, small, completely 2-celled.* — Riverbanks: common from N. New York westward and southward. July-Aug.

5. **A. Coöperi**, Gray. Nearly smooth, erect (1°-2° high); leaflets 11-21, elliptical or oblong, somewhat notched at the end, minutely hoary underneath; *flowers white, rather numerous in a short spike; pods inflated-ovoid (almost 1' long), thin-walled, one-celled, the dorsal suture slightly, the ventral more decidedly projecting inwards.* (Phaca neglecta, Torr. & Gray. Now named for

the discoverer, the late *Wm. Cooper*, there being already an *A. neglectus*.) — Gravelly shores, &c., W. New York to Wisconsin. June, July.

6. *A. distortus*, Torr. & Gray. Low and spreading, branched from the base, smoothish; leaflets 11–23, oblong or obovate; flowers purplish or violet, 10–20 in a short spike; the standard deeply notched at the summit; pods oblong, turgid, incurved ($\frac{3}{4}$ long), coriaceous, incompletely 2-celled. — Mason Co., Illinois, *Dr. Mead*. May. (Also in Arkansas and Texas.)

§ 3. Pod dry and dehiscent, thin-walled, small, stalked in the calyx (stipitate), and with it more or less pubescent with fine blackish hairs, hanging on short pedicels: raceme short, rather many-flowered, long-peduncled: leaflets oval or oblong.

7. *A. alpinus*, L. Smooth or slightly hairy; stem diffuse, 6' to 12' high; leaflets 13–25; corolla violet-purple, or at least the keel tipped with violet or blue (5"–6" long); pods black-hairy, oblong, deeply grooved on the back and partly 2-celled by the intrusion of the dorsal suture, its stipe usually rather exceeding the calyx. — Rocks and banks, Northern Vermont (*Willoughby Mountain, J. Blake*) and Maine (*Dr. Scammon, G. L. Goodale*), and northward. June, July. (Eu.)

8. *A. Robbinsii*, Gray. Nearly smooth and erect (1° high), slender; leaflets 7–11; corolla white (4" long); calyx-teeth short; pods oblong, flattish ($\frac{1}{2}$ long), membranaceous, almost glabrous, the base suddenly contracted into a stipe about equalling the calyx, one-celled, a thin membrane slightly projecting from the dorsal suture. (*Phaca Robbinsii, Oakes*.) — Rocky ledges of Onion River, at Colchester, Vermont, *Dr. Robbins* (1829): the station now obliterated. May.

15. OXYTROPIS, DC. OXYTROPIS.

Keel of the corolla tipped with a sharp projecting point or appendage: otherwise as in *Astragalus*. Pod often partly 2-celled by the intrusion of the ventral suture. — Our species, and most others, are low, nearly acaulescent perennials, with tufts of numerous very short stems from a hard and thick root or rootstock, covered with scaly adnate stipules; pinnate leaves of many leaflets; and naked scapes bearing a head or short spike of flowers. (Name indicates the peculiarity of the flower, from *οξύς*, sharp, and *τρόπις*, keel.)

1. *O. campestris*, DC. Pubescent or smoothish; leaflets lanceolate or oblong: flowers yellowish or white, often tinged or tipped with purple or violet-blue; pods ovate or oblong-lanceolate, of a thin or papery texture. — Northern border of Maine, on the St. John's, near Seven Isles, *G. L. Goodale*, and northward. July. (Eu.)

2. *O. Lamberti*, Pursh. Silky with fine appressed hairs; leaflets mostly linear; flowers larger, purple, violet, or sometimes white; pods cartilaginous or firm coriaceous in texture, strictly erect, cylindraceous-lanceolate and long-pointed, almost 2-celled by intrusion of the ventral suture. — Dry plains, Minnesota and westward. June.

16. GLYCYRRHIZA, Tourn. LIQUORICE.

Calyx with the two upper lobes shorter or partly united. Anther-cells confluent at the apex, the alternate ones smaller. Pod ovate or oblong-linear, com-

pressed, often curved, clothed with rough glands or short prickles, scarcely dehiscent, few-seeded. The flower, &c., otherwise as in *Astragalus*. — Long perennial root sweet (whence the name, from *γλυκὺς*, *sweet*, and *ρίζα*, *root*); herbage glandular-viscid; leaves odd-pinnate, with minute stipules; flowers in axillary spikes, white or bluish.

1. *G. lepidota*, Nutt. (WILD LIQUORICE.) Tall (2°–3° high); leaflets 15–19, oblong-lanceolate, mucronate-pointed, sprinkled with little scales when young, and with corresponding dots when old; spikes peduncled, short; flowers whitish; pods oblong, beset with hooked prickles, so as to resemble the fruit of *Xanthium* on a smaller scale. — Vicinity of Buffalo, New York, on the sands of the shore, probably drifted from its native northwest regions; but perfectly established, *G. W. Clinton*.

17. ÆSCHYNÓMENE, L. SENSITIVE JOINT-VETCH.

Calyx 2-lipped; the upper lip 2-, the lower 3-cleft. Standard roundish: keel boat-shaped. Stamens diadelphous in two sets of 5 each. Pod flattened, composed of several square easily separable joints. — Leaves odd-pinnate, with several pairs of leaflets, sometimes sensitive, as if shrinking from the touch (whence the name, from *αἰσχνομένη*, *being ashamed*).

1. *Æ. hispida*, Willd. Erect, rough-bristly annual; leaflets 37–51, linear; racemes few-flowered; pod stalked, 6–10-jointed. — Along rivers, S. Penn. and southward. Aug. — Flowers yellow, reddish externally.

18. HEDÝSARUM, Tourn. HEDYSARUM.

Calyx 5-cleft, the lobes awl-shaped and nearly equal. Keel nearly straight, obliquely truncate, not appendaged, longer than the wings. Stamens diadelphous, 5 & 1. Pod flattened, composed of several equal-sided separable roundish joints connected in the middle. — Perennial herbs: leaves odd-pinnate. (Name composed of *ἡδύς*, *sweet*, and *ἄρωμα*, *smell*.)

1. *H. boreale*, Nutt. Leaflets 13–21, oblong or lanceolate, nearly glabrous; stipules scaly, united opposite the petiole; raceme of many deflexed purple flowers; standard shorter than the keel; joints of the pod 3 or 4, smooth, reticulated. — Mountain above Willoughby Lake, Vermont, *A. Wood*, &c. St. John's River, Maine, *G. L. Goodale*. Also northward.

19. DESMÒDIUM, DC. TICK-TREFOIL.

Calyx usually more or less 2-lipped. Standard obovate: wings adherent to the straight or straightish and usually truncate keel, by means of a little transverse appendage on each side of the latter. Stamens diadelphous, 9 & 1, or monadelphous below. Pod flat, deeply lobed on the lower margin, separating into few or many flat reticulated joints (mostly roughened with minute hooked hairs, by which they adhere to the fleece of animals or to clothing). — Perennial herbs, with pinnately 3-foliolate (rarely 1-foliolate) leaves, stipellate. Flowers (in summer) in axillary or terminal racemes, often paniced, and 2 or 3 from each bract, purple or purplish, often turning green in withering. Stipules and

bracts scale-like, often striate. (Name from *δεσμός*, a bond or chain, from the connected joints of the pods.)

§ 1. *Pod raised on a stalk within the calyx (stipe) many times longer than the slightly toothed calyx and nearly as long as the pedicel, straightish on the upper margin, deeply sinuate on the lower; the 1-4 joints mostly half-obovate and concave on the back: stamens monadelphous below: plants nearly glabrous: stems erect or ascending: raceme terminal, paniced: stipules bristle-form, deciduous.*

1. **D. nudiflorum**, DC. *Leaves all crowded at the summit of sterile stems; leaflets broadly ovate, bluntish, whitish beneath; raceme elongated on an ascending mostly leafless stalk or scape from the root, 2° long. — Dry woods: common.*

2. **D. acuminatum**, DC. *Leaves all crowded at the summit of the stem from which arises the elongated naked raceme or panicle; leaflets round-ovate, taper-pointed, green both sides, the end one round (4'-5' long). — Rich woods.*

3. **D. pauciflorum**, DC. *Leaves scattered along the low (8'-15' high) ascending stems; leaflets rhombic-ovate, bluntish, pale beneath; raceme few-flowered, terminal. — Woods, W. New York and Pennsylvania to Illinois and southward.*

§ 2. *Pod raised on a stalk (stipe) little if not all surpassing the deeply-cleft calyx: stems long and prostrate or decumbent: racemes axillary and terminal.*

* *Stipules conspicuous, ovate, taper-pointed, striate, persistent: racemes mostly simple.*

4. **D. rotundifolium**, DC. *Soft-hairy all over, truly prostrate; leaflets orbicular, or the odd one slightly rhomboid; flowers purple; pods almost equally sinuate on both edges, 3-5-jointed; the joints rhomboid-oval. — Dry rocky woods: rather common.*

Var. **glabratum**: almost glabrous, otherwise nearly as the ordinary form. (*Hedysarum humifusum*, Muhl. in part, Bigel., &c.) — Mass., New York, &c.

5. **D. ochroleucum**, M. A. Curtis. *Stems sparsely hairy, decumbent; leaflets nearly glabrous, ovate, acute or obtuse, transversely reticulated beneath, the lateral ones smaller or sometimes wanting; racemes much elongated; corolla whitish; pods twisted, 2-4-jointed, the large rhomboid joints smooth and reticulated but the margins downy. (Perhaps Muhlenberg's *H. humifusum* from "Carolina.") — Woodlands, Maryland (*W. M. Canby*) and southward.*

* * *Stipules smaller, lanceolate and awl-shaped, less persistent: racemes paniced.*

6. **D. humifusum**, Beck (as to syn.). *Glabrous or nearly so, procumbent; leaflets ovate or ovate-oblong, rather obtuse, much smaller than in the two preceding (1½'-2' long); corolla purple; pods 2-4-jointed, flat, the oval-rhomboid joints minutely scabrous throughout. (*Hedysarum humifusum*, Muhl. *Fl. Lancaster. herb., ex Canby.*) — Dry sandy soil (Lancaster, Penn., Muhlenberg), Salisbury, Maryland, *W. M. Canby*.*

§ 3. *Pod slightly if at all stalked in the calyx: racemes paniced.*

* *Stems tall (3°-5°) and erect; the persistent stipules and deciduous bracts large and conspicuous, ovate or ovate-lanceolate, taper-pointed: pods of 4-7 unequal-sided rhombic joints, which are considerably longer than broad, about ½' long. (Flowers rather large.)*

7. **D. canescens**, DC. *Stem loosely branched, hairy; leaflets ovate, bluntish, about the length of the petioles, whitish and reticulated beneath, both sides*

roughish with a close fine pubescence; joints of the pod very adhesive. — Moist grounds, W. Vermont to Wisconsin and southward, chiefly westward. — Branches clothed with both minute and hooked, and longer, spreading, rather glutinous hairs.

8. **D. cuspidatum**, Torr. & Gray. *Very smooth* except the panicle; stem straight; *leaflets lanceolate-ovate and taper-pointed*, green both sides; longer than the petiole (3'–5'); joints of the pod rhomboid-oblong, smoothish. — Thickets: common. — The conspicuous bracts and stipules $\frac{3}{4}$ ' long.

* * *Stems* (2°–5° high) *erect*: stipules as well as the bracts mostly deciduous, small and inconspicuous: pods of 3–5 triangular or half-rhombic or very unequal-sided rhomboidal joints, which are longer than broad, $\frac{1}{4}$ ' or less in length. (*Flowers middle-sized.*)

9. **D. lævigatum**, DC. *Smooth* or nearly so throughout: stem straight; *leaflets ovate*, bluntish, pale beneath (2'–3' long); panicles minutely rough-pubescent. — Pine woods, New Jersey and southward.

10. **D. viridiflorum**, Beck. Stem *very downy*, rough at the summit; *leaflets broadly ovate*, very obtuse, rough above, whitened with a soft velvety down underneath (2'–3' long). — S. New York and southward.

11. **D. Dillénii**, Darlingt. Stem pubescent; *leaflets oblong or oblong-ovate*, commonly bluntish, pale beneath, *softly and finely pubescent* (mostly thin, 2'–3' long). — Open woodlands: common.

12. **D. paniculatum**, DC. *Nearly smooth* throughout; stem slender, tall; *leaflets oblong-lanceolate, or narrowly lanceolate, tapering to a blunt point*, thin (3'–5' long); racemes much paniced. — Copses, common.

13. **D. strictum**, DC. Stem very straight and slender, simple (2°–3° high), the upper part and narrow panicle rough-glandular; *leaflets linear, blunt, strongly reticulated, thickish, very smooth* (1'–2' long, $\frac{1}{4}$ ' wide); joints of the pod 1–3, semi-obovate or very gibbous (only 2'' long). — Pine woods of New Jersey, and southward.

* * * *Stipules small and inconspicuous, mostly deciduous*: pods of few roundish or obliquely oval or sometimes roundish-rhomboidal joints, $1\frac{1}{2}$ ''– $2\frac{1}{2}$ '' long.

+ *Stems erect*: bracts before flowering conspicuous: racemes densely flowered.

14. **D. Canadense**, DC. Stem hairy (3°–6° high); *leaflets oblong-lanceolate, or ovate-lanceolate*, obtuse, with numerous straightish veins, *much longer than the petiole* ($1\frac{1}{2}$ '–3' long); *flowers showy*, larger than in any other species ($\frac{1}{2}$ '– $\frac{1}{4}$ ' long). — Dry, rich woods: common, especially northward.

15. **D. sessilifolium**, Torr. & Gray. Stem pubescent (2°–4° high); *leaves nearly sessile*; *leaflets linear or linear-oblong, blunt, thickish, reticulated*, rough above, downy beneath; branches of the panicle long; *flowers small*. — Copses, Pennsylvania and Michigan to Illinois and southward.

+ + *Stems ascending* (1°–3° high): bracts small; racemes or panicles elongated and loosely flowered: flowers small.

16. **D. rigidum**, DC. Stem branching, somewhat hoary, like the lower surface of the leaves, with a close roughish pubescence; *leaflets ovate-oblong, blunt, thickish, reticulated-veiny*, rather rough above, *the lateral ones longer than the petiole*. — Dry hillsides, Mass. to Michigan, Illinois, and southward. — Intermediate, as it were, between No. 17 and No. 11.

17. **D. ciliare**, DC. Stem slender, *hairy or rough-pubescent*; leaves crowded, on very short hairy petioles; leaflets round-ovate or oval, thickish, more or less hairy on the margins and underneath ($\frac{1}{2}$ '-1' long).—Dry hills and sandy fields: common, especially southward.

18. **D. Marilándicum**, Boott. Nearly smooth throughout, slender; leaflets ovate or roundish, very obtuse, thin, the lateral ones about the length of the slender petiole: otherwise resembling the preceding. (*D. obtusum*, DC.)—Copses: common.

+++ Stems reclining or prostrate: racemes loosely flowered.

19. **D. lineatum**, DC. Stem minutely pubescent, striate-angled; leaflets orbicular, smoothish ($\frac{1}{2}$ '-1' long), much longer than the petiole; pod scarcely stalked in the calyx.—Dry soil, Maryland (*W. M. Canby*), Virginia and southward.

20. LESPEDEZA, Mich. BUSH-CLOVER.

Calyx 5-cleft; the lobes nearly equal, slender. Stamens diadelphous (9 & 1): anthers all alike. Pods of a single 1-seeded joint (sometimes 2-jointed, with the lower joint empty and stalk-like), oval or roundish, flat, reticulated.—Perennials with pinnately 3-foliolate leaves, not stipellate. Stipules and bracts minute. Flowers often polygamous, in summer and autumn. (Dedicated to *Lespedez*, the Spanish governor of Florida in the time of Michaux.)

* *Flowers of two sorts, the larger (violet-purple) perfect, but seldom fruitful, paniced or clustered; with smaller pistillate and fertile but mostly apetalous ones intermixed, or in sessile little clusters.*

1. **L. procumbens**, Michx. *Soft-downy*, except the upper surface of the leaves, trailing, slender; leaflets oval or elliptical; peduncles slender, mostly simple, few-flowered.—Sandy soil: commonest southward.—The apetalous fertile flowers, as in the rest, have short hooked styles.

2. **L. rèpens**, Torr. & Gray. *Smooth, except minute close-pressed scattered hairs, prostrate*, spreading, very slender; leaflets oval or obovate-elliptical ($\frac{1}{2}$ ' long); peduncles slender and few-flowered; pods roundish.—Dry sandy soil, S. New York to Kentucky and southward.—Much like the last.

3. **L. violàcea**, Pers. *Stems upright or spreading*, branched; leaflets varying from oval-oblong to linear, whitish-downy beneath with close-pressed pubescence; peduncles or clusters few-flowered; pods ovate.—The principal varieties are, 1. *DIVÉRGENS*, with oval or oblong leaflets and loosely paniced flowers; this runs into, 2. *SESSILIFLORA*, with the flowers principally on peduncles much shorter than the leaves, and clustered; and a more distinct form is 3. *ANGUSTIFOLIA*, with closely clustered flowers on straight branches, crowded leaves, and narrowly oblong or linear leaflets, which are often silky.—Dry copses: common.—Pods ripening from both sorts of flowers.

4. **L. Stùvei**, Nutt. *Stems upright-spreading*, bushy, downy; leaflets oval or roundish, longer than the petiole, silky or white-woolly beneath (and sometimes above); clusters many-flowered, crowded; pods ovate, downy.—Dry hills, and sand, Plymouth, Mass. to Virginia, Michigan, and southward.—Appearing intermediate between No. 3 and No. 5.

* * *Flowers all alike and perfect, in close spikes or heads: corolla whitish or cream-color with a purple spot on the standard, about the length of the downy calyx: stems upright, wand-like (2° - 4° high).*

5. **L. hirta**, Ell. *Peduncles longer than the leaves; petioles slender; leaflets roundish or oval, hairy; spikes cylindrical, rather loose; pods nearly as long as the calyx. (L. polystachia, Michx.)*—Dry hillsides.

6. **L. capitata**, Michx. *Peduncles and petioles short; stems rigid, woolly; leaflets elliptical or oblong, thickish, reticulated and mostly smooth above, silky beneath; spikes or heads short; pods much shorter than the calyx.*—Varies greatly, most of all in var. **ANGUSTIFOLIA**: slender; leaflets linear; peduncles sometimes elongated.—Dry and sandy soil; the narrow variety only found near the coast and southward.

21. STYLOSÁNTHES, Swartz. PENCIL-FLOWER.

Calyx early deciduous; the tube slender and stalk-like; the limb unequally 4-5-cleft, the lower lobe more distinct. Corolla and monadelphous stamens inserted at the summit of the calyx-tube: standard orbicular: keel incurved. Anthers 10, the 5 longer ones fixed near their base, and the 5 alternate shorter ones fixed by the middle. Style filiform, its upper part falling off after flowering, the lower part incurved or hooked, and persistent on the apex of the 1-2-jointed small and short reticulated pod, the lower joint when present empty and stalk-like.—Low perennials, branched from the base, with wiry stems, pinnately 3-foliolate leaves, the sheathing stipules united to the petiole, no stipels, and small, yellow flowers in terminal heads or short spikes. (Name composed of *στύλος*, a column, and *ἄνθος*, a flower, from the stalk-like calyx-tube.)

1. **S. elatior**, Swartz. Tufted; leaflets lanceolate, strongly straight-veined; heads or clusters small and few-flowered.—Pine barrens, Long Island, New York, to Virginia, Illinois, and southward. July—Oct.

22. VÍCIA, Tourn. VETCH. TARE.

Calyx 5-cleft or 5-toothed, the 2 upper teeth often shorter, or the lowest longer. Wings of the corolla adhering to the middle of the keel. Stamens more or less diadelphous (9 & 1); the orifice of the tube oblique. Style filiform, hairy all round or only on the back at the apex. Pod flat, 2-valved, 2-several-seeded. Seeds globular. Cotyledons very thick, remaining under ground in germination.—Herbs, mostly climbing more or less by the tendril at the end of the pinnate leaves. Stipules half-sagittate. Flowers or peduncles axillary. (The classical Latin name.)

* *Annual: flowers 1-2 in the axils, nearly sessile, large, violet-purple.*

1. **V. sativa**, L. (COMMON VETCH OR TARE.) Somewhat pubescent; stem simple; leaflets 5-7 pairs, varying from obovate-oblong to linear, notched and mucronate at the apex; pod linear, several-seeded.—Cultivated fields and waste places; both the common form and the var. **ANGUSTIFOLIA**, which has longer and narrow leaflets. (Adv. from Eu.)

* * *Annual, slender: peduncles elongated: flowers small.* (Species of *Ervum*, L.)

2. **V. TETRASPÉRMA**, L. *Peduncles 1-2-flowered; leaflets 4-6 pairs, linear-oblong, obtuse; calyx-teeth unequal; corolla whitish; pods narrowly oblong, 4-seeded, smooth.* — Waste or open places, near the coast. (Nat. from Eu.)

3. **V. HIRSUTA**, Koch. *Peduncles 3-6-flowered; leaflets 6-8 pairs, truncate; calyx-teeth equal; corolla bluish; pods oblong, 2-seeded, hairy.* — Massachusetts to Virginia. (Nat. from Eu.)

* * * *Perennial: peduncles elongated; calyx-teeth unequal: pod several-seeded.*

4. **V. Cracca**, L. *Downy-pubescent; leaflets 20-24, oblong-lanceolate, strongly mucronate; peduncles densely many-flowered; calyx-teeth shorter than the tube.* — Borders of thickets, New England to Kentucky and northward: rather rare. July. — Flowers blue, turning purple, 6" long, one-sided in the spike, reflexed. (Eu.)

5. **V. Caroliniana**, Walt. *Nearly smooth; leaflets 8-24, oblong, obtuse, scarcely mucronate; peduncles loosely flowered; calyx-teeth very short.* — Riverbanks, &c. May. — Flowers small, more scattered than in the preceding, whitish, the keel tipped with blue.

6. **V. Americana**, Muhl. *Glabrous; leaflets 10-14, elliptical or ovate-oblong, very obtuse, many-veined; peduncles 4-8-flowered.* — Moist soil, New York to Kentucky and northward. June. — Flowers purplish, 8" long.

23. LÁTHYRUS, L. VETCHLING. EVERLASTING PEA.

Style flattish, dilated and flattish (not grooved) above, hairy along the inner side (next the free stamen). Sheath of the filaments scarcely oblique at the apex. Otherwise nearly as in *Vicia*. (*Λάθυρος*, a leguminous plant of Theophrastus.) — Our species are perennial and mostly smooth plants.

1. **L. maritimus**, Bigelow. (BEACH PEA.) *Stem stout (1° high); leaflets 4-8 pairs, crowded, oval or obovate; stipules broadly halberd-shaped, nearly as large as the leaflets; peduncles 6-10-flowered.* — Sea-coast, from New Jersey northward, and shore of the Great Lakes. June-Aug. — Flowers large, purple. Leaflets very veiny, as also in the other species. (Eu.)

2. **L. venosus**, Muhl. *Stem climbing (2°-5° high); leaflets 5-7 pairs, scattered, oblong-ovate, often downy beneath; stipules very small and usually slender, half arrow-shaped, rarely larger and broader; peduncles many-flowered; corolla purple.* — Shady banks, Penn. to Wisconsin, and southward. June.

3. **L. ochroleucus**, Hook. *Stem slender (1°-3° high); leaflets 3-4 pairs, ovate or oval, smooth, glaucous, thin; stipules half heart-shaped, about half as large as the leaflets; peduncles 7-10-flowered; corolla yellowish-white, smaller than in the last.* — Hillsides, W. Vermont to Pennsylvania, Wisconsin, and northward. July.

4. **L. palustris**, L. (MARSH VETCHLING.) *Stem slender (1°-2° high), often winged-margined; leaflets 2-4 pairs, lanceolate, linear, or narrowly oblong, mucronate-pointed; stipules small, lanceolate, half arrow-shaped, sharp-pointed at both ends; peduncles 3-5-flowered; corolla blue-purple.* — Moist places, N. England to Penn., Illinois, and northward. July. (Eu.)

Var. **myrtifolius**. *Taller, climbing 2°-4° high; leaves oblong or ovate-*

elliptical; upper stipules much larger: corolla pale purple. (*L. myrtifolius*, *Muhl.*)—W. New England to Virginia and northward, July.—Ordinarily appears quite distinct from *L. palustris*; but intermediate specimens occur.

5. **L. PRATENSIS**, L. Low and straggling; leaflets a single pair, narrow-lanceolate; stipules large; peduncles several-flowered; corolla yellow.—Spontaneous and abundant along the Connecticut at West Springfield, Mass., *A. P. Foster*. July. (Adv. from Eu.)

24. **ÁPIOS**, Boerhaave. GROUND-NUT. WILD BEAN.

Calyx somewhat 2-lipped, the 2 lateral teeth being nearly obsolete, the upper very short, the lower one longest. Standard very broad, reflexed: the long scythe-shaped keel strongly incurved, at length coiled. Stamens diadelphous. Pod straight or slightly curved, linear, elongated, thickish, many-seeded.—A perennial herb (with some milky juice!), bearing edible tubers on underground shoots, twining and climbing over bushes. Leaflets 5-7, ovate-lanceolate, obscurely stipellate. Flowers in dense and short, often branching racemes. (Name from *ἄπιον*, a pear, from the shape of the tubers.)

1. **A. tuberosa**, Mœnch. (*Glycine Apios*, L.)—Low grounds; common. Aug.—Sept.—Flowers brown purple, or chocolate-color, violet-scented.

25. **PHASÈOLUS**, L. KIDNEY BEAN.

Calyx 5-toothed or 5-cleft, the two upper teeth often higher united. Keel of the corolla, with the included stamens and style, spirally coiled or curved into a ring. Stamens diadelphous. Style bearded along the upper side: stigma oblique or lateral. Pod linear or scythe-shaped, several-many-seeded, tipped with the hardened base of the style. Cotyledons thick and fleshy, rising out of the ground nearly unchanged in germination.—Twining or prostrate herbs, with pinnately 3-foliolate stipellate leaves. Flowers often clustered on the knotty joints of the raceme, produced in summer and autumn. (The ancient name of the Kidney Bean.)

* *Pods scymetar-shaped: racemes long and loose, paniced.*

1. **P. perennis**, Walt. (WILD BEAN.) Stem climbing high from a perennial root; leaflets roundish-ovate, short-pointed; pods drooping, strongly curved, 4-5-seeded.—Copses, Connecticut to Illinois, and southward.—Flowers purple, handsome, but small.

* * *Pods long and straight, linear, rather terete: flowers few in a short clustered and long-peduncled raceme.* (*Strophóstyles*, Ell.)

2. **P. diversifolius**, Pers. Annual; stem prostrate, spreading, rough-hairy; leaflets ovate-3-lobed, or angled towards the base, or some of them oblong-ovate and entire; peduncles at length twice the length of the leaves.—Sandy fields and banks, coast of Massachusetts and along the Great Lakes to Illinois and southward.—Corolla greenish-white, tinged with red or purple. Pod thickish.

3. **P. hélvolus**, L. Perennial, hairy; stems diffuse, slender; leaflets ovate or oblong, entire or obscurely angled; peduncles 3-6 times the length of the leaves.—Sandy fields, S. New York to Illinois and southward.—More slender than the last: pods narrower: flowers as large and similar.

* * *Pods straight and linear, flat : peduncles short, 1 - few-flowered at the summit : flowers small : keel less incurved.*

4. **P. pauciflorus**, Benth. Annual; stems diffuse, but twining, slender, pubescent; leaflets varying from oblong-lanceolate or ovate-oblong to linear. (*P. leiospermus*, Torr. & Gr.) — River-banks, Illinois (*Mead*) and southwestward. July — Sept. — Flowers 3" long, purple. Pod 1' long, pubescent.

26. CENTROSEMA, DC. SPURRED BUTTERFLY-PEA.

Calyx short, 5-cleft. Corolla, &c. much as in *Clitoria*, but the spreading standard with a spur-shaped projection on the back near the base: keel broad. Style bearded at the apex around the terminal stigma. Pod long and linear, flat, pointed with the awl-shaped style, many-seeded, thickened at the edges, the valves marked with a raised line on each side next the margin. — Twining perennials, with 3-foliolate stipellate leaves, and large showy flowers. Stipules, bracts, and bractlets striate, the latter longer than the calyx. (Name from *κέντρον*, a spur, and *σῆμα*, the standard.)

1. **C. Virginianum**, Benth. Rather rough with minute hairs; leaflets varying from oblong-ovate to lanceolate and linear, very veiny, shining; peduncles 1 - 4-flowered; calyx-teeth linear-awl-shaped. — Sandy woods, from Maryland southward. July. — Corolla 1' long, violet. Pods straight, 4' - 5' long.

27. CLITÒRIA, L. BUTTERFLY-PEA.

Calyx tubular, 5-toothed. Standard much larger than the rest of the flower, erect, rounded, notched at the top, not spurred on the back: keel small, shorter than the wings, incurved, acute. Stamens monadelphous below. Style bearded down the inner face. Pod linear-oblong, flattish, knotty, several-seeded, pointed with the base of the style. — Erect or twining perennials, with mostly pinnately 3-foliolate stipellate leaves, and very large flowers. Peduncles 1 - 3-flowered: bractlets opposite, striate. (Derivation *recondite*.)

1. **C. Mariàna**, L. Smooth; leaflets oblong-ovate or ovate-lanceolate; stipules and bracts awl-shaped; peduncles short, 1 - 3-flowered. — Dry banks, E. New York to Virginia and southward. July. — Low, ascending or twining; the showy pale-blue flowers 2' long.

28. AMPHICARPÆA, Ell. HOG PEA-NUT.

Flowers of 2 kinds; those of the racemes from the upper branches perfect, but seldom ripening fruit; those near the base and on creeping branches with the corolla none or rudimentary, and few free stamens, but fruitful. Calyx about equally 4- (rarely 5-) toothed: bractlets none or minute. Keel and wing-petals similar, almost straight; the standard partly folded round them. Stamens diadelphous. Style beardless. Pods of the upper flowers, when formed, somewhat scymetar-shaped, 3 - 4-seeded; of the lower ones commonly subterranean, obovate or pear-shaped, fleshy, ripening usually but one large seed. — Low and slender perennials; the twining stems clothed with brownish hairs. Leaves pinnately 3-foliolate: leaflets rhombic-ovate, stipellate. Flowers small, in clus-

tered or compound racemes, purplish. Bracts persistent, round, partly clasping, striate, as well as the stipules. (Name from *ἀμφί*, *both*, and *καρπός*, *fruit*, in allusion to the two kinds of pods.)

1. **A. monóica**, Nutt. Racemes nodding; bracts each supporting 2 or more flowers, shorter than the pedicels; subterranean pods hairy. — Rich woodlands. Aug., Sept.

29. GALÁCTIA, P. Browne. MILK-PEA.

Calyx 4-cleft; the lobes acute, the upper one broadest, entire. Keel scarcely incurved. Stamens diadelphous or nearly so. Style beardless. Pod linear, flat, several-seeded (some few of them rarely partly subterranean and fleshy or deformed). — Low, mostly prostrate or twining perennial herbs. Leaflets usually 3, stipellate. Flowers in somewhat interrupted or knotty racemes, purplish; in summer. (Name from *γάλα*, *-aktos*, *milk*; some species being said to yield a milky juice, which is unlikely.)

1. **G. glabélla**, Michx. *Stems nearly smooth*, prostrate; leaflets elliptical or ovate-oblong, sometimes slightly hairy beneath; racemes short, 4–8-flowered; *Pods somewhat hairy*. — Sandy woods, S. New York, New Jersey and Penn. to Virginia and southward. — Flowers large for the genus, rose-purple.

2. **G. móllis**, Michx. *Stems* (decumbent and somewhat twining) and *leaves beneath soft-downy and hoary*; leaflets oval; racemes many-flowered; *Pods very downy*. — S. Pennsylvania, Maryland, and southward. July.

30. RHYNCHÓSIA, Lour., DC. RHYNCHOSIA.

Calyx somewhat 2-lipped, or deeply 4–5-parted. Keel scythe-shaped, or incurved at the apex. Stamens diadelphous. Ovules only 2. Pod 1–2-seeded, short and flat, 2-valved. — Usually twining or trailing perennial herbs, pinnately 3-foliolate, or with a single leaflet, not stipellate. Flowers yellow, racemose or clustered. (Name from *ῥύγχος*, *a beak*, from the shape of the keel.)

1. **R. tomentósa**, Torr. & Gray. More or less downy; leaflets roundish; racemes short or capitate; calyx about as long as the corolla, 4-parted, the upper lobe 2-cleft; pod oblong. — Very variable: or perhaps the following are distinct species.

Var. **monophýlla**, Torr. & Gray. Dwarf and upright (3'–6' high); leaves mostly of one round leaflet 1'–2' wide. (*R. renifórmis*, DC.) — Virginia and southward, in dry sandy soil. — Flowers earlier than the following.

Var. **volúbilis**, Torr. & Gray. Trailing and twining, less downy; leaflets 3, roundish; racemes few-flowered, almost sessile in the axils. (*R. diffórmis*, DC.) — S. Virginia and southward.

Var. **erécta**, Torr. & Gray. Upright (1°–2° high), soft-downy; leaflets 3, oval or oblong. (*R. erecta*, DC.) — Maryland and southward.

31. BAPTÍSIA, Vent. FALSE INDIGO.

Calyx 4–5-toothed. Standard not longer than the wings, its sides reflexed: keel-petals nearly separate, and, like the wings, straight. Stamens 10, distinct.

Pod stalked in the persistent calyx, roundish or oblong, inflated, pointed, many-seeded. — Perennial herbs, with palmately 3-foliolate (rarely simple) leaves, which generally blacken in drying, and racemed flowers. (Named from βαρρικόζω, *to dye*, from the economical use of some species, which yield a poor indigo.)

1. **B. tinctoria**, R. Br. (WILD INDIGO.) Smooth and slender (2°–3° high), rather glaucous; leaves almost sessile; leaflets rounded, wedge-obovate ($\frac{3}{4}$ ' long); *stipules and bracts minute and deciduous*; *racemes few-flowered*, terminating the bushy branches; pods oval-globose, on a stalk longer than the calyx. — Sandy dry soil: common. June–Aug. — Corolla yellow, $\frac{1}{2}$ ' long.

2. **B. australis**, R. Br. (BLUE FALSE-INDIGO.) Smooth, tall and stout (4°–5°); leaflets oblong-wedge-form, obtuse; *stipules lanceolate, as long as the petioles, rather persistent*; *raceme elongated* (1°–2°) and many-flowered, erect; bracts deciduous; *stalk of the oval-oblong pods about the length of the calyx*. — Alluvial soil, from Pennsylvania westward and southward: often cultivated. June. — Flowers 1' long, indigo-blue. Pods 2'–3' long.

3. **B. leucantha**, Torr. & Gr. Smooth; stems, leaves, and racemes as in the foregoing; *stipules early deciduous*; *Pods oval-oblong, raised on a stalk fully twice the length of the calyx*. — Alluvial soil, Ohio to Wisconsin and southwestward. July. — Flowers white; the standard short. Pods 2' long.

4. **B. alba**, R. Br. Smooth (1°–3° high); *the branches slender and widely spreading*; *petioles slender*; *stipules and bracts minute and deciduous*; leaflets oblong or oblanceolate; racemes slender on a long naked peduncle; *Pods linear-oblong* (1'–1 $\frac{1}{2}$ ' long), *short-stalked*. — Dry soil, Virginia and southward. May, June. — Flowers white, 6''–9'' long.

5. **B. leucophæa**, Nutt. *Hairy, low* (1° high), with *divergent branches*; *leaves almost sessile*; leaflets narrowly oblong-obovate or spatulate; *stipules and bracts large and leafy, persistent*; *racemes long, reclined*; *flowers on elongated pedicels*; pods ovoid, hoary. — Michigan to Wisconsin and southward. April, May. — Raceme often 1°, pedicels 1'–2', the cream-colored corolla 1', in length.

32. CLADRÁSTIS, Raf. YELLOW-WOOD.

Calyx 5-toothed. Standard large, roundish, reflexed: the distinct keel-petals and wings straight, oblong. Stamens 10, distinct: filaments slender, incurved above. Pod short-stalked above the calyx, linear, flat, thin, marginless, 4–6-seeded, at length 2-valved. — A small and handsome tree, with yellow wood, smooth bark, nearly smooth pinnate leaves of 7–11 oval or ovate leaflets, and ample paniced racemes (10''–20'' long) of showy white flowers drooping from the end of the branches. Stipules obsolete. Base of the petioles hollow, and enclosing the leaf-buds of the next year. Bracts minute and fugacious. (Name of obscure derivation.)

1. **C. tinctoria**, Raf. (*Virgilia lutea, Michx. f.*) Rich hillsides, E. Kentucky and southward along the western base of the Alleghanies. May, June.

33. CÉRCIS, L. RED-BUD. JUDAS-TREE.

Calyx 5-toothed. Corolla imperfectly papilionaceous: standard smaller than the wings, and enclosed by them in the bud: the keel-petals larger and not

united. Stamens 10, distinct, declined. Pod oblong, flat, many-seeded, the upper suture with a winged margin. Embryo straight. — Trees, with rounded-heart-shaped simple leaves, caducous stipules, and red-purple flowers in umbel-like clusters along the branches of the last or preceding years, appearing before the leaves, acid to the taste. (The ancient name of the Oriental *Judas-tree*.)

1. **C. Canadensis**, L. (RED-BUD.) Leaves pointed; pods nearly sessile; above the calyx. — Rich soil, New York to Ohio, Illinois, and southward. March–May. — A small ornamental tree, often cultivated: the blossoms smaller than in the European species.

34. CASSIA, L. SENNA.

Sepals 5, scarcely united. Petals 5, unequal, not papilionaceous, spreading. Stamens 5–10, unequal, and some of them often imperfect, spreading: anthers opening by 2 pores or chinks at the apex. Pod many-seeded, often with cross partitions. — Herbs (in the United States), with simply and abruptly pinnate leaves, and mostly yellow flowers. (An ancient name of obscure derivation.)

* *Leaflets large: stipules deciduous: the 3 upper anthers deformed and imperfect: flowers in short axillary racemes, the upper ones panicled: herbage glabrous.*

1. **C. Marilandica**, L. (WILD SENNA.) *Leaflets 6–9 pairs, lanceolate-oblong, obtuse; petiole with a club-shaped gland near the base; pods linear, slightly curved, flat, at first hairy (2'–4'); root perennial. — Alluvial soil. July. — Stem 3°–4° high. Leaves used as a substitute for the officinal Senna.*

2. **C. OCCIDENTALIS**, L. *Leaflets 4–6 pairs, ovate-lanceolate, acute; an ovate gland at the base of the petiole; pods long-linear (5' long) with a tumid border, glabrous. — Virginia and southward. Aug. (Adv. from Trop. Amer.)*

3. **C. obtusifolia**, L. *Leaflets 3 or rarely 2 pairs, obovate, obtuse, with an elongated gland between those of the lower pairs or lowest pair; pods slender, 6' long, curved; root annual. — Banks of the Ohio River, Illinois (Dr. Vasey), S. Virginia and southward.*

* * *Leaflets small, somewhat sensitive to the touch: stipules striate, persistent: a cup-shaped gland beneath the lowest pair of leaflets: anthers all perfect: flowers in small clusters above the axils: pods flat: root annual.*

4. **C. Chamæcrista**, L. (PARTRIDGE PEA.) *Leaflets 10–15 pairs, linear-oblong, oblique at the base; flowers (large) on slender pedicels; anthers 10, elongated, unequal (4 of them yellow, the others purple) style slender. — Sandy fields: common, especially southward. Aug. — Stems spreading, 1° long: 2 or 3 of the showy yellow petals often with a purple spot at the base.*

5. **C. nictitans**, L. (WILD SENSITIVE-PLANT.) *Leaflets 10–20 pairs, oblong-linear; flowers (very small) on very short pedicels; anthers 5, nearly equal; style short. — Sandy fields, New England, near the coast, to Illinois and southward. Aug.–Sept.*

35. GYMNOCLADUS, Lam. KENTUCKY COFFEE-TREE.

Flowers diœcious or polygamous, regular. Calyx elongated-tubular below, 5-cleft. Petals 5, oblong, equal, inserted on the summit of the calyx-tube. Sta-

mens 10, distinct, short, inserted with the petals. Pod oblong, flattened, hard, pulpy inside, several-seeded. Seeds flattish. — A tall large tree, with rough bark, stout branchlets, not thorny, and large unequally twice-pinnate leaves; the leaflets standing vertically. — Flowers whitish, in axillary racemes. (Name from *γυμνός*, *naked*, and *κλάδος*, *a branch*, alluding to the stout branches destitute of spray.)

1. **G. Canadensis**, Lam. Rich woods, along rivers, W. New York and Penn. to Illinois and southwestward. June. — Cultivated as an ornamental tree: timber valuable. Leaves 2°–3° long, with several large partial leafstalks bearing 7–13 ovate stalked leaflets, the lowest pair with single leaflets. Pod 6'–10' long, 2' broad; the seeds over $\frac{1}{2}$ ' across.

36. GLEDÍTSCHIA, L. HONEY-LOCUST.

Flowers polygamous. Calyx short, 3–5-cleft, the lobes spreading. Petals as many as the sepals and equalling them, the 2 lower sometimes united. Stamens 3–10, distinct, inserted with the petals on the base of the calyx. Pod flat, 1–many-seeded. Seeds flat. — Thorny trees, with abruptly once or twice pinnate leaves, and inconspicuous greenish flowers in small spikes. Thorns above the axils. (Named in honor of *J. G. Gleditsch*, a botanist contemporary with Linnæus.)

1. **G. triacanthos**, L. (THREE-THORNED ACACIA, OR HONEY-LOCUST.) Thorns stout, often triple or compound; *leaflets lanceolate-oblong*, somewhat serrate; *Pods linear, elongated* (1°–1½° long), often twisted, filled with sweet pulp between the seeds. — Rich woods, Penn. to Virginia, Illinois, and southwestward. June. — Common in cultivation as an ornamental tree, and for hedges.

2. **G. monosperma**, Walt. (WATER-LOCUST.) Thorns slender, mostly simple; *leaflets ovate or oblong*; *Pods oval, 1-seeded*, pulpless. — Swamps, Illinois and southwestward. July. — A small tree.

37. DESMANTHUS, Willd. DESMANTHUS.

Flowers perfect or polygamous, regular. Calyx campanulate, 5-toothed. Petals 5, distinct. Stamens 5 or 10. Pod flat, membranaceous or somewhat coriaceous, several-seeded, 2-valved, smooth. — Herbs, with twice-pinnate leaves of numerous small leaflets, and with one or more glands on the petiole, setaceous stipules, and axillary peduncles bearing a head of small greenish-white flowers. (Name composed of *δέσμα*, *a bond*, and *άνθος*, *flower*.)

1. **D. brachylobus**, Benth. Nearly glabrous perennial, erect (1°–4° high); partial petioles 6–15 pairs; leaflets 20–30 pairs; stamens 5; pods oblong or lanceolate, curved, scarcely 1' long, 2–6-seeded. (*Darlingtonia brachyloba* & *glandulosa*, DC.) — Prairies and alluvial banks, Illinois and southwestward.

38. SCHRÁNKIA, Willd. SENSITIVE BRIAR.

Flowers polygamous, regular. Calyx minute, 5-toothed. Petals united into a funnel-form 5-cleft corolla. Stamens 10–12, distinct, or the filaments united at the base. Pods long and narrow, rough-prickly, several-seeded, 4-valved, i. e.

the two narrow valves separating on each side from a thickened margin. — Perennial herbs, nearly related to the true Sensitive Plants (*Mimosa*); the procumbent stems and petioles prickly, with twice-pinnate sensitive leaves of many small leaflets, and axillary peduncles bearing round heads of small rose-colored flowers. (Named for *F. P. Schrank*, a German botanist.)

1. **S. uncinata**, Willd. Prickles hooked; partial petioles 4–6 pairs; leaflets elliptical, reticulated with strong veins beneath; pods oblong-linear, nearly terete, short-pointed, densely prickly (2' long). — Dry sandy soil, Virginia, Illinois? and southward. June–Aug.

2. **S. angustata**, Torr. & Gray. Leaflets oblong-linear, scarcely veined; pods slender, taper-pointed, sparingly prickly (about 4' long). — With the preceding.

ORDER 33. ROSACEÆ. (ROSE FAMILY.)

Plants with regular flowers, numerous (rarely few) distinct stamens inserted on the calyx, and 1–many pistils, which are quite distinct, or (in the Pear tribe) united and combined with the calyx-tube. Seeds (anatropous) 1–few in each ovary, almost always without albumen. Embryo straight, with large and thick cotyledons. Leaves alternate, with stipules, these sometimes caducous, rarely obsolete or wanting. — Calyx of 5 or rarely 3–4–8 sepals (the odd one superior), united at the base, often appearing double by a row of bractlets outside. Petals as many as the sepals (rarely wanting), mostly imbricated in the bud, and inserted with the stamens on the edge of a disk that lines the calyx-tube. Trees, shrubs, or herbs. — A large and important order, almost destitute of noxious qualities, and producing the most valuable fruits. Very intimately connected with Leguminosæ on one hand, and with Saxifragaceæ on the other.

SUBORDER I. AMYGDALÆÆ. (ALMOND FAMILY.)

Calyx entirely free from the (usually) solitary ovary, deciduous. Style terminal or nearly so. Fruit a drupe (stone-fruit), 1-seeded, or rarely 2-seeded. — Trees or shrubs, with simple leaves, the bark exuding gum, and the bark, leaves, and kernels yielding the peculiar flavor of prussic acid. Stipules free, often deciduous.

1. **Prunus**. Flowers perfect. Lobes of the calyx and petals 5. Stone of the drupe bony.

SUBORDER II. ROSACEÆ PROPER.

Calyx free from the ovaries (but sometimes enclosing them in its tube), mostly persistent with the fruit. Pistils few or many, distinct, occasionally single. Stipules commonly united with the petiole.

Tribe I. SPIRÆÆÆ. Pistils mostly 5 and forming follicles in fruit: styles terminal.

Calyx without bractlets. Seeds not rarely with some albumen!

2. **Spiræa**. Calyx 5-cleft, short. Petals obovate, equal, chiefly imbricated in the bud.

3. **Gillenia**. Calyx elongated, 5-toothed. Petals slender, unequal, convolute in the bud.

Tribe II. POTERIEÆ. Pistils 1-4, one-ovuled, becoming achenia, and enclosed in the urn-shaped tube of the dry persistent calyx, which is constricted or nearly closed at the throat. Petals often wanting.

4. **Poterium.** Petals none. Lobes of the calyx 4, petal-like. Style terminal: stigma tufted.
5. **Alchemilla.** Petals none. Stamens and pistils 1-4: style lateral.
6. **Agrimonia.** Petals 5. Stamens 12-15. Pistils 2: style terminal.

Tribe III. DRYADEÆ. Pistils numerous, rarely few or single, one-ovuled, becoming dry achenia; the calyx open, not fleshy in fruit. Petals present, usually conspicuous.

7. **Dryas.** Petals and calyx-lobes 8 or 9. Stamens and carpels numerous: persistent styles becoming long plumose tails in fruit.
8. **Geum.** Petals and calyx-lobes 5, the latter usually with 5 alternating small bractlets. Stamens and carpels numerous: persistent styles becoming long plumose or hairy, or naked and straight or jointed, tails. Radicle inferior.
9. **Waldsteinia.** Petals and calyx-lobes 5; no bractlets. Stamens numerous. Achenia 2-6: styles deciduous from the base. Radicle inferior.
10. **Sibbaldia.** Petals minute: stamens and achenia 5-10: otherwise same as *Potentilla*.
11. **Potentilla.** Petals 5 (rarely 4), conspicuous. Calyx-lobes as many, and also with an alternating set of accessory lobes or bractlets. Stamens and achenia numerous; the latter heaped on a dry receptacle. Styles commonly more or less lateral, deciduous or not enlarging in fruit. Radicle superior.
12. **Fragaria.** Flower as in *Potentilla*. Receptacle much enlarged and pulpy in fruit.

Tribe IV. RUBEÆ. Pistils numerous or several, 2-ovuled, becoming berry-like or drupelets in fruit; the 5-cleft calyx not bracteolate, open, persistent or withering beneath the fruit. Petals conspicuous.

13. **Dalibarda.** Carpels 5-10, in the bottom of the calyx, almost dry.
14. **Rubus.** Carpels numerous, heaped on the receptacle.

Tribe V. ROSEÆ. Pistils numerous, one-ovuled, becoming achenia, contained in the urn-shaped or globular and almost closed fleshy tube of the calyx, or hip: no bractlets. Petals conspicuous.

15. **Rosa.** Character of the Tribe.

SUBORDER III. POMEÆ. (PEAR FAMILY.)

Calyx-tube thick and fleshy in fruit, including and combined with the 2-5 ovaries (forming a *pome*). Stipules free.

* Cells of the compound ovary as many as the styles (2-5), each 2- (rarely several-) ovuled.

16. **Cratægus.** Pome drupe-like, with 1-5 bony stones or kernels. Usually thorny.
17. **Pyrus.** Pome containing 2-5 papery or cartilaginous carpels.

* Cells of the compound ovary becoming twice as many as the styles, each 1-ovuled.

18. **Amelanchier.** Pome usually of 5 carpels; each becomes incompletely 2-celled by a projection from its back: otherwise as *Pyrus*.

1. PRÛNUS, Tourn. PLUM, CHERRY, &c.

Calyx 5-cleft; the tube bell-shaped, urn-shaped or tubular-obconical, deciduous after flowering. Petals 5, spreading. Stamens 15-20. Ovary solitary, with 2 pendulous ovules. Drupe fleshy, with a bony stone.—Small trees or shrubs, with mostly edible fruit. (The ancient Latin name.)

- § 1. PRUNUS & CÉRASUS, Tourn. *Drupe smooth, and the stone smooth or somewhat rugged: flowers (usually white) from separate lateral scaly buds in early spring, preceding or coetaneous with the leaves; the pedicels few or several in simple umbel-like clusters.*—The PLUMS of the Old World have the leaves

convolute in the bud, the fruit with a bloom; its stone oblong, flattened or flattish and acute at both ends; but our wild Plums are like CHERRIES in having the leaves folded before expansion, little or no bloom, and some of them in the thicker or globular stone, thus confounding the distinctions.

1. **P. Americana**, Marshall. (WILD YELLOW OR RED PLUM.) *Leaves ovate or somewhat obovate, conspicuously pointed, coarsely or doubly serrate, very veiny, glabrous when mature; fruit nearly destitute of bloom, roundish-oval, yellow, orange, or red, $\frac{1}{2}$ ' - $\frac{3}{8}$ ' in diameter, with the turgid stone more or less acute on both margins, or in cultivated states 1' or more in diameter, the flattened stone with broader margins: pleasant-tasted, but with a tough and acerb skin.* — Woodlands and river-banks: common. — Tree thorny, 8° - 20° high.

2. **P. maritima**, Wang. (BEACH PLUM.) *Low and straggling (2° - 5°); leaves ovate or oval, finely serrate, softly pubescent underneath; pedicels short, pubescent; fruit globular, purple or crimson with a bloom ($\frac{1}{2}$ ' - 1' in diameter); the stone very turgid, acute on one edge, rounded and minutely grooved on the other. (P. littoralis, Bigelow.) — Varies, when at some distance from the coast, with the leaves smoother and thinner, and the fruit smaller. (P. pygmæa, Willd.) — Sea-beach and the vicinity, Maine to Virginia; the variety, New Jersey and southward.*

3. **P. Chicasa**, Michx. (CHICKASAW PLUM.) *Stem scarcely thorny (8' - 15' high); leaves nearly lanceolate, finely serrulate, glabrous; fruit globular, red, nearly destitute of bloom ($\frac{1}{2}$ ' - $\frac{3}{8}$ ' in diameter); the ovoid stone almost as thick as wide, rounded at both sutures, one of them minutely grooved.* — Maryland to Illinois (probably not indigenous) and southwestward.

4. **P. spinosa**, L. (SLOE. BLACK THORN.) *Branches thorny; leaves obovate-oblong or ovate-lanceolate, sharply serrate, at length glabrous; pedicels glabrous; fruit small, globular, black with a bloom, the stone turgid, acute on one edge.* — Var. **INSITITIA** (BULLACE-PLUM), is less spiny, the pedicels and lower side of the leaves pubescent. (P. insititia, L.) — Roadsides and waste places, New England, to Penn., &c. (Adv. from Eu.)

5. **P. pumila**, L. (DWARF CHERRY.) *Smooth, depressed, and trailing (6' - 18' high); leaves obovate-lanceolate, tapering to the base, somewhat toothed near the apex, pale underneath; flowers 2 - 4 together; fruit ovoid, dark red, without bloom; stone ovoid, marginless, of the size of a large pea.* — Rocks or sandy banks, Massachusetts northward to Wisconsin, and south to Virginia along the mountains.

6. **P. Pennsylvanica**, L. (WILD RED CHERRY.) *Leaves oblong-lanceolate, pointed, finely and sharply serrate, shining, green and smooth both sides; flowers many in a cluster, on long pedicels; fruit globose, light red, very small, with thin and sour flesh; stone globular.* — Rocky woods: common, especially northward. May. — Tree 20° - 30° high, with light red-brown bark.

§ 2. **PADUS**, Mill. (CHERRY.) *Drupe small, globose, without bloom; the stone turgid-ovate, marginless: flowers in racemes terminating leafy branches, therefore appearing after the leaves, late in spring.*

7. **P. Virginiana**, L. (CHOKE-CHERRY.) *Leaves oval, oblong, or obovate, abruptly pointed, very sharply (often doubly) serrate with slender teeth, thin; racemes*

short and close; petals roundish; fruit red turning to dark crimson; stone smooth. — River-banks: common, especially northward. May. — A tall shrub, seldom a tree, with grayish bark; the fruit very austere and astringent till perfectly ripe. (*P. obovata*, *Bigelow*. *P. serotina*, of several authors.)

8. **P. serotina**, Ehrhart. (WILD BLACK CHERRY.) *Leaves oblong or lanceolate-oblong, taper-pointed, serrate with incurved short and callous teeth*, thickish, shining above; racemes elongated; petals obovate; fruit purplish-black. — Woods: common. June. — A fine large tree, with reddish-brown branches, furnishing valuable timber to the cabinet-maker: also abounding eastward as a shrub. Fruit slightly bitter, but with a pleasant vinous flavor.

2. SPIRÆA, L. MEADOW-SWEET.

Calyx 5-cleft, short, persistent. Petals 5, obovate, equal, imbricated in the bud (except in No. 6). Stamens 10–50. Pods (follicles) 2–12, several- (2–15-) seeded. — Flowers white or rose-color, sometimes diœcious: rarely the parts are 4 instead of 5. (Name probably from *σπειράω*, to wind, alluding to the fitness of the plants to be formed into garlands.)

§ 1. **PHYSOCÁRPOS**, Camb. (*NEILLIA*, Don.) *Shrubs with simple palmately-lobed leaves and umbel-like corymbs: pods inflated and diverging when grown, 2–4-seeded: seeds pretty large, roundish, bitter, with a thick crustaceous seed-coat, and rather copious albumen!*

1. **S. opulifolia**, L. (NINE-BARK.) *Leaves roundish, somewhat 3-lobed and heart-shaped; pods 2–5.* — Rocky river-banks, especially westward. June. — Shrub 4°–10° high, with long recurved branches, and white flowers, succeeded by membranaceous purplish pods: the old bark loose and separating in numerous thin layers.

§ 2. **SPIRÆA** proper. *Shrubs, with simple leaves; the stipules obsolete: pods (mostly 5) not inflated, several-seeded: seeds mostly linear and with a thin or loose coat and no albumen, in this and the following sections.*

2. **S. corymbosa**, Raf. *Nearly smooth (1°–2° high); leaves oval or ovate, cut-toothed towards the apex; corymbs large, flat, several times compound; flowers white.* — Alleghanies of Pennsylvania to Virginia and Kentucky. Also Morris Co., New Jersey, *C. F. Austin*. June. — A form of *S. betulæfolia*, *Pallas*.

3. **S. salicifolia**, L. (COMMON MEADOW-SWEET.) *Nearly smooth (2°–3° high); leaves wedge-lanceolate, simply or doubly serrate; flowers in a crowded panicle, white or flesh-color; pods smooth.* — Wet or low grounds: also cultivated. July. (Eu.)

4. **S. tomentosa**, L. (HARDHACK. STEEPLE-BUSH.) *Stems and lower surface of the ovate or oblong serrate leaves very woolly; flowers in short racemes crowded in a dense panicle; pods woolly.* — Low grounds: commonest in New England. July. — Flowers rose-color, rarely white.

§ 3. **ULMÁRIA**, Mœnch. *Perennial herbs, with pinnate leaves and panicled cymose flowers: calyx reflexed: pods 5–8 in number, 1–2-seeded.*

5. **S. lobata**, Murr. (QUEEN OF THE PRAIRIE.) *Glabrous (2°–8° high); leaves interruptedly pinnate; the terminal leaflet very large, 7–9-parted, the lobes incised and toothed; stipules kidney-form; panicle compound-clustered,*

on a long naked peduncle. — Meadows and Prairies, Penn. to Michigan, Illinois, and Kentucky. June. — Flowers deep peach-blossom color, handsome, the petals and sepals often in fours! The bruised foliage exhales the odor of Sweet Birch.

§ 4. **ARÚNCUS**, Seringe. *Perennial herbs, with diœcious whitish flowers, in many slender spikes disposed in a long compound panicle: leaves thrice-pinnate: stipules obsolete: pods 3-5, several-seeded: pedicels reflexed in fruit.*

6. **S. Arúncus**, L. (GOAT'S-BEARD.) Smooth, tall; leaflets thin, lanceolate-oblong, or the terminal ones ovate-lanceolate, taper-pointed, sharply cut and serrate. — Rich woods, Catskill and Alleghany Mountains and westward. June. (Eu.)

3. GILLÈNIA, Mœnch. INDIAN PHYSIC.

Calyx narrow, somewhat constricted at the throat, 5-toothed; teeth erect. Petals 5, rather unequal, linear-lanceolate, inserted in the throat of the calyx; convolute in the bud. Stamens 10-20, included. Pods 5, included, at first lightly cohering with each other, 2-4-seeded. Seeds ascending, with a close coriaceous coat, and some albumen. — Perennial herbs, with almost sessile 3-foliolate leaves; the thin leaflets doubly serrate and incised. Flowers loosely paniculate-corymbed, pale rose-color or white. (Dedicated to an obscure German botanist or gardener, *A. Gille*, or *Gillenius*.)

1. **G. trifoliata**, Mœnch. (BOWMAN'S ROOT.) Leaflets ovate-oblong, pointed, cut-serrate; stipules small, awl-shaped, entire. — Rich woods, from W. New York southward, and sparingly in the Western States. July.

2. **G. stipulacea**, Nutt. (AMERICAN IPECAC.) Leaflets lanceolate, deeply incised; stipules large and leaf-like, doubly incised. — From W. Pennsylvania and New York to Illinois and Kentucky. June.

4. POTÈRIUM, L. (including SANGUISORBA.) BURNET.

Calyx with a top-shaped tube, constricted at the throat, persistent; the 4 broad petal-like spreading lobes imbricated in the bud, deciduous. Petals none. Stamens 4-12 or more, with flaccid filaments and short anthers. Pistils 1-3: the slender terminal style tipped with a tufted or brush-like stigma. Achenium (commonly solitary) enclosed in the 4-angled dry and thickish closed calyx-tube. Seed suspended. — Chiefly perennial herbs, with unequally pinnate leaves, stipules coherent with the petiole, and small, often polygamous or diœcious flowers crowded in a dense head or spike at the summit of a long and naked peduncle, each bracteate and 2-bracteolate. (Name *ποτήριον*, a *drinking-cup*, the foliage of Burnet having been used in the preparation of some medicinal drink.)

1. **P. Canadense**. (CANADIAN BURNET.) Stamens 4, long-exserted, club-shaped, white, as is the whole of the elongated and cylindrical spike; stem 3°-6° high, leaflets numerous, ovate or oblong-lanceolate, coarsely serrate, obtuse, heart-shaped at the base, as if stipellate; stipules serrate. (*Sanguisorba Canadensis*, L., and former edition.) — Bogs and wet meadows; chiefly northward. Aug. - Oct.

5. **ALCHEMÍLLA**, Tourn. LADY'S MANTLE.

Calyx-tube inversely conical, contracted at the throat; limb 4-parted with as many alternate accessory lobes outside. Petals none. Stamens 1-4. Pistils 1-4; the slender style arising from near the base of the ovary; the achenia included in the tube of the persistent calyx. — Low herbs, with palmately lobed or compound leaves, and small corymbed greenish flowers. (From *Alkemelyeh*, the Arabic name.)

1. **A. ARVÉNSIS**, L. (PARSLEY PIERT.) Small annual (3'-8' high), leafy; leaves 3-parted, with the wedge-shaped lobes 2-3-cleft, pubescent; flowers sessile in the axils. — Eastern Virginia. (Adv. from Eu.)

A. ALPÍNA, L., is said by Pursh to grow on the Green and White Mountains, New England: but there is most probably some mistake about it.

6. **AGRIMÒNIA**, Tourn. AGRIMONY.

Calyx-tube top-shaped, contracted at the throat, beset with hooked bristles above, indurated in fruit and enclosing the 2 achenia; the limb 5-cleft, closed after flowering. Petals 5. Stamens 12-15. Styles terminal. Seed suspended. — Perennial herbs, with interruptedly pinnate leaves, and yellow flowers in slender spiked racemes: bracts 3-cleft. (Name a corruption of *Argemonia*, of the same derivation as *Argemone*, p. 59.)

1. **A. EUPATÒRIA**, L. (COMMON AGRIMONY.) *Leaflets 5-7 with minute ones intermixed, oblong-obovate, coarsely toothed; petals twice the length of the calyx.* — Borders of woods: common. July-Sept. (Eu.)

2. **A. PARVIFLÒRA**, Ait. (SMALL-FLOWERED A.) *Leaflets crowded, 11-19, with smaller ones intermixed, lanceolate, acute, deeply and regularly cut-serrate, as well as the stipules; petals small.* — Woods and glades, S. New York (*C. F. Austin*) to Virginia, Kentucky, and southward. July.

7. **DRÝAS**, L. DRYAS.

Calyx open, flattish, 8-9-parted. Petals 8-9, large. Otherwise like *Geum* § *Sieversia*. — Dwarf and matted slightly shrubby plants, with simple toothed leaves, and solitary large flowers. (Name from *Dryades*, the nymphs of the Oaks, the foliage of the original species resembling oak-leaves in miniature.)

1. **D. INTEGRIFÒLIA**, Vahl. Leaves oblong-ovate, slightly heart-shaped, with revolute margins, nearly entire, white-downy beneath, flowers white. — White Mountains, New Hampshire, *Prof. Peck*, according to *Pursh*; not since met with: but it grows in Lower Canada. (Eu.)

8. **GÈUM**, L. AVENS.

Calyx bell-shaped or flattish, deeply 5-cleft, usually with 5 small bractlets at the sinuses. Petals 5. Stamens many. Achenia numerous, heaped on a conical or cylindrical dry receptacle, the long persistent styles forming hairy or naked and straight or jointed tails. Seed erect. — Perennial herbs, with pinnate or lyrate leaves. (Name from *γεύω*, to give an agreeable flavor, the roots being rather aromatic.)

§ 1. **GEUM** proper. *Styles jointed and bent near the middle; the lower portion persistent, naked, hooked at the end after the deflexed and mostly somewhat hairy upper joint falls away: head of fruit sessile in the calyx: calyx-lobes reflexed. (Flowers somewhat panicked at the summit of a leafy stem: achenia in our species glabrous or nearly so below, more or less bristly at the top or along the base of the style.)*

* *Petals white or pale greenish-yellow, small, spatulate or oblong: stipules small.*

1. **G. álbum**, Gmelin. *Smoothish or softly pubescent; stem slender (2° high); root-leaves of 3-5 leaflets, or simple and rounded, with a few minute leaflets on the petiole below; those of the stem 3-divided or lobed, or only toothed; petals about the length of the calyx; receptacle of the fruit densely bristly-hirsute. — Borders of woods, &c.: common. May-Aug. — Too near the European G. urbanum; probably a white or whitish-flowered form of it.*

2. **G. Virginianum**, L. *Bristly-hairy, especially the stout stem; lower and root-leaves pinnate, very various, the upper mostly 3-parted or divided, incised; petals inconspicuous, shorter than the calyx; receptacle of the fruit glabrous or nearly so. — Borders of woods and low grounds: common. June-Aug. — Heads of fruit larger than in the preceding, on stouter hirsute peduncles.*

** *Petals golden-yellow, conspicuous, broadly obovate, exceeding the calyx: stipules larger and all deeply cut.*

3. **G. macrophýllum**, Willd. *Bristly-hairy, stout (1°-3° high); root-leaves lyrate and interruptedly pinnate, with the terminal leaflet very large and round-heart-shaped; lateral leaflets of the stem-leaves 2-4, minute, the terminal roundish, 3-cleft, the lobes wedge-form and rounded; receptacle of the fruit nearly naked. — Around the base of the White Mountains, New Hampshire, also Northern Michigan, Illinois, and northwestward. June. (Eu.)*

4. **G. strictum**, Ait. *Somewhat hairy (3°-5° high); root-leaves interruptedly pinnate, the leaflets wedge-obovate; leaflets of the stem-leaves 3-5, rhombic-ovate or oblong, acute; receptacle of fruit downy. — Moist meadows: common, especially northward. July, Aug. (Eu.)*

§ 3. **STÝLIPUS**, Raf. *Styles smooth: head of fruit conspicuously stalked in the calyx: bractlets of the calyx none: otherwise nearly as § 1.*

5. **G. vérnum**, Torr. & Gr. *Somewhat pubescent; stems ascending, few-leaved, slender; root-leaves roundish-heart-shaped, 3-5-lobed, or some of them pinnate, with the lobes cut; petals yellow, about the length of the calyx; receptacle smooth. (Stylipus vernus, Raf.) — Thickets, Ohio to Illinois and Kentucky. April-June.*

§ 3. **CARYOPHYLLÀTA**, Tourn. *Style jointed and bent in the middle, the upper joint plumose: flowers large: calyx erect or spreading: petals erect.*

6. **G. rivale**, L. (WATER, OR PURPLE AVENS.) *Stems nearly simple, several-flowered (2° high); root-leaves lyrate and interruptedly pinnate; those of the stem few, 3-foliolate or 3-lobed; petals dilated-obovate, retuse, contracted into a claw, purplish-orange; head of fruit stalked in the calyx. — Bogs and wet meadows, New England to Penn., Wisconsin, and northward. May. — Blossoms nodding, but the feathery fruiting heads upright. Calyx brown-purple. (Eu.)*

§ 4. **SIEVÉRSIA**, Willd. *Style not jointed, wholly persistent and straight: head of fruit sessile: flowers large: calyx erect or spreading. (Flowering stems simple, and bearing only bracts or small leaves.)*

7. **G. triflorum**, Pursh. Low, softly-hairy; root-leaves interruptedly pinnate; the leaflets very numerous and crowded, oblong-wedge-form, deeply cut-toothed; flowers 3 or more on long peduncles; bractlets linear, longer than the purple calyx, as long as the oblong purplish erect petals; styles very long (2'), strongly plumose in fruit. — Rocks, N. New Hampshire and N. New York to Wisconsin and westward: rare. April–June.

8. **G. radiatum**, Michx. Hirsutely hairy or smoothish; root-leaves rounded-kidney-shaped, radiate-veined (2'–5' broad), doubly or irregularly cut-toothed and obscurely 5–7-lobed, also a set of minute leaflets down the long petiole; stems (8'–18' high) 1–5-flowered; bractlets minute; petals yellow, round-obovate and more or less obcordate, exceeding the calyx ($\frac{1}{2}$ ' long), spreading; styles naked except the base. (High mountains of Carolina.)

Var. **Péckii**. Nearly glabrous, or the stalks and veins of the leaves sparsely hirsute. (*G. Peckii*, Pursh.) — Alpine tops of the White Mountains of New Hampshire. July–Sept.

9. WALDSTEINIA, Willd. (COMAROPSIS, DC.)

Calyx-tube inversely conical; the limb 5-cleft, with 5 often minute and deciduous bractlets. Petals 5. Stamens many, inserted into the throat of the calyx. Achenia 2–6, minutely hairy; the terminal slender styles deciduous from the base by a joint. Seed erect. — Low perennial herbs, with chiefly radical 3–5-lobed or divided leaves, and small yellow flowers on bracted scapes. (Named in honor of *Francis von Waldstein*, a German botanist.)

1. **W. fragarioides**, Tratt. (BARREN STRAWBERRY.) Low; leaflets 3, broadly wedge-form, cut-toothed; scapes several-flowered; petals longer than the calyx. (*Dalibarda fragarioides*, Michx.) — Wooded hillsides, common northward, and southward along the Alleghanies. June.

10. SIBBÁLDIA, L. SIBBALDIA.

Calyx flattish, 5-cleft, with 5 bractlets. Petals 5, linear-oblong, minute. Stamens 5, inserted alternate with the petals into the margin of the woolly disk which lines the base of the calyx. Achenia 5–10: styles lateral. — Low and depressed mountain perennials, — in fact only reduced *Potentillas*. (Dedicated to *Dr. Robert Sibbald*, Prof. at Edinburgh at the close of the 17th century.)

1. **S. procumbens**, L. Leaflets 3, wedge-shaped, 3-toothed at the apex; petals yellow. — Alpine summits of the White Mountains of New Hampshire, and northward. (Eu.)

11. POTENTÍLLA, L. CINQUE-FOIL. FIVE-FINGER.

Calyx flat, deeply 5-cleft, with as many bractlets at the sinuses, thus appearing 10-cleft. Petals 5, usually roundish. Stamens many. Achenia many, collected in a head on the dry mostly pubescent or hairy receptacle: styles lateral or terminal, deciduous. Radicle superior. — Herbs, or rarely shrubs, with

compound leaves, and solitary or cymose flowers: their parts rarely in fours. (Name a kind of diminutive from *potens*, powerful, alluding to the reputed medicinal power, of which in fact these plants possess very little, being merely mild astringents, like the rest of the tribe.)

§ 1. *Style terminal, or attached above the middle of the ovary: achenia glabrous.*

* *Annuals or biennials: petals pale yellow, small, not exceeding the calyx: receptacle globular, ovoid, or even oblong in fruit.*

1. **P. Norvègica**, L. Hairy, erect, branched above; leaves palmately 3-foliolate; leaflets obovate-oblong, cut-toothed. — Fields: common, especially northward. A homely weed. (Eu.)

2. **P. paradóxa**, Nutt. Somewhat pubescent, spreading or decumbent, branched; leaves pinnate; leaflets 5–9, obovate-oblong, cut-toothed; achenia with a thick appendage at the base. — Banks of the Ohio and Mississippi. Shore of L. Ontario, *J. A. Paine*; probably an immigrant from the West.

* * *Perennial herbs: petals yellow, commonly longer than the calyx.*

+ *Low: leaves palmate, of 3 or 5 leaflets.*

3. **P. frígida**, Vill. Dwarf (1'–3' high), tufted, villous when young, stems or scapes mostly 1-flowered; leaflets 3, broadly wedge-obovate, deeply cut into 5–7 oblong approximate teeth. (*P. Robbinsiana*, *Oakes*.) — Alpine summits of the White Mountains of New Hampshire (*Robbins*, *Tuckerman*, &c.). July. — Less villous with age and smaller-flowered than *P. frigida* of the Alps, but agreeing closer with it than with *P. minima*, which probably is only another form of the same species. (Eu.)

4. **P. Canadénsis**, L. (COMMON CINQUE-FOIL OR FIVE-FINGER.) Low or dwarf, silky-hairy; stems decumbent, prostrate, or at length creeping; peduncles axillary, 1-flowered; leaflets 5, obovate-wedge-form, cut-toothed towards the apex. — Dry soil. April–July: producing summer runners (*P. sarmentosa*, *Muhl.*).

Var. **simplex**, Torr. & Gr. Less hairy and greener, larger, the ascending stems (1°–2° long, seldom if ever creeping) from a thicker and harder caudex: leaflets obovate-oblong, sometimes almost glabrous. (*P. simplex*, *Michx.*) — Meadows or moister soil. May–Sept. — Usually appearing distinct.

5. **P. argétea**, L. (SILVERY CINQUE-FOIL.) Stems ascending, paniculately branched at the summit, many-flowered, white-woolly; leaflets 5, wedge-oblong, almost pinnatifid, entire towards the base, with revolute margins, green above, white with silvery wool beneath. — Dry barren fields, &c. June–Sept. (Eu.)

+ + *Leaves pinnate, of 3–9 leaflets.*

6. **P. Pennsylvánica**, L. Stems erect (5'–18' high), hairy or woolly; cymose at the summit, many-flowered; leaflets 5–9, oblong, obtuse, pinnatifid, silky or downy with white hairs, especially beneath, the upper ones larger and crowded; petals scarcely longer than the calyx. — Pennsylvania? New Hampshire (Isle of Shoals, *Robbins*), Maine (Cape Elizabeth, *C. J. Sprague*), N. Wisconsin, and northwestward. July–Aug.

§ 2. *Style deeply lateral: petals yellow or white, deciduous.*

* *Achenia glabrous: style thickened upwards: receptacle conical in fruit.*

7. **P. argúta**, Pursh. Stem erect and stout (1°–4° high,) brownish-hairy, clammy above; leaves pinnate, of 3–9 oval or ovate cut-serrate leaflets, downy

underneath; flowers cymose-clustered; *petals yellowish or whitish*; disk thick and glandular. — Rocky hills: common northward and westward. July.

* * *Achenia* (at least below) and the convex receptacle villous.

8. **P. Anserina**, L. (SILVER-WEED.) Herbaceous, creeping with slender runners; leaves all radical, pinnate; leaflets 9–19, with minute pairs interposed, oblong, pinnatifid-serrate, mostly green and nearly smooth above, silvery-white with silky down underneath; stipules many-cleft; flowers solitary (yellow), on long scape-like peduncles. — Brackish marshes, river-banks, &c., New England to Penn., Wisconsin, and northward. June–Sept. (Eu.)

9. **P. fruticosa**, L. (SHRUBBY CINQUE-FOIL.) Stem erect, shrubby (2°–4° high), very much branched; leaves pinnate; leaflets 5–7, closely crowded, oblong-lanceolate, entire, silky, especially beneath; stipules scale-like; flowers numerous (yellow), terminating the branchlets. — Wet grounds: common northwards. June–Sept. (Eu.)

10. **P. tridentata**, Ait. (THREE-TOOTHED C.) Stems low (4'–6' high), rather woody at the base, tufted, ascending, cymosely several-flowered; leaves palmate; leaflets 3, wedge-oblong, nearly smooth, thick, coarsely 3-toothed at the apex; petals white; achenia and receptacle very hairy. — Coast of New England, from Cape Cod, and mountain-tops from the Alleghanies, northward; also shores of the upper Great Lakes. June.

§ 3. *Styles moderately lateral: petals (shorter than the calyx, ovate-lanceolate) and filaments more or less persistent: disk thick and hairy: achenia glabrous: receptacle hairy, convex, at length large and spongy.* (Comarum, L.)

11. **P. palustris**, Scop. (MARSH FIVE-FINGER.) Stems ascending from a creeping perennial base (1°–2° high); leaves pinnate, of 5–7 lanceolate or oblong crowded serrate leaflets, whitish beneath; flowers somewhat cymose; calyx (1' broad) dark purple inside; petals purple. (Comarum palustre, L.) — Cool bogs, New England to Penn., Illinois, and northward. June–Aug. (Eu.)

12. FRAGÀRIA, Tourn. STRAWBERRY.

Flowers nearly as in *Potentilla*. Styles deeply lateral. Receptacle in fruit much enlarged and conical, becoming pulpy and scarlet, bearing the minute dry achenia scattered over its surface. — Stemless perennials, with runners, and with white cymose flowers on scapes. Leaves radical: leaflets 3, obovate-wedge-form, coarsely serrate. Stipules cohering with the base of the petioles, which with the scapes are usually hairy. (Name from the fragrance of the fruit.) — Flowering in spring. (The species are indiscriminately called WILD STRAWBERRY.)

1. **F. Virginiana**, Ehrhart. *Achenia imbedded in the deeply pitted fruiting receptacle*, which usually has a narrow neck; calyx becoming erect after flowering and connivent over the hairy receptacle when sterile or unfructified; leaflets of a firm or coriaceous texture. (F. Canadensis, Michx.) — Moist or rich woodlands, fields, &c. — In the true *F. Virginiana*, the hairs of the scapes, and especially of the pedicels, are silky and appressed. It is the original of the Virginian Scarlet strawberries.

Var. **Illinoensis** (F. Grayana, Vilmorin, F. Illinoensis & F. Iowensis,

Prince) is a coarser or larger plant, perhaps a distinct species, the flowers more inclined to be polygamo-dioecious, the *villous hairs of the scape and pedicels widely spreading*, as in *F. elatior* and *F. collina*, which it seems to represent in this country. — Common in richer soil, from W. New York to Illinois and beyond the Rocky Mountains. The supposed original of Hovey's Seedling, Boston Pine, and other cultivated varieties.

2. **F. véscá**, L. *Achenia superficial on the glabrous conical or hemispherical fruiting receptacle* (not sunk in pits); calyx remaining spreading or reflexed; hairs on the scape mostly widely spreading, on the pedicels appressed; *leaflets thin*, even the upper face strongly marked by the veins. — Fields and rocky places: common; certainly indigenous northward. (Eu.)

3. **F. ÍNDICA**, L. (or *Duchésnea fragarioides*, *Smith*), — which differs from the true Strawberries in having leafy runners, a calyx with incised leafy bractlets larger than the sepals, *yellow petals*, and *insipid fruit*, — has sparingly established itself in copses around Philadelphia (*Charles E. Smith*, &c.), and in the Southern States. (Adv. from Ind.)

13. DALIBÁRDA, L. DALIBARDA.

Calyx deeply 5-6-parted, 3 of the divisions larger and toothed. Petals 5, sessile, deciduous. Stamens many. Ovaries 5-10, becoming nearly dry seed-like drupes: styles terminal, deciduous. — Low perennials, with creeping and densely tufted stems or rootstocks, and roundish-heart-shaped crenate leaves on slender petioles. Flowers 1 or 2, white, on scape-like peduncles. (Named in honor of *Thomas Dalibard*, a French botanist of the time of Linnæus.)

1. **D. répens**, L. Downy; sepals spreading in the flower, converging and enclosing the fruit. — Wooded banks: common northward. June-Aug. — In aspect and foliage resembling a stemless Violet.

14. RUBÛS, Tourn. BRAMBLE.

Calyx 5-parted, without bractlets. Petals 5, deciduous. Stamens numerous. Achenia usually many, collected on a spongy or succulent receptacle, becoming small drupes: styles nearly terminal. — Perennial herbs, or somewhat shrubby plants, with white (rarely reddish) flowers, and edible fruit. (The Roman name, kindred with *ruber*, red.)

§ 1. RASPBERRY. *Fruit, or collective mass of drupes, falling off whole from the dry receptacle when ripe, or of few grains which fall separately.*

* *Leaves simple: flowers large: prickles none: fruit and receptacle flat and broad.*

1. **R. odorátus**, L. (PURPLE FLOWERING-RASPBERRY.) *Stem shrubby* (3°-5° high); *branches, stalks, and calyx bristly with glandular clammy hairs*; leaves 3-5-lobed, the lobes pointed and minutely toothed, the middle one prolonged; peduncles many-flowered; calyx-lobes tipped with a long narrow appendage; *petals rounded, purple rose-color*; fruit reddish. — Dells, &c.: common northward. June-Aug. — Flowers showy, 2' broad.

2. **R. Nutkánus**, Moçino. (WHITE F.) *Glandular, scarcely bristly*; leaves almost equally 5-lobed, coarsely toothed; peduncles few-flowered; *petals oval, white*. (*R. parviflorus*, *Nutt.*) — Upper Michigan, and westward.

3. **R. Chamæmorus**, L. (CLOUD-BERRY.) *Herbaceous, low, dioecious; stem simple, 2-3-leaved, 1-flowered; leaves roundish-kidney-form, somewhat 5-lobed, serrate, wrinkled; calyx-lobes pointless; petals obovate, white; fruit of few grains, amber-color.* — White Mountains of New Hampshire at the limit of trees: also on the coast at Lubeck, Maine, and northward. (Eu.)

* * Leaflets (pinnately) 3-5: petals small, erect, white.

+ Stems annual, herbaceous, not prickly: fruit of few separate grains.

4. **R. triflorus**, Richardson. (DWARF RASPBERRY.) Stems ascending (6'-12' high) or trailing; leaflets 3 (or pedately 5), rhombic-ovate or ovate-lanceolate, acute at both ends, coarsely doubly serrate, thin, smooth; peduncle 1-3-flowered. — Wooded hillsides, New England to Pennsylvania, Wisconsin, and northward. June. — Sepals and petals often 6 or 7.

+ + Stems biennial and woody, prickly: receptacle oblong: fruit hemispherical.

5. **R. strigosus**, Michx. (WILD RED RASPBERRY.) *Stems upright, and with the stalks, &c. beset with stiff straight bristles (or a few becoming weak hooked prickles), glandular when young, somewhat glaucous; leaflets 3-5, oblong-ovate, pointed, cut-serrate, whitish-downy underneath; the lateral ones sessile; petals as long as the sepals; fruit light red.* — Thickets and hills: common everywhere, especially northward. June, July. — Fruit ripening all summer, more tender than that of the Garden or European Raspberry (*R. IDÆUS*), which it too closely resembles.

6. **R. occidentalis**, L. (BLACK RASPBERRY. THIMBLEBERRY.) *Glaucous all over; stems recurved, armed like the stalks, &c., with hooked prickles, not bristly; leaflets 3 (rarely 5), ovate, pointed, coarsely doubly serrate, whitened-downy underneath; the lateral ones somewhat stalked; petals shorter than the sepals; fruit purple-black (rarely a whitish variety), ripe early in July.* — Very common northward, especially where ground has been burned over.

§ 2. BLACKBERRY. *Fruit, or collective drupes, not separating from the juicy prolonged receptacle, mostly ovate or oblong, blackish.*

7. **R. villosus**, Ait. (COMMON OR HIGH BLACKBERRY.) *Shrubby (10°-60° high), furrowed, upright or reclining, armed with stout curved prickles; branchlets, stalks, and lower surface of the leaves hairy and glandular; leaflets 3 (or pedately 5), ovate, pointed, unequally serrate; the terminal ones somewhat heart-shaped, conspicuously stalked; flowers racemed, numerous, bracts short; sepals linear-pointed, much shorter than the obovate-oblong spreading petals.* — Var. 1. FRONDOSUS: smoother and much less glandular; flowers more corymbose, with leafy bracts; petals roundish. Var. 2. HUMIFUSUS: trailing, smaller; peduncles few-flowered. — Borders of thickets, &c.: common. May, June: the pleasant large fruit ripe in Aug. and Sept. — Plant very variable in size, aspect, and shape of the fruit; — the varieties connecting with

8. **R. Canadensis**, L. (LOW BLACKBERRY. DEWBERRY.) *Shrubby, extensively trailing, slightly prickly; leaflets 3 (or pedately 5-7), oval or ovate-lanceolate, mostly pointed, thin, nearly smooth, sharply cut-serrate; flowers racemed, with leaf-like bracts.* (*R. trivialis*, Pursh, Bigel., &c.; not of Michx.) — Rocky hills and copses: common. May; ripening its excellent fruit earlier than No. 7.

9. **R. hispídus**, L. (RUNNING SWAMP-BLACKBERRY.) *Stems slender, scarcely woody, extensively procumbent, beset with small reflexed prickles; leaflets 3 (or rarely pedately 5), smooth, thickish, mostly persistent, obovate, obtuse, coarsely serrate, entire towards the base; peduncles leafless, several-flowered, often bristly; flowers small.* (R. obovális, Michx. R. sempérvirens and R. scòsus, Bigelow.) — Low woods: common northward. June. — Flowering shoots short, ascending; sterile ones forming long runners. Fruit of few grains, red or purple, sour.

10. **R. cuneifólius**, Pursh. (SAND BLACKBERRY.) *Shrubby (1°-3° high), upright, armed with stout recurved prickles; branchlets and lower surface of the leaves whitish-woolly; leaflets 3-5, wedge-obovate, thickish, serrate above; peduncles 2-4-flowered; petals large.* — Sandy woods, S. New York, Penn. and southward. May-July; ripening its well-flavored black fruit in August.

11. **R. trivialis**, Michx. (LOW BUSH-BLACKBERRY.) *Shrubby, procumbent, bristly and prickly; leaves evergreen, coriaceous, nearly glabrous; leaflets 3 (or pedately 5), ovate-oblong or lanceolate, sharply serrate; peduncles 1-3-flowered; petals large.* — Sandy soil, Virginia and southward. March-May.

15. RÒSA, Tourn. ROSE.

Calyx-tube urn-shaped, contracted at the mouth, becoming fleshy in fruit. Petals 5, obovate or obcordate, inserted, with the many stamens, into the edge of the hollow thin disk that lines the calyx-tube and within bears the numerous pistils below. Ovaries hairy, becoming bony achenia in fruit. — Shrubby and prickly, with odd-pinnate leaves, and stipules cohering with the petiole: stalks, foliage, &c. often bearing aromatic glands. (The ancient Latin name.)

* *Styles cohering in a protruding column, as long as the stamens.*

1. **R. setigera**, Michx. (CLIMBING OR PRAIRIE ROSE.) *Stems climbing, armed with stout nearly straight prickles, not bristly; leaflets 3-5, ovate, acute, sharply serrate, smooth or downy beneath; stalks and calyx glandular; flowers corymbed; sepals pointed; petals deep rose-color changing to white; fruit (hip) globular.* — Borders of prairies and thickets, W. New York (indigenous?) to Wisconsin and southwestward: also cultivated. July. — The only American climbing rose, or with united protruding styles: strong shoots growing 10°-20° in a season.

** *Styles separate, included in the calyx-tube: petals rose-color.*

2. **R. Carolina**, L. (SWAMP ROSE.) *Stems tall (4°-7° high), armed with stout hooked prickles, not bristly; leaflets 5-9, elliptical, often acute, dull above and pale beneath; stipules narrow; flowers numerous, in corymbs; peduncles and calyx (with leaf-like appendages) glandular-bristly; fruit (hip) depressed-globular, somewhat bristly.* — Low grounds: common. June-Sept.

3. **R. lúcida**, Ehrhart. (DWARF WILD-ROSE.) *Stems (1°-2° high), armed with unequal bristly prickles, which are mostly deciduous, the stouter persistent ones nearly straight, slender; leaflets 5-9, elliptical or oblong-lanceolate, shining above, sharply serrate; stipules broad; peduncles 1-3-flowered, and with the appendaged calyx-lobes glandular-bristly; fruit depressed-globular, smooth when ripe.* — Common in dry soil, or along the borders of swamps. May-July. — R. nítida, Willd., is a smooth and narrow-leaved form.

4. **R. blánda**, Ait. (EARLY WILD-ROSE.) *Nearly unarmed*, or with scattered straight deciduous prickles (1° – 3° high); leaflets 5–7, oval or oblong, obtuse, pale on both sides and minutely downy or hoary beneath, serrate; stipules large; flowers 1–3, the peduncles and calyx-tube smooth and glaucous; fruit globose, crowned with the persistent erect and connivent entire calyx-lobes. — Rocks and banks, Vermont to Penn. and Wisconsin, chiefly northward. May, June. — Petals light rose-color.

5. **R. RUBIGINOSA**, L. (SWEET-BRIER.) Climbing high; prickles numerous, the larger ones strong and hooked, the smaller awl-shaped; leaflets doubly serrate, rounded at the base; downy and clothed with fragrant russet glands beneath; fruit pear-shaped or obovate, crowned with the mostly persistent calyx-lobes. — Roadsides and thickets. June–Aug. (Nat. from Eu.)

6. **R. MICRÁNTHA**, Smith. (SMALLER-FL. S.) Prickles uniform and hooked; fruit elliptical and ovate; calyx-lobes deciduous; flowers smaller: otherwise as the last: probably a mere variety. — E. New England. (Nat. from Eu.)

16. CRATÆGUS, L. HAWTHORN. WHITE THORN.

Calyx-tube urn-shaped, the limb 5-cleft. Petals 5, roundish. Stamens many, or only 10–5. Styles 1–5. Pome drupe-like, containing 1–5 bony 1-seeded stones. — Thorny shrubs or small trees, with simple and mostly lobed leaves, and white (rarely rose-colored) blossoms. (Name from *κράτος*, strength, on account of the hardness of the wood.)

* *Corymbs many-flowered.*

+ Fruit very small, depressed-globose (not larger than peas), bright red: flowers small: calyx-teeth short and broad: styles 5: plants glabrous (except No. 1) and glandless.

1. **C. PYRACÁNTHA**, Pers. (EVERGREEN THORN.) Leaves evergreen, shining (1' long), oblong or spatulate-lanceolate, crenulate; the short petioles and young branchlets pubescent; corymbs small. — Shrub, spontaneous at Washington, and near Philadelphia, *Isaac Martindale*. (Adv. from Eu.)

2. **C. spathuláta**, Michx. Leaves thickish, shining, but deciduous, spatulate or oblanceolate, with a long tapering base, crenate above, rarely cut-lobed, nearly sessile. — Virginia and southward. May. — Shrub 10° – 15° high.

3. **C. cordáta**, Ait. (WASHINGTON THORN.) Leaves broadly ovate or triangular, mostly truncate or a little heart-shaped at the base, on a slender petiole, variously 3–5-cleft or cut, serrate. — Virginia, Kentucky, and southward. June. — Trunk 15° – 25° high.

+ + Fruit small ($\frac{1}{4}$ – $1\frac{1}{2}$ ' long), ovoid, deep red: flowers rather large: styles 1–3.

4. **C. OXYACÁNTHA**, L. (ENGLISH HAWTHORN.) Smooth; leaves obovate, cut-lobed and toothed, wedge-form at the base; calyx not glandular. May. — More or less spontaneous as well as cultivated. (Adv. from Eu.)

5. **C. apiifolia**, Michx. Softly pubescent when young; leaves roundish, with a broad truncate or slightly heart-shaped base, pinnately 5–7-cleft, the crowded divisions cut-lobed and sharply serrate; petioles slender; calyx-lobes glandular-toothed, slender. — Virginia and southward. March, April.

+ + + *Fruit large* ($\frac{1}{2}$ ' - $\frac{3}{8}$ ' long), red: *flowers large*: styles and stones of the fruit even in the same species 1-3 (when the fruit is ovoid or pear-shaped), or 4-5 (when the fruit is globular): stipules, calyx-teeth, bracts, &c. often beset with glands.

6. **C. coccinea**, L. (SCARLET-FRUITED THORN.) *Glabrous throughout*; leaves thin, roundish-ovate, sharply toothed and cut, or somewhat cut-lobed, usually abrupt at the base, on slender petioles; flowers white, often with a rosy tinge ($\frac{3}{8}$ ' broad); fruit bright coral-red, ovoid ($\frac{1}{2}$ ' broad), scarcely edible. — Thickets and rocky banks: common. May. — A low tree.

7. **C. tomentosa**, L. (BLACK OR PEAR THORN.) *Downy or villous-pubescent* at least when young on the peduncles, calyx, and lower side of the leaves; leaves thickish, rather large, oval or ovate-oblong, sharply toothed and often cut, abruptly narrowed at the base into a somewhat margined petiole, the upper surface more or less furrowed along the veins; flowers large (often 1, broad), white; fruit scarlet or orange, large ($\frac{3}{8}$ ' - $\frac{3}{4}$ ' broad), globular or somewhat pear-shaped, edible. — Thickets: common. May, June. — A tall shrub or low tree, of many varieties, of which the following are the most marked.

Var. **pyrifolia**. Leaves sparingly pubescent beneath when young, soon glabrous, smooth above, and shining often slightly cut-lobed; fruit large, bright-colored, sparingly dotted, of a pleasant flavor. (*C. pyrifolia*, Ait.)

Var. **punctata**. Leaves rather small, mostly wedge-obovate, with a longer tapering and entire base, unequally toothed above, rarely cut, villous-pubescent when young, smooth but dull when old, the numerous veins more strongly impressed on the upper surface and prominent underneath; fruit globose, usually dull red and yellowish with whitish dots. (*C. punctata*, Jacq.)

Var. **mollis**. Leaves rounded, abrupt or somewhat heart-shaped at the base, soft-downy both sides, or at least beneath, very sharply doubly-toothed and cut; fruit often downy, dull red. (*C. subvillosa*, Schrader. *C. coccinea*, var. ? *mollis*. Torr. & Gray.) — Michigan, Illinois, and southwestward.

8. **C. Crus-galli**, L. (COCKSPUR THORN.) *Glabrous*; leaves thick, shining above, wedge-obovate and oblanceolate, tapering into a very short petiole, serrate above the middle; fruit globular, bright-red ($\frac{1}{3}$ ' broad). — Thickets. June. — Shrub or tree 10° - 20° high, with firm dark-green leaves very shining above, and slender thorns often 2' long. This is our best species for hedges.

* *Corymbs simple, few- (1-6-) flowered: calyx, bracts, &c. glandular.*

9. **C. flava**, Ait. (SUMMER HAW.) Somewhat pubescent or glabrous; leaves wedge-obovate or rhombic-obovate, narrowed into a glandular petiole, unequally toothed and somewhat cut above the middle, rather thin, the teeth glandular; styles 4-5; fruit somewhat pear-shaped, yellowish, greenish, or reddish ($\frac{1}{2}$ ' - $\frac{3}{8}$ ' broad). — Sandy soil, Virginia and southward. May. — Tree 15° - 20° high, with rather large flowers, 2-6 in a corymb.

Var. **pubescens**. Downy or villous-pubescent when young; leaves thickish, usually obtuse or rounded at the summit. (*C. elliptica*, Ait. *C. glandulosa*, Michx. *C. Virginica*, Lodd.) — Virginia and southward.

10. **C. parvifolia**, Ait. (DWARF THORN.) Downy; leaves thick, obovate-spatulate, crenate-toothed ($\frac{1}{2}$ ' - 1 $\frac{1}{2}$ ' long), almost sessile, the upper surface at length

shining; flowers solitary or 2-3 together on *very short peduncles*; *calyx-lobes as long as the petals*; styles 5; fruit globular or pear-shaped, yellowish. — Sandy soil, New Jersey to Virginia and southward. May. — Shrub 3°-6° high.

17. PÿRUS, L. PEAR. APPLE.

Calyx-tube urn-shaped, the limb 5-cleft. Petals roundish or obovate. Stamens numerous. Styles 2-5. Pome fleshy or berry-like; the 2-5 carpels or cells of a papery or cartilaginous texture, 2-seeded. — Trees or shrubs, with handsome flowers in corymbed cymes. (The classical name of the Pear-tree.)

§ 1. MĀLUS, Tourn. (APPLE.) *Leaves simple: cymes simple and umbel-like: pome fleshy, globular, sunk in at the attachment of the stalk.*

1. **P. coronāria**, L. (AMERICAN CRAB-APPLE.) *Leaves ovate, often rather heart-shaped, cut-serrate or lobed, soon glabrous; styles woolly and united at the base.* — Glades, W. New York to Wisconsin and southward. May. — Tree 20' high, with large, rose-colored, fragrant blossoms, few in the corymb, and fragrant, greenish fruit.

2. **P. angustifolia**, Ait. (NARROW-LEAVED C.) *Leaves oblong or lanceolate, often acute at the base, mostly toothed, glabrous; styles distinct.* — Glades, from Pennsylvania southward. April. — Perhaps a variety of No. 1.

§ 2. ADENŌRHACHIS, DC. *Leaves simple, the mid-rib glandular along the upper side: cymes compound: styles united at the base: fruit berry-like, small.*

3. **P. arbutifolia**, L. (CHOKE-BERRY.) *Leaves oblong or obovate, finely serrate; fruit pear-shaped, or when ripe globular.* — Var. 1. **ERYTHROCĀRPA**, has the cyme and leaves beneath woolly, and red or purple fruit. Var. 2. **MELANOCĀRPA**, is nearly smooth, with black fruit. — Damp thickets: common. May, June. — Shrub 2°-10° high. Flowers white, or tinged with purple.

§ 3. SŌRBUS, Tourn. *Leaves odd-pinnate, with rather numerous leaflets: cymes compound: styles separate: pome berry-like, small.*

4. **P. Americana**, DC. (AMERICAN MOUNTAIN-ASH.) *Nearly glabrous or soon becoming so; leaflets 13-15, lanceolate, taper-pointed, sharply serrate with pointed teeth, bright green; cymes large and flat; berries globose, not larger than peas; leaf-buds pointed, glabrous and somewhat glutinous.* — Swamps and mountain-woods, Maine to Penn. and Michigan, and southward along the whole length of the Alleghanies. June. (*P. microcarpa*, DC.) — Tree or tall shrub, with leaflets rather shining above and scarcely pale underneath, the rachis and petiole reddish and elongated: prized in cultivation for the autumnal clusters of bright-red berries.

5. **P. sambucifolia**, Cham. & Schlecht. *Leaflets oblong, oval, or lanceolate, mostly obtuse or abruptly short-pointed, serrate (mostly doubly) with more spreading teeth, often pale beneath; cymes smaller; flowers and berries larger, the latter (4" broad) when young ovoid, at length globose; leaf-buds sparingly hairy: otherwise nearly as the preceding.* (*Sorbus aucuparia*, var. β . Michx.) — Along the northern frontiers of the United States, northward and westward to the Pacific, &c. Perhaps passes into No. 4: it is sometimes cultivated for it, and nearly connects it with.

P. aucuparia, Gærtn., the EUROPEAN MOUNTAIN ASH or ROWAN-TREE, the one more commonly planted in grounds: it has paler, oblong, and obtuse leaflets, their lower surface downy, larger globose berries, and blunter and tomentose leaf-buds.

18. AMELANCHIER, Medic. JUNE-BERRY.

Calyx 5-cleft. Petals oblong, elongated. Stamens numerous, short. Styles 5, united below. Ovary 5-celled, each cell 2-ovuled; but a projection grows from the back of each, and forms a false partition; the berry-like pome thus 10-celled, with one seed in each cell (when all ripen): partitions cartilaginous. — Small trees or shrubs, with simple sharply serrated leaves, and white flowers in racemes. (*Amelancier* is the popular name of *A. vulgaris* in Savoy.)

1. **A. Canadensis**, Torr. & Gray. (SHAD-BUSH. SERVICE-BERRY.) Calyx-lobes triangular-lance-form; fruit globular, purplish, edible (sweet, ripe in June). — Along streams, &c. common, especially northward. April, May. — Varies exceedingly; the leading forms are, —

Var. **Botryapium**; a tree 10° – 30° high, nearly or soon glabrous; leaves ovate-oblong, sometimes heart-shaped to the base, pointed, very sharply serrate; flowers in long drooping racemes; the oblong petals 4 times the length of the calyx. (*Pyrus Botryapium*, *Willd.*)

Var. **oblongifolia**; a smaller tree or shrub; leaves oblong, beneath, like the branchlets white downy when young; racemes and petals shorter.

Var. **rotundifolia**; with broader leaves and smaller petals than in the first variety; racemes 6 – 10-flowered.

Var. **alnifolia**; shrub, with the roundish leaves blunt or notched at both ends, serrate towards the summit; racemes dense and many-flowered. — Chiefly in the Western States and westward.

Var. **oligocarpa**; shrub, with thin and smooth narrowly oblong leaves, and 2–4-flowered racemes, the broader petals scarcely thrice the length of the calyx. — Cold and deep mountain swamps, northward.

ORDER 34. CALYCANTHACEÆ. (CALYCANTHUS FAMILY.)

Shrubs with opposite entire leaves, no stipules, the sepals and petals similar and indefinite, the anthers adnate and extrorse, and the cotyledons convolute: the fruit like a rose-hip. Chiefly represented by the genus

1. CALYCANTHUS, L. CAROLINA ALLSPICE. SWEET-SCENTED SHRUB.

Calyx of many sepals, united below into a fleshy inversely conical cup (with some leaf-like bractlets growing from it); the lobes lanceolate, mostly colored like the petals; which are similar, in many rows, thickish, inserted on the top of the closed calyx-tube. Stamens numerous, inserted just within the petals, short; some of the inner ones sterile (destitute of anthers). Pistils several or many, enclosed in the calyx-tube, inserted on its base and inner face, resembling those of the Rose; but the enlarged hip dry when ripe, enclosing the achenia.

— The lurid purple flowers terminating the leafy branches. Bark and foliage aromatic; the crushed flowers exhaling more or less the fragrance of strawberries. (Name composed of *κάλυξ*, a cup or calyx, and *ἄνθος*, flower, from the closed cup which contains the pistils.)

1. **C. flóridus**, L. *Leaves oval, soft-downy underneath.* — Virginia? and southward, on hillsides in rich soil. Common in gardens. April – Aug.

2. **C. lævigátus**, Willd. *Leaves oblong, thin, either blunt or taper-pointed, bright green and glabrous or nearly so on both sides, or rather pale beneath; flowers smaller.* — Mountains of Franklin Co., Penn. (*Prof. Porter*), and southward along the Alleghanies. May – Aug.

3. **C. glaucus**, Willd. *Leaves oblong-ovate or ovate-lanceolate; conspicuously taper-pointed, glaucous-white beneath, roughish above, glabrous, large (4' – 7' long); probably a variety of the preceding.* — Virginia? near the mountains and southward. May – Aug.

ORDER 35. SAXIFRAGACEÆ. (SAXIFRAGE FAMILY.)

Herbs or shrubs, of various aspect, distinguishable from Rosaceæ by having copious albumen in the seeds, opposite as well as alternate leaves, and usually no stipules when the leaves are alternate; the stamens mostly definite, and the carpels commonly fewer than the sepals, either separate or partly so, or all combined into one compound pistil. Calyx either free or adherent, usually persistent or withering away. Stamens and petals almost always inserted on the calyx. Ovules anatropous. — A large family, to which Parnassia, formerly associated with Drosera, is commonly referred, — now made to include Ribes also.

Tribe I. GROSSULARIÆ. Shrubs, with alternate and palmately veined and lobed leaves: stipules none or united with the base of the petiole. Calyx-tube coherent with the one-celled ovary, which has 2 parietal placentæ and forms a many-seeded berry. Seed-coat externally gelatinous. Embryo minute at the base of the hard albumen.

1. **Ribes.** Character of the tribe. Stamens and small petals 5.

Tribe II. ESCALLONIÆ. Shrubs or trees, with alternate and simple pinnately veined leaves, and no stipules. Ovary 2 – 5-celled.

2. **Itea.** Calyx 5-cleft, free from the 2-celled ovary, which becomes a septicidal pod.

Tribe III. HYDRANGIÆ. Shrubs or trees, with opposite simple leaves, and no stipules. Ovary 2 – 5-celled; the calyx coherent at least with its base. Fruit (in the following) a many-seeded pod.

3. **Hydrangea.** Lobes of the calyx minute in complete flowers. Petals valvate in the bud. Stamens 8 or 10.

4. **Philadelphus.** Lobes of the calyx and petals conspicuous; the former valvate, the latter convolute in the bud. Stamens 20 – 40.

Tribe IV. SAXIFRAGEÆ. Herbs, without stipules, except perhaps a membranous dilatation of the base of the petiole. Petals imbricated or rarely convolute in the bud. Fruit dry, capsular or follicular.

* A cluster of sterile or gland-tipped filaments at the base of each petal. Stigmas 3 or 4, situated directly over as many parietal placentæ!

5. **Parnassia.** Sepals, petals, and proper stamens 5. Peduncle a scape or scape-like, 1-flowered.

* * No sterile stamens or bodies resembling them.

→ Pod 2-3-celled and 2-3-beaked, or of as many distinct follicles.

6. **Astilbe.** Flowers polygamous. Stamens twice as many as the small petals. Seeds few. Leaves decompound.
7. **Saxifraga.** Flowers perfect. Stamens twice as many as the petals. Seeds numerous, with a close coat.
8. **Boykinia.** Flowers perfect. Stamens only as many as the petals, which are convolute in the bud and deciduous. Calyx-tube adherent to the ovary. Seed-coat close.
9. **Sullivantia.** Flowers perfect. Stamens 5. Calyx nearly free. Seeds wing-margined.
 + + Pod one-celled with 2 parietal placentæ.
 ++ Stamens as many as the lobes of the calyx, namely 5.
10. **Heuchera.** Calyx bell-shaped, coherent with the ovary below. Petals small, entire.
 ++ ++ Stamens twice as many as the lobes of the calyx, namely 8 or 10.
11. **Mitella.** Calyx partly cohering with the depressed ovary. Petals small, pinnatifid.
12. **Tiarella.** Calyx nearly free from the slender ovary. Petals entire.
13. **Chrysosplenium.** Calyx-tube coherent with the ovary. Petals none.

1. RĪBES, L. CURRANT. GOOSEBERRY.

Calyx 5-lobed, often colored; the tube coherent with the ovary. Petals 5, inserted in the throat of the calyx, small. Stamens 5, alternate with the petals. Ovary 1-celled, with 2 parietal placentæ and 2 distinct or united styles. Berry crowned with the shrivelled remains of the calyx; the surface of the numerous seeds swelling into a gelatinous outer coat investing a crustaceous one. Embryo minute at the base of hard albumen. — Low, sometimes prickly shrubs, with alternate and palmately-lobed leaves, which are plaited in the bud (except in one species), often clustered in the axils; the small flowers from the same clusters, or from separate lateral buds. (An Arabic name, properly belonging to a species of Rheum. Grossularia was the proper name to have been adopted for the genus.)

§ 1. GROSSULĀRIA, Tourn. (GOOSEBERRY.) *Stems mostly bearing thorns at the base of the leafstalks or clusters of leaves, and often with scattered bristly prickles: berries prickly or smooth. (Our species are indiscriminately called WILD GOOSEBERRY: the flowers greenish.)*

* *Peduncles 1-3-flowered: leaves roundish-heart-shaped, 3-5-lobed.*

1. **R. Cynósbati, L.** Leaves pubescent; *peduncles slender, 2-3-flowered; stamens and undivided style not longer than the broad calyx.* — Rocky woods: common, especially northward. May. — Spines small or obsolete. Berry large, armed with long prickles like a burr, or rarely smooth.

2. **R. hirtéllum, Michx.** Leaves somewhat pubescent beneath; *peduncles very short, 1-2-flowered, deflexed; stamens and 2-cleft style scarcely longer than the bell-shaped calyx; fruit smooth, small, purple, sweet.* — Moist grounds, New England to Illinois, common. May. — Stems either smooth or prickly, and with very short thorns, or none. — This yields the commonest smooth gooseberry of New England, &c., and usually passes for *R. triflorum, Willd.*, which name belongs to the next.

3. **R. rotundifólium, Michx.** Leaves smooth or downy; *peduncles slender, 1-3-flowered; stamens and 2-parted style slender, longer than the narrow cylindrical calyx; fruit smooth, pleasant.* — Rocks, W. Massachusetts to Wisconsin, and southward along the mountains to Virginia, &c. June.

* * *Racemes 4-9-flowered, slender, nodding.*

4. **R. lacustre**, Poir. Young stems clothed with bristly prickles, and with weak thorns; leaves heart-shaped, 3-5-parted, with the lobes deeply cut; calyx broad and flat; stamens and style not longer than the petals; fruit bristly (small, unpleasant).—Cold woods and swamps, New England to Wisconsin and northward. June.

§ 2. **RIBËSIA**, Berl. (CURRANT.) *Stems neither prickly nor thorny: flowers (greenish) in racemes: berries never prickly.*

5. **R. prostratum**, L'Her. (FETID CURRANT.) Stems reclined; leaves deeply heart-shaped, 5-7-lobed, smooth; the lobes ovate, acute, doubly serrate; racemes erect, slender; calyx flattish; pedicels and the (pale red) fruit glandular-bristly.—Cold damp woods and rocks, from N. England and Penn. northward. May.—The bruised plant and berries exhale an unpleasant odor.

6. **R. flóridum**, L. (WILD BLACK CURRANT.) *Leaves sprinkled with resinous dots, slightly heart-shaped, sharply 3-5-lobed, doubly serrate; racemes drooping, downy; bracts longer than the pedicels; calyx tubular-bell-shaped, smooth; fruit round-ovoid, black, smooth.*—Woods: common. May.—Much like the *Black Currant* of the gardens, which the berries resemble in smell and flavor. Flowers large, whitish.

7. **R. rúbrum**, L. (RED CURRANT.) Stems straggling or reclined; leaves somewhat heart-shaped, obtusely 3-5-lobed, serrate, downy beneath when young; racemes from lateral buds distinct from the leaf-buds, drooping; calyx flat (green or purplish); fruit globose, smooth, red; on our wild plant apt to turn upwards in the drooping raceme: the veins of the leaves are whitish beneath (whence the name *R. albinervium*, Michx.): but apparently not distinct from the garden *Red Currant* of the Old World.—Cold bogs and damp woods, New Hampshire to Minnesota and northward. May, June. (Eu.)

R. aúreum, Pursh, the **BUFFALO** or **MISSOURI CURRANT**, remarkable for the spicy fragrance of its yellow blossoms in early spring, is widely cultivated for ornament. Its leaves are convolute (instead of plaited) in the bud.

2. ÍTEA, L. ÍTEA.

Calyx 5-cleft, free from the ovary or nearly so. Petals 5, lanceolate, much longer than the calyx, and longer than the 5 stamens. Pod oblong, 2-grooved, 2-celled, tipped with the 2 united styles, 2-parted (septicidal) when mature, several-seeded.—Shrubs, with simple, alternate, petioled leaves, without stipules, and small white flowers in simple racemes. (Greek name of the Willow.)

1. **I. Virgínica**, L. Leaves deciduous, oblong, pointed, minutely serrate; seeds oval, flattish, with a crustaceous coat.—Wet places, New Jersey and southward, near the coast. June.

3. HYDRÁNGEA, Gronov. HYDRANGEA.

Calyx-tube hemispherical, 8-10-ribbed, coherent with the ovary; the limb 4-5-toothed. Petals ovate, valvate in the bud. Stamens 8-10, slender. Pod crowned with the 2 diverging styles, 2-celled below, many-seeded, opening by a hole between the styles.—Shrubs, with opposite petioled leaves, no stipules,

and numerous flowers in compound cymes. The marginal flowers are usually sterile and radiant, consisting merely of a membranaceous and colored flat and dilated calyx, and showy. (Name from ὕδωρ, *water*, and ἄγγος, *a vase*.)

1. **H. arboréscens**, L. (WILD HYDRANGEA.) Glabrous or nearly so; leaves ovate, rarely heart-shaped, pointed, serrate, green both sides; cymes flat. — Rocky banks, N. Penn. to Illinois, and southward. July. — Flowers often all fertile, rarely all radiant, like the *Garden Hydrangea*.

4. PHILADÉLPHUS, L. MOCK ORANGE OF SYRINGA.

Calyx-tube top-shaped, coherent with the ovary; the limb 4–5-parted, spreading, persistent, valvate in the bud. Petals rounded or obovate, large, convolute in the bud. Stamens 20–40. Styles 3–5, united below or nearly to the top. Stigmas oblong or linear. Pod 3–5-celled, splitting at length into as many pieces. Seeds very numerous, on thick placentæ projecting from the axis, pendulous, with a loose membranaceous coat prolonged at both ends. — Shrubs, with opposite often toothed leaves, no stipules, and solitary or cymose-clustered showy white flowers. (An ancient name, applied by Linnæus to this genus for no obvious reason.)

1. **P. inodorus**, L. Glabrous; leaves ovate or ovate-oblong, pointed, entire or with some spreading teeth; flowers single or few at the ends of the diverging branches, pure white, scentless; calyx-lobes acute, scarcely longer than the tube. — Mountains of Virginia and southward.

Var. **grandiflorus**. Somewhat pubescent; flowers larger; calyx-lobes longer and taper-pointed. — Virginia and southward, near the mountains. May–July. — A tall shrub, with long and recurved branches: often cultivated.

P. coronarius, L., the common MOCK ORANGE OF SYRINGA of the gardens, with cream-colored, odorous flowers, in full clusters, the crushed foliage with the odor and taste of cucumbers, — has sometimes escaped from grounds.

5. PARNÁSSIA, Tourn. GRASS OF PARNASSUS.

Sepals 5, imbricated in the bud, slightly united at the base, and sometimes also with the base of the ovary, persistent. Petals 5, veiny, spreading, at length deciduous, imbricated in the bud: a cluster of somewhat united gland-tipped sterile filaments at the base of each. Proper stamens 5, alternate with the petals: filaments persistent: anthers opening inwards. Ovary 1-celled, with 4 projecting parietal placentæ: stigmas 4, sessile, directly over the placentæ. Pod 4-valved, the valves bearing the placentæ on their middle. Seeds very numerous, anatropous, with a thick wing-like seed-coat and little if any albumen. Embryo straight: cotyledons very short. — Perennial smooth herbs, with the leaves entire and chiefly radical, and the solitary flowers terminating the long naked stems. Petals white, with greenish or yellowish veins. (Named from Mount Parnassus: called Grass of Parnassus by Dioscorides.) In former edition placed between Droseraceæ and Hypericaceæ.

1. **P. parviflora**, DC. *Petals sessile*, little longer than the calyx; *sterile filaments about 5 in each set, slender*; leaves ovate or oblong, tapering at the base. — N. W. shore of L. Michigan (White-Fish Bay, Wisconsin, *Henry Gillman*),

and westward. July. — More slender than the next, and the flower only half the size.

2. **P. palústris**, L. *Petals sessile*; rather longer than the calyx, few-veined; *sterile filaments* 9–15 in each set, slender; leaves heart-shaped. — Shore of Lake Superior, and northward. Aug. — Stalks 3'–10' high. Flower nearly 1' broad. (Eu.)

3. **P. Caroliniána**, Michx. *Petals sessile*, more than twice the length of the calyx, many-veined; *sterile filaments* 3 in each set, stout, distinct almost to the base. — Wet banks, New England to Wisconsin and southward, especially along the mountains. July–Sept. — Leaves thickish, ovate or rounded, often heart-shaped, usually but one on the stalk, and that low down and clasping. Stalk 9'–2° high. Flower 1'–1½' broad.

4. **P. asarifólia**, Vent. *Petals abruptly contracted into a claw* at the base; *sterile filaments* 3 in each set; leaves rounded kidney-shaped: otherwise as in the foregoing. — High Alleghanies of Virginia and southward.

6. **ASTÍLBE**, Don. FALSE GOATBEARD.

Flowers diœciously polygamous. Calyx 4–5-parted, small. Petals 4–5, spatulate, small, withering-persistent. Stamens 8 or 10. Ovary 2-celled, almost free, many-ovuled: styles 2, short. Pod 2-celled, separating into 2 follicles, each ripening few seeds. Seed-coat loose and thin, tapering at each end. — Perennial herbs, with twice or thrice ternately-compound ample leaves, cut-lobed and toothed leaflets, and small white or yellowish flowers in spikes or racemes, which are disposed in a compound panicle. (Name composed of $\acute{\alpha}$ privative and $\sigma\tau\acute{\iota}\lambda\beta\eta$, a bright surface, because the foliage is not shining.)

1. **A. decándra**, Don. Somewhat pubescent; leaflets mostly heart-shaped; petals minute or wanting in the fertile flowers; stamens 10. — Rich woods, Alleghanies of S. W. Virginia and southward. July. — Plant imitating *Spiræa Aruncus*, but coarser, 3°–5° high.

7. **SAXÍFRAGA**, L. SAXIFRAGE.

Calyx either free from or cohering with the base of the ovary, 5-cleft or parted. Petals 5, entire, imbricated in the bud, commonly deciduous. Stamens 10. Styles 2. Pod 2-beaked, 2-celled, opening down or between the beaks; or sometimes 2 almost separate follicles. Seeds numerous, with a close coat. — Chiefly perennial herbs, with the root-leaves clustered, those of the stem mostly alternate. (Name from *saxum*, a rock, and *frango*, to break; many species rooting in the clefts of rocks.)

* *Stems prostrate, in tufts, leafy: leaves opposite: calyx free from the pod.*

1. **S. oppositifólia**, L. (MOUNTAIN SAXIFRAGE.) Leaves fleshy, ovate, keeled, ciliate, imbricated on the sterile branches (1"–2" long); flowers solitary, large; petals purple, obovate, much longer than the 5-cleft-calyx. — Rocks, Willoughby Mountain, Vermont, and northward. (Eu.)

** *Stems ascending, leafy: stem-leaves alternate: calyx coherent below with the pod.*

← *Petals all alike.*

2. **S. rivuláris**, L. (ALPINE BROOK-S.) Small; stems weak, 3–5-

flowered; lower leaves rounded, 3-5-lobed, on slender petioles, the upper lanceolate; petals white, ovate. — Alpine region of Mount Washington, New Hampshire (*Oakes, &c.*): rare. (Eu.)

3. **S. aizoides**, L. (YELLOW MOUNTAIN-S.) Low (3'-5' high), in tufts, with few or several corymbose flowers; leaves linear-lanceolate, entire, fleshy, distantly spinulose-ciliate; petals yellow, spotted with orange, oblong. — Willoughby Mountain, Vermont; near Oneida Lake, New York; N. Michigan; and northward. June. (Eu.)

4. **S. tricuspidata**, Retz. Stems tufted (4'-8' high), naked above; flowers corymbose; leaves oblong or spatulate, with 3 rigid sharp teeth at the summit; petals obovate-oblong, yellow. — Shore of L. Superior and northward. (Eu.)

* * Leaves clustered at the root: scape many-flowered, erect, clammy-pubescent.

5. **S. Aizoon**, Jacq. Leaves persistent, thick, spatulate, with white cartilaginous toothed margins; calyx partly adherent; petals obovate, cream-color, often spotted at the base. — Moist rocks, Upper Michigan and Wisconsin; Willoughby Mountain, Vermont (*Mr. Blake*), and northward. — Scape 5'-10' high. (Eu.)

6. **S. Virginiensis**, Michx. (EARLY S.) Low (4'-9' high); leaves obovate or oval-spatulate, narrowed into a broad petiole, crenate-toothed, thickish; flowers in a clustered cyme, which is at length open and loosely paniced; lobes of the nearly free calyx erect, not half the length of the oblong obtuse (white) petals; pods 2, united merely at the base, divergent, purplish. — Exposed rocks: common, especially northward. April-June.

7. **S. Pennsylvanica**, L. (SWAMP S.) Large (1°-2° high); leaves oblanceolate, obscurely toothed (4'-8' long), narrowed at the base into a short and broad petiole; cymes in a large oblong panicle, at first clustered; lobes of the nearly free calyx recurved, about the length of the linear-lanceolate (greenish) small petals; filaments awl-shaped: pods at length divergent. — Bogs: common, especially northward. May, June. — A homely species.

8. **S. erosa**, Pursh. (LETTUCE S.) Leaves oblong or oblanceolate, obtuse, sharply toothed, tapering into a margined petiole (8'-12' long); scape slender (1°-3° high); panicle elongated, loosely flowered; pedicels slender; calyx reflexed, entirely free, nearly as long as the oval obtuse (white) petals; filaments club-shaped; pods 2, nearly separate, diverging. — Cold mountain brooks, Pennsylvania (near Bethlehem, *Mr. Wolfe*), and throughout the Alleghanies, southward. June.

←← Petals unequal, with claws, white, all or some of them with a pair of yellow spots near the base: leaves oblong, wedge-shaped or spatulate; calyx free and reflexed.

9. **S. leucanthemifolia**, Lapeyrouse, Michx. Leaves coarsely toothed or cut, tapering into a petiole; scapes (5'-18' high) bearing one or more leaves or leafy bracts and a loose, spreading corymbose or paniculate cyme; petals lanceolate; the 3 larger ones with a heart-shaped base and a pair of spots; the 2 smaller with a tapering base and no spots. — Salt Pond Mountain, Virginia (*Wm. M. Canby*), and southward in the Alleghanies.

10. **S. stellaris**, L., var. **comosa**, Willd. Leaves wedge-shaped, more or less toothed; scape (4'-5' high) bearing a small contracted panicle; many

or most of the flowers changed into little tufts of green leaves; *petals all lanceolate and tapering into the claw*. — Mount Katahdin, Maine (*Rev. J. Blake*) and high northward. (Eu.)

8. BOYKÍNIA, Nutt. BOYKINIA.

Calyx-tube top-shaped, coherent with the 2-celled and 2-beaked pod. Stamens 5, as many as the deciduous petals, these mostly convolute in the bud. Otherwise as in *Saxifraga*. — Perennial herbs, with alternate palmately 5-7-lobed or cut petioled leaves, and white flowers in cymes. (Dedicated to the late *Dr. Boykin* of Georgia.)

1. *B. aconitifolia*, Nutt. Stem glandular (6'-20' high); leaves deeply 5-7-lobed. — Mountains of S. W. Virginia, and southward. July.

9. SULLIVÁNTIA, Torr. & Gray. SULLIVANTIA.

Calyx bell-shaped, cohering below only with the base of the ovary, 5-cleft. Petals 5, entire, acutish, withering-persistent. Stamens 5, shorter than the petals. Pod 2-celled, 2-beaked, many-seeded, opening between the beaks: the seeds wing-margined, imbricated upwards. — A low and reclined-spreading perennial herb, with rounded and cut-toothed, or slightly lobed smooth leaves, on slender petioles, and small white flowers in a branched loosely cymose panicle, raised on a nearly leafless slender scape (6'-12' long). Peduncles and calyx glandular: pedicels recurved in fruit. (Dedicated to the distinguished bryologist who discovered the only species.)

1. *S. Ohionis*, Torr. & Gr. (*Gray, Chloris Bor.-Am., pl. 6.*) — Limestone cliffs, Highland County, Ohio, *Sullivant*; Wisconsin River, *Lapham*. June.

10. HEÛCHERA, L. ALUM-ROOT.

Calyx bell-shaped; the tube cohering at the base with the ovary 5-cleft. Petals 5, spatulate, small, entire. Stamens 5. Styles 2, slender. Pod 1-celled, with 2 parietal many-seeded placentæ, 2-beaked, opening between the beaks. Seeds oval, with a rough and close seed-coat. — Perennials, with the round heart-shaped leaves principally from the rootstock; those on the scapes, if any, alternate. Petioles with dilated margins or adherent stipules at their base. Flowers in small clusters disposed in a prolonged and narrow panicle, greenish or purplish. (Named in honor of *John Henry Heucher*, a German botanist of the beginning of the 18th century.)

* *Flowers small, loosely paniced: stamens and styles exerted: calyx regular.*

1. *H. villósa*, Michx. Scapes (1°-3° high), petioles, and veins of the acutely 7-9-lobed leaves beneath *villous with rusty hairs*; calyx 1½" long; *petals spatulate-linear, about as long as the stamens, soon twisted*. — Rocks, Maryland, Kentucky and southward, in and near the mountains. Aug. - Sept.

2. *H. Americana*, L. (COMMON ALUM-ROOT.) Scapes (2°-3° high), &c. *glandular* and more or less *hirsute with short hairs*; leaves roundish, with short rounded lobes and crenate teeth; calyx broad, 2" long, the *spatulate petals not longer than its lobes*. — Rocky woodlands, Connecticut to Wisconsin and southward. June.

* * *Flowers larger*: calyx (3'' - 4'' long) more or less oblique: stamens short: panicle very narrow: leaves rounded, slightly 5 - 9-lobed.

3. **H. hispida**, Pursh. *Hispid* or *hirsute* with long spreading hairs (occasionally almost glabrous), scarcely glandular; *stamens soon exerted, longer than the spatulate petals.* (*H. Richardsonii*, R. Br.) — Mountains of Virginia. Also Illinois (*Dr. Mead*) and northwestward. May - July. — Scapes 2° - 4° high.

4. **H. pubescens**, Pursh. Scape (1° - 3° high) and petioles *granular-pubescent* or *glandular above*, not hairy, below often glabrous; *stamens shorter than the lobes of the calyx* and the spatulate petals. — Rich woods, Lancaster, Penn. to Virginia and Kentucky, along the mountains. June, July.

11. MITÉLLA, Tourn. MITRE-WORT. BISHOP'S-CAP.

Calyx short, coherent with the base of the ovary, 5-cleft. Petals 5, slender, pinnatifid. Stamens 10, included. Styles 2, very short. Pod short, 2-beaked, 1-celled, with 2 parietal or rather basal several-seeded placenta, 2-valved at the summit. Seeds smooth and shining. — Low and slender perennials, with round heart-shaped alternate leaves on the rootstock or runners, on slender petioles; those on the scapes, opposite, if any. Flowers small, in a simple slender raceme or spike. (Name a diminutive from *μίτρα*, a mitre or cap, alluding to the form of the young pod.)

1. **M. diphýlla**, L. *Hairy, leaves heart-shaped, acute*, somewhat 3 - 5-lobed, toothed, *those on the many-flowered-scape 2, opposite, nearly sessile.* — Hillsides in rich woods: common, especially westward and northward. May. — Flowers white, in a raceme 6' - 8' long.

2. **M. nuda**, L. Small and slender; *leaves rounded or kidney-form*, deeply and doubly crenate; *scape usually leafless, few-flowered*, very slender (4' - 6' high). (*M. cordifolia*, Lam. *M. prostrata*, Michx.) — Deep moist woods in moss, Maine to Wisconsin and northward. May - July. — A delicate little plant, sending forth runners in summer. Blossoms greenish.

12. TIARÉLLA, L. FALSE MITRE-WORT.

Calyx bell-shaped, nearly free from the ovary, 5-parted. Petals 5, with claws, entire. Stamens 10, long and slender. Styles 2. Pod membranaceous, 1-celled, 2-valved; the valves unequal. Seeds few, at the base of each parietal placenta, globular, smooth. — Perennials: flowers white. (Name a diminutive from *τιάρα*, a tiara, or turban, from the form of the pod, or rather pistil, which is like that of *Mitella*, to which the name of *Mitre-wort* properly belongs.)

1. **T. cordifolia**, L. Leaves from the rootstock or summer runners heart-shaped, sharply lobed and toothed, sparsely hairy above, downy beneath; scape leafless (5' - 12' high); raceme simple; petals oblong. — Rich rocky woods: common northward, and southward along the mountains. April, May.

13. CHRYSOSPLÈNIUM, Tourn. GOLDEN SAXIFRAGE.

Calyx-tube coherent with the ovary; the blunt lobes 4 - 5, yellow within. Petals none. Stamens 8 - 10, very short, inserted on a conspicuous disk.

Styles 2. Pod inversely heart-shaped or 2-lobed, flattened, very short, 1-celled, with 2 parietal placentæ, 2-valved at the top, many-seeded. — Low and small smooth herbs, with tender succulent leaves, and small solitary or leafy-cymed flowers. (Name compounded of χρυσός, *golden*, and σπλήν, *the spleen*, probably from some reputed medicinal qualities.)

1. **C. Americanum**, Schwein. Stems slender, spreading and forking; leaves principally opposite, roundish or somewhat heart-shaped, obscurely crenate-lobed; flowers distant, inconspicuous, nearly sessile (greenish tinged with yellow or purple.) — Cold wet places: common northward. April, May.

ORDER 36. CRASSULACEÆ. (ORPINE FAMILY.)

Succulent herbs, with perfectly symmetrical flowers: viz. the petals and pistils equalling the sepals in number (3–20), and the stamens the same or double their number, — technically different from Saxifrageæ only in this complete symmetry, and in the carpels being quite distinct from each other, but even this does not hold in two N. American genera. Also, instead of a perigynous disk, there are usually little scales on the receptacle, one behind each carpel. Fruit dry and dehiscent; the pods (follicles) opening down the ventral suture, many- rarely few-seeded. — Stipules none. Flowers usually cymose, small. Leaves mostly sessile, in Penthorum not at all fleshy.

* Not succulent: the carpels united, forming a 5-celled pod. Transition to the Saxifrage Family.

1. **Penthorum**. Sepals 5. Petals none. Stamens 10. Pod 5-beaked, many-seeded.

* * Leaves, &c., thick and succulent. Carpels distinct.

2. **Tillæa**. Sepals, petals, stamens, and pistils 3 or 4. Seeds few or many.

3. **Sedum**. Sepals, petals, and pistils 4 or 5. Stamens 10–8. Seeds many.

1. PENTHORUM, Gronov. DITCH STONE-CROP.

Sepals 5. Petals rare, if any. Stamens 10. Pistils 5, united below, forming a 5-angled, 5-horned, and 5-celled pod, which opens by the falling off of the beaks, many-seeded. — Upright weed-like perennials (not fleshy like the rest of the family), with scattered leaves, and yellowish-green flowers loosely spiked along the upper side of the naked branches of the cyme. (Name from πέντε, *five*, and ῥῆσος, *a rule or mode*, probably from the quinary order of the flower.)

1. **P. sedoides**, L. Leaves lanceolate, acute at both ends. — Open wet places, everywhere. July–Oct. — Parts of the flower rarely in sixes or sevens.

2. TILLÆA, L. TILLÆA.

Sepals, petals, stamens, and pistils 3 or 4. Pods 2–many-seeded. — Very small tufted annuals, with opposite entire leaves and axillary flowers. (Named in honor of *Michael Angelo Tilli*, an early Italian botanist.)

1. **T. simplex**, Nutt. Rooting at the base (1'–2' high); leaves linear-oblong; flowers solitary, nearly sessile; calyx half the length of the (greenish-white) petals and the narrow 8–10-seeded pods, the latter with a scale at the base of each. (*T. ascéndens*, *Eaton*.) — Muddy river-banks, Nantucket to Maryland. July–Sept.

3. SÈDUM, Tourn. STONE-CROP. ORPINE.

Sepals and petals 4 or 5. Stamens 8 or 10. Pods many-seeded; a little scale at the base of each. — Chiefly perennial, smooth, and thick-leaved herbs, with the flowers cymose or one-sided. Petals almost always narrow and acute or pointed. (Name from *sedeo*, to sit, alluding to the manner in which these plants fix themselves upon rocks and walls.)

* *Flowers perfect and sessile, as it were spiked along one side of spreading flowering branches or of the divisions of a scorpioid cyme, the first or central flower mostly 5-merous and 10-androus, the others often 4-merous and 8-androus.*

1. **S. ACRE**, L. (MOSSY STONE-CROP.) Spreading on the ground, moss-like; leaves very small, alternate, almost imbricated on the branches, ovate, very thick; petals yellow. — Escaped from cultivation to rocky roadsides, &c. July. (Adv. from Eu.)

2. **S. pulchellum**, Michx. Stems ascending or trailing (4'–12' high); leaves *terete, linear-filiform*, much crowded; spikes of the cyme several, densely flowered; petals *rose-purple*. — Virginia to S. Illinois, Kentucky, and southward; also cultivated in gardens. July.

3. **S. Névii**, Gray. Stems spreading, simple (3'–5' high); leaves *all alternate*, those of the sterile shoots *wedge-obovate or spatulate*, on flowering stems *linear-spatulate* and flattish; cyme about 3-spiked, densely flowered; petals *white*, more pointed than in the next; the flowering 3 or 4 weeks later; leaves and blossoms smaller. — Mountains of Virginia (Salt Pond Mountain, *W. M. Canby*) to Alabama (*R. D. Nevius*).

4. **S. ternatum**, Michx. Stems spreading (3'–6' high); leaves *flat*; the lower *whorled in threes, wedge-obovate*, the upper scattered, *oblong*; cyme 3-spiked, leafy; petals *white*. — Rocky woods, Penn. to Illinois and southward: common in gardens. May, June.

** *Flowers in a terminal naked and regular cyme or cluster, more or less peduncled: leaves flat, obovate or oblong, mostly alternate.*

+ *Flowers perfect, 5-merous, 10-androus.*

5. **S. telephioides**, Michx. Stems ascending (6'–12' high), stout, leafy to the top; leaves oblong or oval, entire or sparingly toothed; cyme small; petals *flesh-color*, ovate-lanceolate, taper-pointed; pods *tapering into a slender style*. — Dry rocks, Alleghany Mountains, from Maryland southward, and sparingly in New Jersey? W. New York? and Indiana. June. — Too near the next.

6. **S. TELÈPHICUM**, L. (GARDEN ORPINE OR LIVE-FOR-EVER.) Stems erect (2° high), stout; leaves oval, obtuse, toothed; cymes compound; petals *purple*, oblong-lanceolate; pods *abruptly pointed with a short style*. — Rocks and banks, escaped from cultivation in some places. July. (Adv. from Eu.)

+ + *Flowers dioecious, mostly 4-merous and 8-androus.*

7. **S. Rhodiola**, DC. (ROSEROOT.) Stems erect (5'–10' high); leaves oblong or oval, smaller than in the preceding; flowers in a close cyme, greenish-yellow, or the fertile turning purplish. — Pennsylvania, on cliffs of Delaware River, above Easton! (*Professors Porter & Green*); Quoddy Head, Maine (*Prof. Verrill*), and northward. May, June. (Eu.)

ORDER 37. **HAMAMELACEÆ.** (WITCH-HAZEL FAMILY.)

Shrubs or trees, with alternate simple leaves and deciduous stipules; flowers in heads or spikes, often polygamous or monœcious; the calyx cohering with the base of the ovary; which consists of 2 pistils united below, and forms a 2-beaked, 2-celled woody pod, opening at the summit, with a single bony seed in each cell, or several, only one or two of them ripening. — Petals inserted on the calyx, narrow, valvate, or involute in the bud, or often none at all. Stamens twice as many as the petals, and half of them sterile and changed into scales, or numerous. Seeds anatropous. Embryo large and straight, in sparing albumen: cotyledons broad and flat.

Tribe I. HAMAMELEÆ. Flowers with a manifest calyx or calyx and corolla, and a single ovule suspended from the summit of each cell.

1. **Hamamelis.** Petals 4, strap-shaped. Stamens and scales each 4, short.

2. **Fothergilla.** Petals none. Stamens about 24, long: filaments thickened upwards.

Tribe II. BALSAMIFLUÆ. Flowers naked, with barely rudiments of a calyx, and no corolla, crowded into catkin-like heads. Ovules several or many in each cell.

3. **Liquidambar.** Monœcious or polygamous. Stamens very numerous. Pods consolidated by their bases in a dense head.

1. **HAMAMELIS, L.** WITCH-HAZEL.

Flowers in little axillary clusters or heads, usually surrounded by a scale-like 3-leaved involucre. Calyx 4-parted, and with 2 or 3 bractlets at its base. Petals 4, strap-shaped, long and narrow, spirally involute in the bud. Stamens 8, very short; the 4 alternate with the petals anther-bearing, the others imperfect and scale-like. Styles 2, short. Pod opening loculicidally from the top; the outer coat separating from the inner, which encloses the single large and bony seed in each cell, but soon bursts elastically into two pieces. — Tall shrubs, with straight-veined leaves, and yellow, perfect or polygamous flowers. (From *ἄμα*, like to, and *μηλís*, an apple-tree; a name anciently applied to the Medlar, or some other tree resembling the Apple, which the Witch-Hazel does not.)

1. **H. Virginica, L.** Leaves obovate or oval, wavy-toothed, somewhat downy when young. — Damp woods: blossoming late in autumn, when the leaves are falling, and maturing its seeds the next summer.

2. **FOTHERGILLA, L. f.** FOTHERGILLA.

Flowers in a terminal catkin-like spike, mostly perfect. Calyx bell-shaped, the summit truncate, slightly 5-7-toothed. Petals none. Stamens about 24, borne on the margin of the calyx in one row, all alike: filaments very long, thickened at the top (white). Styles 2, slender. Pod cohering with the base of the calyx, 2-lobed, 2-celled, with a single bony seed in each cell. — A low shrub; the oval or obovate leaves smooth, or hoary underneath, toothed at the summit; the flowers appearing rather before the leaves, each partly covered by a scale-like bract. (Dedicated to the distinguished *Dr. John Fothergill.*)

1. **F. alnifolia, L. f.** — Low grounds, Virginia and southward. April, May.

3. LIQUIDÁMBAR, L. SWEET-GUM TREE.

Flowers usually monœcious, in globular heads or catkins; the sterile arranged in a conical cluster, naked: stamens very numerous, intermixed with minute scales: filaments short. Fertile flowers consisting of many 2-celled 2-beaked ovaries, subtended by minute scales in place of a calyx, all more or less cohering together and hardening in fruit, forming a spherical catkin or head; the pods opening between the 2 awl-shaped beaks. Styles 2, stigmatic down the inner side. Ovules many, but only one or two perfecting. Seeds with a wing-angled seed-coat. — Catkins racemed, nodding, in the bud enclosed by a 4-leaved deciduous involucre. (A mongrel name, from *liquidus*, fluid, and the Arabic *ambar*, amber; in allusion to the fragrant terebinthine juice which exudes from the tree.)

1. **L. Styraciflua**, L. (SWEET GUM. BILSTED.) Leaves rounded, deeply 5-7-lobed, smooth and shining, glandular-serrate, the lobes pointed. — Moist woods, from Connecticut to Illinois, and southward. April. — A large and beautiful tree, with fine-grained wood, the gray bark commonly with corky ridges on the branchlets. Leaves fragrant when bruised, turning deep crimson in autumn. The woody pods filled mostly with abortive seeds, resembling sawdust.

ORDER 38. HALORÀGEÆ. (WATER-MILFOIL FAMILY.)

Aquatic or marsh plants (at least in northern countries), with the inconspicuous symmetrical flowers sessile in the axils of leaves or bracts, calyx-tube coherent with the ovary, which consists of 2-4 more or less united carpels (or in Hippuris of only one carpel), the styles or sessile stigmas distinct. Limb of the calyx obsolete or very short in fertile flowers. Petals small or none. Stamens 1-8. Fruit indehiscent, 1-4-celled, with a single anatropous seed suspended from the summit of each cell. Embryo in the axis of fleshy albumen: cotyledons minute. — Formerly attached as a suborder to Onagraceæ, but now deemed quite distinct.

1. **Myriophyllum**. Flowers monœcious or polygamous, the parts in fours, with or without petals. Stamens 4 or 8. Immersed leaves pinnately dissected.
2. **Proserpinaca**. Flowers perfect, the parts in threes. Petals none. Immersed leaves pinnately dissected.
3. **Hippuris**. Flowers usually perfect. Petals none. Stamen, style, and cell of the ovary only one. Leaves entire.

1. MYRIOPHÝLLUM, Vaill. WATER-MILFOIL.

Flowers monœcious or polygamous. Calyx of the sterile flowers 4-parted, of the fertile 4-toothed. Petals 4, or none. Stamens 4-8. Fruit nut-like, 4-celled, deeply 4-lobed: stigmas 4, recurved. — Perennial aquatics. Leaves crowded, often whorled; those under water pinnately parted into capillary divisions. Flowers sessile in the axils of the upper leaves, usually above water in summer; the uppermost staminate. (Name from *μυρίος*, a thousand, and *φύλλον*, a leaf, i. e. Milfoil.)

* *Stamens 8; petals deciduous: carpels even: leaves whorled in threes or fours.*

1. **M. spicatum**, L. Leaves all pinnately parted and capillary, except the floral ones or bracts; these ovate, entire or toothed, and chiefly shorter than the flowers, which thus form an interrupted spike. — Deep water: common. (Eu.)

2. **M. verticillatum**, L. Floral leaves much longer than the flowers, pectinate-pinnatifid: otherwise nearly as No. 1. — Ponds, &c. northward. (Eu.)

* * *Stamens 4: petals rather persistent: carpels 1-2-ridged and roughened on the back: leaves whorled in fours and fives, the lower with capillary divisions.*

3. **M. heterophyllum**, Michx. Stem stout; floral leaves ovate and lanceolate, thick, crowded, sharply serrate, the lowest pinnatifid; fruit obscurely roughened. — Lakes and rivers, from N. New York westward and southward.

4. **M. scabratum**, Michx. Stem rather slender; lower leaves pinnately parted with few capillary divisions; floral leaves linear (rarely scattered), pectinate-toothed or cut-serrate: carpels strongly 2-ridged and roughened on the back. — Shallow ponds, from S. New England and Ohio southward.

* * * *Stamens 4: petals rather persistent: carpels even on the back: leaves chiefly scattered, or wanting on the flowering stems.*

5. **M. ambiguum**, Nutt. Immersed leaves pinnately parted into about 10 very delicate capillary divisions; the emerging ones pectinate, or the upper floral linear and sparingly toothed or entire; flowers mostly perfect; fruit (minute) smooth. — Var. 1. NĀTANS: stems floating, prolonged. Var. 2. CAPILLACEUM: stems floating, long and very slender; leaves all immersed and capillary. Var. 3. LIMOSUM: small, rooting in the mud; leaves all linear, incised, toothed, or entire. — Ponds and ditches, Massachusetts to New Jersey, Pennsylvania, and southward, near the coast.

6. **M. tenellum**, Bigelow. Flowering stems nearly leafless and scape-like, (3'-10' high), erect, simple; the sterile shoots creeping and tufted; bracts small, entire; flowers alternate, monœcious; fruit smooth. — Borders of ponds, N. New York, New England, and northward.

2. PROSERPINACA, L. MERMAID-WEED.

Flowers perfect. Calyx-tube 3-sided, the limb 3-parted. Petals none. Stamens 3. Stigmas 3, cylindrical. Fruit bony, 3-angled, 3-celled, 3-seeded, nut-like. — Low, perennial herbs, with the stems creeping at the base (whence the name, from *proserpo*, to creep), alternate leaves, and small flowers sessile in the axils, solitary or 3-4 together, in summer.

1. **P. palustris**, L. Leaves lanceolate, sharply serrate, the lower pectinate when under water; fruit sharply angled. — Wet swamps: not rare.

2. **P. pectinacea**, Lam. Leaves all pectinate, the divisions linear-awl-shaped; fruit rather obtusely angled. — Sandy swamps, near the coast.

9. HIPPURIS, L. MARE'S TAIL.

Flowers perfect or polygamous. Calyx entire. Petals none. Stamen one, inserted on the edge of the calyx. Style single, thread-shaped, stigmatic down one side, received in the groove between the lobes of the large anther. Fruit

nut-like, 1-celled, 1-seeded. — Perennial aquatics, with simple entire leaves in whorls, and minute flowers sessile in the axils in summer. (Name from *ἵππος*, a horse, and *οὐρά*, a tail.)

1. **H. vulgaris**, L. Leaves in whorls of 8 or 12, linear, acute. — Ponds and springs, New York to Kentucky and northward: rare. Stems simple, 1°-2° high. Flowers very inconspicuous. (Eu.)

ORDER 39. ONAGRACEÆ. (EVENING-PRIMROSE FAMILY.)

Herbs, with 4-merous (sometimes 2-3- or 5-6-merous) perfect and symmetrical flowers; the tube of the calyx cohering with the 2-4-celled ovary, its lobes valvate in the bud, or obsolete; the petals convolute in the bud, sometimes wanting; and the stamens as many or twice as many as the petals or calyx-lobes, inserted on the summit of the calyx-tube. Style single, slender: stigma 2-4-lobed or capitate. Pollen grains often connected by cobwebby threads. Seeds anatropous, small, without albumen.

* Parts of the flower in twos.

1. **Circæa**. Petals 2, obcordate or 2-lobed. Stamens 2. Fruit 1-2-seeded, bristly.

** Parts of the flowers in fours or more.

— Fruit dry and indehiscent, mostly becoming 1-celled, 1-4-seeded.

2. **Gaura**. Petals 4. Stamens 8 and with the long style turned downwards.

— — Fruit a many-seeded pod, usually loculicidal.

3. **Epilobium**. Stamens 8. Petals 4. Seeds with a large downy tuft at the apex.

4. **Oenothera**. Stamens 8 and petals 4 on the prolonged calyx-tube. Seeds naked.

5. **Jussiaea**. Stamens 8-12. Petals 4-6. Calyx-tube not prolonged beyond the ovary.

6. **Ludwigia**. Stamens 4. Petals 4 or more. Calyx-tube not prolonged beyond the ovary.

1. CIRCÆA, Tourn. ENCHANTER'S NIGHTSHADE.

Calyx-tube slightly prolonged, the end filled by a cup-shaped disk, deciduous; lobes 2, reflexed. Petals 2, inversely heart-shaped. Stamens 2. Fruit indehiscent, small and bur-like, bristly with hooked hairs, 1-2-celled: cells 1-seeded. — Low and inconspicuous perennials, in cool or damp woods, with opposite thin leaves on slender petioles, and small whitish flowers in racemes, produced in summer. (Named from *Circe*, the enchantress.)

1. **C. Lutetiàna**, L. Taller (1°-2° high); *leaves ovate*, slightly toothed; *bracts none*; hairs of the *roundish 2-celled fruit bristly*. — Very common. (Eu.)

2. **C. alpina**, L. *Low* (3'-8' high), *smooth and weak*; *leaves heart-shaped, thin, shining, coarsely toothed*; *bracts minute*; hairs of the *obovate-oblong 1-celled fruit soft and slender*. — Deep woods: common northward. July. (Eu.)

2. GAÛRA, L. GAURA.

Calyx-tube much prolonged beyond the ovary, deciduous; the lobes 4 (rarely 3), reflexed. Petals clawed, unequal or turned to the upper side. Stamens mostly 8, often turned down, as is also the long style. A small scale-like appendage before the base of each filament. Stigma 4-lobed, surrounded by a ring or cup-like border. Fruit hard and nut-like, 3-4-ribbed or angled, inde-

hiscent or nearly so, usually becoming 1-celled and 1-4-seeded. Seeds naked. — Leaves alternate, sessile. Flowers rose-color or white, changing to reddish in fading, in wand-like spikes or racemes; in our species quite small (so that the name, from γαῦρος, *superb*, does not seem appropriate).

1. **G. biennis**, L. *Soft-hairy or downy* (3°-8° high); leaves *oblong-lanceolate*, acute, denticulate; fruit oval or oblong, *nearly sessile*, ribbed, downy. — Dry banks, from New York westward and southward: common. Aug.

2. **G. filipes**, Spach. *Nearly smooth*; stem slender (2°-4° high); leaves *linear*, mostly toothed, tapering at the base; branches of the panicle very slender, naked; fruit obovate-club-shaped, 4-angled at the summit, *slender pedicelled*. — Open places, Virginia to Ohio, Illinois, and southward. Aug.

3. EPILÒBIUM, L. WILLOW-HERB.

Calyx-tube not prolonged beyond the ovary; the limb 4-cleft, deciduous. Petals 4. Stamens 8: anthers short. Pod linear, many-seeded. Seeds with a tuft of long hairs at the end. — Perennials, with nearly sessile leaves, and violet, purple, or white flowers; in summer. (Name composed of ἐπὶ λοβοῦ ἴον, viz. *a violet on a pod*.)

* *Flowers large in a long spike or raceme: petals widely spreading, on claws, entire: stamens and style turned downwards: stigma of 4 long lobes: leaves scattered.*

1. **E. angustifolium**, L. (GREAT WILLOW-HERB.) Stem simple, tall (4°-7°); leaves lanceolate. — Low grounds, especially in newly cleared land: common northward. — Flowers pink-purple, very showy. (Eu.)

* * *Flowers rather large, regular: petals obcordate: stamens and style erect: stigma of 4 long linear lobes: leaves mostly opposite.*

2. **E. hirsutum**, L. *Soft-hairy, branching* (3°-5° high); leaves lance-oblong, serrulate; flowers in the upper axils or in a leafy short raceme; petals rose-purple, 6" long. — Spontaneous in waste grounds, New Bedford, Mass. (*T. A. Greene*) and Roxbury (*D. Murray*); and in a ravine near Albany, New York (*C. H. Peck*). (Nat. from Eu.)

* * *Flowers small, corymbed or paniced: petals (mostly notched at the end), stamens, and style erect: stigma club-shaped, nearly entire: lower leaves opposite, entire or denticulate.*

3. **E. alpinum**, L. *Low* (2'-6' high), *nearly glabrous*; stems ascending from a stoloniferous base, *simple*; leaves elliptical or ovate-oblong, obtuse, nearly entire, on short petioles; flowers few or solitary, drooping in the bud; petals purple; pods long, glabrous. — Alpine summits of the White Mountains of New Hampshire, and Adirondack Mountains, New York. (Eu.)

Var. **majus**, Wahl. Taller; upper leaves more or less acute and toothed, pod glabrous or somewhat pubescent. (*E. alsinifolium*, *Vill.* *E. organifolium*, *Lam.*) — With the typical form: also upper Wisconsin and Michigan. (Eu.)

4. **E. palustre**, L., var. lineare. *Erect and slender* (1°-2° high), branched above, *minutely hoary-pubescent*; stem roundish; leaves *narrowly lanceolate or linear*, nearly entire; flower-buds somewhat nodding; petals purplish or white; pods hoary. (*E. lineare*, *Muhl.* *E. squamatum*, *Nutt.*) — Bogs, N.

England to Penn., Illinois, and northward. There is also a small an simple 1-few-flowered form (4'-9' high), less hoary or nearly glabrous, with shorter leaves (*E. oligánthum*, Michx.), found in N. New York, White Mountains of New Hampshire, and northward. This is *E. nutans*, *Sommerf.* & *E. lineare*, *Fries*, but the pods are usually a little hoary. (Eu.)

5. **E. mólle**, Torr. *Soft-downy all over*, strictly erect (1° - $2\frac{1}{2}^{\circ}$ high), at length branching; *leaves crowded; linear-oblong or lanceolate, blunt*, mostly petioled; petals rose-color, notched (2"-3" long). — Bogs, Rhode Island and Pennsylvania to Michigan, and northward. Sept.

6. **E. colorátum**, Muhl. *Glabrous or nearly so; stem roundish, not angled, much branched* (1° - 3° high), many-flowered; *leaves lanceolate or ovate-oblong, acute, denticulate, often petioled*, not at all decurrent, thin, usually purple-veined; flower-buds erect; petals purplish, 2-cleft at the summit ($1\frac{1}{2}$ "-2" long). — Wet places: common. July-Sept.

4. CENOThÈRA, L. EVENING PRIMROSE.

Calyx-tube prolonged beyond the ovary, deciduous; the lobes 4, reflexed. Petals 4. Stamens 8: anthers mostly linear and versatile. Pod 4-valved, many-seeded. Seeds naked. — Leaves alternate. (Name from *οἶνος*, wine, and *θήρα*, a hunt or eager pursuit, given to some plant the roots of which were eaten to provoke a relish for wine.)

§ 1. *Tube of the calyx filiform or cylindrical and much prolonged beyond the ovary: stamens nearly equal: anthers linear: stigma of 4 filiform or linear divergent lobes.*

* *Annuals or biennials: flowers nocturnal, odorous, yellow, withering the next day: pods fusiform or cylindrical, closely sessile.*

1. **Ce. biénnis**, L. (COMMON EVENING-PRIMROSE.) *Erect*, mostly hairy; leaves ovate-lanceolate, acute, obscurely toothed; flowers in a terminal rather leafy spike; calyx-tube much prolonged; *petals inversely heart-shaped* (light yellow); *Pods oblong*, somewhat tapering above. — Varies greatly; as Var. 1. MURICATA, with rough-bristly stem and pods, and petals rather longer than the stamens. Var. 2. GRANDIFLORA, with larger and more showy petals. Var. 3. PARVIFLORA, with petals about the length of the stamens. Var. 4. CRUCIATA, with singularly small and narrow linear-oblong petals, shorter than the stamens, and smooth pods. Var. 5. OAKESIANA (*Ce. Oakesiana*, *Robbins*), with a minute and wholly appressed pubescence, or almost glabrous: otherwise as Var. 3. — Fields and waste places: very common. June-Sept.

2. **Ce. rhombipétala**, Nutt. *Petals rhombic-ovate, acute; calyx-tube very slender, extended an inch longer than the short cylindrical ovary: otherwise resembling a smoothish or rather hoary narrow-leaved state of No. 1.* — Sandy soil, Illinois, Wisconsin, and westward. July-Sept.

3. **Ce. sinuáta**, L. *Hairy, low, ascending, or at length procumbent; leaves oblong or lanceolate, sinuate-toothed, often pinnatifid, the lower petioled; flowers (small) axillary; petals not longer than the stamens (pale yellow, rose-color in fading); pods cylindrical, elongated.* — Sandy fields, New Jersey and southward, principally a dwarf state. June.

* * *Biennials or perennials: flowers diurnal (opening in sunshine), yellow: pods club-shaped, with 4 strong or winged angles and 4 intermediate ribs.*

4. **Œ. glauca**, Michx. Very glabrous, glaucous; leaves ovate or ovate-lanceolate; pods obovoid-oblong, 4-winged, almost sessile; root perennial. — Mountains of Virginia, Kentucky, and southward. May — July. — Leaves broader and flowers larger than in the next.

5. **Œ. fruticosa**, L. (SUNDRUPS.) Hairy or nearly smooth (1° – 3° high); leaves lanceolate or oblong; raceme corymbed, naked below; petals broadly obcordate, longer than the calyx-lobes and stamens; pods oblong-club-shaped, 4-winged, longer than the pedicels; root perennial. — Open places, S. New England to Illinois, and southward. June — Aug. — Corolla $1\frac{1}{2}'$ broad.

6. **Œ. riparia**, Nutt. Scarcely pubescent; leaves linear-lanceolate, elongated, tapering below and somewhat stalked; flowers (large) in a rather leafy at length elongated raceme; petals slightly obcordate; pods oblong-club-shaped, slender-pedicelled, scarcely 4-winged; root biennial. — River-banks and swamps, Quaker Bridge, New Jersey, to Virginia and southward.

7. **Œ. linearis**, Michx. Slender, minutely hoary-pubescent; leaves linear; flowers (rather large) somewhat corymbed at the end of the branches; pods obovate, hoary, scarcely 4-winged at the summit, tapering into a slender pedicel. — Montauk Point, Long Island, to Virginia and southward. June, July. — Plant 1° high, bushy-branched: flowers $1'$ wide.

8. **Œ. chrysantha**, Michx. Slender, smooth or pubescent; leaves lanceolate, rather blunt; flowers crowded or at first corymbed; petals obovate, notched at the end (orange-yellow), longer than the stamens: pods all pedicelled, oblong-club-shaped, scarcely wing-angled; root biennial? — Banks, Oswego, New York, to Wisconsin and northward. July. — Stem $12'$ – $15'$ high; flowers larger than in No. 9, from which it may not be distinct.

9. **Œ. pumila**, L. Almost smooth, small; leaves lanceolate or oblanceolate, mostly obtuse; flowers in a loose and prolonged leafy raceme; petals obcordate (pale yellow), scarcely longer than the stamens; pods almost sessile, oblong-club-shaped, strongly wing-angled; root perennial or biennial? — Dry fields: common northward, and southward along the Alleghanies. June. — Stems mostly simple, $5'$ – $12'$ high: the corolla $\frac{1}{2}'$ broad.

§ 2. *Tube of the calyx funnel-form, strongly 4-nerved, and shorter than the cylindrical ovary, its lobes keeled with the midrib: filaments opposite the petals shorter: anthers oblong, versatile: stigma disk-shaped, almost entire: flowers opening in sunshine or daylight.*

10. **Œ. serrulata**, Nutt. Stems low, slightly woody at the base; leaves lance-linear, oblanceolate or linear-spatulate, sharply serrulate or toothed; flowers axillary mostly small; petals yellow, obovate, wavy-crenulate, much longer than the stamens; pods cylindrical, puberulent. — Falls of St. Anthony, Wisconsin (*Lesquereux*, *T. J. Hale*), and westward.

5. JUSSIÆA, L. JUSSIÆA.

Calyx-tube elongated, not at all prolonged beyond the ovary; the lobes 4–6, herbaceous and persistent. Petals 4–9. Stamens twice as many as the petals.

Pod 4-6-celled, usually long, opening between the ribs. Seeds very numerous. — Herbs (ours glabrous perennials), with mostly entire and alternate leaves, and axillary yellow flowers, in summer. (Dedicated to *Bernard de Jussieu*, the founder of the Natural System of Botany, as further developed by his illustrious nephew.)

1. **J. decurrens**, DC. *Stem erect* (1° - 2° high), branching, *winged* by the decurrent lanceolate leaves; calyx-lobes 4, as long as the petals; pod oblong-club-shaped, wing-angled. — Wet places, Virginia to Illinois, and southward.

2. **J. repens**, L. *Stem creeping, or floating and rooting*; leaves oblong, tapering into a slender petiole; flowers large, long-peduncled; calyx-lobes and obovate petals 5; pod cylindrical, with a tapering base. — In water, Illinois, Kentucky, and southward. Also nat. near Philadelphia.

6. LUDWIGIA, L. FALSE LOOSESTRIFE.

Calyx-tube not at all prolonged beyond the ovary; the lobes 4, usually persistent. Petals 4, often small or wanting. Stamens 4. Pod short or cylindrical, many-seeded. Seeds minute, naked. — Perennial herbs, with axillary (rarely capitate) flowers, produced through summer and autumn. (Named in honor of *Christian G. Ludwig*, Professor of Botany at Leipsic, contemporary with Linnaeus.)

* *Leaves all alternate, sessile or nearly so.*

← *Flowers peduncled in the upper axils, with conspicuous yellow petals (4" - 8" long), equalling the ovate or lanceolate foliaceous lobes of the calyx.*

1. **L. alternifolia**, L. (SEED-BOX.) *Smooth* or nearly so, branched (3° high); leaves lanceolate, acute or pointed at both ends; pods cubical, rounded at the base, wing-angled. — Swamps: common, especially near the coast. — Pods opening first by a hole where the style falls off, afterwards splitting in pieces.

2. **L. hirtella**, Raf. *Hairy* all over; stems nearly simple (1° - 2° high); leaves oblong, or the upper lanceolate, blunt at both ends; pods nearly as in the last, but scarcely wing-angled. — Moist pine barrens, New Jersey to Virginia, and southward. — Fascicled roots often tuberous-thickened.

← ← *Flowers small, sessile (solitary or sometimes clustered or crowded) in the axils, with very small greenish petals (in No. 5) or mostly none: leaves mostly lanceolate or linear on the erect stems (1° - 3° high) and numerous branches; but prostrate or creeping sterile shoots or stolons are often produced from the base of the stem, these are thickly beset with shorter obovate or spatulate leaves. (Our species glabrous, except No. 3.)*

3. **L. sphærocárpa**, Ell. Minutely pubescent, especially the calyx, or nearly glabrous; leaves lanceolate or linear, acute, tapering at the base; those of the runners obovate with a wedge-shaped base and glandular-denticulate; bractlets minute, obsolete, or none; pods globular or depressed (sometimes acute at the base), not longer than the calyx-lobes (less than 2" long). — Water or wet swamps, E. Mass., S. New York, New Jersey, Penn., and southward. — Bark of lower part of the stem often spongy-thickened.

4. **L. polycárpa**, Short & Peter. Leaves narrowly lanceolate, acute at both ends; those of the runners oblong-spatulate, acute, entire; bractlets linear-

awl-shaped, and conspicuous on the base of the 4-sided somewhat top-shaped pod, which is longer than the calyx-lobes.—Swamps, Michigan to Illinois, Kentucky, and southward.

5. **L. linearis**, Walt. Slender, mostly low; leaves narrowly linear, those of the short runners obovate; minute petals usually present; *bractlets minute at the base of the elongated top-shaped 4-sided pod, which is 3" long and much longer than the calyx-lobes.*—Bogs, pine barrens of New Jersey and southward.

6. **L. cylíndrica**, Ell. Much branched; leaves oblong- or spatulate-lanceolate, much tapering at the base, or even petioled; *bractlets very minute at the base of the cylindrical pod, which is 3" long, and several times exceeds the calyx-lobes.*—Mound City, S. Illinois, *Dr. Vasey*, and southward.

* * *Leaves all opposite: stems creeping or floating.*

7. **L. palústris**, Ell. (WATER PURSLANE.) Smooth: leaves ovate or oval, tapering into a slender petiole; petals none, or small and reddish when the plant grows out of water; calyx-lobes very short; pods oblong, 4-sided, not tapering at the base, sessile in the axils (2" long). (*Isnárdia palustris, L.*)—Ditches: common. (Eu.)

8. **L. arcuáta**, Walt. Smooth, small and creeping; leaves oblanceolate, nearly sessile; flowers solitary, long-peduncled; petals yellow, exceeding the calyx (3" long); pods oblong-club-shaped, somewhat curved ($\frac{1}{3}$ ' long).—Swamps, Eastern Virginia and southward.

ORDER 40. MELASTOMACEÆ. (MELASTOMA FAMILY.)

Plants with opposite 3-7-ribbed leaves, and definite stamens, the anthers opening by pores at the apex; otherwise much as in the Evening-Primrose Family.—All tropical, except the genus

1. RHÉXIA, L. DEER-GRASS. MEADOW-BEAUTY.

Calyx-tube urn-shaped, coherent with the ovary below, and continued above it, persistent, 4-cleft at the apex. Petals 4, convolute in the bud, oblique, inserted, along with the 8 stamens, on the summit of the calyx-tube. Anthers long, 1-celled, inverted in the bud. Style 1. stigma 1. Pod invested by the permanent calyx, 4-celled, with 4 many-seeded placentæ projecting from the central axis. Seeds coiled like a snail-shell, without albumen.—Low perennial herbs, often bristly, with sessile 3-5-nerved and bristle-edged leaves, and large showy cymose flowers; in summer; the petals falling early. (Name from $\rho\eta\chi\iota\varsigma$, a rupture, applied to this genus for no obvious reason.)

* *Anthers linear, curved, with a minute spur on the back at the attachment of the filament above its base: flowers cymose, peduncled.*

1. **R. Virginica**, L. *Stem square, with wing-like angles; leaves oval-lanceolate, acute; petals bright purple.*—Sandy swamps, E. Massachusetts to Penn., Wisconsin, Illinois, and southward.

2. **R. Mariána**, L. *Stems cylindrical; leaves linear-oblong, narrowed below; petals paler.*—Sandy swamps, New Jersey, Kentucky, and southward.

* * *Anthers oblong, straight, without any spur: flowers few, sessile.*

3. **R. ciliōsa**, Michx. Stem square, glabrous; leaves broadly ovate, ciliate with long bristles; calyx glabrous. — Maryland and southward.

ORDER 41. LYTHRÀCÆ. (LOOSESTRIFE FAMILY.)

Herbs, with mostly opposite entire leaves, no stipules, the calyx enclosing but free from the 1-4-celled many-seeded ovary and membranous pod, and bearing the 4-7 deciduous petals and 4-14 stamens on its throat; the latter lower down. Style 1: stigma capitate, or rarely 2-lobed. — Flowers axillary or whorled, rarely irregular, perfect, sometimes dimorphous or even trimorphous, those on different plants with filaments and style reciprocally longer and shorter. Petals sometimes wanting. Pod often 1-celled by the early breaking away of the thin partitions: placentæ in the axis. Seeds anatropous, without albumen. — Branches usually 4-sided.

* Flowers regular, or nearly so.

1. **Ammannia**. Calyx short, 4-angled, not striate. Petals 4, or none. Stamens 4, rarely 2.
2. **Lythrum**. Calyx tubular-cylindrical, striate. Petals 5-7. Stamens 5-14.
3. **Nesaea**. Calyx short-campanulate or hemispherical. Stamens 10-14, exserted.

* * Flowers irregular: petals unequal.

4. **Cuphea**. Calyx spurred or enlarged on one side at the base. Stamens 12.

1. AMMÁNIA, Houston. AMMANNIA.

Calyx globular or bell-shaped, 4-angled, 4-toothed, usually with a little horn-shaped appendage at each sinus. Petals 4 (purplish), small and deciduous, sometimes wanting. Stamens 4, rarely 2, short. Pod globular, 2-4-celled. — Low and inconspicuous smooth herbs, with opposite narrow leaves, and small greenish flowers in their axils, produced all summer. (Named after *Paul Ammann*, a German botanist anterior to Linnaeus.)

§ 1. *Calyx with manifest tooth-like or horn-shaped appendages at the sinuses: pod 4-celled: plants of low or wet ground; ours are annuals.*

1. **A. humilis**, Michx. *Leaves tapering at the base or into a short petiole, linear-oblongate or somewhat spatulate; flowers solitary or 3 together in the axils of the leaves, sessile; style very short.* — Massachusetts to Michigan, Illinois and southward.

2. **A. latifolia**, L. *Leaves linear-lanceolate (2'-3' long), with a broad auricled sessile base; style sometimes very short, sometimes slender.* — Ohio, Illinois, and southward. Ship-yards, Philadelphia, an immigrant from the south, *C. F. Parker*.

§ 2. **HYPOBRÝCHIA**, M. A. Curtis. *Appendages at the sinuses of the calyx mere callous points or none: petals none: pod 2-celled.*

3. **A. Nuttállii**, Gray. Submersed aquatic, or sometimes terrestrial, rooting in the mud; leaves linear, when immersed elongated, thin, and closely sessile by a broad base, when out of water shorter and contracted at the base; flowers mostly solitary in the axils, sessile, small; calyx with broad triangular lobes; style very short. (*Péplis diandra*, *Nutt.*, but stamens usually 4. *Hypo-*

brichia Nuttallii, *M. A. Curtis.*) — Wisconsin and Minnesota (*T. J. Hale*), Illinois (*Buckley, Vasey, Hall, &c.*), and southward. — When in deep water the stems are 1°–3° long, very leafy, the flowers and pods not larger than a pin's head: when terrestrial 2'–6' long, larger-flowered, resembling depauperate specimens of No. 1, with obsolete projections at the sinuses.

2. LÿTHRUM, L. LOOSESTRIFE.

Calyx cylindrical, striate, 5–7-toothed, with as many little processes in the sinuses. Petals 5–7. Stamens as many as the petals or twice the number, inserted low down on the calyx, commonly nearly equal. Pod oblong, 2-celled. — Slender herbs, with opposite or scattered mostly sessile leaves, and purple (rarely white) flowers; produced in summer. (Name from *λύθρον*, *blood*; perhaps from the crimson blossoms of some species.)

* *Stamens and petals 5–7: flowers small, solitary and nearly sessile in the axils of the mostly scattered upper leaves: proper calyx-teeth often shorter than the intermediate processes: plants smooth.*

1. **L. Hyssopifolia**, L. Low annual (6'–10' high), pale; leaves oblong-linear, obtuse, longer than the inconspicuous flowers; petals (pale-purple) 5–6. — Marshes, coast of New England and New Jersey. (Eu.)

2. **L. alatum**, Pursh. Tall and wand-like perennial; branches with margined angles; leaves from oblong-ovate to lanceolate, the upper not longer than the flowers; petals (deep purple) 6. — Michigan, Wisconsin, and southward.

3. **L. lineare**, L. Stem slender and tall (3°–4° high), bushy at the top, two of the angles margined; leaves linear, short, chiefly opposite, obtuse, or the upper acute and scarcely exceeding the flowers; calyx obscurely striate; petals (whitish) 6. — Brackish marshes, New Jersey and southward.

* * *Stamens 12, twice the number of the petals, 6 longer and 6 shorter: flowers large, crowded and whorled, in an interrupted wand-like spike.*

4. **L. Salicaria**, L. (SPIKED LOOSESTRIFE.) Leaves lanceolate, heart-shaped at the base, sometimes whorled in threes. — Wet meadows, Eastern New England, and Orange County, New York: also cultivated. — Plant more or less downy, tall: flowers large, purple, trimorphous, as to respective length of style and filaments in 3 different kinds of individuals.

3. NESÆA, Commerson, Juss. SWAMP LOOSESTRIFE.

Calyx short, broadly bell-shaped or hemispherical, with 5–7 erect teeth, and as many longer and spreading horn-like processes at the sinuses. Petals 5. Stamens 10–14, exserted, of two lengths. Pod globose, 3–5-celled. — Perennial herbs or slightly shrubby plants, with opposite or whorled leaves, and axillary flowers (these probably dimorphous or trimorphous).

1. **N. verticillata**, H. B. K. Smooth or downy; stems recurved (2°–8° long), 4–6-sided; leaves lanceolate, nearly sessile, opposite or whorled, the upper with clustered flowers in their axils on short pedicels; petals 5, wedge-lanceolate, rose-purple ($\frac{1}{2}$ ' long); stamens 10, half of them shorter. (*Décodon verticillatum*, *Gmelin.*) — Swampy grounds: common eastward. July–Sept.

4. CÛPHEA, Jacq. CUPHEA.

Calyx tubular, 12-ribbed, somewhat inflated below, gibbous or spurred at the base on the upper side, 6-toothed at the apex, and usually with as many little processes in the sinuses. Petals 6, very unequal. Stamens mostly 12, approximate in 2 sets, included, unequal. Ovary with a curved gland at the base next the spur of the calyx, 1-2-celled: style slender: stigma 2-lobed. Pod oblong, few-seeded, early ruptured through one side. — Flowers solitary or racemose, stalked. (Name from *κυφός*, *gibbous*, from the shape of the calyx, &c.)

1. *C. viscosissima*, Jacq. (CLAMMY CUPHEA.) Annual, very viscid-hairy, branching: leaves ovate-lanceolate; petals ovate, short-clawed, purple. — Dry fields, from Connecticut to Illinois, and southward. Aug. — Seeds flat, borne on one side of the placenta, which is early forced out the ruptured pod.

ORDER 42. LOASÀCEÆ. (LOASA FAMILY.)

Herbs, with a rough or stinging pubescence, no stipules, the calyx-tube adherent to a 1-celled ovary with 2 or 3 parietal placenta; — represented here only by the genus

1. MENTZÈLIA, Plumier. (BARTONIA, Nutt.)

Calyx-tube cylindrical or club-shaped; the limb 5-parted, persistent. Petals 5 or 10, regular, spreading, flat, convolute in the bud, deciduous. Stamens indefinite, rarely few, inserted with the petals on the throat of the calyx. Styles 3, more or less united into one: stigmas terminal, minute. Pod at length dry and opening irregularly, few-many-seeded. Seeds flat, anatropous, with little albumen. — Stems erect. Leaves alternate. Flowers terminal, solitary or cymose-clustered. (Dedicated to *C. Mentzel*, an early German botanist.)

1. *M. oligosperma*, Nutt. Rough and adhesive (1°-3° high), much branched, the brittle branches spreading; leaves ovate and oblong, cut-toothed or angled; flowers yellow (7"-10" broad), opening in sunshine; petals wedge-oblong, pointed; stamens 20 or more: filaments filiform: pod small, about 9-seeded. — Prairies and plains, Illinois, and on the western plains; — where *M. ORNATA* and *M. NUDA*, with large white flowers, are showy representatives of the genus.

ORDER 43. CACTÀCEÆ. (CACTUS FAMILY.)

Fleshy and thickened mostly leafless plants, of peculiar aspect, globular, or columnar and many-angled, or flattened and jointed, usually with prickles. Flowers solitary, sessile; the sepals and petals numerous, imbricated in several rows, adherent to the 1-celled ovary. — Stamens numerous, with long and slender filaments, inserted on the inside of the tube or cup formed by the union of the sepals and petals. Style 1. stigmas numerous. Fruit a 1-celled berry, with numerous campylotropous seeds on several parietal placenta. — Represented east of the Mississippi only by the genus

1. **OPÚNTIA**, Tourn. PRICKLY PEAR. INDIAN FIG.

Sepals and petals not united into a prolonged tube, spreading, regular, the inner roundish. Berry often prickly. Seeds flat and margined. Embryo coiled around albumen: cotyledons large, foliaceous in germination.—Stem composed of joints, bearing very small awl-shaped and usually deciduous leaves arranged in a spiral order, with clusters of barbed bristles and often spines also in their axils. Flowers in our species yellow, opening in sunshine for more than one day. (A name of Theophrastus, originally belonging to some different plant.)

1. **O. vulgaris**, Mill. (Cactus *Opuntia*, L.) Low, prostrate or spreading, pale, with flat and broadly obovate joints; the minute leaves ovate-subulate and appressed; the axils bristly, rarely with a few small spines; flowers sulphur-yellow; berry nearly smooth, pulpy, eatable.—Sandy fields and dry rocks, from Nantucket, Mass., southward, usually near the coast. June.

2. **O. Rafinesqui**, Engelm. Joints (*deep green*) and flowers larger than in the preceding, the latter often with a red centre, and with more numerous (10–12) petals; leaves spreading, longer and narrower (3"–4"); axils some of them bearing a few small spines and a single strong one (9"–12" long).—Wisconsin to Kentucky and westward. June.

3. **O. Missouriënsis**, DC. Prostrate; the joints broadly obovate and flat (2'–4' long), tuberculate; leaves minute; axils armed with a tuft of straw-colored bristles and 5–10 slender radiating spines (1'–2' long); flowers light yellow; berry dry, prickly.—Borders of Wisconsin and westward. May–July.

ORDER 44. **PASSIFLORACEÆ**. (PASSION-FLOWER FAMILY.)

Herbs or woody plants, climbing by tendrils, with perfect flowers, 5 monadelphous stamens, and a stalked 1-celled ovary free from the calyx, with 3 or 4 parietal placentæ, and as many club-shaped styles;—represented by the typical genus

1. **PASSIFLÒRA**, L. PASSION-FLOWER.

Calyx of 5 sepals united at the base into a short cup, imbricated in the bud, usually colored like the petals, at least within; the throat crowned with a double or triple fringe. Petals 5, on the throat of the calyx. Stamens 5: filaments united in a tube which sheathes the long stalk of the ovary, separate above: anthers large, fixed by the middle. Berry (often edible) many-seeded; the anatropous albuminous seeds invested by a pulpy covering. Seed-coat brittle, grooved.—Leaves alternate, generally palmately lobed, with stipules. Peduncles axillary, jointed. Ours are perennial herbs. (Name, from *passio*, passion, and *flos*, a flower, given by the early missionaries in South America to these blossoms, in which they fancied a representation of the implements of the crucifixion.)

1. **P. lutea**, L. Smooth, slender; leaves obtusely 3-lobed at the summit, the lobes entire; petioles glandless; flowers greenish-yellow (1' broad).—Damp thickets, S. Penn. to Ill. and southward. July–Sept.—Fruit $\frac{1}{2}$ ' in diameter.

2. **P. incarnata**, L. Nearly smooth; leaves 3-cleft; the lobes serrate; petiole bearing 2 glands; flower large (2' broad), nearly white, with a triple purple and flesh-colored crown; involucre 3-leaved. — Dry soil, Virginia, Kentucky, and southward. May–July. — Fruit of the size of a hen's egg, oval, called *Maypops*.

ORDER 45. CUCURBITACEÆ. (GOURD FAMILY.)

Mostly succulent herbs with tendrils, dioecious or monoecious (often monopetalous) flowers, the calyx-tube cohering with the 1–3-celled ovary, and the 5 or usually 2½ stamens (i. e. one with a one-celled and 2 with two-celled anthers) commonly united by their often tortuous anthers, and sometimes also by the filaments. Fruit (pepo) fleshy, or sometimes membranaceous. — Limb of the calyx and corolla usually more or less combined. Stigmas 2 or 3. Seeds large, usually flat, anatropous, with no albumen. Cotyledons leaf-like. Leaves alternate, palmately lobed or veined. — Mostly a tropical or subtropical order; represented in cultivation by the GOURD (LAGENARIA VULGARIS), PUMPKIN and SQUASH (species of CUCURBITA), MUSKMELON (CUCUMIS MELO), CUCUMBER (C. SATIVUS), and WATERMELON (CITRULLUS VULGARIS); while as wild plants, there are only the three following:—

1. **Sicyos**. Corolla of the sterile flowers flat and spreading, 5-lobed. Fruit prickly, indehiscent, 1-celled, 1-seeded.
2. **Echinocystis**. Corolla of the sterile flowers flat and spreading, 6-parted. Pod prickly, 2-celled, 4-seeded, bursting at the top.
3. **Melothria**. Corolla of the sterile flowers somewhat campanulate, 5-cleft. Berry smooth, many-seeded.

1. SÍCYOS, L. ONE-SEEDED STAR-CUCUMBER.

Flowers monoecious. Petals 5, united below into a bell-shaped or flattish corolla. Anthers cohering in a mass. Ovary 1-celled, with a single suspended ovule: style slender: stigmas 3. Fruit ovate, dry and indehiscent, filled by the single seed, covered with barbed prickly bristles which are readily detached. — Climbing annuals, with 3-forked tendrils, small whitish flowers; the sterile and fertile mostly from the same axils, the former corymbed, the latter in a capitate cluster, long-peduncled. (Greek name for the Cucumber.)

1. **S. angulatus**, L. Leaves roundish heart-shaped, 5-angled or lobed, the lobes pointed; plant clammy-hairy. — River-banks; and a weed in damp yards. July–Sept.

2. ECHINOCÝSTIS, Torr. & Gray. WILD BALSAM-APPLE.

Flowers monoecious. Petals 6, lanceolate, united at the base into an open spreading corolla. Anthers more or less united. Ovary 2-celled, with 2 erect ovules in each cell: stigma broad. Fruit fleshy, at length dry, clothed with weak prickles, bursting at the summit, 2-celled, 4-seeded, the inner part fibrous-netted. Seeds large, with a thickish hard coat. Tall climbing plants, nearly smooth, with 3-forked tendrils, thin leaves, and very numerous small greenish-

white flowers; the sterile in compound racemes often 1° long, the fruitful in small clusters or solitary, from the same axils. (Name composed of *ἐχίνος*, a *hedgehog*, and *κύστις*, a *bladder*, from the prickly covering of the at length bladdery fruit.)

1. **E. lobata**, Torr. & Gr. Root annual; leaves deeply and sharply 5-lobed; fruit oval (2' long); seeds flat, dark-colored. (*Sicyos lobatus*, *Michx.* *Momórdica echinata*, *Muhl.*) — Rich soil along rivers, W. New England to Wisconsin and Kentucky: also cult. for arbors. July — Oct.

3. MELÔTHRIA, L. MELOTHRIA.

Flowers polygamous or monœcious; the sterile campanulate, the corolla 5-lobed; the fertile with the calyx-tube constricted above the ovary, then campanulate. Anthers more or less united. Berry small, pulpy, filled with many flat and horizontal seeds. — Tendrils simple. Flowers very small. (Altered from *Μήλωθρον*, an ancient name for a sort of white grape.)

1. **M. péndula**, L. Slender, from a perennial root, climbing; leaves small, roundish and heart-shaped, 5-angled or lobed, roughish; sterile flowers few in small racemes; the fertile solitary, greenish or yellowish; berry oval, green. — Copses, Virginia and southward. June — Aug.

ORDER 46. UMBELLIFERÆ. (PARSLEY FAMILY.)

Herbs, with small flowers in umbels (or rarely in heads), the calyx entirely adhering to the 2-celled and 2-ovuled ovary, the 5 petals and 5 stamens inserted on the disk that crowns the ovary and surrounds the base of the 2 styles. Fruit consisting of 2 seed-like dry carpels. Limb of the calyx obsolete, or a mere 5-toothed border. Petals either imbricated in the bud or valvate with the point inflexed. The two carpels (called mericarps) cohering by their inner face (the commissure), when ripe separating from each other and usually suspended from the summit of a slender prolongation of the axis (carpophore): each carpel marked lengthwise with 5 primary ribs, and often with 5 intermediate (secondary) ones; in the interstices or intervals between them are commonly lodged the oil-tubes (vitæ), which are longitudinal canals in the substance of the fruit, containing aromatic oil. (These are best seen in slices made across the fruit.) Seed suspended from the summit of the cell, anatropous, with a minute embryo in hard albumen. — Stems usually hollow. Leaves alternate, mostly compound, the petioles expanded or sheathing at the base: rarely with true stipules. Umbels usually compound; when the secondary ones are termed umbellets: each often subtended by a whorl of bracts (that under the umbel is the involucre; that of the umbellet, involucl). — In many the flowers are *dichogamous*, i. e. the styles are protruded from the bud some time before the anthers develop, — an arrangement for cross-fertilization. — A large family, some of the plants innocent and aromatic, others with very poisonous (acid-narcotic) properties; the flowers much alike in all,

—therefore to be studied by their fruits, inflorescence, &c., which likewise exhibit comparatively small diversity. The family is therefore difficult for the young student.

I. Inner face of each seed flat or nearly so (not hollowed out).

- * Umbels or heads simple or irregularly compound, sometimes proliferous (i. e. one from the summit of another).
1. **Hydrocotyle.** Fruit smooth, orbicular or shield-shaped, flattened laterally. Leaves with an orbicular or roundish blade.
 2. **Crantzia.** Fruit smooth, globular, corky. Leaves are thread-shaped or awl-shaped petioles, with no true blade.
 3. **Sanicula.** Fruit clothed with hooked prickles, globular. Flowers polygamous, capitate in the umbellets.
 4. **Eryngium.** Fruit clothed with appressed scales or tubercles, top-shaped. Flowers perfect, in dense heads.
 - * * Umbels compound and perfect ; i. e. its rays regularly bearing umbellets.
 - ← Fruit beset with bristly prickles, not flat.
 5. **Daucus.** Fruit beset with weak prickles in single rows on the ribs.
 - ← ← Fruit smooth, strongly flattened on the back, and single-winged or margined at the junction of the 2 carpels (next to the commissure).
 6. **Polytaenia.** Fruit surrounded with a broad and tumid corky margin thicker than the fruit itself, which is nearly ribless on the back.
 7. **Heracleum.** Fruit broadly wing-margined : the carpels minutely 5-ribbed on the back : lateral ribs close to the margin. Flowers white, the marginal ones radiant.
 8. **Pastinaca.** Fruit wing-margined : ribs of the carpels as in No. 7. Flowers yellow, the marginal ones perfect, not radiant.
 9. **Archemora.** Fruit broadly winged : the 5 ribs on the back equidistant ; the 2 lateral ones close to the wing. Flowers white. Leaves pinnate or 3-foliolate.
 10. **Tiedemannia.** Fruit winged, much as in No. 9. Leaves simple, long and cylindrical, hollow, with some cross partitions.
 - ← ← ← Fruit smooth, flat or flattish on the back, and double-winged or margined at the edge, each carpel also 3-ribbed or sometimes 3-winged on the back.
 11. **Angelica.** Carpels with 3 slender ribs on the back ; a single oil-tube in each interval. Seed not loose.
 12. **Archangelica.** Carpels with 3 rather stout ribs on the back, and 2-3 or more oil-tubes in each interval adhering to the loose seed.
 13. **Conioselinum.** Carpels with 3 wings on the back narrower than those of the margins.
 - ← ← ← ← Fruit smooth, not flattened either way, or only slightly so, the cross-section nearly orbicular or quadrate ; the carpels each with 5 wings or strong ribs.
 14. **Æthusa.** Fruit ovate-globose : carpels with 5 sharply keeled ridges, and with single oil-tubes in the intervals.
 15. **Ligusticum.** Fruit elliptical : carpels with 5 sharp almost winged ridges, and with several oil-tubes in each interval.
 16. **Thasptum.** Fruit elliptical or ovoid : carpels 5-winged or 5-ribbed, and with single oil-tubes in each interval. Flowers yellow or dark purple.
 - ← ← ← ← ← Fruit smooth, flattened laterally or contracted at the sides, wingless.
 17. **Zizia.** Flowers yellow. Fruit oval, somewhat twin : the carpels narrowly 5-ribbed : oil-tubes 3 in each interval. Leaves compound.
 18. **Bupleurum.** Flowers yellow. Fruit ovoid-oblong : the carpels somewhat 5-ribbed. Leaves all simple.
 19. **Discopleura.** Flowers white. Fruit ovoid : the lateral ribs united with a thick corky margin. Leaves cut into capillary divisions.

20. **Cicuta.** Flowers white. Fruit subglobose, twin: the carpels strongly and equally 5-ribbed. Leaves twice or thrice ternate.
21. **Stum.** Flowers white. Fruit ovate or globular: the carpels 5-ribbed. Leaves all simply pinnate.
22. **Cryptotaenia.** Flowers white. Fruit oblong. Leaves 3-parted. Umbel irregular.

II. Inner face of the seed hollowed out lengthwise, or the margins involute, so that the cross-section is semilunar. (Umbels compound.)

23. **Chærophyllum.** Fruit linear or oblong, narrowed or beaked at the apex.
24. **Osmorrhiza.** Fruit linear-club-shaped, tapering below: ribs bristly.
25. **Conium.** Fruit ovate, flattened at the sides: ribs prominent, wavy.
26. **Eulophus.** Fruit ovoid, somewhat twin, nearly destitute of ribs.

III. Inner face of the seed hollowed in the middle, or curved inwards at the top and bottom, so that the section lengthwise is semilunar.

27. **Erigenia.** Fruit twin: carpels nearly kidney-form. Umbellets few-flowered.

1. HYDROCÓTYLE, Tourn. WATER PENNYWORT.

Calyx-teeth obsolete. Fruit flattened laterally, orbicular or shield-shaped; the carpels 5-ribbed, two of the ribs enlarged and often forming a thickened margin: oil-tubes none. — Low, mostly smooth, marsh or aquatic perennials, with slender creeping stems, and round shield-shaped or kidney-form leaves, with scale-like stipules. Flowers small, white, in simple umbels or clusters, which are either single or proliferous, appearing all summer. (Name from ὕδωρ, *water*, and κοτύλη, *a flat cup*, the peltate leaves of several species being somewhat cup-shaped.)

* *Peduncles much shorter than the petioles: pedicels short or none: leaves not peltate.*

1. **H. repánda**, Pers. Petioles (2'–9' long) and peduncles (1'–2' long) clustered on the creeping stems or runners; *leaves ovate-heart-shaped* with a shallow open sinus, *repand-toothed*, thickish; flowers 2–4 in a head or cluster, with a conspicuous 2-leaved involucre; ripe fruit ribbed, reticulated between the ribs. — Maryland (*W. M. Canby*) and southward. — Probably a variety of *H. Asiática*, L.

2. **H. ranunculoides**, L. Petioles (2'–9' long) and peduncles ($\frac{1}{2}$ '–1' long, in fruit reflexed) from long commonly floating creeping stems; *leaves orbicular or kidney-form*, 3–7-cleft, the lobes broad and crenate; flowers 5–10 in a capitate umbel; fruit smooth, scarcely ribbed. — Pennsylvania? Virginia, and southward.

3. **H. Americana**, L. *Stems filiform*, branching, spreading and creeping; *leaves rounded kidney-form*, *crenate-lobed and the lobes crenate*, thin, very smooth and shining, short-petioled; the few-flowered umbels of minute flowers in their axils almost sessile. — Shady damp places: common northward.

* * *Peduncles scape-like, as long as the slender petioles, all from slender runners or rootstocks creeping in the mud: leaves orbicular, centrally peltate, simply or doubly crenate: fruit sharp-margined.*

4. **H. umbelláta**, L. Umbel many-flowered and simple or sometimes proliferous (2 or 3, above one another); pedicels slender $1\frac{1}{2}$ "–3" long; fruit

notched at base and apex. — Massachusetts on the coast, to Penn. (on the Juniata River, *Prof. Porter*), and southward. — Petioles and peduncles 3'–8'' high; leaves 1'–2' wide.

Var. ? **ambigua**. Umbels 3–4; pedicels only once or twice the length of the fruit. — Maryland, *W. M. Canby*. Intermediate between *H. umbellata* and *H. vulgaris*: differs from the next by the distinctly pedicelled fruit.

5. **H. interrúpta**, Muhl. Umbels or rather little heads few-flowered, proliferous and forming an interrupted spike; pedicels scarcely any, the broadly margined fruit acutish at the base. — Massachusetts to Virginia and southward, along the coast. — Usually smaller than No. 4.

2. CRÁNTZIA, Nutt. CRANTZIA.

Calyx-teeth obsolete. Fruit globose; the carpels corky, 5-ribbed: an oil-tube in each interval. — Minute perennials, creeping and rooting in the mud, like *Hydrocotyle*, but with fleshy and hollow cylindrical or awl-shaped petioles, in place of leaves, marked with cross divisions. Umbels few-flowered, simple. Flowers white. (Named for *Prof. Henry John Crantz*, an Austrian botanist of the 18th century.)

1. **C. lineata**, Nutt. Leaves somewhat club-shaped, very obtuse (1'–2' long); lateral ribs of the fruit projecting, forming a corky margin. — Brackish marshes, from Massachusetts southward along the coast. July.

3. SANÍCULA, Tourn. SANICLE. BLACK SNAKEROOT.

Calyx-teeth manifest, persistent. Fruit globular; the carpels not separating spontaneously, ribless, thickly clothed with hooked prickles, each with 5 oil-tubes. — Perennial rather tall herbs, with palmately-lobed or parted leaves, those from the root long-petioled. Umbels irregular or compound, the flowers (greenish or yellowish) capitate in the umbellets, perfect, and with staminate ones intermixed. Involucre and involuclers few-leaved. (Name from *sano*, to heal.)

1. **S. Canadénsis**, L. Leaves 3–5- (the upper only 3-) parted; *sterile flowers few, scarcely pedicelled*, shorter than the fertile ones; *styles shorter than the prickles of the fruit*. — Copses. June–Aug. — Plant 1°–2° high, with thin leaves; their divisions wedge-obovate or oblong, sharply cut and serrate, the lateral mostly 2-lobed. Fruits few in each umbellet.

2. **S. Marilándica**, L. Leaves all 5–7-parted; *sterile flowers numerous, on slender pedicels*, about the length of the fertile; *styles elongated and conspicuous, recurved*. — Woods and copses: common. — Stem 2°–3° high; the leaves more rigid and with narrower divisions than in the former, with almost cartilaginous teeth. Fruits several in each umbellet.

4. ERÝNGIUM, Tourn. ERYNGO.

Calyx-teeth manifest, persistent. Styles slender. Fruit top-shaped, covered with little scales or tubercles, with no ribs, and scarcely any oil-tubes. — Chiefly perennials, with coriaceous, toothed, cut, or prickly leaves, and blue or white bracted flowers closely sessile in dense heads. (A name used by *Dioscorides*, of uncertain origin.)

1. **E. yuccæfolium**, Michx. (RATTLESNAKE-MASTER. BUTTON SNAKE-ROOT.) *Leaves* linear, taper-pointed, rigid, grass-like, nerved, bristly-fringed; leaflets of the involucre mostly entire and shorter than the heads; root perennial. (*E. aquaticum*, L., in part, but never aquatic.) — Dry or damp pine-barrens or prairies, New Jersey to Wisconsin, and southward. July, Aug.

2. **E. Virginianum**, Lam. *Leaves* linear-lanceolate, serrate with hooked or somewhat spiny teeth, veiny; leaflets of the involucre cleft or spiny-toothed, longer than the cymose whitish or bluish heads; root biennial. — Swamps, New Jersey and southward near the coast. July.

5. DAÛCUS, Tourn. CARROT.

Calyx 5-toothed. Corolla irregular. Fruit ovoid or oblong; the carpels scarcely flattened on the back, with 5 primary slender bristly ribs, two of them on the inner face, also with 4 equal and more or less winged secondary ones, each bearing a single row of slender bristly prickles: an oil-tube under each of these ribs. — Biennials, with finely 2-3-pinnate or pinnatifid leaves, cleft involucre, and concave umbels, dense in fruit. (The ancient Greek name.)

1. **D. CAROTA**, L. (COMMON CARROT.) Stem bristly; involucre pinnatifid, equalling the umbel. — Spontaneous in old fields. July-Sept. — Flowers white or cream-color, the central one of each umbellet abortive and dark-purple. Umbel in fruit dense and concave, resembling a bird's nest. (Adv. from Eu.)

6. POLYTÆNIA, DC. POLYTÆNIA.

Calyx 5-toothed. Fruit oval, very flat, with an entire broad and thick corky margin, the impressed back very obscurely ribbed: oil-tubes 2 in each interval, and many in the corky margin. — A smooth herb (2°-3° high), resembling a Parsnip, with twice-pinnate leaves, the uppermost opposite and 3-cleft, no involucre, bristly involucre, and bright yellow flowers. (Name from *πολύς*, many, and *ταβία*, a fillet, alluding to the numerous oil-tubes.)

1. **P. Nuttallii**, DC. — Barrens, Michigan, Wisconsin, and southwestward. May.

7. HERACLËUM, L. COW-PARSNIP.

Calyx-teeth minute. Fruit as in *Pastinaca*, but the oil-tubes shorter than the carpels (reaching from the summit to the middle). Petals (white) inversely heart-shaped, those of the outer flowers commonly larger and radiant, appearing 2-cleft. — Stout perennials, with broad sheathing petioles and large flat umbels. Involucre deciduous: involucre many-leaved. (Dedicated to *Hercules*.)

1. **H. lanatum**, Michx. Woolly; stem grooved; leaves 1-2-ternately compound; leaflets somewhat heart-shaped; fruit obovate or orbicular. — Moist rich ground: most common northward. June. — A very large, strong-scented plant, 4°-8° high, in some places wrongly called *Masterwort*.

8. PASTINÀCA, Tourn. PARSNIP.

Calyx-teeth obsolete. Fruit oval, flat, with a thin single-winged margin; the carpels minutely 5-ribbed; three of the ribs equidistant on the back, the lateral

ones distant from them and near the margin: an oil-tube in each interval, of the length of the fruit. Petals yellow, roundish, entire; none of the flowers radiant. — Chiefly biennials, with spindle-shaped roots, and pinnately-compound leaves. Involucre and involucels small or none. (The Latin name, from *pastus*, food.)

1. **P. SATIVA**, L. (COMMON PARSNIP.) Stem grooved, smooth; leaflets ovate or oblong, obtuse, cut-toothed, somewhat shining above. — Fields, &c. July. (Adv. from Eu.)

9. ARCHÉMORA, DC. COWBANE.

Calyx 5-toothed. Fruit with a broad single-winged margin, oval, flattish; the carpels with 5 obtuse and approximated equidistant ribs on the convex back: oil-tubes one in each interval, and 4–6 on the inner face. — Smooth perennials, with rather rigid leaves of 3–9 lanceolate or linear leaflets. Involucre nearly none: involucels of numerous small leaflets. Flowers white. (Name applied to this poisonous umbelliferous plant in fanciful allusion to *Archemorus*, who is said to have died from eating parsley. DC.)

1. **A. rigida**, DC. Leaves simply pinnate; leaflets 3–9, varying from lanceolate to ovate-oblong, entire or remotely toothed, or, in Var. *AMBIGUA*, linear, long and narrow. — Sandy swamps, New Jersey and W. New York to Michigan, Illinois, and southward. Aug. — Stem 2°–5° high.

10. TIEDEMÁNIA, DC. FALSE WATER-DROPWORT.

Calyx 5-toothed. Fruit with a single-winged margin, obovate, flattish; the carpels with 5 equidistant slender ribs on the convex back: oil-tubes one in each interval, and 2 on the inner face. — A smooth and erect aquatic herb, with a hollow stem (2°–6° high), and cylindrical pointed and hollow petioles (the cavity divided by cross partitions) in place of leaves. Involucre and involucels of few subulate leaflets. Flowers white. (Dedicated to the distinguished anatomist the late *Prof. Tiedemann*, of Heidelberg.)

1. **T. teretifolia**, DC. Virginia (Harper's Ferry) and southward. Aug.

11. ANGÉLICA, L. ANGELICA.

Calyx-teeth obsolete. Fruit flattened, with a double-winged margin at the commissure; i. e. the lateral rib of each oval carpel expanded into a wing; their flattish backs each strongly 3-ribbed: an oil-tube in each interval, and 2–4 on the inner face. Seed adherent to the pericarp. — Stout perennial herbs, more or less aromatic, with first ternately, then once or twice pinnately or ternately divided leaves, toothed and cut ovate or oblong leaflets, large terminal umbels, scanty or no involucre, and small many-leaved involucels. Flowers white or greenish. Petioles membranaceous at the base. (Named *angelic* from its cordial and medicinal properties.)

1. **A. Curtisii**, Buckley. Nearly glabrous; leaves twice ternate or the divisions quinate; leaflets thin, ovate or ovate-lanceolate, pointed, sharply cut and toothed; involucels of small subulate leaflets; wings of the fruit broad. — Mountains of Pennsylvania (*Prof. Porter*), Virginia, and southward in the Alleghanies. Aug.

12. **ARCHANGÉLICA**, Hoffm. ARCHANGELICA.

Calyx-teeth short. Seed becoming loose in the pericarp, coated with numerous oil-tubes which adhere to its surface. Otherwise as in *Angelica*, from which the species have been separated, with hardly sufficient reason.

1. **A. hirsuta**, Torr. & Gr. *Woolly or downy* at the top (2° – 5° high), *rather slender*; leaves twice pinnately or ternately divided; leaflets thickish, ovate-oblong, often blunt, serrate; involuclers as long as the umbellets; peduncles and *fruit downy, broadly winged*. (*Angélica triquinata*, Nutt.)—Dry open woods, New York to Michigan, and southward. July. — Flowers white.

2. **A. atropurpurea**, Hoffm. (GREAT ANGELICA.) *Smooth*; stem dark purple, *very stout* (4° – 6° high), hollow; leaves 2–3-ternately compound; the leaflets pinnate, 5–7, sharply cut serrate, acute, pale beneath; petioles much inflated; involuclers very short; *fruit smooth, winged*. (*Angélica triquinata*, Michx.)—Low river-banks, New England to Penn., Wisconsin, and northward. June. — Flowers greenish-white. Plant strong-scented.

3. **A. Gmélini**, DC. Stem a little downy at the summit (1° – 3° high); leaves 2–3-ternately divided; the leaflets ovate, acute, cut-serrate, glabrous; involuclers about as long as the umbellets; *fruit oblong with 5 thick and corky wing-like ribs* to each carpel, *the marginal ones little broader than the others*. (*A. peregrina*, Nutt., & ed. 2.)—Rocky coast of Massachusetts Bay and northward. July. — Flowers greenish-white. Plant little aromatic. Fruit so thick and so equally ribbed, rather than winged, that it might be taken for a *Ligusticum*.

13. **CONIOSELINUM**, Fischer. HEMLOCK-PARSLEY.

Calyx-teeth obsolete. Fruit oval: the carpels convex-flattish and narrowly 3-winged on the back, and each more broadly winged at the margins: oil-tubes in the substance of the pericarp, 1–3 in each of the intervals, and several on the inner face. — Smooth perennials, with finely 2–3-pinnately compound thin leaves, inflated petioles, and white flowers. Involucre scarcely any: leaflets of the involuclers awl-shaped. (Name compounded of *Conium*, the Hemlock, and *Selinum*, Milk-Parsley, from its resemblance to these two genera.)

1. **C. Canadense**, Torr. & Gr. Leaflets pinnatifid; fruit longer than the pedicels. — Swamps, Vermont to Wisconsin northward, and southward through the Alleghanies. Aug. — Herbage resembling the Poison Hemlock.

14. **ÆTHÛSA**, L. FOOL'S PARSLEY.

Calyx-teeth obsolete. Fruit ovate-globose; the carpels each with 5 thick sharply-keeled ridges: intervals with single oil-tubes. — Annual, poisonous herbs, with 2–3-ternately compound and many-cleft leaves, the divisions pinnate, and white flowers. (Name from *αἶθω*, to burn, from the acrid taste.)

1. **Æ. CYNAPIUM**, L. Divisions of the leaves wedge-lanceolate; involucre none: involuclers 3-leaved, long and narrow. — About cultivated grounds, New England to Penn. July. — A fetid, poisonous herb, with much the aspect of Poison Hemlock, but with dark-green foliage, long hanging involuclers, and unspotted stem. (Adv. from Eu.)

15. **LIGŪSTICUM**, L. **LOVAGE.**

Calyx-teeth small or minute. Fruit elliptical, round on the cross-section, or slightly flattened on the sides; the carpels each with 5 sharp and projecting or narrowly winged ridges: intervals and inner face with many oil-tubes. — Perennials, with aromatic roots and fruit, 2-3-ternately compound leaves, and white flowers. (Named from the country *Liguria*, where the officinal *Lovage* of the gardens, *L. Levisticum*, abounds.)

1. **L. Scoticum**, L. (SCOTCH LOVAGE.) Very smooth; stem (2° high), simple; leaves 2-ternate; leaflets rhombic-ovate, coarsely toothed or cut; leaflets of the involucre and involucels linear; calyx-teeth distinct; fruit narrowly oblong. — Salt marshes, from Rhode Island northward. Aug. — (Eu.)

2. **L. actæifolium**, Michx. (NONDO. ANGELICO.) Smooth; stem (3°-6° high) branched above; the numerous umbels forming a loose and naked somewhat whorled panicle, the lateral ones mostly barren; leaves 3-ternate; leaflets broadly ovate, equally serrate, the end ones often 3-parted; calyx-teeth minute; ribs of the short fruit wing-like. — Rich woods, Virginia, Kentucky, and southward along the mountains. July, Aug. — Root large, with the strong aromatic odor and taste of *Angelica*. (Michaux's habitat, "Banks of the St. Lawrence," is probably a mistake.)

16. **THÁSPIUM**, Nutt. **MEADOW-PARSNIP.**

Calyx-teeth obsolete or short. Fruit ovoid or oblong, somewhat flattish or contracted at the sides (the cross-section of each seed orbicular and somewhat angled or 5-angular); the carpels each with 5 strong and equal ribs or wings, the lateral ones marginal: oil-tubes single in each interval. — Perennial herbs, with 1-2-ternately divided leaves (or the root-leaves simple), umbels with no involucre, minute few-leaved involucels, and yellow or sometimes dark-purple flowers. (Name a play upon *Thapsia*, a genus so called from the island of *Thapsus*.) I include in this genus *Zizia*, Koch, — because the same species has fruit either ribbed or winged, — and retain the name of *Zizia* for *Z. integerrima*, DC.

* Stems loosely branched, 2°-5° high, mostly pubescent on the joints: calyx short but manifest: corolla light yellow: leaves all ternately compound.

1. **T. barbinode**, Nutt. Leaves 1-3-ternate; leaflets ovate or lance-ovate and acute, mostly with a wedge-shaped base, above deeply cut-serrate, often 2-3-cleft or parted, the terminal one long-stalked (1'-2' long); fruit oblong, 6-10-winged (3" long), some of the dorsal wings often narrow or obsolete. — River-banks, W. New York to Wisconsin, and southward. July.

2. **T. pinnatifidum**, Gray. Branchlets, umbels, &c. roughish-puberulent; leaves 1-3-ternate; leaflets 1-2-pinnatifid, the lobes linear or oblong; fruit oblong, narrowly 8-10-winged (1½' long), the intervals minutely scabrous. (*Zizia pinnatifida*, Buckley. *Thaspium Walteri*, Shuttlew., excl. syn. Walt.) — Barrens of Kentucky (*Short*), and southward in the mountains.

** Stems somewhat branched; the whole plant glabrous: calyx-teeth obscure.

3. **T. aureum**, Nutt. Leaves all 1-2-ternately divided or parted (or rarely some of the root-leaves simple and heart-shaped); the divisions or leaflets oblong-

lanceolate, very sharply cut-serrate, with a wedge-shaped entire base; flowers deep yellow; fruit oblong-oval, with 10 winged ridges. — Moist river-banks, &c. June.

Var. **ápterum**. Fruit with strong and sharp ribs in place of wings. (*Smyrniium aureum*, L. *Zizia aurea*, Koch.) — With the winged form.

4. **T. trifoliátum**. *Root-leaves* or some of them round and *heart-shaped; stem-leaves simply ternate or quinate, or 3-parted; the divisions or leaflets ovate-lanceolate or roundish, mostly abrupt or heart-shaped at the base, crenately toothed; flowers deep yellow; fruit globose-ovoid, with 10 winged ridges.* — Rocky thickets, Vermont to Wisconsin, and southward: rare eastward. June.

Var. **atropurpúreum**, Torr. & Gr. Petals deep dark-purple. (*Thápsia trifoliata*, L. *Smyrniium cordatum*, Walt. *Thaspium atropurpureum*, Nutt.) — From New York westward and southward.

Var. **ápterum**. Petals yellow: fruit with sharp ribs in place of wings. (*Zizia cordata*, Koch, Torr.) With the preceding form.

17. ZÍZIA, DC. partly. (ZIZIA § TÆNÍDIA, Torr. & Gr.)

Calyx-teeth obsolete. Fruit ovoid-oblong, contracted at the junction of the carpels so as to become twin, the cross-section of each seed nearly orbicular: carpels somewhat fleshy when fresh, with 5 slender ribs (which are more conspicuous when dry): oil-tubes 3 in each interval and 4 on the inner face. — A perennial smooth and glaucous slender herb (2°–3° high), with 2–3-ternately compound leaves, the leaflets with entire margins; umbels with long and slender rays, no involucre, and hardly any involucels. Flowers yellow. (Named for *I. B. Ziz*, a Rhenish botanist.)

1. **Z. integrérrima**, DC. — Rocky hillsides: not rare. May, June.

18. BUPLEÛRUM, Tourn. THOROUGH-WAX.

Calyx-teeth obsolete. Fruit ovate-oblong, flattened laterally or somewhat twin, the carpels 5-ribbed, with or without oil-tubes. Plants with simple entire leaves and yellow flowers. (Name from βούρ, an ox, and πλευρόν, a rib.)

1. **B. rotundifólium**, L. Leaves ovate, perfoliate; involucre none; involucels of 5 ovate leaflets. — Fields, New York to Virginia: rare. (Adv. from Eu.)

19. DISCOPLÉÛRA, DC. MOCK BISHOP-WEED.

Calyx-teeth awl-shaped. Fruit ovoid; the carpels each with 3 strong ribs on the back, and 2 broad lateral ones united with a thickened corky margin: intervals with single oil-tubes. — Smooth and slender branched annuals, with the leaves finely dissected into bristle-form divisions, and white flowers. Involucre and involucels conspicuous. (Name from δίσκος, a disk, and πλευρόν, a rib.)

1. **D. capillácea**, DC. Umbel few-rayed; leaflets of the involucre 3–5-cleft; involucels longer than the umbellets; fruit ovate in outline. — Brackish swamps, Massachusetts to Virginia, and southward. July–Oct.

Var. ? **costáta**, DC. Larger; rays and divisions of the involucre numerous; ribs of the fruit stronger. — S. Illinois (*Vasey*) and southwestward.

2. **D. Nuttállii**, DC. Umbel many-rayed; leaflets of the involucre entire and shorter; fruit globular. — Wet prairies, Kentucky and southward.

20. CÍCÛTA, L. WATER-HEMLOCK.

Calyx minutely 5-toothed. Fruit subglobose, a little contracted at the sides, the carpels with 5 flattish and strong ribs: intervals with single oil-tubes. — Marsh perennials, very poisonous, smooth, with thrice pinnately or ternately compound leaves, the veins of the lanceolate or oblong leaflets terminating in the notches. Involucre few-leaved: involucels many-leaved. Flowers white; in summer. (The ancient Latin name of the Hemlock.)

1. **C. maculàta**, L. (SPOTTED COWBANE. MUSQUASH ROOT. BEAVER-POISON.) Stem streaked with purple, stout; leaflets oblong-lanceolate, coarsely serrate, sometimes lobed, pointed. — Swamps: common. Aug. — Plant 3' – 6° high, coarse; the root a deadly poison.

2. **C. bulbifera**, L. Leaflets linear, remotely toothed or cut-lobed; upper axils bearing clustered bulblets. — Swamps: common northward: seldom fruiting.

21. SÌUM, L. WATER-PARSNIP.

Calyx-teeth small or none. Fruit oblong, ovate or globular, flattish or contracted at the sides; the carpels with 5 mostly strong ribs: intervals with 1 – several oil-tubes. — Marsh or aquatic perennials, smooth, poisonous, with sharp-angled or grooved stems, simply pinnate leaves, and serrate or incised leaflets, or the immersed leaves more compound. Involucre several-leaved. Flowers white; in summer. (Name supposed to be from the Celtic *siu*, water.)

* *Pericarp thin between the strong projecting ribs: lateral ribs marginal.*

1. **S. lineàre**, Michx. Tall (2° – 4°); leaflets linear, lanceolate, or oblong-lanceolate, tapering to a sharp point, very sharply serrate; calyx-teeth none or obsolete; fruit globular, with corky and very salient ribs, or rather wings; oil-tubes 1 – 3 in the very narrow intervals. (*S. cicutæfolium*, Gmelin, of Siberia is probably the same, and if so, the older name.) — In water or wet places: common.

S. LATIFOLIUM, L., of Europe (for which broad-leaved forms of our plant have been mistaken), if ever found in this country, will be known by its blunt or less tapering leaves, slender calyx-teeth, and much less projecting or wing-like ribs to the more oval fruit.

2. **S. Carsonii**, Durand, ined. Smaller (1° – 2° high), branched; leaflets 3 – 7, thin, varying from linear to oblong, acute, sharply serrate; calyx-teeth none or obsolete; fruit ovoid-globular, with strong filiform ribs, the broad intervals with 1 – 3 conspicuous oil-tubes. — Wet banks of streams, or in flowing water (when the submersed or floating leaflets are very thin, sometimes reduced to the terminal one, ovate or oblong, and usually laciniate-toothed or dissected: in this state it was doubtfully referred to *Helosciadium nodiflorum*, Koch, in addend. to ed. 2). — Pennsylvania, around the Pocono Mountain, Prof. Traill Green, Prof. T. C. Porter, Dr. Carson. Connecticut, Prof. D. C. Eaton.

* *Pericarp of a thick texture, concealing the oil-tubes: ribs inconspicuous, the lateral ones not quite marginal.* (Bérula, Koch.)

2. **S. angustifolium**, L. Low (9' – 20' high); leaflets varying from oblong to linear, mostly cut-toothed and cleft or even pinnatifid; peduncles short; fruit ovate; calyx-teeth scarcely any. — Wet places, Massachusetts (*M. A. Curtis* in herb Durand), Michigan, Illinois (*Dr. Vasey*), and westward. (Eu.)

22. CRYPTOTÆNIA, DC. HONEWORT.

Calyx-teeth obsolete. Fruit oblong, contracted at the sides; the carpels equally and obtusely 5-ribbed: oil-tubes very slender, one in each interval and one under each rib. Seed slightly concave on the inner face. — A perennial smooth herb, with thin 3-foliolate leaves, the umbels and umbellets with very unequal rays, no involucre, and few-leaved involucels. Flowers white. (Name composed of *κρυπτός*, *hidden*, and *ταινία*, *a fillet*, from the concealed oil-tubes.)

1. **C. Canadensis**, DC. — Thickets: common. June – Sept. — Plant 2° high. Leaflets large, ovate, pointed, doubly serrate, the lower ones lobed.

23. CHÆROPHÝLLUM, L. CHERVIL.

Calyx-teeth obsolete. Fruit linear or oblong, pointed or short-beaked, contracted at the sides; the carpels 5-ribbed, at least at the apex: inner face of the seed deeply grooved lengthwise: intervals usually with single oil-tubes. — Chiefly annuals or biennials: leaves ternately decomposed; the leaflets lobed or toothed: involucre scarcely any: involucels many-leaved. Flowers chiefly white. (Name from *χαίρω*, *to gladden*, and *φύλλον*, *a leaf*, alluding to the agreeable aromatic odor of the foliage.)

1. **C. procumbens**, Lam. Stems slender (6'–18'), spreading, a little hairy; lobes of the pinnatifid leaflets obtuse, oblong; umbels few-rayed (sessile or peduncled); fruit narrowly oblong, with conspicuous narrow ribs, not beaked. — Moist copses, New Jersey to Illinois and southward. May, June.

2. **C. SATIVUM**, Lam. (GARDEN CHERVIL.) Taller; lobes of the leaves ovate and pinnatifid; fruit oblong-linear, ribless, but tapering into a ribbed beak shorter than the seed. (*Anthriscus Cerefolium*, *Hoffm.*) — Rarely seen in gardens (as a sweet herb), but thoroughly spontaneous in fields and copses near Lancaster, Penn., *Prof. T. C. Porter*. (Nat. from Eu.)

24. OSMORRHÌZA, Raf. SWEET CICELY.

Calyx-teeth obsolete. Fruit linear-oblong, angled, tapering downwards into a stalk-like base, contracted at the sides, crowned with the styles; the carpels with sharp upwardly bristly ribs: inner face of the nearly terete seed with a deep longitudinal channel: oil-tubes none. — Perennials, with thick sweet-aromatic roots, and large 2–3-ternately compound leaves; the leaflets ovate, pinnatifid-toothed. Involucre and involucels few-leaved. Flowers white. (Name from *ὄσμή*, *a scent*, and *ρίζα*, *a root*, from the anise-like flavor of the latter.)

1. **O. longistylis**, DC. (SMOOTHER SWEET CICELY.) *Styles slender, nearly as long as the ovary; leaflets sparingly pubescent or smooth when old, short-pointed, cut-toothed, sometimes lobed.* — Rich moist woods: commonest northward. May, June. — Plant 3° high, branching: stem reddish.

2. **O. brevistylis**, DC. (HAIRY SWEET CICELY.) *Styles conical and not longer than the breadth of the ovary; fruit somewhat tapering at the summit; leaflets downy-hairy, taper-pointed, pinnatifid-cut.* — Common: root less sweet.

25. CONIUM, L. POISON HEMLOCK.

Calyx-teeth obsolete. Fruit ovate, flattened at the sides, the carpels with 5 prominent wavy ribs, and no oil-tubes: inner face of the seed with a deep nar-

row longitudinal groove. — Biennial poisonous herbs, with large decomposed leaves. Involucre and involucels 3 - 5-leaved, the latter 1-sided. Flowers white. (*Κόνηιον*, the Greek name of the Hemlock, by which criminals and philosophers were put to death at Athens.)

1. *C. MACULATUM*, L. Smooth; stem spotted; leaflets lanceolate, pinnatifid: involucels shorter than the umbellets. — Waste places. July. — A large branching herb: the pale green leaves exhale a disagreeable odor when bruised. A virulent narcotico-acrid poison, used in medicine. (Nat. from Eu.)

26. EULOPHUS, Nutt. EULOPHUS.

Calyx-teeth small. Fruit ovoid, contracted at the sides and somewhat twin; the carpels smooth, indistinctly ribbed, and with a close row of oil-tubes: inner face of the seed longitudinally channelled, the cross-section semilunar. — A slender, smooth perennial: leaves 2-ternately divided into narrow linear leaflets. Involucre scarcely any: involucels bristle-form. Flowers white. (Name from *εὖ*, *well*, and *λόφος*, *a crest*, not well applied to a plant with no crest at all.)

1. *E. Americanus*, Nutt. — Darby plains, near Columbus, Ohio (*Sullivan*), Illinois, and southwestward. July. — Root a cluster of small tubers.

27. ERIGÈNIA, Nutt. HARBINGER-OF-SPRING.

Calyx-teeth obsolete. Petals obovate or spatulate, flat, entire. Fruit twin; the carpels incurved at top and bottom, nearly kidney-form, with 5 very slender ribs, and several small oil-tubes in the interstices: inner face of the seed hollowed into a broad deep cavity. — A small and smooth vernal plant, producing from a deep round tuber a simple stem, bearing one or two 2-3-ternately divided leaves, and a somewhat imperfect and leafy-bracted compound umbel. Flowers few, white. (Name from *ἡριγενής*, *born in the spring*.)

1. *E. bulbosa*, Nutt. — Alluvial soil, Western New York and Penn. to Wisconsin, Kentucky, &c. March, April. — Stem 3' - 9' high.

ORDER 47. ARALIACEÆ. (GINSENG FAMILY.)

Herbs, shrubs, or trees, with much the same characters as Umbelliferæ, but with usually more than 2 styles, and the fruit a few-several-celled drupe. (Albumen mostly fleshy. Petals not inflexed.) — Represented here only by the genus

1. ARÀLIA, Tourn. GINSENG. WILD SARSAPARILLA.

Flowers more or less polygamous. Calyx-tube coherent with the ovary, the teeth very short or almost obsolete. Petals 5, epigynous, oblong or obovate, lightly imbricated in the bud, deciduous. Stamens 5, epigynous, alternate with the petals. Styles 2 - 5, mostly distinct and slender, or in the sterile flowers short and united. Ovary 2 - 5-celled, with a single anatropous ovule suspended from the top of each cell, ripening into a berry-like drupe, with as many seeds as cells. Embryo minute. — Leaves compound or decomposed. Flowers white or greenish, in umbels. Roots (perennial), bark, fruit, &c. warm and aromatic. (Derivation obscure.)

§ 1. **ARALIA**, L. *Flowers monœciously polygamous or perfect, the umbels usually in corymbs or panicles: styles and cells of the (black or dark purple) fruit 5: stems herbaceous or woody: ultimate divisions of the leaves pinnate.*

* *Umbels very numerous in a large compound panicle: leaves very large, quinately or pinnately decomposed.*

1. **A. spinosa**, L. (ANGELICA-TREE. HERCULES' CLUB.) *Shrub, or a low tree; the stout stem and stalks prickly; leaflets ovate, pointed, serrate, pale beneath.* — River-banks, Pennsylvania to Kentucky and southward: common in cultivation. July, August.

2. **A. racemosa**, L. (SPIKENARD.) *Herbaceous; stem widely branched; leaflets heart-ovate, pointed, doubly serrate, slightly downy; umbels racemose; styles united.* — Rich woodlands. July. — Well known for its spicy-aromatic large roots. There are traces of stipules at the dilated base of the leafstalks.

* * *Umbels 2-7, corymbed: stem short, somewhat woody.*

3. **A. hispida**, Michx. (BRISTLY SARSAPARILLA. WILD ELDER.) *Stem (1°-2° high), bristly, leafy, terminating in a peduncle bearing several umbels; leaves twice pinnate; leaflets oblong-ovate, acute, cut-serrate.* — Rocky places: common northward, and southward along the mountains. June.

4. **A. nudicaulis**, L. (WILD SARSAPARILLA.) *Stem scarcely rising out of the ground, smooth, bearing a single long-stalked leaf and a shorter naked scape, with 2-7 umbels; leaflets oblong-ovate or oval, pointed, serrate, 5 on each of the 3 divisions.* — Moist woodlands; with the same range as No. 3. May, June. — The aromatic horizontal roots, which are several feet long, are employed as a substitute for the officinal Sarsaparilla. Leafstalk 1° high.

§ 2. **GINSENG**, Decaisne & Planchon. (Panax, L.) *Flowers diœciously polygamous: styles and cells of the (red or reddish) fruit 2 or 3: stem herbaceous, low, simple, bearing at its summit a whorl of 3 palmately 3-7-foliolate leaves (or perhaps rather a single and sessile twice-compound leaf), and a single umbel on a slender naked peduncle.*

5. **A. quinquefolia**. (GINSENG.) *Root large and spindle-shaped, often forked (4'-9' long, aromatic); stem 1° high; leaflets long-stalked, mostly 5, large and thin, obovate-oblong, pointed; styles mostly 2; fruit bright red.* (Panax quinquefolium, L.) — Rich and cool woods: becoming rare. July.

6. **A. trifolia**. (DWARF GINSENG. GROUND-NUT.) *Root or tuber globular, deep in the ground (pungent to the taste, not aromatic); stems 4'-8' high; leaflets 3-5, sessile at the summit of the leafstalk, narrowly oblong, obtuse; styles usually 3; fruit yellowish.* — Rich woods: common northward. April, May.

HÈDERA HÈLIX, the European IVY, is almost the only other representative of this family in the northern temperate zone.

ORDER 48. CORNACEÆ. (DOGWOOD FAMILY.)

Shrubs or trees (rarely herbaceous), with opposite or alternate simple leaves, the calyx-tube coherent with the 1-2-celled ovary, its limb minute, the petals (valvate in the bud) and as many stamens borne on the margin of an epigynous disk in the perfect flowers; style one; a single anatropous ovule hang-

ing from the top of the cell; the fruit a 1-2-seeded drupe; embryo nearly the length of the albumen, with large and foliaceous cotyledons.—A small family represented by *Cornus*, and by a partly apetalous genus, *Nyssa*. (Bark bitter and tonic.)

1. C. CORNUS, Tourn. CORNEL. DOGWOOD.

Flowers perfect (or in some foreign species diœcious). Calyx minutely 4-toothed. Petals 4, oblong, spreading. Stamens 4: filaments slender. Style slender: stigma terminal, flat or capitate. Drupe small, with a 2-celled and 2-seeded stone.—Leaves opposite (except in one species), entire. Flowers small, in open naked cymes, or in close heads which are surrounded by a corolla-like involucre. (Name from *cornu*, a horn; alluding to the hardness of the wood.)

§ 1. *Flowers greenish, in a head or close cluster, which is surrounded by a large and showy, 4-leaved, corolla-like, white or rarely pinkish involucre: fruit bright red.*

1. **C. Canadensis**, L. (DWARF CORNEL. BUNCH-BERRY.) *Stems low and simple (5'-7' high) from a slender creeping and subterranean rather woody trunk; leaves scarcely petioled, the lower scale-like, the upper crowded into an apparent whorl in sixes or fours, ovate or oval, pointed; leaves of the involucre ovate; fruit globular.—Damp cold woods: common northward. June.*

2. **C. florida**, L. (FLOWERING DOGWOOD.) *Leaves ovate, pointed, acutish at the base; leaves of the involucre inversely heart-shaped or notched (1½' long); fruit oval.—Rocky woods: more common southward. May, June.—Tree 12°-30° high, very showy in flower, scarcely less so in fruit.*

§ 2. *Flowers white, in open and flat spreading cymes: involucre none: fruit spherical.*

* *Leaves all opposite: shrubs.*

3. **C. circinata**, L'Her. (ROUND-LEAVED CORNEL OR DOGWOOD.) *Branches greenish, warty-dotted; leaves round-oval, abruptly pointed, woolly beneath (2'-5' broad); cymes flat; fruit light blue.—Copses; in rich soil. June.—Shrub 6°-10° high. Leaves larger than in any other species.*

4. **C. sericea**, L. (SILKY CORNEL. KINNIKINNIK.) *Branches purplish; the branchlets, stalks, and lower surface of the narrowly ovate or elliptical pointed leaves silky-downy (often rusty), pale and dull; cymes flat, close; calyx-teeth lanceolate; fruit pale blue.—Wet places: common. June.—Shrub 3°-10° high.*

5. **C. stolonifera**, Michx. (RED-OSIER DOGWOOD.) *Branches, especially the osier-like annual shoots, bright red-purple, smooth; leaves ovate, rounded at the base, abruptly short-pointed, roughish with a minute close pubescence on both sides, whitish underneath; cymes small and flat, rather few-flowered, smooth; fruit white or lead-color.—Wet places: common, especially northward. Multiplies freely by prostrate or subterranean suckers, and forms broad clumps, 3°-6° high. June.*

6. **C. asperifolia**, Michx. (ROUGH-LEAVED DOGWOOD.) *Branches brownish; the branchlets, &c. rough-pubescent; leaves oblong or ovate, on very short petioles, pointed, rough with a harsh pubescence above, and downy beneath; calyx-teeth minute.—Dry or sandy soil, Illinois and southward. May, June.*

7. **C. stricta**, Lam. (STIFF CORNEL.) *Branches brownish or reddish, smooth; leaves ovate or ovate-lanceolate, taper-pointed, acutish at the base, glabrous,*

of nearly the same hue both sides; cymes loose, flattish; anthers and fruit pale blue. — Swamps, Virginia and southward. April, May. — Shrub 8°–15° high.

8. **C. paniculata**, L'Her. (PANICLED CORNELL.) Branches gray, smooth; leaves ovate-lanceolate, taper-pointed, acute at the base, whitish beneath but not downy; cymes convex, loose, often paniced; fruit white, depressed-globose. — Thickets and river-banks. June. — Shrub 4°–8° high, very much branched, bearing a profusion of pure white blossoms.

* * Leaves mostly alternate, crowded at the ends of the branches.

9. **C. alternifolia**, L. (ALTERNATE-LEAVED CORNELL.) Branches greenish streaked with white, alternate; leaves ovate or oval, long-pointed, acute at the base, whitish and minutely pubescent underneath; fruit deep blue on reddish stalks. — Hillsides in copses. May, June. — Shrub or tree 8°–20° high, with flattish top, and very open, broad cymes.

2. NYSSA, L. TUPELO. PEPPERIDGE. SOUR-GUM TREE.

Flowers diœciously polygamous, clustered or rarely solitary at the summit of axillary peduncles. Stam. Fl. numerous in a simple or compound dense cluster of fascicles. Calyx small, 5-parted. Petals as in fertile flower or none. Stamens 5–12, oftener 10, inserted on the outside of a convex disk: filaments slender: anthers short. No pistil. Pist. Fl. solitary, or 2–8, sessile in a bracted cluster, much larger than the staminate flowers. Calyx with a very short repand-truncate or minutely 5-toothed limb. Petals very small and fleshy, deciduous, or often wanting. Stamens 5–10, with perfect or imperfect anthers. Style elongated, revolute, stigmatic down one side. Ovary one-celled. Drupe ovoid or oblong, with a bony and grooved or striate 1-celled and 1-seeded stone. — Trees with entire or sometimes angulate-toothed leaves, which are alternate, but mostly crowded at the end of the branchlets, and greenish flowers appearing with the leaves. (The name of a Nymph: “so called because it [the original species] grows in the water.”)

1. **N. multiflora**, Wang. (TUPELO. PEPPERIDGE. BLACK or SOUR GUM.) Leaves oval or obovate, commonly acuminate, glabrous or villous-pubescent when young, at least on the margins and midrib, shining above when old (2'–5' long); fertile flowers 3–8, at the summit of a slender peduncle; fruit ovoid, bluish-black (about $\frac{1}{2}$ ' long). (N. sylvatica, Marsh. N. villosa, Willd., &c.) — Rich soil, either moist or nearly dry, Massachusetts to Illinois, and southward. April, May. — A middle-sized tree, with horizontal branches and a light flat spray, like the Beech: the wood firm, close-grained and very unwedgeable, on account of the oblique direction and crossing of the fibre of different layers. Leaves turning bright crimson in autumn.

2. **N. uniflora**, Walt. (LARGE TUPELO.) Leaves oblong or ovate, sometimes slightly cordate at the base, long-petioled, entire or angulate-toothed, pale and downy-pubescent beneath, at least when young (4'–12' long); fertile flower solitary on a slender peduncle; fruit oblong, blue (1' or more in length). (N. denticulata, Ait. N. tomentosa, and angulisans, Michx. N. grandidentata, Michx. f.) — In water or wet swamps, Virginia, Kentucky, and southward. April. — Wood soft: that of the roots very light and spongy, used for corks.

DIVISION II. MONOPÉTALOUS EXÓGENOUS PLANTS.

Floral envelopes consisting of both calyx and corolla, the latter composed of more or less united petals, that is, monopetalous.*

ORDER 49. CAPRIFOLIACEÆ. (HONEYSUCKLE FAMILY.)

Shrubs, or rarely herbs, with opposite leaves, no (genuine) stipules, the calyx-tube coherent with the 2-5-celled ovary, the stamens as many as (or one fewer than) the lobes of the tubular or wheel-shaped corolla, and inserted on its tube.—Fruit a berry, drupe, or pod, 1-several-seeded. Seeds anatropous, with a small embryo in fleshy albumen.

Tribe I. LONICEREÆ. Corolla tubular, often irregular, sometimes 2-lipped. Style slender: stigma capitate.

1. **Linnæa.** Stamens 4, one fewer than the lobes of the corolla. Fruit dry, 3-celled, but only 1-seeded; two of the cells sterile.
2. **Symphoricarpus.** Stamens 4 or 5, as many as the lobes of the bell-shaped regular corolla. Berry 4-celled, but only 2-seeded; two of the cells sterile.
3. **Lonicera.** Stamens 5, as many as the lobes of the tubular and more or less irregular corolla. Berry several-seeded; all the 2 or 3 cells fertile.
4. **Diervilla.** Stamens 5. Corolla funnel-form, nearly regular. Pod 2-celled, 2-valved, many-seeded, slender.
5. **Triosteum.** Stamens 5. Corolla gibbous at the base. Fruit a 3-celled drupe.

Tribe II. SAMBUCEÆ. Corolla wheel-shaped or urn-shaped, regular, deeply 5-lobed. Stigmas 1-3, rarely 5, sessile. Flowers in broad cymes.

6. **Sambucus.** Fruit berry-like containing the small seed-like nutlets. Leaves pinnate.
7. **Viburnum.** Fruit a 1-celled 1-seeded drupe, with a compressed stone. Leaves simple.

1. LINNÆA, Gronov. LINNÆA. TWIN-FLOWER.

Calyx-teeth 5, awl-shaped, deciduous. Corolla narrow bell-shaped, almost equally 5-lobed. Stamens 4, two of them shorter, inserted toward the base of the corolla. Ovary and the small dry pod 3-celled, but only 1-seeded, two of the cells having only abortive ovules.—A slender creeping and trailing little evergreen, somewhat hairy, with rounded-oval sparingly crenate leaves contracted at the base into short petioles, and thread-like upright peduncles forking into 2 pedicels at the top, each bearing a delicate and fragrant nodding flower. Corolla purple and whitish, hairy inside. (Dedicated to the immortal *Linnaeus*, who first pointed out its characters, and with whom this pretty little plant was a special favorite.)

1. **L. borealis**, Gronov.—Moist mossy woods and cold bogs: common northward, but towards the south of rare occurrence as far as New Jersey, and along the mountains to Maryland. Junc. (Eu.)

* In certain families, such as Ericaceæ, &c., the petals in some genera are nearly or quite separate. In Compositæ and some others, the calyx is mostly reduced to a pappus, or to scales, or a mere border, or even to nothing more than a covering of the surface of the ovary. The student might look for these in the first or the third division. But the *artificial analysis* prefixed to the volume provides for such anomalies, and will lead the student to the proper order.

2. SYMPHORICÁRPUS, Dill. SNOWBERRY.

Calyx-teeth short, persistent on the fruit. Corolla bell-shaped, regularly 4-5-lobed, with as many short stamens inserted into its throat. Ovary 4-celled, only 2 of the cells with a fertile ovule; the berry therefore 4-celled but only 2-seeded. Seeds bony. — Low and branching upright shrubs, with oval short-petioled leaves, which are downy underneath and entire, or wavy-toothed or lobed on the young shoots. Flowers white tinged with rose-color, in close short spikes or clusters. (Name composed of *συμφορέω*, to bear together, and *καρπός*, fruit; from the clustered berries.)

1. **S. occidentális**, R. Brown. (WOLFBERRY.) Flowers in dense terminal and axillary spikes; corolla much bearded within; the *stamens and style protruded*; berries white. — Northern Michigan, Illinois, and westward. — Flowers larger and more funnel-form, and stamens longer, than in the next.

2. **S. racemósus**, Michx. (SNOWBERRY.) *Flowers in a loose and somewhat leafy interrupted spike* at the end of the branches; corolla bearded inside; berries large, bright white. — Rocky banks, W. Vermont to Pennsylvania and Wisconsin: common in cultivation. June-Sept. — Berries ripe in autumn.

Var. **pauciflorus**, Robbins. Low, diffusely branched and spreading; leaves smaller (about 1' long), the spike reduced to one or two flowers in the axils of the uppermost. — Rocky woods of L. Superior, Dr. Robbins, and north-westward. Alleghanies of Pennsylvania, J. R. Lowrie, Mr. Böcking.

3. **S. vulgáris**, Michx. (INDIAN CURRANT. CORAL-BERRY.) *Flowers in small close clusters in the axils of nearly all the leaves*; corolla sparingly bearded; berries small, dark red. — Rocky banks, W. New York and Penn. to Illinois and southward: also cultivated. July.

3. LONICÈRA, L. HONEYSUCKLE. WOODBINE.

Calyx-teeth very short. Corolla tubular or funnel-form, often gibbous at the base, irregularly or almost regularly 5-lobed. Stamens 5. Ovary 2-3-celled. Berry several-seeded. — Leaves entire. Flowers often showy and fragrant. (Named in honor of Adam Lonitzer, latinized *Lonicerus*, a German herbalist of the 16th century.)

§ 1. CAPRIFOLIUM, Juss. *Twining shrubs, with the flowers in sessile whorled clusters from the axils of the (often connate) upper leaves, and forming interrupted terminal spikes: calyx-teeth persistent on the (red or orange) berry.*

* *Corolla trumpet-shaped, almost regularly and equally 5-lobed.*

1. **L. sempervirens**, Ait. (TRUMPET HONEYSUCKLE.) Flowers in somewhat distant whorls; leaves oblong, smooth; the lower petioled, the uppermost pairs united round the stem. — Copses, New York (near the city) to Virginia, and southward: common also in cultivation. May-Oct. — Leaves deciduous at the North. Corolla scentless, nearly 2' long, deep red outside, yellowish within or rarely throughout.

** *Corolla ringent: the lower lip narrow, the upper broad and 4-lobed.*

2. **L. gràta**, Ait. (AMERICAN WOODBINE.) *Leaves smooth, glaucous beneath, obovate, the 2 or 3 upper pairs united; flowers whorled in the axils of*

the uppermost leaves or leaf-like connate bracts; *corolla smooth* (whitish with a purple tube, fading yellowish), not gibbous at the base, fragrant. — Rocky woodlands, New York, Penn., and westward: also cultivated. May.

3. **L. flava**, Sims. (YELLOW HONEYSUCKLE.) *Leaves smooth, very pale and glaucous both sides, thickish*, obovate or oval, the 2-4 upper pairs united into round cup-like disks; flowers in approximate whorls; tube of the smooth (light yellow) *corolla* somewhat gibbous; filaments almost or quite smooth. — Rocky banks. Catskill mountains (*Pursh*), Ohio to Wisconsin (a form with rather short flowers), and southward along the Alleghany Mountains. June.

4. **L. parviflora**, Lam. (SMALL HONEYSUCKLE.) *Leaves smooth, oblong, green above, very glaucous beneath*, the upper pairs united, all closely sessile; flowers in 2 or 3 closely approximate whorls raised on a peduncle; *corolla gibbous at the base, smooth outside (greenish-yellow tinged with dull purple)*, short (9" long); filaments rather hairy below. — Rocky banks, mostly northward. May, June. — Stem commonly bushy, only 2°-4° high.

Var. **Douglasii**. Leaves greener, more or less downy underneath when young, or ciliate; *corolla* crimson or deep dull purple. (*L. Douglasii*, DC.) — Northern Ohio to Wisconsin and northward.

5. **L. hirsuta**, Eaton. (HAIRY HONEYSUCKLE.) *Leaves not glaucous, downy-hairy beneath*, as well as the branches, and slightly so above, veiny, dull, broadly oval; the uppermost united, the lower short-petioled; flowers in approximate whorls; tube of the (orange-yellow) clammy-pubescent *corolla* gibbous at the base, slender. — Damp copses and rocks, Maine to Wisconsin northward. July. — A coarse, large-leaved species.

§ 2. **XYLÓSTEON**, Juss. *Upright bushy shrubs: leaves all distinct at the base: peduncles axillary, single, 2-flowered at the summit; the two berries sometimes united into one: calyx-teeth not persistent.*

* *The two flowers involucrate by 4 conspicuous and broad foliaceous bracts.*

6. **L. involucrata**, Banks. Pubescent, or becoming glabrous; branches 4-angular; leaves (3' - 6' long) ovate-oblong, mostly pointed, petioled, and with a strong midrib, exceeding the peduncle; *corolla* yellowish, viscid-pubescent, cylindraceous (6" - 8" long); ovaries and globose berries distinct. — Deep woods, Lake Superior (*C. G. Loring, Jr., Dr. Robbins*) and westward. June.

** *Bracts (2 or sometimes 4) at the base of the ovaries minute.*

7. **L. ciliata**, Muhl. (FLY-HONEYSUCKLE.) Branches straggling (3° - 5° high); *leaves oblong-ovate, often heart-shaped, petioled, thin, downy beneath; filiform peduncles shorter than the leaves; corolla funnel-form, almost spurred at the base (greenish-yellow, ¾ long), the lobes nearly equal; berries separate (red).* — Rocky woods, Mass. to Penn., Wisconsin, and northward. May.

8. **L. cærulea**, L. (MOUNTAIN F.) Low (1° - 2° high); branches upright; *leaves oval, downy when young; peduncles very short; bracts awl-shaped, longer than the ovaries of the two (yellowish) flowers, which are united into one (blue) berry.* (*Xylósteum villosum, Michx.*) — Mountain woods and bogs, Massachusetts to Wisconsin, northward. May. (Eu.)

9. **L. oblongifolia**, Muhl. (SWAMP F.) Branches upright; *leaves oblong, downy when young, smooth when old; peduncles long and slender; bracts*

minute or deciduous; corolla deeply 2-lipped ($\frac{1}{2}$ long, yellowish-white); berries (purple) united or nearly distinct.—Bogs, Northern New York to Wisconsin. June.—Shrub 2°–5° high. Leaves 2'–3' long.

4. DIERVILLA, Tourn. BUSH-HONEYSUCKLE.

Calyx-tube tapering at the summit; the lobes slender, awl-shaped, persistent. Corolla funnel-form, 5-lobed, almost regular. Stamens 5. Pod ovoid-oblong, pointed, 2-celled, 2-valved, septical, many-seeded.—Low, upright shrubs, with ovate or oblong pointed serrate leaves, and cymosely 3–several-flowered peduncles, from the upper axils, or terminal. (Named in compliment to *M. Dierville*, who brought it from Canada to Tournefort.)

1. *D. trifida*, Mœnch. Leaves oblong-ovate, taper-pointed, petioled; peduncles mostly 3-flowered; pod long-beaked. (*D. Canadensis*, *Muhl.*)—Rocks: common, especially northward. June–Aug.—Flowers honey-color, not showy, like the Japanese species cultivated under the name of WEIGELA.

D. sessilifolia, Buckley, of the mountains of North Carolina, may occur in those of S. W. Virginia.

5. TRIÓSTEUM, L. FEVER-WORT. HORSE-GENTIAN.

Calyx-lobes linear-lanceolate, leaf-like, persistent. Corolla tubular, gibbous at the base, somewhat equally 5-lobed, scarcely longer than the calyx. Stamens 5. Ovary mostly 3-celled, in fruit forming a rather dry drupe, containing as many angled and ribbed 1-seeded bony nutlets.—Coarse, hairy, perennial herbs, leafy to the top; with the ample entire pointed leaves tapering to the base, but connate round the simple stem. Flowers sessile, and solitary or clustered in the axils. (Name from *τρεις*, three, and *ὄστέον*, a bone, alluding to three bony seeds, or rather nutlets.)

1. *T. perfoliatum*, L. Softly hairy (2°–4° high); leaves oval, abruptly narrowed below, downy beneath; flowers brownish-purple, mostly clustered.—Rich woodlands: not rare. June.—Fruit orange-color, $\frac{1}{2}$ long.

2. *T. angustifolium*, L. Smaller; bristly-hairy; leaves lanceolate, tapering to the base; flowers greenish-cream-color, mostly single in the axils.—S. Pennsylvania to Illinois, and southward. May.

6. SĀMBŪCUS, Tourn. ELDER.

Calyx-lobes minute or obsolete. Corolla open urn-shaped, with a broadly spreading 5-cleft limb. Stamens 5. Stigmas 3. Fruit a berry-like juicy drupe, containing 3 small seed-like nutlets.—Shrubby plants, with a rank smell when bruised, pinnate leaves, serrate pointed leaflets, and numerous small and white flowers in compound cymes. (Name from *σαμβύκη*, an ancient musical instrument, supposed to have been made of Elder-wood.)

1. *S. Canadensis*, L. (COMMON ELDER.) Stems scarcely woody (5°–10° high); leaflets 7–11, oblong, mostly smooth, the lower often 3-parted; cymes flat; fruit black-purple.—Rich soil, in open places. June.—Pith white.

2. *S. pūbens*, Michx. (RED-BERRIED ELDER.) Stems woody (2°–18° high), the bark warty; leaflets 5–7, ovate-lanceolate, downy underneath; cymes

panicled, convex or pyramidal; fruit bright red (rarely white). — Rocky woods; chiefly northward, and southward in the mountains. May: the fruit ripening in June. — Pith brown. — Var. with dissected leaves, Lake Superior, *Lewis Foote, Dr. Robbins.*

7. VIBURNUM, L. ARROW-WOOD. LAURESTINUS.

Calyx 5-toothed. Corolla spreading, deeply 5-lobed. Stamens 5. Stigmas 1-3. Fruit a 1-celled, 1-seeded drupe, with soft pulp and a thin-crustaceous (flattened or tumid) stone. — Shrubs, with simple leaves, and white flowers in flat compound cymes. Petioles sometimes bearing little appendages which are evidently stipules. Leaf-buds naked, or with a pair of scales. (The classical Latin name, of unknown meaning.)

§ 1. *Flowers all alike and perfect.*

* *Leaves finely serrate or entire, bright green; veins not prominent: no stipular appendages: whole plant glabrous or with some minute rusty scurf: fruit black or with a blue bloom, sweet; the stone very flat and even, broadly oval or orbicular.*

1. **V. Lentago**, L. (SWEET VIBURNUM. SHEEP-BERRY.) *Leaves ovate, strongly pointed, closely and very sharply serrate; petioles long and margined; cyme sessile; fruit oval, $\frac{1}{2}$ ' or more long, ripe in autumn, edible; tree 15°-30° high.* — Copses, &c. common, especially northward. May, June.

2. **V. prunifolium**, L. (BLACK HAW.) *Leaves oval, obtuse or slightly pointed, finely and sharply serrate, smaller than in the preceding (1'-2' long); fruit similar or rather smaller; cyme sessile.* — Dry copses; Connecticut to Illinois, and common southward. May. — A tall shrub or small tree.

3. **V. nudum**, L. (WITHE-ROD.) *Leaves thickish, oval, oblong or lanceolate, not shining, the margins entire, repand, or crenate; cyme short-peduncled; fruit round-ovoid (3" long).* — Var. 1. **CLAYTONI** has the leaves nearly entire, the veins somewhat prominent underneath, and grows in swamps from Massachusetts, near the coast, to Virginia and southward. Var. 2. **CASSINOIDES** (**V. pyriformium**, Pursh, &c.) has more opaque, often toothed leaves; and grows in cold swamps from Pennsylvania northward. May, June. — Shrub 6°-10° high.

4. **V. obovatum**, Walt. *Leaves obovate or spatulate, obtuse, entire or denticulate, thickish, small (1'-1½' long), shining; cymes sessile, small; fruit ovoid-oblong.* — River-banks, Virginia and southward. May. — Shrub 2°-8° high.

** *Leaves (with base inclined to heart-shaped) coarsely toothed, prominently pinnately veined, the veins straight and simple or sparingly forked: no rusty scurf: fruit small, ovoid, blue or purple; the stone tumid and grooved: cymes peduncled.*

5. **V. dentatum**, L. (ARROW-WOOD.) *Smooth; leaves broadly ovate, very numerous sharp-toothed and strongly veined, on slender petioles; fruit bright blue; the turgid stone deeply excavated on one face; cross section of the seed between kidney- and horseshoe-shaped.* — Wet places, common northward. June. — Shrub 5°-10° high, with ash-colored bark; the pale leaves often with hairy tufts in the axils of the straight veins.

6. **V. molle**, Michx. *Leaves broadly oval, obovate or ovate, scarcely pointed, coarsely crenate or repand-toothed, the lower surface, rather slender petioles, branchlets and cymes soft-downy, the latter with stellate pubescence; fruit oily (Eu-*

gelm.), the stone as in No. 5, but less deeply excavated on the face. (*V. dentatum*, var. *scabrèllum*, Torr. & Gr.) — Rich woods, Kentucky and southward.

7. **V. pubescens**, Pursh. (DOWNY A.) *Leaves ovate or oblong-ovate*, acute or taper-pointed, the veins and teeth fewer and less conspicuous than in No. 5, the lower surface and very short petioles soft-downy, at least when young; fruit dark-purple; the stone plano-convex and 3-4-grooved on the flat face. — Rocks, &c., W. Vermont to New Jersey, Kentucky, Wisconsin, and northward. June. — A low, straggling shrub.

* * * *Leaves coarsely toothed and somewhat 3-lobed, roundish, the base mostly truncate or somewhat heart-shaped, 3-5-ribbed from the base, the ribs and veins prominent beneath: stipular appendages bristle-shaped: cymes small, slender-peduncled: fruit red; the stone flattened.*

8. **V. acerifolium**, L. (MAPLE-LEAVED A. DOCKMACKIE.) *Leaves soft-downy beneath, 3-ribbed, the pointed lobes diverging, unequally toothed; stamens exserted; fruit crimson turning purple; the lenticular stone undulately 2-grooved on one face and 3-grooved on the other.* — Rocky woods: common. May, June. Shrub 3°-6° high.

9. **V. pauciflorum**, Pylaie. *Leaves glabrous or loosely pubescent beneath, 5-ribbed at base, unequally serrate nearly all round, with 3 short lobes at the summit; cyme few-flowered; stamens shorter than the corolla; fruit red, sour, globular; the stone very flat and even.* (*V. Oxycoccus*, var. *eradiatum*, Oakes.) — Cold woods, Northern N. Hampshire, New York, Wisconsin, and northward. — A low straggling shrub, most related to the next; the leaf-buds similar.

§ 2. *Marginal flowers of the cyme destitute of stamens and pistils, and with corollas many times larger than the others, forming a kind of ray, as in Hydrangea: stipular appendages conspicuous on the petiole.*

* *Fruit spherical, pleasantly acid, bright red; the stone very flat, smooth and even, nearly orbicular: leaf-buds enclosed in one or two pairs of scales.*

10. **V. Ópulus**, L. (CRANBERRY-TREE.) Nearly smooth, upright (5°-10° high); leaves 3-5-ribbed, strongly 3-lobed, broadly wedge-shaped or truncate at the base, the spreading lobes pointed, mostly toothed on the sides, entire in the sinuses; petioles bearing 2 glands at the apex, cymes peduncled. (*V. Oxycoccus* and *V. édule*, Pursh.) — Low grounds, along streams: common northward, and southward in the Alleghanies to the borders of Maryland. June, July. — The acid fruit is a substitute for cranberries, whence the name *High Cranberry-bush*, &c. — The well-known SNOW-BALL TREE, or GUELDER-ROSE, is a cultivated state, with the whole cyme turned into showy sterile flowers. (Eu.)

* * *Fruit ovoid, red, turning darker; the stone tumid, 6-grooved: buds wholly naked.*

11. **V. lantanoides**, Michx. (HOBBLE-BUSH. AMERICAN WAYFARING-TREE.) *Leaves round-ovate, abruptly pointed, heart-shaped at the base, closely serrate, pinnately many-veined; the veins and veinlets underneath along with the stalks and branchlets very rusty-scurfy; cymes sessile, very broad and flat.* — Cold moist woods, New England to Penn. and northward, and southward in the Alleghanies. May. — A straggling shrub; the reclining branches often taking root. Flowers handsome. Leaves 4'-8' across.

ORDER 50. **RUBIACEÆ.** (MADDER FAMILY.)

Shrubs or herbs, with opposite entire leaves connected by interposed stipules, or in whorls without apparent stipules, the calyx coherent with the 2-4-celled ovary, the stamens as many as the lobes of the regular corolla (3-5), and inserted on its tube. — Flowers perfect, but often dimorphous (as in Mitchellia and Houstonia). Fruit various. Seeds anatropous or amphitropous. Embryo commonly pretty large, in copious hard albumen. — A very large family, the greater part, and all its most important plants (such as the Coffee and Peruvian-Bark trees) tropical; not sufficiently represented in our district to render it worth while to note the tribes and the larger systematic divisions.

I. **STELLATÆ.** Leaves in whorls: no apparent stipules.

1. **Galium.** Corolla wheel-shaped, 4- (or rarely 3-) parted. Calyx-teeth obsolete. Fruit twin, separating into 2 indehiscent one-seeded carpels.

II. **CINCHONÆ,** &c. Leaves rarely in whorls, with stipules.

* Ovules and seeds solitary in each cell.

— Flowers axillary, separate. Fruit dry when ripe. Herbs.

2. **Spermacece.** Corolla funnel-form or salver-form: lobes 4. Fruit separating when ripe into 2 carpels, one or both of them opening.

3. **Diodia.** Fruit separating into 2 or 3 closed and indehiscent carpels: otherwise as No. 2.

— — Flowers in a close and globose long-peduncled head. Fruit dry. Shrubs.

4. **Cephalanthus.** Corolla tubular: lobes 4. Fruit inversely pyramidal, 2-4-seeded.

— — — Flowers twin; their ovaries united into one. Fruit a 2-eyed berry.

5. **Mitchella.** Corolla funnel-form; its lobes 4. — A creeping herb.

* * Ovules and seeds many or several in each cell of the (loculicidal) pod.

6. **Oldenlandia.** Corolla wheel-shaped in our species, 4-lobed. Seeds very numerous and minute, angular.

7. **Houstonia.** Corolla salver-form or funnel-form, 4-lobed. Seeds rather few, thimble-shaped or saucer-shaped.

1. **GÀLIUM,** L. BEDSTRAW. CLEAVERS.

Calyx-teeth obsolete. Corolla 4-parted, rarely 3-parted, wheel-shaped, valvate in the bud. Stamens 4, rarely 3, short. Styles 2. Fruit dry or fleshy, globular, twin, separating when ripe into the 2 seed-like, indehiscent, 1-seeded carpels. — Slender herbs, with small cymose flowers (produced in summer), square stems, and whorled leaves: the roots often containing a red coloring matter. (Name from γάλα, *milk*, which some species are used to curdle.)

* Annual: leaves about 8 in a whorl: peduncles 1-2-flowered, axillary.

1. **G. Aparine,** L. (CLEAVERS. GOOSE-GRASS.) Stem weak and reclining, bristle-prickly backwards, hairy at the joints; leaves lanceolate, tapering to the base, short-pointed, rough on the margins and midrib (1' - 2' long); flowers white; fruit (large) bristly with hooked prickles. — Moist thickets. Doubtful if truly indigenous in our district. (Eu.)

* * Perennial, ascending; leaves 4-6 or 8 in a whorl, with prominent midrib, but no lateral nerves: flowers white, few or numerous, on slender pedicels: fruit smooth.

+ *Flowers very abundant, the small clusters or cymes paniced on the branches.*

2. **G. MOLLÛGO**, L. *Stems* (1° – 3° long) very smooth; *leaves mostly in whorls of 8, oblanceolate or oblong-linear, barely rough on the margins, slender-pointed; flowers forming a long panicle.* — Washington Heights, near New York, *W. W. Denslow.* (Adv. from Eu.)

3. **G. asprëllum**, Michx. (ROUGH BEDSTRAW.) *Stem weak, much branched, rough backwards with hooked prickles, leaning on bushes* (3° – 5° high); *leaves in whorls of 6, or 4–5 on the branchlets, oval-lanceolate, pointed, with almost prickly margins and midrib; peduncles short, 2–3 times forked.* — Low thickets: common northward.

4. **G. concinnum**, Torr. & Gr. *Stems low and slender* ($6'$ – $12'$ high), with minutely roughened angles; *leaves all in whorls of 6, linear, slightly pointed, veinless, the margins upwardly roughened; peduncles slender, 2–3 times forked, diffusely paniced at the summit; pedicels short.* — Dry ground, Pennsylvania from the Susquehanna, to Virginia, Michigan, Illinois, and Kentucky. — Leaves not blackening in drying.

+ + *Flowers few, lateral or terminating the branches, not paniced.*

5. **G. trifidum**, L. (SMALL BEDSTRAW.) *Stems weak, ascending* ($5'$ – $20'$ high), branching, mostly roughened backwards on the angles; *leaves in whorls of 4 to 6, linear or oblanceolate, obtuse, the margins and midrib rough; corolla-lobes and stamens often only 3.* — Var. 1. **PUSILLUM**: *stems rather simple, 5'–8' high, nearly smooth; leaves only 3''–4'' long, all in fours, soon reflexed; peduncles 1–3-flowered.* (In deep sphagnous swamps, northward.) Var. 2. **TINCTORIUM**: *stem taller and stouter, and with nearly smooth angles; peduncles 3–7-flowered, the corolla-lobes and stamens 4.* Var. 3. **LATIFOLIUM** (*G. obtusum, Bigel.*): *stem smooth and widely branched; leaves oblong or elliptical, quite rough on the midrib and margins.* — Swamps: common, and very variable. (Eu.)

* * * *Perennial, procumbent: leaves 6 or rarely 5 in a whorl, with prominent midrib and no lateral nerves: flowers greenish: fruit bur-like, beset with hooked bristles.*

6. **G. triflorum**, Michx. (SWEET-SCENTED BEDSTRAW.) *Stem* (1° – 3° long) bristly-roughened backwards on the angles; *leaves elliptical-lanceolate, bristle-pointed, with slightly roughened margins* ($1'$ – $2'$ long); *peduncles 3-flowered, the flowers all pedicelled.* — Rich woodlands: common. — Sweet-scented in drying. (Eu.)

* * * * *Perennial, ascending or upright: leaves all in fours, more or less 3-nerved: peduncles loosely or remotely 3–several-flowered: corolla dull-purple, brownish, or rarely cream-color; the lobes pointed or bristle-tipped: fruit, except in No. 10, bur-like, beset with hooked bristles.*

7. **G. pilosum**. *Stem hairy; leaves oval, dotted, hairy* ($1'$ long), scarcely 3-nerved; *peduncles twice or thrice 2–3-forked, the flowers all pedicelled.* — Dry copses, Rhode Island and Vermont to Illinois and southward. — Var. **PUNCTICULOSUM** is a nearly smooth form (*G. punctulosum, Michx.*): Virginia and southward.

8. **G. circæzans**, Michx. (WILD LIQUORICE.) *Smooth or downy* (1° high); *leaves oval, varying to ovate-oblong, mostly obtuse, 3-nerved, ciliate* ($1'$ – $1\frac{1}{2}'$ long); *peduncles usually once forked, the branches elongated and widely diverging in*

fruit, bearing several remote flowers on very short lateral pedicels, reflexed in fruit; lobes of the corolla hairy outside above the middle. — Rich woods: common. — The var. *MONTANUM* is a dwarf, broad-leaved form, from mountain woods.

9. *G. lanceolatum*, Torr. (WILD LIQUORICE.) Leaves (except the lowest) lanceolate or ovate-lanceolate, tapering to the apex (2' long); corolla glabrous: otherwise like the last. — Woodlands: common northward.

10. *G. latifolium*, Michx. Smooth (1°-2° high); leaves lanceolate or ovate-lanceolate, acute, 3-nerved below; the midrib and margins rough; cymes paniced, loosely many-flowered, the purple flowers all on slender spreading pedicels; fruit smooth. — Dry woods, Mercersberg, Penn. (*Prof. Porter*), Maryland, and southward in the Alleghanies. (Also Arkansas, *Engelmann*.)

* * * * Perennial, erect: leaves 4 or 8 in a whorl; flowers very numerous and crowded in a narrow and compact terminal panicle, white or yellow.

11. *G. boreale*, L. (NORTHERN BEDSTRAW.) Smooth (1°-2° high); leaves in fours, linear-lanceolate, 3-nerved; flowers white; fruit minutely bristly, sometimes smooth. — Rocky banks of streams: common, especially northward. (Eu.)

12. *G. verum*, L. (YELLOW BEDSTRAW.) Leaves in eights (or some in sixes), linear, grooved above, roughish, deflexed; flowers yellow; fruit smooth. — Dry fields, E. Massachusetts. (Adv. from Eu.)

2. SPERMACÔCE, L. BUTTON-WEED.

Calyx-tube short; the limb parted into 4 teeth. Corolla funnel-form or salver-form; the lobes valvate in the bud. Stamens 4. Stigma or style 2-cleft. Fruit small and dry, 2-celled, 2-seeded, splitting when ripe into 2 carpels, one of them usually carrying with it the partition, and therefore closed, the other open on the inner face. — Small herbs, the bases of the leaves or petioles connected by a bristle-bearing stipular membrane. Flowers small, crowded into sessile axillary whorled clusters or heads. Corolla whitish. (Name compounded of *σπέρμα*, seed, and *ἄκωκή*, a point, probably from the pointed calyx-teeth on the fruit.)

1. *S. glabra*, Michx. Glabrous perennial; stems spreading (9' - 20' long); leaves oblong-lanceolate; whorled heads many-flowered; corolla little exceeding the calyx, bearded in the throat, bearing the anthers at its base; filaments and style hardly any. — River-banks, S. Ohio, Illinois, and southward. Aug.

3. DIÓDIA, L. BUTTON-WEED.

Calyx-teeth 2-5, often unequal. Fruit 2- (rarely 3-) celled; the crustaceous carpels into which it splits all closed and indehiscent. Otherwise resembling *Spermacoe*. Flowering all summer. (Name from *δίodos*, a thoroughfare; the species often growing by the wayside.)

1. *D. Virginia*, L. Smooth or hairy perennial; stems spreading (1°-2° long); leaves lanceolate or oblong-lanceolate, sessile; flowers 1-3 in each axil; corolla white ($\frac{1}{2}$ ' long), the slender tube abruptly expanded into the large limb; style 2-parted; fruit oblong, strongly furrowed, crowned mostly with 2 slender calyx-teeth. — River-banks, Maryland, and southward. Also naturalized near Philadelphia, *C. F. Parker*.

2. **D. tères**, Walt. Hairy or minutely pubescent annual; stem spreading 3'-9' long), nearly terete; leaves linear-lanceolate, closely sessile, rigid; flowers 1-3 in each axil; *corolla funnel-form* (2''-3'' long, whitish), with short lobes, not exceeding the long bristles of the stipules; *style undivided*; *fruit* obovate-turbinate, *not furrowed*, crowned with 4 short calyx-teeth. — Sandy fields, from New Jersey and Illinois southward.

4. CEPHALÁNTHUS, L. BUTTON-BUSH.

Calyx-tube inversely pyramidal, the limb 4-toothed. Corolla tubular, 4-toothed; the teeth imbricated in the bud. Style thread-form, much protruded. Stigma capitate. Fruit dry and hard, small, inversely pyramidal, 2-4-celled, at length splitting from the base upward into 2-4-closed 1-seeded portions. — Shrubs, with the flowers densely aggregated in spherical peduncled heads. Flowers white. (Name composed of *κεφαλή*, a head, and *ἄνθος*, a flower.)

1. **C. occidentális**, L. Smooth or pubescent; leaves petioled, ovate or lanceolate-oblong, pointed, opposite or whorled in threes, with short intervening stipules. — Wet places: common. July, Aug.

5. MITCHÉLLA, L. PARTRIDGE-BERRY.

Flowers in pairs, with their ovaries united. Calyx 4-toothed. Corolla funnel-form, 4-lobed; the lobes spreading, densely bearded inside, valvate in the bud. Stamens 4. Style 1: stigmas 4, linear. Fruit a berry-like double drupe, crowned with the calyx-teeth of the two flowers, each with 4 small and seed-like bony nutlets. — A smooth and trailing small evergreen herb, with round-ovate and shining petioled leaves, minute stipules, white fragrant flowers often tinged with purple, and scarlet edible (but nearly tasteless) dry berries, which remain over winter. Flowers occasionally 3-6-merous, always dimorphous; all those of some individuals having exerted stamens and included stigmas; of others, included stamens and exerted style. (This very pretty plant commemorates *Dr. John Mitchell*, an early correspondent of Linnæus, and an excellent botanist, who resided in Virginia.)

1. **M. repens**, L. — Dry woods, creeping about the foot of trees: common. June, July. — Leaves often variegated with whitish lines. Rarely the two flowers are completely confluent into one, with a 10-lobed corolla.

6. OLDENLÁNDIA, Plumier, L. OLDENLANDIA.

Calyx 4- (rarely 5-) lobed, persistent. Corolla short, in our species wheel-shaped; the limb 4- (rarely 5-) parted, valvate in the bud. Stamens 4 (rarely 5): anthers short. Style 1 or none: stigmas 2. Pod thin, 2-celled, many-seeded, opening loculicidally across the summit. Seeds concave, very numerous, minute and angular. — Low herbs, with small stipules united to the petioles. (Dedicated, in 1703, to the memory of *Oldenland*, a German physician and botanist, who died early at the Cape of Good Hope.)

1. **O. glomerata**, Michx. An inconspicuous, pubescent or smoothish, branched and spreading annual (2'-12' high); leaves oblong; flowers in sessile

axillary clusters; corolla nearly wheel-shaped (white), much shorter than the calyx. (*O. uniflora*, *L.* *Hedyotis glomerata*, *Ell.*) — Wet places, S. New York to Virginia near the coast, and southward.

7. HOUSTÔNIA, L. HOUSTONIA.

Calyx 4-lobed, persistent; the lobes in fruit distant. Corolla salver-form or funnel-form, usually much longer than the calyx-lobes, 4-lobed, the lobes valvate in the bud. Stamens 4: anthers linear or oblong. Style 1: stigmas 2. Ovary 2-celled. Pod top-shaped, globular, or didymous, thin, its summit or upper half free from and projecting beyond the tube of the calyx, loculicidal across the top. Seeds rather few (4–20 in each cell), peltate and saucer-shaped or globular-thimble-shaped, pitted. — Small herbs, with short entire stipules connecting the petioles or narrowed bases of the leaves, and cymose or solitary and peduncled flowers. These are dimorphous, in some individuals with the anthers borne high up on the tube of the corolla and projecting from its throat, while the style is short and the stigma therefore included: in the other sort the anthers are low down in the corolla and the style long, the stigmas therefore protruding; — an arrangement for cross-fertilization. (Named for *Dr. Wm. Houston*, an English botanist who collected in Central America.) The genus, formerly merged in *Oldenlandia*, merits restoration.

* *Erect, mostly perennial herbs (6'–20' high), with stem-leaves sessile, and flowers in terminal small cymes or clusters: corolla funnel-form, purplish, often hairy inside: seeds meniscoidal, with a ridge across the hollowed inner face.*

1. **H. purpurea**, *L.* Pubescent or smooth (8'–15' high); leaves varying from roundish-ovate to lanceolate, 3–5-ribbed; calyx-lobes longer than the half free globular pod. (*Houstonia purpurea*, *L.* *H. varians*, *Michx.* *Oldenlandia purpurea*, ed. 2.) — Woodlands, W. Pennsylvania to Illinois and southward. May–July. — Varying wonderfully, as into: —

Var. **longifolia**. Leaves varying from oblong-lanceolate to linear, narrowed at the base, 1-ribbed; calyx-lobes scarcely as long as the pod; stems 5'–12' high. (*Houstonia longifolia*, *Willd.*) — Maine to Wisconsin and southward. — A narrow-leaved, slender form is *H. tenuifolia*, *Nutt.*

Var. **ciliolata**. More tufted stems 3'–6' high; root-leaves in rosettes, thickish and ciliate; calyx-lobes about as long as the pod. (*Houstonia ciliolata*, *Torr.*) — Along the Great Lakes and rivers, from N. New York to Wisconsin.

2. **H. angustifolia**, *Michx.* Stems tufted from a hard or woody root; leaves narrowly linear, acute, 1-ribbed, many of them fascicled; flowers crowded, short-pedicelled; lobes of the corolla densely bearded inside; pod obovoid, acute at the base, only its summit free from the calyx, opening first across the top, at length through the partition. (*Oldenlandia angustifolia*, ed. 2. *Hedyotis stenophylla*, *Torr. & Gray.*) — Plains and banks, from Illinois southward. June–Aug.

** *Small and delicate, chiefly annuals or biennials, vernal-flowering: peduncles 1-flowered: corolla salver-form: upper half of the broad and somewhat 2-lobed pod free: seeds globular, with a very deep round cavity occupying the inner face.*

3. **H. minima**, *Beck.* Scabrous; stems at length much branched and spreading (1'–4' high); lowest leaves ovate or spatulate, the upper oblong or

nearly linear; earlier peduncles elongated and spreading in fruit, the later ones short; *tube of the purplish corolla not longer than its lobes nor than the ample calyx-lobes* ($1\frac{1}{2}$ ' long). — Dry hills, W. Illinois and southwestward. March — May.

4. **H. cærulea**, L. (BLUETS.) *Glabrous*; stems erect, slender, sparingly branched from the base (3' — 5' high); leaves oblong-spatulate (3' — 4' long); peduncle filiform, erect; *corolla with tube much longer than its lobes or than those of the calyx*. (Oldenlandia cærulea, ed. 2.) Moist and grassy places; producing from early spring to midsummer its delicate little flowers, light blue, pale lilac, or nearly white with a yellowish eye.

H. serpyllifolia, Michx., — with similar flowers, but with slender creeping stems, abounding in the mountains of N. Carolina, — may occur in those of Virginia.

H. rotundifolia, Michx., — also creeping, but with much larger roundish leaves, and axillary peduncles nodding in fruit, — belongs to the low country of the Southern States, and may occur in S. E. Virginia.

ORDER 51. VALERIANACEÆ. (VALERIAN FAMILY.)

Herbs, with opposite leaves and no stipules; the calyx-tube coherent with the ovary, which has one fertile 1-ovuled cell and two abortive or empty ones; the stamens distinct, 1 — 3, fewer than the lobes of the corolla, and inserted on its tube. — Corolla tubular or funnel-form, often irregular, mostly 5-lobed, the lobes imbricated in the bud. Style slender: stigmas 1 — 3. Fruit indehiscent, 1-celled (the two empty cells of the ovary disappearing), or 3-celled, two of them empty, the other 1-seeded. Seed suspended, anatropous, with a large embryo and no albumen. — Flowers in paniced or clustered cymes. (Roots often odorous and antispasmodic.) — Represented by only two genera.

1. VALERIANA, Tourn. VALERIAN.

Limb of the calyx of several plumose bristles (like a pappus) which are rolled up inwards in flower, but unroll and spread as the seed-like 1-celled fruit matures. Corolla commonly gibbous near the base, the 5-lobed limb nearly regular. Stamens 3. — Perennial herbs, with thickened strong-scented roots, and simple or pinnate leaves. Flowers in many species imperfectly dioecious, or dimorphous. (Name from *valere*, to have efficacy, alluding to the medicinal qualities.)

* *Root fibrous: leaves thin. (Stems 1° — 3° high.)*

1. **V. pauciflora**, Michx. Smooth, slender, surculose; *root-leaves ovate, heart-shaped, toothed*, pointed, sometimes with 2 small lateral divisions; stem-leaves pinnate, with 3 — 7 ovate toothed leaflets; branches of the paniced cyme few-flowered; *tube of the (pale pink) corolla long and slender* ($\frac{1}{2}$ ' long). — Woodlands, and alluvial banks, Penn. (near Lancaster, Prof. Porter) and Ohio to S. Illinois and southward. June.

2. **V. sylvatica**, Richards. Smooth or minutely pubescent; *root-leaves ovate or oblong, entire*, rarely with 2 small lobes; stem-leaves pinnate, with 5 — 11

oblong-ovate or lanceolate nearly entire leaflets; cyme at first close, many-flowered; *corolla inversely conical* (3" long, rose-color or white). — Cedar swamps, Western Vermont to Wisconsin and northward. June. (Probably a form of *V. dioica*, L.)

* *Root spindle-shaped, large and deep* (6'–12' long): *leaves thickish*.

3. **V. édulis**, Nutt. Smooth, or minutely downy when very young; stem straight (1°–4° high), few-leaved; leaves commonly minutely and densely ciliate, those of the root spatulate and lanceolate, of the stem pinnately parted into 3–7 long and narrow divisions; flowers in a long and narrow interrupted panicle, nearly diœcious; corolla whitish, obconical (2" long). (*V. ciliata*, Torr. & Gr.) — Alluvial ground, Ohio to Wisconsin, and westward. June.

2. FÈDIA, Gærtl. CORN SALAD. LAMB-LETTUCE.

Limb of the calyx obsolete or merely toothed. Corolla funnel-form, equally or unequally 5-lobed. Stamens 3, rarely 2. Fruit 3-celled, two of the cells empty and sometimes confluent into one, the other 1-seeded. — Annuals and biennials, usually smooth, with forking stems, tender and rather succulent leaves (entire or cut-lobed towards the base), and white or whitish cymose-clustered and bracted small flowers. (Name of uncertain derivation.) — Our species all have the limb of the calyx obsolete, and are so much alike in aspect, flowers, &c., that good characters are only to be taken from the fruit. They all have a rather short corolla, the limb of which is nearly regular, and therefore belong to the section (by many botanists taken as a genus) VALERIANÉLLA.

1. **F. OLITÒRIA**, Vahl. *Fruit compressed, oblique, at length broader than long, with a corky or spongy mass at the back of the fertile cell nearly as large as the (often confluent) empty cells; flowers bluish.* — Fields, New York and Penn. to Virginia: rare. (Adv. from Eu.)

2. **F. Fagopyrum**, Torr. & Gr. *Fruit ovate-triangular, smooth, not grooved between the (at length confluent) empty cells, which form the anterior angle, and are much smaller than the broad and flat fertile one; flowers white.* — Low grounds, from Western New York to Wisconsin and Kentucky. May, June. — Plant 1°–2° high.

3. **F. radiata**, Michx. *Fruit ovoid, downy (rarely smooth), obtusely and unequally somewhat 4-angled; the empty cells parallel and contiguous, but with a deep groove between them, rather narrower than the flattish fertile cell.* — Low grounds, Penn. to Michigan, and southward. — Plant 6'–15' high.

4. **F. umbilicata**, Sulliv. *Fruit globular-ovate, smooth; the much inflated sterile cells wider and many times thicker than the flattish fertile one, contiguous, and when young with a common partition, when grown, indented with a deep circular depression in the middle, opening into the confluent sterile cells; bracts not ciliate.* — Moist grounds, Columbus, Ohio, Sullivant. (*Sill. Jour.* Jan. 1842.)

5. **F. patellaria**, Sulliv. *Fruit smooth, circular, platter-shaped or disk-like, slightly notched at both ends, the flattened-concave sterile cells widely divergent, much broader than the fertile one, and forming a kind of wing around it when ripe.* — Low grounds, Columbus, Ohio, Sullivant. — Plant 1°–2° high, resembling the last, but with a very different fruit.

ORDER 52. **DIPSACÆ.** (TEASEL FAMILY.)

Herbs, with opposite or whorled leaves, no stipules, and the flowers in dense heads, surrounded by an involucre, as in the Composite Family; but the stamens are distinct, and the suspended seed has albumen. — Represented by the Scabious (cultivated) and the genus

1. **DÍPSACUS**, Tourn. TEASEL.

Involucre many-leaved, longer than the chaffy leafy-tipped and pointed bracts among the densely capitate flowers: each flower with a 4-leaved calyx-like involucre investing the ovary and fruit (achenium). Calyx-tube coherent with the ovary, the limb cup-shaped, without a pappus. Corolla nearly regular, 4-cleft. Stamens 4, inserted on the corolla. Style slender. — Stout and coarse biennials, hairy or prickly, with large oblong heads. (Name from *διψάω*, to thirst, probably because the united cup-shaped bases of the leaves in some species hold water.)

1. **D. SYLVÉSTRIS**, Mill. (WILD TEASEL.) Prickly; leaves lance-oblong; leaves of the involucre slender, longer than the head; bracts (chaff) tapering into a long flexible awn with a straight point. — Roadsides: rather rare. (Nat. from Eu.) Suspected to be the original of

2. **D. FULLÓNUM**, L., the cultivated FULLER'S TEASEL, which has a shorter involucre, and stiff chaff to the heads, with hooked points, used for raising a nap upon woollen cloth: it has escaped from cultivation in some places. (Adv. from Eu.)

ORDER 53. **COMPÓSITÆ.** (COMPOSITE FAMILY.)

Flowers in a close head (the compound flower of the older botanists), on a common receptacle, surrounded by an involucre, with 5 (rarely 4) stamens inserted on the corolla, their anthers united in a tube (syngenesious). — Calyx-tube united with the 1-celled ovary, the limb (called a pappus) crowning its summit in the form of bristles, awns, scales, teeth, &c., or cup-shaped, or else entirely absent. Corolla either strap-shaped or tubular; in the latter chiefly 5-lobed, valvate in the bud, the veins bordering the margins of the lobes. Style 2-cleft at the apex. Fruit seed-like (achenium), dry, containing a single erect anatropous seed, with no albumen. — An immense family, in temperate regions chiefly herbs, without stipules, with perfect, polygamous, monœcious, or diœcious flowers. The flowers with a strap-shaped (ligulate) corolla are called rays or ray-flowers: the head which presents such flowers, either throughout or at the margin, is radiate. The tubular flowers compose the disk; and a head which has no ray-flowers is said to be discoid. When the head contains two sorts of flowers it is said to be heterógamous; when only one sort, homógamous. The leaves of the involucre, of whatever form or texture, are termed scales. The bracts or scales, which often grow on the recep-

tacle among the flowers, are called the *chaff*: when these are wanting, the receptacle is said to be *naked*. — The largest order of Phænogamous plants. It is divided by the corolla into three suborders, only two of which are represented in the Northern United States. The first is much the larger.

SUBORDER I. TUBULIFLORÆ.

Corolla tubular in all the perfect flowers, regularly 5- (rarely 3-4-) lobed, ligulate only in the marginal or ray-flowers, which when present are either pistillate only, or neutral (with neither stamens nor pistil).

The technical characters of the five tribes of the vast suborder *Tubulifloræ*, taken from the styles, require a magnifying-glass to make them out, and will not always be clear to the student. The following artificial analysis, founded upon other and more obvious distinctions, will be useful to the beginner. (The numbers are those of the genera.)

Artificial Key to the Genera of this Suborder.

§ 1. Rays or ligulate flowers none: corollas all tubular (or rarely none).

* Flowers of the head all perfect and alike.

Pappus composed of bristles;

Double, the outer of very short, the inner of longer bristles. No. 1.

Simple, the bristles all of the same sort.

Heads few-flowered, themselves aggregated into a compound or dense cluster. 2.

Heads separate, few-flowered or many-flowered.

Receptacle (when the flowers are pulled off) bristly-hairy 67, 68, 70.

Receptacle deeply honeycomb-like. 69.

Receptacle naked.

Pappus of plumose or bearded stiff bristles. Flowers purple. 4.

Pappus of very plumose bristles. Flowers whitish. 5.

Pappus of slender but rather stiff rough bristles. 6, 7, 8, 20.

Pappus of very soft and weak naked bristles. 62, 63.

Pappus composed of scales or chaff.

Receptacle naked. Leaves in whorls. 3.

Receptacle naked. Leaves alternate. 45.

Receptacle bearing chaff among the flowers. 49.

Pappus of 2 or few barbed awns or teeth. 41, 42.

Pappus none, or a mere crown-like margin to the fruit. 55, 57.

** Flowers of two kinds in the same head.

Marginal flowers neutral and sterile, either conspicuous or inconspicuous. 65, 66.

Marginal flowers pistillate and fertile.

Receptacle elongated and bearing broad chaff among the flowers. 60.

Receptacle naked or bearing no conspicuous chaff.

Pappus of capillary bristles. Involucre imbricated. 23, 58, 59

Pappus of capillary bristles. Involucre merely one row of scales. 14, 61.

Pappus obsolete or none.

Achenia becoming much longer than the involucre. 11.

Achenia not exceeding the involucre. 29, 56, 57.

*** Flowers of two kinds in separate heads, the one pistillate, the other staminate.

Heads dioecious; in both kinds many-flowered. Pappus capillary. 24, 59.

Heads monoecious; the fertile 1-2-flowered and closed. Pappus none. 30, 31.

§ 2. Rays present; i. e. the marginal flowers or some of them with ligulate corollas.

* Pappus of capillary bristles. (Rays all pistillate.)

| | |
|--|--------------|
| Rays occupying several rows. | 9, 10, 14. |
| Rays in one marginal row, and | |
| White, purple or blue, never yellow. | 12-15. |
| Yellow, of the same color as the disk. | |
| Pappus double, the outer short and minute. | 21. |
| Pappus simple. | |
| Scales of the involucre equal and all in one row. Leaves alternate. | 63. |
| Scales of the involucre in two rows. Leaves opposite. | 64. |
| Scales of the involucre imbricated. Leaves alternate. | 19, 22. |
| * * Pappus a circle of chaffy scales, dissected into bristles. | 44. |
| * * * Pappus a circle of thin chaffy scales or short chaffy bristles. | |
| Heads several-flowered. Receptacle chaffy. | 50. |
| Heads 8-10-flowered. Receptacle naked. | 18. |
| Heads many-flowered. Receptacle deeply honeycombed. | 48. |
| Heads many-flowered. Receptacle naked. | 45½, 46, 47. |
| * * * Pappus none, or a cup or crown, or of 2 or 3 awns, teeth, or chaffy scales corresponding with the edges or angles of the achenium, often with intervening minute bristles or scales. | |
| + Receptacle naked. | |
| Achenia flat, wing-margined. Pappus of separate little bristles or awns. | 16. |
| Achenia flat, marginless. Pappus none. Receptacle conical. | 17. |
| Achenia terete or angled. Pappus none. Receptacle flattish. | 54. |
| Achenia angled. Pappus a little cup or crown. Receptacle conical. | 55. |
| + + Receptacle chaffy. | |
| Rays neutral (rarely pistillate but sterile); the disk-flowers perfect and fertile. | |
| Receptacle elevated (varying from strongly convex to columnar), and | |
| Chaffy only at the summit; the chaff deciduous. Pappus none. | 51. |
| Chaffy throughout. Achenia flattened laterally if at all. | 36-40. |
| Receptacle flat or flattish. Achenia flattened parallel with the scales or chaff. | 41, 42. |
| Rays pistillate and fertile; the disk-flowers also perfect and fertile. | |
| Achenia much flattened laterally, 1-2-awned. | 43. |
| Achenia flattened parallel with the scales and chaff. Pappus none. | 53. |
| Achenia 3-4-angular, terete or laterally flattish, awnless. | |
| Receptacle convex or conical. Leaves alternate, dissected. | 52. |
| Receptacle conical. Leaves opposite simple. | |
| Achenia obovoid. Involucre a leafy cup. | 32. |
| Achenia 4-angular. Involucre of separate scales. | 35. |
| Receptacle flat. Leaves opposite and simple. | 33, 34. |
| Rays pistillate and fertile: the disk-flowers staminate and sterile (pistil imperfect). | |
| Receptacle chaffy. | 25-28. |

Systematic Synopsis.

Tribe I. VERNONIACEÆ. Heads discoid; the flowers all alike, perfect and tubular. Branches of the style long and slender, terete, thread-shaped, minutely bristly-hairy all over. — Leaves alternate or scattered.

1. **Vernonia.** Heads several — many-flowered, separate. Involucre of many scales. Pappus of many capillary bristles.

2. **Elephantopus.** Heads 3-5-flowered, several crowded together into a compound head. Involucre of 8 scales. Pappus of several chaffy bristles.

Tribe II. EUPATORIACEÆ. Heads discoid, the flowers all alike, perfect and tubular; or in a few cases dissimilar, and the outer ones ligulate. Branches of the style thickened upwards or club-shaped, obtuse, very minutely and uniformly pubescent; the stigmatic lines indistinct.

Subtribe 1. Eupatoriæ. Flowers all perfect and tubular, never truly yellow.

* Pappus a row of hard scales.

3. **Sclerolepis.** Head many-flowered. Scales of the involucre equal. Leaves whorled.

** Pappus of slender bristles.

4. **Liatris.** Achenia many-ribbed. Bristles of the pappus plumose or barbellate. Corollas red-purple, strongly 5-lobed.

5. **Kuhnia.** Achenia many-ribbed. Bristles of the pappus very strongly plumose. Corollas whitish, 5-toothed.

6. **Eupatorium.** Achenia 5-angled. Bristles of the pappus roughish. Scales of the involucre many or several. Receptacle of the 5-many flowers flat or barely convex.

7. **Mikania.** Achenia and pappus as No. 6. Scales of the involucre and flowers only 4.

8. **Conoclinium.** Achenia, pappus, &c. as No. 6. Receptacle conical.

Subtribe 2. Tussilagines. Flowers (sometimes yellow, more or less monœcious or dioecious) of 2 sorts in the same head.

* Outer flowers of each (many-flowered) head pistillate and ligulate. Scape leafless.

9. **Nardosmia.** Heads corymbed. Flowers somewhat diœcious. Pappus capillary.

10. **Tussilago.** Head single; the outer pistillate flowers in many rows. Pappus capillary.

** Flowers all tubular. Stem leafy.

11. **Adenocaulon.** Head few-flowered; the outer flowers pistillate. Pappus none.

Tribe III. ASTEROIDEÆ. Heads discoid, with the flowers all alike and tubular; or else radiate, the outer ones ligulate and pistillate. Branches of the style in the perfect flowers flat, smooth up to where the conspicuous marginal stigmatic lines abruptly terminate, and prolonged above this into a flattened lance-shaped or triangular appendage which is evenly hairy or pubescent outside. — Leaves alternate. Receptacle naked (destitute of chaff) in all our species.

Subtribe 1. Asterinæ. Flowers of the head all alike and perfect, or the marginal ones ligulate and pistillate. Anthers without tails at the base.

* Ray-flowers white, blue, or purple, never yellow.

+ Pappus of numerous long and capillary bristles: receptacle flat.

12. **Sericocarpus.** Heads 12-15 flowered: rays 4 or 5. Involucre oblong or club-shaped, imbricated, cartilaginous. Achenia short, narrowed downwards, silky.

13. **Aster.** Heads many-flowered. Involucre loosely or closely imbricated. Achenia flattish. Pappus simple, copious.

14. **Erigeron.** Heads many-flowered. Involucre of narrow scales, little imbricated. Achenia flattened. Pappus simple and rather scanty, or with some outer minute scales.

15. **Diplopappus.** Heads many-flowered. Involucre imbricated. Pappus double; the outer obscure, of minute stiff bristles.

+ + Pappus of very short rigid bristles, or none: receptacle conical or hemispherical.

16. **Boltonia.** Achenia flat and wing-margined. Pappus very short.

17. **Bellis.** Achenia marginless. Pappus none. Receptacle conical.

** Ray-flowers yellow (in one species of *Solidago* whitish), or sometimes none at all.

18. **Brachyœta.** Heads 8-10 flowered, clustered: rays 4 or 5. Pappus a row of minute bristles shorter than the achenium.

19. **Solidago.** Heads few-many-flowered: rays 1-16. Pappus simple, of numerous slender and equal capillary bristles.

20. **Bigelovia.** Heads 3-4 flowered: rays none. Receptacle awl-shaped. Pappus simple, a single row of capillary bristles.

21. **Chrysopsis.** Heads many-flowered: rays numerous. Pappus double; the outer of very small chaffy bristles, much shorter than the inner of copious capillary bristles.

Subtribe 2. Inulæ. Anthers with tails at their base: otherwise as Subtribe 1.

22. **Inula.** Heads many-flowered. Rays many. Pappus capillary and copious.

Subtribe 3. Baccharidæ & Tarchonanthes. Flowers of the head all tubular, either diœcious or monœcious, namely, the staminate and pistillate flowers either in different heads on distinct plants, or in the same head. Corolla of the pistillate fertile flowers a very slender tube sheathing the style, and truncate at the summit.

23. **Pluchea.** Heads containing a few perfect but sterile flowers in the centre, and many pistillate fertile ones around them. Anthers tailed at the base. Pappus capillary.

24. **Baccharis.** Heads diœcious, some all pistillate, others all staminate, on different plants. Anthers tailless. Pappus capillary.

Tribe IV. SENECTIONIDÆ. Heads various. Branches of the style in the fertile flowers linear, thickish or convex externally, flat internally, hairy or pencil-tufted at the apex (where the stigmatic lines terminate abruptly), and either truncate, or continued beyond into a bristly-hairy appendage. — Leaves either opposite or alternate.

Subtribe 1. Melampodinæ. Flowers none of them truly perfect, but either staminate or pistillate; the two sorts either in the same or in different heads. Anthers tailless. Pappus, if any, never of bristles.

* Heads containing two kinds of flowers, the marginal ones pistillate, the central and tubular staminate flowers having a style, but always sterile.

+ Fertile flowers with a ligulate corolla (rarely wanting in No. 25): receptacle chaffy.

25. **Polymnia.** Achenia thick and turgid, roundish. Pappus none.

26. **Chrysogonum.** Achenia flattish. Pappus a one-sided 2-3-toothed chaffy crown.

27. **Silphium.** Achenia flat, wing-margined, numerous in several rows: rays deciduous.

28. **Parthenium.** Achenia flat, slightly margined: rays very short, persistent.

+ + Fertile flowers with tubular or no corolla: no pappus.

29. **Iva.** Achenia short and thick: receptacle with narrow chaff.

57. **Artemisia,** in part. Achenia short and small: receptacle naked.

11. **Adenocaulon.** Achenia elongated, bearing stalked glands: receptacle naked.

* * Heads of two sorts, one containing staminate, the other pistillate flowers, both borne on the same plant; the pistillate only 1 or 2. in a closed involucre resembling an achenium or a bur; the staminate several, in an open cup-shaped involucre.

30. **Ambrosia.** Fertile involucre (fruit) small, 1-flowered, pointed and often tubercled.

31. **Xanthium.** Fertile involucre (fruit) an oblong prickly bur, 2-celled, 2-flowered.

Subtribe 2. Helianthes. Heads radiate, or rarely discoid; the rays ligulate, the disk-flowers all perfect and fertile. Receptacle chaffy. Anthers blackish, tailless. Pappus none, or a crown or cup, or of one or two chaffy awns, never capillary, nor of several uniform chaffy scales. — Leaves more commonly opposite.

* Rays pistillate and fertile: achenia 3-4-sided, slightly if at all flattened.

+ Involucre double; the outer forming a cup.

32. **Tetragonotheca.** Outer involucre 4-leaved. Achenia obovoid. Pappus none.

+ + Involucre of one or more rows of separate scales.

33. **Eclipta.** Receptacle flat; its chaff bristle-shaped. Pappus obsolete or none.

34. **Borrchia.** Receptacle flat, its chaff scale-like and rigid. Pappus an obscure crown.

35. **Hellopsis.** Receptacle conical; its chaff linear. Pappus none or a mere border.

* * Rays sterile (either entirely neutral or with an imperfect style), or occasionally none; achenia 4-angular or flattened laterally, i. e. their edges directed inwards and outwards, the chaff of the receptacle embracing their outer edge.

+ Receptacle elevated, conical or columnar. Pappus none or a short crown.

36. **Echinacea.** Rays (very long) pistillate, but sterile. Achenia short, 4-sided.

37. **Rudbeckia.** Rays neutral. Achenia 4-sided, flat at the top, marginless.

38. **Lepachys.** Rays few, neutral. Achenia flattened laterally and margined.

+ + Receptacle flattish or conical. Pappus chaffy or awned.

39. **Helianthus.** Achenia flattened, marginless. Pappus of 2 very deciduous chaffy scales.

40. **Actinomeris.** Achenia very flat, wing-margined, bearing 2 persistent awns.

* * * Rays sterile, neutral: achenia obcompressed, i. e. flattened parallel with the scales of the involucre, the faces looking inwards and outwards. Involucre double; the outer spreading and often foliaceous. Receptacle flat.

41. **Coreopsis.** Pappus of 2 (or rarely more) scales, teeth, or awns, which are naked or barbed upwards, sometimes obsolete or a mere crown.

42. **Bidens.** Pappus of 2 or more rigid and persistent downwardly barbed awns or teeth.

* * * * Rays pistillate or fertile (rarely none): achenia laterally flattened, 2-awned.

43. **Verbesina.** Rays few and small, or rarely none. Receptacle convex. Achenia sometimes winged.

Subtribe 3. Tagetinae. Heads commonly radiate; the rays ligulate; the disk-flowers all perfect and fertile. Receptacle naked, flat. Scales of the involucre united into a cup. Pappus various. — Herbage strong-scented (as in *Tagetes* of the gardens), being dotted with large pellucid glands containing a volatile oil.

44. **Dysodia.** Pappus a row of chaffy scales dissected into many bristles.

Subtribe 4. Heleniæ. Heads radiate or sometimes discoid; the disk-flowers perfect. Pappus a circle of several chaffy scales. Anthers tailless.

* Receptacle naked (not chaffy nor honeycombed.)

45. **Hymenopappus.** Rays none. Receptacle flat. Scales of the involucre colored.

45½. **Actinella.** Rays pistillate, merely toothed. Receptacle elevated. Involucre appressed.

46. **Helenium.** Rays pistillate, 3-5-cleft. Receptacle elevated. Involucre small, reflexed.

47. **Leptopoda.** Rays neutral or sterile: otherwise as No. 46.

* * Receptacle deeply pitted, like honeycomb.

48. **Baldwinia.** Rays numerous: neutral, elongated. Involucre much imbricated.

* * * Receptacle chaffy.

49. **Marshallia.** Rays none. Involucre of many narrow and foliaceous scales.

50. **Galinsoga.** Rays 4 or 5, short, pistillate, whitish; the disk yellow. Involucre of 4 or 5 ovate and thin scales.

Subtribe 5. Anthemidæ. Heads radiate or discoid; the perfect flowers sometimes infertile, and the pistillate flowers rarely tubular. Pappus a short crown or none. Otherwise nearly as Subtribe 4.

* Receptacle chaffy, at least in part: rays ligulate.

51. **Naruta.** Rays neutral. Achenia obovoid and many-ribbed. Pappus none.

52. **Anthemis.** Rays pistillate. Achenia terete or 4-angular. Pappus minute or none.

53. **Achillea.** Rays pistillate, short. Achenia flattened and margined.

* * Receptacle naked.

54. **Leucanthemum.** Rays numerous, pistillate. Receptacle flattish. Achenia striate or ribbed. Pappus none.

55. **Matricaria.** Rays pistillate, or none, and then all the flowers perfect. Receptacle conical. Pappus crown-like or none.

56. **Tanacetum.** Rays none, but the marginal flowers pistillate. Achenia broad at the top. Pappus a short crown.

57. **Artemisia.** Rays none; some of the outer flowers often pistillate. Achenia narrow at the top. Pappus none.

Subtribe 6. Gnaphalinae. Heads all discoid, with tubular corollas; those of the fertile flowers filiform. Anthers with tails at their base. Pappus of capillary bristles. Flocculent-woolly herbs: leaves alternate.

58. **Gnaphalium.** Receptacle naked, flat. Heads containing both perfect and pistillate flowers. Bristles of the pappus all slender.

59. **Antennaria.** Receptacle naked, flat. Heads dioecious, or nearly so. Pappus of the staminate flowers thickened or club-shaped at the summit.

60. **Filago.** Receptacle columnar or top-shaped, chaffy. Pappus of the inner flowers capillary, of the outer often none.

Subtribe 7. Senecioneæ. Heads radiate or discoid; the central flowers perfect. Anthers tailless. Pappus capillary. Receptacle naked. (Scales of the involucre commonly in a single row.)

* Heads discoid. Leaves alternate.

61. **Erechtites.** Heads many-flowered: flowers whitish; the marginal ones pistillate and with filiform corollas.

62. **Cacalia.** Heads 5-many-flowered: flowers white or cream-color, all tubular and perfect.

63. **Senecio.** Heads many-flowered: flowers yellow, all perfect.

** Heads radiate, many-flowered.

← Rays conspicuous and in a single row: flowers all fertile, yellow.

63. **Senecio.** Pappus fine and soft. Leaves alternate.

64. **Arnica.** Pappus of more rigid and rough-denticulate bristles. Leaves all opposite.

← ← Rays narrow or small, in more than one row, at least in fertile heads. Leaves all radical.

9. **Nardosmia.** Flowers whitish or purplish. Scape bearing several heads.

10. **Tussilago.** Flowers yellow. Scape bearing a single head.

Tribe V. CYNAREÆ. Heads (in our species) discoid, with the flowers tubular, or some of the outer corollas enlarged and appearing like rays but not truly ligulate. Style thickened or thickish near the summit; the branches stigmatic to the apex, without any appendage, often united below. (Heads large.)

* Marginal flowers mostly neutral or sterile. Pappus not plumose.

65. **Centaurea.** Achenia flat. Pappus of short naked bristles, or none. Marginal neutral flowers commonly enlarged or ray-like.

66. **Cnicus.** Achenia terete, bearing 10 horny teeth and a pappus of 10 long and 10 shorter rigid naked bristles. Marginal flowers inconspicuous.

** Flowers all alike in the (ovoid or globular) head.

67. **Cirsium.** Achenia smooth. Pappus of plumose bristles. Receptacle clothed with long and soft bristles or hairs.

68. **Carduus.** Pappus of naked bristles: otherwise as No. 67.

69. **Onopordon.** Achenia wrinkled transversely, 4-angled. Pappus not plumose. Receptacle honeycombed.

70. **Lappa.** Achenia wrinkled, flattened. Pappus of short and rough bristles. Receptacle bristly.

SUBORDER II. **LIGULIFLORÆ.** **Tribe VI. CICHORACEÆ.**

Corolla ligulate in all the flowers of the head, and all the flowers perfect. — Herbs, with milky juice. Leaves alternate.

* Pappus none.

71. **Lampsana.** Involucre cylindrical, of 8 scales in a single row. 8-12-flowered.

** Pappus chaffy, or of both chaff and bristles.

72. **Cichorium.** Pappus a small crown of many bristle-form scales. Involucre double.

73. **Krigia.** Pappus of 5 broad chaffy scales, and 5 slender bristles.

74. **Cynthia.** Pappus double; the outer short, of many minute chaffy scales, the inner of numerous long capillary bristles.

*** Pappus plumose.

75. **Leontodon.** Bristles of the pappus several, chaffy-dilated at the base, tawny.

**** Pappus composed entirely of capillary bristles, not plumose.

← Achenia not flattened nor distinctly beaked, columnar or terete, often slender: pappus rather stiff, mostly tawny or dirty-white.

76. **Troximon.** Involucre loosely imbricated, many-flowered: corolla yellow. Achenia 10-ribbed. Pappus very copious and unequal.

77. **Hieracium.** Involucre more or less imbricated, 12-many-flowered: corolla yellow. Achenia short. Pappus of rather scanty and tawny roughish bristles.

78. **Nabalus**. Involucre cylindrical, of 5-14 linear equal scales in a single row and a few little scales at base, 5-many-flowered: corolla whitish, cream-color, or purplish. Achenia rather short and blunt. Pappus of very copious tawny or brown roughish bristles.
79. **Lygodesmia**. Involucre as the preceding, 5-10-flowered: corolla rose-purple. Achenia long and slender, tapering at the summit. Pappus of copious whitish bristles.
- + + Achenia terete or nearly so, ribbed, roughish above, abruptly slender-beaked. Pappus soft, fine, and flaccid. Involucre cylindrical, of several linear scales in a single row and some small short ones at the base. Corolla yellow.
80. **Chondrilla**. Involucre few-flowered. Pappus white. Stems branching, leafy.
81. **Pyrrhopappus**. Involucre many-flowered. Pappus reddish or rusty. Stems branching, leafy below.
82. **Taraxacum**. Involucre many-flowered. Pappus whitish. Scape naked, simple.
- + + + Achenia flat or flattish. Involucre somewhat imbricated, mostly many-flowered.
83. **Lactuca**. Achenia abruptly long and slender-beaked, very flat: pappus bright white.
84. **Mulgedium**. Achenia flattish, and with a short and thick beak.
85. **Sonchus**. Achenia flattish, beakless. Pappus white. Flowers yellow.

1. VERNONIA, Schreb. IRON-WEED.

Heads 15-many-flowered, in corymbose cymes; flowers all perfect. Involucre shorter than the flowers, of many appressed closely imbricated scales. Receptacle naked. Achenia cylindrical, ribbed. Pappus double; the outer of minute scale-like bristles; the inner of copious capillary bristles. — Perennial herbs, with alternate leaves and mostly purple flowers. (Named in honor of a *Mr. Vernon*, an early English botanist who travelled in this country.)

1. **V. noveboracensis**, Willd. *Scales of the involucre tipped with a long bristle-form or awl-shaped spreading appendage or awn*; in some varieties merely pointed. — Low grounds near the coast, Maine to Virginia; and river-banks in the Western States, from Wisconsin southward. Aug. — A tall coarse weed with lanceolate or oblong leaves.

2. **V. fasciculata**, Michx. *Scales of the involucre (all but the lowest) rounded and obtuse, without appendage*. — Prairies and river-banks, Ohio to Wisconsin and southward. Aug. — Leaves narrowly or broadly lanceolate: heads mostly crowded. Very variable, and manifestly passing into No. 1.

2. ELEPHANTOPUS, L. ELEPHANT'S-FOOT.

Heads 3-5-flowered, several together clustered into a compound head: flowers perfect. Involucre narrow, flattened, of 8 oblong dry scales. Achenia many-ribbed. Pappus of stout bristles, chaffy-dilated at the base. — Perennials, with alternate leaves and purplish flowers. (Name composed of *ἔλεφας*, *elephant*, and *ποῦς*, *foot*.)

1. **E. Carolinianus**, Willd. Somewhat hairy, corymbose, leafy; leaves ovate-oblong, thin. — Dry soil, Pennsylvania to Illinois and southward.

3. SCLERÓLEPIS, Cas. SCLEROLEPIS.

Head many-flowered: flowers perfect. Scales of the involucre linear, equal, in 1 or 2 rows. Corolla 5-toothed. Achenia 5-angled. Pappus a single row of almost horny oval and obtuse scales. — A smooth perennial, with simple stems, rooting at the base, linear entire leaves in whorls of 5 or 6, and a terminal head

of flesh-colored flowers. (Name of *σκληρός*, *hard*, and *λεπίς*, *a scale*, from the pappus.)

1. **S. verticillata**, Cass. — In water: pine barrens, New Jersey and southward. Aug.

4. **LIÀTRIS**, Schreb. **BUTTON SNAKEROOT. BLAZING-STAR.**

Head several-many-flowered: flowers perfect. Scales of the involucre imbricated, appressed. Receptacle naked. Corolla 5-lobed. Achenia slender, tapering to the base, about 10-ribbed. Pappus of 15-40 capillary bristles, which are manifestly plumose, or only barbellate. — Perennial herbs, often resinous-dotted, with rigid alternate entire leaves (these sometimes twisted so as to become vertical), and heads of handsome rose-purple flowers, spicate, racemose, or paniced-cymose, appearing late in summer or in autumn. (Derivation of the name unknown.)

§ 1. *Stem usually wand-like and simple, from a globular or roundish corm or tuber (impregnated with resinous matter), very leafy: leaves narrow or grass-like, 1-5-nerved: heads spicate or racemed: involucre well imbricated: lobes of the corolla long and slender.*

* *Pappus very plumose; scales of the 5-flowered involucre with ovate or lanceolate spreading petal-like (purple or sometimes white) tips, exceeding the flowers.*

1. **L. élegans**, Willd. Stem (3°-5° high) and involucre hairy; leaves short and spreading; spike or raceme compact (1° long). — Barren soil, Virginia and southward.

** *Pappus very plumose: scales of the cylindrical many-flowered involucre imbricated in many rows, the tips rigid, not petal-like: corolla hairy within.*

2. **L. squarròsa**, Willd. (BLAZING-STAR, &c.) Often hairy (1°-3° high); leaves linear, elongated; heads few (1' long); scales of the involucre mostly with elongated and leaf-like spreading tips. — Dry soil, Pennsylvania to Illinois and southward.

3. **L. cylindràcea**, Michx. Commonly smooth (6'-18' high); leaves linear; heads few ($\frac{1}{2}$ '- $\frac{3}{4}$ ' long); scales of the involucre with short and rounded appressed tips. — Dry open places, Niagara Falls to Wisconsin, and southwestward.

*** *Pappus not obviously plumose to the naked eye: corolla smooth inside.*

4. **L. scariòsa**, Willd. Stem stout (2°-5° high) pubescent or hoary; leaves (smooth, rough, or pubescent) lanceolate; the lowest oblong-lanceolate or obovate-oblong, tapering into a petiole; heads few or many, large, 30-40-flowered; scales of the broad or depressed involucre obovate or spatulate, very numerous, with dry and scarious often colored tips or margins. — Dry soil, New England to Minnesota, and southward. — Widely variable: heads 1' or less in diameter.

5. **L. pilòsa**, Willd. Beset with long scattered hairs; stem stout; leaves linear or linear-lanceolate, elongated; heads few, 10-15-flowered; scales of the top-shaped or bell-shaped involucre slightly margined, the outer narrowly oblong, very obtuse, the innermost linear. — Mountains of Virginia and southward: rare and obscure. Perhaps a remarkable state of the next; but the flowers as large as in the preceding.

6. **L. spicàta**, Willd. Smooth or somewhat hairy; stems very leafy (2° – 5° high); leaves linear, the lower 3–5-nerved; heads 8–12-flowered ($\frac{1}{2}$ '– $\frac{1}{2}$ ' long), crowded in a long spike; scales of the cylindrical-bell-shaped involucre oblong or oval, obtuse, appressed, with slight margins; achenia pubescent or smoothish. — Moist grounds: common from S. New York to Wisconsin and southward. — Involucre somewhat resinous, very smooth.

7. **L. graminifolia**, Willd. Hairy or smoothish; stem (1° – 3° high) slender, leafy; leaves linear, elongated, 1-nerved; heads several or numerous, in a spike or raceme, 7–12-flowered; scales of the obconical or obovoid involucre spatulate or oblong, obtuse, or somewhat pointed, rigid, appressed; achenia hairy. — Virginia and southward. — Inflorescence sometimes paniced, especially in

Var. **dùbia**. Scales of the involucre narrower and less rigid, oblong, often ciliate. (*L. dubia*, Barton.) — Wet pine barrens, New Jersey and southward.

8. **L. pycnostàchya**, Michx. Hairy or smoothish; stem stout (3° – 5° high), very leafy; leaves linear-lanceolate, the upper very narrowly linear; spike thick and dense (6'–20' long); heads about 5-flowered ($\frac{1}{2}$ ' long); scales of the cylindrical involucre oblong or lanceolate, with recurved or spreading colored tips. — Prairies, from Indiana southward and westward.

§ 2. Stem simple or branched above, not from a tuber: heads small, corymbed or paniced, 4–10-flowered: involucre little imbricated: lobes of the corolla ovate: pappus not plumose.

9. **L. odoratissima**, Willd. (VANILLA-PLANT.) Very smooth; leaves pale, thickish, obovate-spatulate, or the upper oval and clasping; heads corymbed. — Low pine barrens, Virginia and southward. — Leaves exhaling the odor of Vanilla when bruised.

10. **L. paniculàta**, Willd. Viscid-hairy; leaves narrowly oblong or lanceolate, smoothish, those of the stem partly clasping, heads paniced. — Virginia and southward.

5. KÙHNIA, L. KUHNIA.

Heads 10–25-flowered: flowers perfect. Scales of the involucre few and loosely imbricated, lanceolate. Corolla slender, 5-toothed. Achenia cylindrical, many-striate. Pappus a single row of very plumose (white) bristles. — A perennial herb, resinous-dotted, with mostly alternate lanceolate leaves, and paniculate-corymbose heads of cream-colored flowers. (Dedicated to *Dr. Kuhn*, of Pennsylvania, who brought the living plant to Linnæus.)

1. **K. eupatorioides**, L. Leaves varying from broadly lanceolate and toothed, to linear and entire. — Dry soil, New Jersey to Wisconsin and southward. Sept.

6. EUPATÒRIUM, Tourn. THOROUGHWORT.

Heads 3–many-flowered: flowers perfect. Involucre cylindrical or bell-shaped. Receptacle flat or barely convex. Corolla 5-toothed. Achenia 5-angled. Pappus a single row of slender capillary barely roughish bristles. — Perennial herbs, often sprinkled with bitter resinous dots, with generally corymbose heads of white, bluish, or purple blossoms, appearing near the close of

summer. (Dedicated to *Eupator Mithridates*, who is said to have used a species of the genus in medicine.)

* *Heads cylindrical, 5-15-flowered; the purplish scales numerous, closely imbricated in several rows, of unequal length, slightly striate: stout herbs, with ample mostly whorled leaves, and flesh-colored flowers.*

1. **E. purpureum**, L. (JOE-PYE WEED. TRUMPET-WEED.) Stems tall and stout, simple; leaves 3-6 in a whorl, oblong-ovate or lanceolate, pointed, very veiny, roughish, toothed; corymbs very dense and compound. — Varies greatly in size (2°-12° high), &c., and with spotted or unspotted, often dotted stems, &c., — including several nominal species. — Low grounds: common.

* * *Heads 3-20-flowered: involucre of 8-15 more or less imbricated and unequal scales, the outer ones shorter: flowers white.*

+ *Leaves all alternate, mostly dissected: heads paniced, very small, 3-5-flowered.*

2. **E. fœniculaceum**, Willd. Smooth or nearly so, paniculately much-branched (3°-10° high); leaves 1-2-pinnately parted, filiform. — Virginia, near the coast, and southward. Adv. near Philadelphia.

+ + *Leaves mostly opposite and sessile: heads 5-8-flowered, corymbed.*

3. **E. hyssopifolium**, L. Minutely pubescent (1°-2° high); leaves narrow, linear or lanceolate, elongated, obtuse, 1-3-nerved, entire, or the lower toothed, often crowded in the axils, acute at the base; scales of the involucre obtuse. — Sterile soil, Massachusetts to Virginia, E. Kentucky and southward.

4. **E. leucolépis**, Torr. & Gr. Minutely pubescent, simple (1°-2° high); leaves linear-lanceolate, closely sessile, 1-nerved, obtuse, serrate, rough both sides; corymb hoary; scales of the involucre with white and scarious acute tips. — Sandy bogs, Long Island, New Jersey, and southward.

5. **E. parviflorum**, Ell. Minutely velvety-pubescent, branching (2°-3° high); leaves lanceolate or oblong, triple-ribbed and veiny, serrate above the middle, tapering to the base, the lower slightly petioled; scales of the short involucre obtuse. (Leaves sometimes whorled in threes, or the upper alternate.) — Damp soil, Virginia and southward.

6. **E. altissimum**, L. Stem stout and tall (3°-7° high), downy; leaves lanceolate, tapering at both ends, conspicuously 3-nerved, entire, or toothed above the middle, the uppermost alternate; corymbs dense; scales of the involucre obtuse, shorter than the flowers. — Dry soil, Pennsylvania to Illinois and Kentucky. — Leaves 3'-4' long, somewhat like those of a Solidago.

7. **E. album**, L. Roughish-hairy (2° high); leaves oblong-lanceolate, coarsely toothed, veiny; heads clustered in the corymb; scales of the involucre closely imbricated, rigid, narrowly lanceolate, pointed, white and scarious above, longer than the flowers. — Sandy and barren places, pine barrens of New Jersey to Virginia and southward.

8. **E. teucrifolium**, Willd. Roughish-pubescent (2°-3° high); leaves ovate-oblong and ovate-lanceolate, obtuse or truncate at the base, slightly triple-nerved, veiny, coarsely toothed or incised towards the base, the upper ones alternate; branches of the corymb few, unequal; scales of the involucre oblong-lanceolate, rather obtuse, at length shorter than the flowers. (*E. verbenæfolium*, Michx.) — Low grounds, Massachusetts to Virginia and southward near the coast.

9. **E. rotundifolium**, L. Downy-pubescent (2° high); leaves roundish-ovate, obtuse, truncate or slightly heart-shaped at the base, deeply crenate-toothed, triple-nerved, veiny, roughish (1'–2' long); corymb large and dense; scales of the (5-flowered) involucre linear-lanceolate, slightly pointed. — Dry soil, Rhode Island to Virginia, near the coast, and southward.

10. **E. pubescens**, Muhl. Pubescent: leaves ovate, mostly acute, slightly truncate at the base, serrate-toothed, somewhat triple-nerved, veiny; scales of the 7–8-flowered involucre lanceolate, acute. (*E. ovatum*, Bigel.) — Massachusetts to New Jersey, near the coast, and Kentucky. — Like the last, but larger.

11. **E. sessilifolium**, L. (UPLAND BONESET.) Stem tall (4°–6° high), smooth, branching; leaves lanceolate or ovate-lanceolate, tapering from near the rounded sessile base to the sharp point, serrate, veiny, smooth (3'–6' long); corymb very compound, pubescent; scales of the 5- (or 5–12-?) flowered involucre oval and oblong, obtuse. — Copses and banks, Massachusetts to Illinois, and southward along the mountains.

+ + + Leaves opposite, clasping or united at the base, long, widely spreading: heads mostly 10–15-flowered: corymbs very compound and large.

12. **E. resinosum**, Torr. Minutely velvety-downy (2°–3° high); leaves linear-lanceolate, elongated, serrate, partly clasping, tapering to the point, slightly veiny beneath (4'–6' long); scales of the involucre oval, obtuse. — Wet pine barrens, New Jersey. — Name from the copious resinous globules of the leaves.

13. **E. perfoliatum**, L. (THOROUGHWORT. BONESET.) Stem stout (2°–4° high), hairy; leaves lanceolate, united at the base around the stem (connate-perfoliate), tapering to a slender point, serrate, very veiny, wrinkled, downy beneath (5'–8' long); scales of the involucre linear-lanceolate. — Low grounds: common and well-known. — Varies with the heads 30–40-flowered.

+ + + + Leaves long-petioled, the upper ones alternate: heads 12–15-flowered, in compound corymbs.

14. **E. serotinum**, Michx. Stem pulverulent-pubescent, bushy-branched (3°–6° high); leaves ovate-lanceolate, tapering to a point, triple-nerved and veiny, coarsely serrate (5'–6' long); involucre very pubescent. — Alluvial ground, Maryland to Illinois and southward.

* * * Heads 8–30-flowered; the scales of the involucre nearly equal and in one row: leaves opposite, ovate, petioled, triple-nerved and veiny, not resinous-dotted: flowers white.

15. **E. ageratoides**, L. (WHITE SNAKE-ROOT.) Smooth, branching (3° high); leaves broadly ovate, pointed, coarsely and sharply toothed, long-petioled, thin (4'–5' long); corymbs compound. — Rich woods: common northward.

16. **E. aromaticum**, L. Smooth or slightly downy; stems nearly simple; leaves on short petioles, ovate, rather obtusely toothed, not pointed, thickish. — Copses, Massachusetts to Virginia and southward, near the coast. — Lower and more slender than No. 15, with fewer, but usually larger heads.

7. MIKÀNIA, Willd. CLIMBING HEMP-WEED.

Heads 4-flowered. Involucre of 4 scales. Receptacle small. Flowers and achenia, &c., as in Eupatorium. — Twining perennials, climbing bushes, with

opposite commonly heart-shaped and petioled leaves, and corymbose-panicled flesh-colored flowers. (Named for *Prof. Mikan*, of Prague.)

1. **M. scândens**, L. Nearly smooth, twining; leaves somewhat triangular-heart-shaped or halberd-form, pointed, toothed at the base. — Copses along streams, E. New England to Kentucky and southward. July – Sept.

8. CONOCLINIUM, DC. MIST-FLOWER.

Heads many-flowered. Involucre bell-shaped, the nearly equal linear-awl-shaped scales somewhat imbricated. Receptacle conical! Otherwise as in *Eupatorium*, of which it is rather a section. — Perennial erect herbs, with opposite petioled leaves, and violet-purple or blue flowers in crowded terminal corymbs. (Name formed of *κῶνος*, a cone, and *κλίνη*, a bed, from the conical receptacle.)

1. **C. cœlestinum**, DC. Somewhat pubescent (1° – 2° high); leaves triangular-ovate and slightly heart-shaped, coarsely and bluntly toothed. — Rich soil, Pennsylvania to Michigan, Illinois, and southward. Sept.

9. NARDÓSMIA, Cass. SWEET COLTSFOOT.

Heads many-flowered, somewhat dioecious: in the sterile plant with a single row of ligulate pistillate ray-flowers, and many tubular ones in the disk; in the fertile plant with many rows of minutely ligulate ray-flowers, and a few tubular perfect ones in the centre. Scales of the involucre in one row. Receptacle flat. Achenia terete. Pappus of soft capillary bristles, longer and copious in the fertile flowers. — Perennial woolly herbs, with the leaves all from the rootstock, the scape with sheathing scaly bracts, bearing heads of purplish or whitish fragrant flowers, in a corymb. (Name from *νάρδος*, *spikenard*, and *ὄσμή*, *odor*.)

1. **N. palmata**, Hook. Leaves rounded, somewhat kidney-form, white-woolly beneath, palmately and deeply 5 – 7-lobed, the lobes toothed and cut. (*Tussilago palmata*, *Ait.* *T. frigida*, *Bigel.*) — Swamps, Maine and Massachusetts to Michigan and northwestward: rare. April, May. — Full-grown leaves (6' – 10' broad).

10. TUSSILÀGO, Tourn. COLTSFOOT.

Head many-flowered; the ray-flowers narrowly ligulate, pistillate, fertile, in many rows; the tubular disk-flowers few, staminate. Scales of the involucre nearly in a single row. Receptacle flat. Fertile achenia cylindrical-oblong. Pappus capillary, copious in the fertile flowers. — A low perennial, with horizontal creeping rootstocks, sending up simple scaly scapes in early spring, bearing a single head, and producing rounded-heart-shaped angled or toothed leaves later in the season, woolly when young. Flowers yellow. (Name from *tussis*, a cough, for which the plant is a reputed remedy.)

1. **T. FÁRFARA**, L. — Wet places, and along brooks, New England, New York, and Pennsylvania; thoroughly wild. (Nat. from Eu.)

11. ADENOCÀULON, Hook. ADENOCAULON.

Heads 5 – 10-flowered; the flowers all tubular and with similar corollas; the marginal ones pistillate, fertile; the others staminate. Scales of the involucre

equal, in a single row. Achenia elongated at maturity, club-shaped, beset with stalked glands above. Pappus none. — Slender perennials, with the alternate thin and petioled leaves smooth and green above, white woolly beneath, and few small (whitish) heads in a loose panicle, beset with glands (whence the name, from ἀδήν, a gland, and καυλός, a stem).

1. **A. bicolor**, Hook. Leaves triangular, rather heart-shaped, with angular-toothed margins; petioles margined. — Moist woods, shore of Lake Superior, and westward.

12. SERICOCÁRPUS, Nees. WHITE-TOPPED ASTER.

Heads 12–15-flowered, radiate; the rays about 5, fertile (white). Involucre somewhat cylindrical or club-shaped; the scales closely imbricated in several rows, cartilaginous and whitish, appressed, with short and abrupt often spreading green tips. Receptacle alveolate-toothed. Achenia short, inversely pyramidal, very silky. Pappus simple, of numerous capillary bristles. — Perennial tufted herbs (1°–2° high), with sessile somewhat 3-nerved leaves, and small heads mostly in little clusters, disposed in a flat corymb. Disk-flowers pale yellow. (Name from σηρικός, silky, and καρπός, fruit.)

1. **S. solidagineus**, Nees. Smooth, slender; leaves linear, rigid, obtuse, entire, with rough margins, tapering to the base; heads narrow (3" long), in close clusters, few-flowered; pappus white. — Thickets, S. New England to Virginia, near the coast. July.

2. **S. conyzoides**, Nees. Somewhat pubescent; leaves oblong-lanceolate or the lower spatulate, mostly serrate towards the apex, ciliate, veiny; heads rather loosely corymbed, obconical (4"–6" long); pappus rusty-color. — Dry ground: common. July.

3. **S. tortifolius**, Nees. Hoary-pubescent; leaves obovate or oblong-spatulate, short, ($\frac{1}{2}$ –1' long), turned edgewise, both sides alike, nearly veinless; heads rather loosely corymbed, obovoid (4"–5" long); pappus white. — Pine woods, Virginia and southward. Aug.

13. ÁSTER, L. STARWORT. ASTER.

Heads many-flowered, radiate; the ray-flowers in a single series, fertile. Scales of the involucre more or less imbricated, usually with herbaceous or leaf-like tips. Receptacle flat, alveolate. Achenia more or less flattened. Pappus simple, of capillary bristles. — Perennial herbs (or annual only in § 6), with corymbed, paniced, or racemose heads; flowering in autumn. Rays white, purple, or blue: the disk yellow, often changing to purple. (Name ἀστήρ, a star, from the appearance of the radiate heads of flowers.)

§ 1. **ΒΙΌΤΙΑ**, DC. Involucre obovoid-bell-shaped; the scales regularly imbricated in several rows, appressed, nearly destitute of herbaceous tips: rays 6–15 (white or nearly so): achenia slender: lower leaves large, heart-shaped, petioled, coarsely serrate: heads in open corymbs.

1. **A. corymbosus**, Ait. Stem slender, somewhat zigzag; leaves thin, smoothish, coarsely and unequally serrate with sharp spreading teeth, sharp-pointed, ovate or ovate-lanceolate, all but the uppermost heart-shaped at the base and on

slender naked petioles; rays 6-9. — Woodlands: common, especially northward. July, Aug. — Plant 1° - 2° high, with smaller heads, looser corymbs, rounder and less rigid exterior involucre scales, and thinner leaves than the next; not rough, but sometimes pubescent.

2. **A. macrophyllus**, L. *Stem stout and rigid* (2° - 3° high); leaves thickish, rough, closely serrate, somewhat pointed; the lower heart-shaped ($4'$ - $10'$ long, $3'$ - $6'$ wide), long-petioled; the upper ovate or oblong, sessile or on margined petioles; heads in ample rigid corymbs; rays 12-25 (white or bluish). — Moist woods; common northward, and southward along the mountains. Aug., Sept. — Involucre $\frac{1}{2}'$ broad; the outer scales rigid, oblong or ovate-oblong, the innermost much larger and thinner.

§ 2. **CALLIÁSTRUM**, Torr. & Gr. *Scales of the involucre imbricated in several rows, coriaceous, with short herbaceous tips: rays 12-30, violet or blue: achenia narrow (smoothish): pappus of rigid bristles of unequal thickness: stem-leaves all sessile; lower ones not heart-shaped: heads few, or when several corymbose, large and showy.* (Allied to § 1, and to Sericocarpus.)

3. **A. Rádula**, Ait. *Stem simple or corymbose at the summit, smooth, many-leaved* (1° - 3° high); leaves oblong-lanceolate pointed, sharply serrate in the middle, very rough both sides and rugose-veined, closely sessile ($2'$ - $3'$ long), nearly equal; scales of the bell-shaped involucre oblong, appressed, with very short and slightly spreading herbaceous tips; achenia smooth. — Bogs and low grounds, Delaware to Maine and northward, near the coast. Also Pocono Mountain, Penn. (*Prof. T. Green*); and a dwarf variety, with linear-lanceolate leaves, at White Mountains, New Hampshire. Aug. — Rays light violet. Involucre nearly smooth, except the ciliate margins.

4. **A. surculósus**, Michx. *Stems slender* ($\frac{1}{2}^{\circ}$ - 1° high), from long and slender, or here and there tuberous-thickened, creeping, subterranean shoots or suckers, roughish-pubescent above, 1-2- or corymbosely several-flowered; leaves roughish, obscurely toothed, lanceolate or the lower spatulate; involucre obconical or bell-shaped ($\frac{1}{2}'$ - $\frac{1}{2}'$ long), the whitish and coriaceous scales with short herbaceous tips, the outer ones shorter; achenia slightly pubescent. — Var. **GRÁCILIS** (*A. gracilis*, Nutt.) is a form with the scales of the narrower obconical involucre successively shorter and with very short and scarcely spreading green tips, resembling a Sericocarpus. — Moist grounds, pine barrens of New Jersey and southward. Sept. — Rays about 12, violet, $6''$ long.

5. **A. spectábilis**, Ait. *Stems* (1° - 2° high) minutely rough and glandular-pubescent at the summit; leaves oblong-lanceolate, roughish, obscurely toothed, tapering to the base; scales of the short and almost hemispherical involucre linear-oblong, with conspicuous spatulate glandular-downy tips, the outermost scarcely shorter; achenia slightly pubescent. — Sandy soil, Massachusetts to New Jersey, near the coast, and southward. Sept. — Nov. — One of the handsomest of the genus, though the heads are few. The rays, about 20, are narrowly lanceolate, nearly $1'$ long, very deep violet-blue. Involucre $\frac{1}{2}'$ long and wide.

6. **A. Hérveyi**, n. sp. *Stem slender* (1° - 2° high), nearly smooth, the summit and peduncles of the several corymbose heads minutely glandular-pubescent; leaves thinnish, roughish, obscurely serrate, oblong-lanceolate, very acute, all but

the uppermost taper-pointed, and also tapering below into a narrowed base or winged petiole; heads small (less than $\frac{1}{2}$ ' long, exclusive of the narrow rays); involucre between bell-shaped and top-shaped; the scales obscurely glandular, linear, or the short outer ones oblong, with greenish appressed tips; achenia linear, slightly pubescent. — Borders of oak woods, in rather moist soil, New Bedford, Mass., *E. W. Hervey*. Sept. — An ambiguous member, and the smallest-flowered, of the section. Pappus whitish, finer than that of the preceding.

§ 3. **ASTER**, proper. Scales of the involucre imbricated in various degrees, with herbaceous or leaf-like summits, or the outer ones entirely foliaceous: rays numerous: pappus soft and nearly uniform: achenia flattened. (All flowering late in summer or in autumn.)

* Leaves whitened, silvery-silky both sides, all sessile and entire, mucronulate: involucre imbricated in 3 to several rows: rays showy, purple-violet.

7. **A. sericeus**, Vent. Stems slender, branched; leaves silver-white, lanceolate or oblong; heads mostly solitary, terminating the short branchlets; scales of the globular involucre similar to the leaves, spreading, except the short coriaceous base; achenia smooth, many-ribbed. — Prairies and dry banks, Wisconsin to Kentucky and southward. — Heads large: rays 20–30.

8. **A. cóncolor**, L. Stems wand-like, nearly simple; leaves crowded, oblong or lanceolate, appressed, the upper reduced to little bracts; heads in a simple or compound wand-like raceme; scales of the obovoid involucre closely imbricated in several rows, appressed, rather rigid, silky, lanceolate; achenia silky. — Dry sandy soil, pine barrens of New Jersey and southward. — Plant 1° – 3° high, with the short leaves 1' or less in length, grayish-silky both sides.

* * Lower leaves not heart-shaped; the upper all sessile and more or less clasping by a heart-shaped or auricled base: heads showy: scales of the inversely conical or bell-shaped involucre regularly imbricated in several rows, the outer successively shorter, appressed, coriaceous, whitish, with short herbaceous tips: rays large, purple or blue.

9. **A. pátens**, Ait. Rough-pubescent; stem loosely paniced above (1° – 3° high), with widely spreading branches, the heads mostly solitary, terminating the slender branchlets; leaves oblong-lanceolate or ovate-oblong, often contracted below the middle, all clasping by a deep auricled-heart-shaped base, rough, especially above and on the margins, entire; scales of the minutely roughish involucre with spreading pointed tips; achenia silky. — Var. **PHLOGIFÓLIUS** is a form which the plant assumes in shady moist places, with larger and elongated thin scarcely rough leaves, downy underneath, sometimes a little toothed above, mostly much contracted below the middle. — Dry ground: common, especially southward. Heads $\frac{1}{2}$ ' broad, and with showy deep blue-purple rays.

10. **A. lævis**, L. Very smooth throughout; heads in a close panicle; leaves thickish, lanceolate or ovate-lanceolate, chiefly entire, the upper more or less clasping by an auricled or heart-shaped base; scales of the short-obovoid or hemispherical involucre with appressed green points; rays sky-blue; achenia smooth. A variable and elegant species, of which the two best-marked forms are: —

Var. **lævigátus**. Scarcely if at all glaucous; leaves lanceolate or oblong; involucre nearly hemispherical; the scales lanceolate or linear, with narrow and

acute green tips tapering down on the midnerve. (*A. lævis*, *L.* *A. lævigatus*, *Willd.*) — Dry woodlands : rather common.

Var. **cyaneus**. Very smooth, but pale or glaucous ; leaves thicker ; the upper often oblong or ovate-lanceolate, clasping by a heart-shaped base ; involucre narrowed at the base, of broader and more coriaceous scales with shorter and abrupt tips. (*A. cyaneus*, *Hoffm.*, &c.) — Border of woodlands : common, especially northward.

11. **A. turbinellus**, *Lindl.* *Very smooth* ; stem slender, paniculately branched ; leaves lanceolate, tapering to each end, entire, with rough margins ; involucre elongated-obconical or almost club-shaped ($\frac{1}{2}$ long) ; the scales linear, with very short and blunt green tips ; rays violet-blue ; achenia nearly smooth. — Dry hills, &c., Illinois and southwestward. — Well-marked and handsome.

* * * *Lower leaves all heart-shaped and petioled, the upper sessile or petioled : involucre imbricated much as in the last division, but the heads smaller, very numerous, racemosed or panicled.*

+ *Leaves entire or slightly serrate : heads middle-sized : rays bright-blue.*

12. **A. azureus**, *Lindl.* Stem rather rough, erect, racemose-compound at the summit, the branches slender and rigid ; leaves rough ; the lower ovate-lanceolate or oblong, heart-shaped, on long often hairy petioles ; the others lanceolate or linear, sessile, on the branches awl-shaped ; involucre inversely conical. — Copses and prairies, Niagara Falls (*Clinton*), and Ohio to Wisconsin and southward. — Involucre much as in *A. lævis*, but much smaller, slightly pubescent ; the rays bright blue.

13. **A. Shörtii**, *Boott.* Stem slender, spreading, nearly smooth, bearing very numerous heads in racemose panicles ; leaves smooth above, minutely pubescent underneath, lanceolate or ovate-lanceolate, elongated, tapering gradually to a sharp point, all but the uppermost more or less heart-shaped at the base, and on naked petioles ; involucre bell-shaped. — Cliffs and banks, Ohio to Wisconsin and southward. — A pretty species, 2°–4° high ; the leaves 3'–5' long.

14. **A. undulatus**, *L.* Pale or somewhat hoary with close pubescence ; stem spreading, bearing numerous heads in racemose panicles ; leaves ovate or ovate-lanceolate, with wavy or slightly toothed margins, roughish above, downy underneath, the lowest heart-shaped on margined petioles, the others abruptly contracted into short broadly winged petioles which are dilated and clasping at the base, or directly sessile by a heart-shaped base ; involucre obovoid. (*A. diversifolius*, *Michx.*) — Dry copses : common.

+ + *Leaves conspicuously serrate : heads small : rays pale blue or nearly white.*

15. **A. cordifolius**, *L.* Stem much branched above, the spreading or diverging branches bearing very numerous panicled heads ; lower leaves all heart-shaped, on slender and mostly naked ciliate petioles ; scales of the inversely conical involucre all appressed and tipped with short green points, obtuse or acutish. — Woodlands : very common. — Varies with the stem and leaves either smooth, roughish, or sometimes hairy. Heads profuse, but quite small.

16. **A. sagittifolius**, *Willd.* Stem rigid, erect, with ascending branches bearing numerous racemose heads ; leaves ovate-lanceolate, pointed ; the lower heart-shaped at the base, on margined petioles ; the upper lanceolate or linear,

pointed at both ends; *scales of the oblong involucre linear, tapering into awl-shaped slender and loose tips.* — Dry ground, New York and Penn. to Kentucky and northward. — Usually more or less hairy or downy; the heads rather larger than in the last, almost sessile. — *A. Drummondii*, Lindl., which probably grows on the Illinois side of the Mississippi, is a downy-leaved variety of this.

* * * * *Leaves none of them heart-shaped; those of the stem sessile, narrow, rigid, entire: involucre imbricated in several rows: the coriaceous scales appressed and whitish at the base, with abrupt and conspicuous spreading herbaceous tips: heads small and very numerous, paniculate-racemose; rays white.*

17. *A. ericoides*, L. *Smooth or sparingly hairy (1° - 1½° high); the simple branchlets or peduncles racemose along the upper side of the wand-like spreading branches; lowest leaves oblong-spatulate, sometimes toothed; the others linear-lanceolate or linear-awl-shaped, acute at both ends; scales of the involucre broadest at the base, with acute or awl-shaped green tips.* — Var. *VILLOSUS* is a hairy form, often with broader leaves; chiefly in the Western States. — Dry open places, S. New England to Wisconsin and southward.

18. *A. multiflorus*, Ait. *Pale or hoary with minute close pubescence (1° high), much branched and bushy; the heads much crowded on the spreading racemose branches; leaves crowded, linear, spreading, with rough or ciliate margins, the upper somewhat dilated and partly clasping at the base; scales of the involucre with spatulate spreading green tips broader than the lower portion, the outer obtuse.* — Dry gravelly or sandy soil: common.

* * * * * *Leaves none of them heart-shaped; those of the stem tapering at the base, sessile; involucre imbricated; the scales unequal, with short and narrow appressed or rather loose greenish tips: heads small or middle-sized: rays white or bluish-purple.*

← *Heads small. (Involucre 2" - 4" long.)*

19. *A. dumosus*, L. *Smooth or nearly so, racemosely compound, the scattered heads mostly solitary at the end of the spreading branchlets; leaves linear or the upper oblong, crowded, entire or slightly serrate, with rough margins; scales of the closely imbricated involucre linear-spatulate, obtuse, in 4 - 6 rows.* — Thickets: common. — A variable species, 1° - 3° high, loosely branched, with small leaves, especially the upper, and an inversely conical or bell-shaped involucre, with more abrupt green tips than any of the succeeding. Rays pale purple or blue, larger than in the next. Runs into several peculiar forms.

20. *A. Tradescanti*, L. *Smooth or smoothish; the numerous heads closely racemed along one side of the erect-spreading or diverging branches; leaves lanceolate-linear, elongated, the larger ones remotely serrate in the middle with fine sharp teeth; scales of the involucre narrowly linear, acute or acutish, in 3 or 4 rows.* — Var. *FRAGILIS* has the leaves entire or nearly so, except the lowest, the heads more scattered. — Moist banks: very common. — Stems 2° - 4° high, bushy: heads very numerous, smaller than in the last. Rays white or nearly so.

21. *A. miser*, L., Ait. *More or less hairy, much branched; the branches usually diverging, bearing racemose often scattered heads; leaves lanceolate or oblong-lanceolate, tapering or pointed at each end, sharply serrate in the middle;*

scales of the involucre linear, acute or rather obtuse, imbricated in 3 or 4 rows. — Thickets, fields, &c.: very common, and extensively variable. — Leaves larger than in either of the preceding (2'–5'); the involucre intermediate between them, as to the form of the scales. Rays mostly short, pale bluish-purple or white.

← ← Heads middle-sized. (Involucre 3"–5" long.)

22. **A. simplex**, Willd. Smooth or nearly so (3°–6° high), much branched; the branches and scattered heads somewhat corymbose at the summit; leaves lanceolate, pointed, the lower serrate; scales of the involucre linear-awl-shaped, loosely and sparingly imbricated. — Shady moist banks: common. — Rays pale. Approaches in its different forms the preceding and the two following.

23. **A. tenuifolius**, L. Nearly smooth; stem much branched (2°–3° high); the heads somewhat paniced or racemed; leaves narrowly lanceolate, tapering into a long slender point (2'–6' long), with rough margins, the lower somewhat serrate in the middle; scales of the hemispherical involucre linear-awl-shaped, very slender-pointed, numerous, closely imbricated. — Low grounds, New York to Wisconsin, and southward. Rays short and narrow, pale purple or whitish.

24. **A. carneus**, Nees. Smooth, or the branches rough or pubescent; leaves lanceolate, somewhat pointed, or the upper short and partly clasping; heads racemose along the ascending leafy branches; scales of the obovate involucre lanceolate, abruptly acute, closely imbricated. — Moist soil: common. Leaves firm in texture, smooth, or rough above. Rays rather large, bluish, purplish, violet-purple, or almost white. — On a thorough revision of the genus, older names will be found and verified for this and No. 22, which here cover a multitude of forms. *A. mutabilis*, L., is probably one of them.

* * * * * Stem-leaves sessile, the upper more or less clasping; scales of the hemispherical involucre loosely more or less imbricated, somewhat equal, with herbaceous tips, or the outer often entirely herbaceous: heads middle-sized or large: rays blue or purple. (The species of this group are still perplexing.)

25. **A. æstivus**, Ait. Stem slender, rough, bushy-branched; leaves narrowly lanceolate-linear, elongated, taper-pointed, entire, with rough margins; heads corymbose, loose; scales of the involucre linear, loose; rays large, apparently light blue. (*A. laxifolius*, Nees.) — Var. **LÆTIFLORUS** has very slender branches and leaves, and the scales of the involucre unequal and more appressed. — Moist shady places, Ohio to Wisconsin and northward. — Heads about as large as in *A. puniceus*, in some forms appearing more like *A. carneus*. Leaves 4'–7' long, $\frac{1}{4}$ ' to $\frac{1}{2}$ ' wide.

26. **A. Novi-Belgii**, L. Nearly smooth; stem stout; leaves oblong-lanceolate, pale or somewhat glaucous, serrate in the middle, acute, tapering to each end; scales of the involucre rather closely imbricated, with broadish acute herbaceous tips; rays pale blue or purplish. — Low grounds, not clearly known in a wild state. The plant here in view is intermediate between No. 24 and No. 27. — Heads smaller and less showy than in the next.

27. **A. longifolius**, Lam. Smooth or nearly so; stem branched, corymbose-paniced at the summit; leaves lanceolate or linear, or the lower ovate-lanceolate, entire or sparingly serrate in the middle, taper-pointed, shining above; scales

of the involucre imbricated in 3-5 rows, linear, with acute or awl-shaped spreading or recurved green tips; rays large and numerous, bright purplish-blue. — Moist places, along streams, &c.: common eastward. — Plant 1°-5° high, with large and showy heads; very variable in the foliage, involucre, &c.; its multiform varieties including *A. thyrsoiflorus*, *Hoffm.*, *A. laxus*, *Willd.* (a form with more leafy involucre), *A. prææltus*, *Poir.*, *A. elodes*, *Torr. & Gr.* also *A. salicifolius*, *Scholler*, the oldest name of all.

28. **A. puniceus**, L. *Stem tall and stout, rough-hairy* all over or in lines, usually purple below, paniced above; leaves oblong-lanceolate, clasping by an auricled base, sparingly serrate in the middle with appressed teeth, rough above, nearly smooth underneath, pointed; scales of the involucre narrowly linear, acute, loose, equal, in about 2 rows; rays long and showy (lilac-blue, paler or whitish in shade). — Low thickets and swamps, very common. — Stems 3°-6° high, in open grounds rough with rigid bristly hairs.

Var. **vimineus** (*A. vimineus*, *Willd.*) is a variety nearly smooth throughout; growing in shade.

29. **A. prenanthoides**, Muhl. *Stem low*, (1°-3° high), corymbose-paniced, hairy above in lines; leaves rough above, very smooth underneath, ovate-lanceolate, sharply cut-toothed in the middle, conspicuously taper-pointed, and tapering below in a long contracted entire portion, which is abruptly dilated into an auricled-heart-shaped clasping base; scales of the involucre narrowly linear, with recurved-spreading tips; rays light blue. — Borders of rich woods, W. New York to Wisconsin, and southwards to Virginia.

* * * * * *Leaves entire, those of the stem sessile, the base often clasping: heads solitary terminating the branches or somewhat corymbed, showy: scales of the involucre very numerous, with loose and spreading or recurved mostly foliaceous tips, usually more or less glandular or viscid, as are the branchlets, &c.*

+ *Involucre imbricated, the scales in several or many ranks.*

30. **A. grandiflorus**, L. *Rough with minute hispid hairs*; stems slender, loosely much branched (1°-3° high); leaves very small ($\frac{1}{4}$ '-1' long), oblong-linear, obtuse, rigid, the uppermost passing into scales of the hemispherical squarrose many-ranked involucre; rays bright violet (1' long); achenia hairy. — Dry open places, Virginia and southward. — Heads large and very showy.

31. **A. oblongifolius**, Nutt. *Minutely glandular-puberulent*, much branched above, rigid, paniculate-corymbose (1°-2° high); leaves narrowly oblong or lanceolate, mucronate-pointed, partly clasping, thickish (1'-2' long by 2''-5'' wide); scales of the involucre broadly linear, appressed at the base; rays violet-purple; achenia canescent. — Banks of rivers, from Pennsylvania (Huntingdon County, *Prof. Porter!*) and Virginia to Wisconsin, Kentucky, and westward. — Heads middle-sized or smaller.

32. **A. amethystinus**, Nutt. *Taller than the foregoing* (2°-5° high), more upright, with more hirsute and not at all glandular or viscid pubescence; heads more racemose on the branches and smaller; involucre only about 3'' in diameter, the tips of the scales less spreading; rays light clear blue. — Moist grounds, Massachusetts (near Salem, Cambridge, &c., *Nuttall, Dr. Robbins, &c.*: Amherst (*Prof. Tuckerman*), Illinois (*E. Hall*), and Wisconsin.

+ + *Involucre of many very slender equal scales appearing like a single row.*

33. **A. Novæ-Ángliæ**, L. *Stem stout, hairy (3° - 8° high), corymbed at the summit; leaves very numerous, lanceolate, entire, acute, auriculate-clasping, clothed with minute pubescence; scales of the involucre linear-awl-shaped, loose, glandular-viscid, as well as the branchlets; rays violet-purple, in var. ROSEUS rose-purple (A. roseus, Desf.), very numerous; achenia hairy. — Moist grounds: common. — Heads large, corymbed. Var. ? with white rays, Carroll Co., Ill., H. Shimer.*

* * * * * *Heads, &c., as in the preceding group; but foliage as in * * *.*

34. **A. anómalus**, Engelm. *Somewhat hoary-pubescent; stems slender (2° - 4° high), simple or racemose-branched above; leaves ovate or ovate-lanceolate, pointed, entire or nearly so, the lower cordate and long-petioled, the upper small and almost sessile; scales of the hemispherical involucre imbricated in several rows, appressed, with linear spreading leafy tips; achenia smooth. — Limestone cliffs, W. Illinois (and Missouri, near St. Louis), Engelmann. — Heads as large as those of No. 31: rays violet-purple.*

§ 4. **ORITRÓPHIUM**, Kunth. *Scales of the involucre narrow, nearly equal and almost in a single row, more or less herbaceous: pappus of soft and uniform capillary bristles: mostly low perennials, bearing solitary or few heads.*

35. **A. graminifólius**, Pursh. *Slightly pubescent, slender (6' - 12' high); leaves very numerous, narrowly linear; branches prolonged into slender naked peduncles, bearing solitary small heads; rays rose-purple or whitish. — Northern borders of New England, Lake Superior, and northward.*

§ 5. **ORTHÓMERIS**, Torr. & Gr. *Scales of the involucre regularly imbricated, unequal, often carinate, with membranaceous margins, entirely destitute of herbaceous tips: pappus of soft and unequal capillary bristles.*

36. **A. acuminátus**, Michx. *Somewhat hairy; stem (about 1° high) simple, zigzag, paniced-corymbose at the summit; peduncles slender; leaves oblong-lanceolate, conspicuously pointed, coarsely toothed above, wedge-form and entire at the base; scales of the involucre few and loosely imbricated, linear-lanceolate, pointed, thin (3'' - 5'' long); heads few or several; rays 12 - 18, white, or slightly purple. — Cool rich woods: common northward and southward along the Alleghanies. Aug. — There is a depauperate narrow-leaved variety on the White Mountains of New Hampshire.*

37. **A. nemorális**, Ait. *Minutely roughish-pubescent; stem slender, simple or corymbose at the summit, very leafy (1° - 2° high); leaves small (1' - 1½' long), rather rigid, lanceolate, nearly entire, with revolute margins; scales of the inversely conical involucre narrowly linear-lanceolate, the outer passing into awl-shaped bracts; rays lilac-purple, elongated. — Bogs, pine barrens of New Jersey to Maine along the coast, and northward. Also White Mountains of New Hampshire; a small form, with solitary heads. Sept.*

38. **A. ptarmicoides**, Torr. & Gr. *Smooth or roughish; stems clustered (6' - 15' high), simple; leaves linear-lanceolate, acute, rigid, entire, tapering to the base, 1 - 3-nerved, with rough margins (2' - 4' long); heads small, in a flat corymb; scales of the involucre imbricated in 3 or 4 rows, short; rays white (2'' - 3'' long). — Dry rocks, W. Vermont to Illinois and Wisconsin along the Great Lakes, and northward. Aug.*

§ 6. **OXYTRIPOLIUM**, DC. *Scales of the involucre imbricated, without herbaceous tips, usually very acute, the outer passing into scale-like bracts: pappus soft and capillary: achenia striate.*

39. **A. flexuosus**, Nutt. Stem zigzag, rigid, forked (6' - 20' high); the branches bearing large solitary heads; leaves linear, thick and fleshy, pointed, entire; scales of the bell-shaped involucre imbricated in many rows, ovate-lanceolate with awl-shaped points; rays numerous, large, pale purple. — Salt marshes on the coast, Maine to Virginia. Sept.

40. **A. linifolius**, L. Annual; stem much branched (6' - 24' high), the branches bearing numerous racemose or panicked small heads; leaves linear-lanceolate, pointed, entire, flat, on the branches awl-shaped; scales of the oblong involucre linear-awl-shaped, in few rows; rays somewhat in two rows, short, not projecting beyond the disk, more numerous than the disk-flowers, purplish. (*A. subulatus*; Michx.) Salt marshes, on the coast, Maine to Virginia. Aug. - Oct.

41. **A. angustus**, Torr. & Gr. (*Tripolium angustum*, Lindl. *T. frondosum*, Nutt. Also in Siberia, *Conyza Altaica*, DC., and *Brachyaetis ciliata*, Ledeb.) An annual species related to the last, but with broader leaves, very many minute rays, and more copious pappus: comes from the northwest to near the borders of Wisconsin.

14. ERIGERON, L. FLEABANE.

Heads many-flowered, radiate, mostly flat or hemispherical; the narrow rays very numerous, pistillate. Scales of the involucre narrow, nearly equal and little imbricated. Receptacle flat or convex, naked. Achenia flattened, usually pubescent and 2-nerved. Pappus a single row of capillary bristles, with minuter ones intermixed, or with a distinct short outer pappus of little bristles or chaffy scales. — Herbs, with entire or toothed and generally sessile leaves, and solitary or corymbed heads. Disk yellow; ray white or purple. (Name from ἔρις, *spring*, and γέρων, *an old man*, suggested by the hoary appearance of some of the vernal species.)

§ 1. **CÆNÔTUS**, Nutt. *Rays inconspicuous, in several rows, scarcely longer than the pappus: pappus simple: chiefly annuals and biennials.*

1. **E. Canadense**, L. (HORSE-WEED. BUTTER-WEED.) Bristly-hairy; stem erect, wand-like (3° - 5° high); leaves linear, mostly entire; those from the root cut-lobed; heads very numerous and small, cylindrical, panicked. — Waste places; a common weed, now widely diffused over the world. July - Oct. — Ligules much shorter than their tube, white. (Nat. in Eu. &c.)

2. **E. divaricatum**, Michx. *Diffuse and decumbent (3' - 6' high); leaves linear or awl-shaped; heads loosely corymbed; rays purple: otherwise like No. 1.* — Illinois, Kentucky, and southward.

3. **E. acre**, L. Sparsely hairy or smoothish; stem erect (10' - 20' high); leaves lanceolate or the lower spatulate-oblong, entire; heads several or rather numerous, racemose or at length corymbose, nearly hemispherical (4" - 5" long); rays purplish or bluish, equalling or a little exceeding the copious pappus. — Shores of Lake Superior (*Dr. Robbins, Prof. Porter, &c.*), and northward. (Eu.)

§ 2. **EUERIGERON**, Torr. & Gr. *Rays elongated, crowded in one or more rows: pappus simple. (Erect perennials: heads somewhat corymbed, middle-sized.)*

4. **E. bellidifolium**, Muhl. (ROBIN'S PLANTAIN.) Hairy, producing offsets from the base: stem simple, rather naked above, bearing few (1-9) large heads on slender peduncles, root-leaves obovate and spatulate, sparingly toothed; those of the stem distant, lanceolate-oblong, partly clasping, entire; rays (about 50) rather broad, light bluish-purple. — Copses and moist banks: common. May.

5. **E. Philadelphicum**, L. (COMMON FLEABANE.) Hairy; stem leafy, corymbed, bearing several small heads; leaves thin, with a broad midrib, oblong; the upper smoothish, clasping by a heart-shaped base, mostly entire; the lowest spatulate, toothed; rays innumerable and very narrow, rose-purple or flesh-color. (*E. purpureum*, Ait.) — Moist ground: common. June - Aug.

§ 3. **STENÁCTIS**, Cass. *Outermost bristles of the pappus short and minute, or rather chaffy, forming an external series: otherwise as § 2.*

6. **E. glabellum**, Nutt. Stem (6' - 15' high) stout, hairy above, the leafless summit bearing 1-7 large heads; leaves nearly glabrous, except the margins, entire, the upper oblong-lanceolate and pointed, closely sessile or partly clasping, the lower spatulate and petioled; rays (more than 100, purple) more than twice the length of the hoary-hispid involucre. — Plains N. Wisconsin, and westward. June.

§ 4. **PHALACROLÔMA**, Cass. *Rays numerous, but nearly in a single row, conspicuous: pappus plainly double, the outer a crown of minute chaffy-bristle-form scales; the inner of scanty capillary bristles which are deciduous, or entirely wanting in the ray: annuals and biennials.*

7. **E. ánnuum**, Pers. (DAISY FLEABANE. SWEET SCABIOUS.) Stem stout (3° - 5° high), branched, beset with spreading hairs; leaves coarsely and sharply toothed; the lowest ovate, tapering into a margined petiole; the upper ovate-lanceolate, acute and entire at both ends; heads corymbed; rays white, tinged with purple, not twice the length of the bristly involucre. (*E. heterophyllum*, Muhl. *E. strigòsum*, Bigel.) — Fields and waste places: a very common weed. June - Aug. (Nat. in Eu.)

8. **E. strigòsum**, Muhl. (DAISY FLEABANE.) Stem paniced-corymbose at the summit, roughish like the leaves with minute appressed hairs, or almost smooth; leaves entire or nearly so, the upper lanceolate, scattered, the lowest oblong or spatulate, tapering into a slender petiole; rays white, twice the length of the minutely hairy involucre. (*E. integrifolium*, Bigel.) — Fields, &c.: common. June - Aug. — Stem smaller and more simple than the last, with smaller heads but longer rays. — Var. **DISCOFDEUM**, Robbins, has the rays minute, scarcely exceeding the involucre. — Uxbridge, Massachusetts, and adjacent parts of Connecticut and Rhode Island, Dr. Robbins.

§ 5. **ERIGERÍDIUM**, Torr. & Gr. *Rays about 30, in a single row, rather broad: pappus simple: achenia mostly 4-nerved: root perennial.*

9. **E. vérnum**, Torr. & Gr. Glabrous; leaves clustered at the root, oval or spatulate; scape leafless, slender (1° - 2° high), bearing 5-12 small corymbed heads; rays white. (*E. nudicaule*, Michx. *Aster vernus*, L.) — Low grounds, E. Virginia and southward. May.

15. DIPLOPÁPPUS, Cass. DOUBLE-BRISTLED ASTER.

Heads many-flowered, radiate; the rays 8-12, pistillate. Scales of the involucre imbricated, appressed, narrow, 1-nerved or keeled, destitute of herbaceous tips. Receptacle flat, alveolate. Achenia flattish. Pappus double; the outer of very short and small stiff bristles, the inner of capillary bristles as long as the disk-corolla. — Perennials with corymbose or simple heads: disk-flowers yellow; rays white or violet. (Name composed of διπλός, *double*, and πάππος, *pappus*, the character which distinguishes the genus from Aster.)

§ 1. *Rays violet, showy: head solitary, pretty large: involucre much imbricated: achenia silky: bristles of the inner pappus all alike.*

1. **D. linariifolius**, Hook. Stems (6'-20' high) several from the same woody root, mostly simple, very leafy; leaves rigid, spreading, linear, strongly 1-nerved, smooth, rough-margined. — Dry soil: common. Sept., Oct.

§ 2. *Rays white: heads small, corymbed: involucre shorter than the disk, imbricated in about 3 rows: achenia smoothish: bristles of the inner pappus unequal, some of them thickened at the tip: leaves rather large, scattered, membranaceous, veiny, entire.*

2. **D. umbellatus**, Torr. & Gr. Smooth, leafy to the top (2°-6° high); leaves lanceolate, elongated, taper-pointed and tapering at the base (3'-6' long); heads very numerous in compound flat corymbs; scales of the involucre rather close, obtusish. — Moist thickets: common, especially northward. Aug.

3. **D. amygdalinus**, Torr. & Gr. Smooth or roughish above, leafy; leaves ovate-lanceolate, acute, abruptly narrowed at the base; scales of the involucre loose, obtuse. — Low grounds, New Jersey, Penn., and southward. Aug. — Too near the last; but lower, rougher, and with broader and shorter leaves.

4. **D. cornifolius**, Darl. Stem (1°-2° high) pubescent, bearing few heads on divergent peduncles; leaves elliptical or ovate-lanceolate, conspicuously pointed at both ends, ciliate, hairy on the veins underneath. — Woodlands, E. Massachusetts to Kentucky, and southward along the mountains. July-Sept.

16. BOLTÓNIA, L'Her. BOLTONIA.

Heads many-flowered, radiate; the rays numerous, pistillate. Scales of the hemispherical involucre imbricated somewhat in 2 rows, appressed, with narrow membranaceous margins. Receptacle conical or hemispherical, naked. Achenia very flat, obovate or inversely heart-shaped, margined with a callous wing, or in the ray 3-winged, crowned with a pappus of several minute bristles and usually 2-4 longer awns. — Perennial and bushy-branched smooth herbs, pale green, with the aspect of Aster: the thickish leaves chiefly entire, often turned edgewise. Flowers autumnal: disk yellow: rays white or purplish. (Dedicated to James Bolton, an English botanist of the last century.)

* *Heads middle-sized, loosely corymbed.*

1. **B. asteroides**, L'Her. Leaves lanceolate; achenia broadly oval; pappus of few minute bristles and no awns. — Moist places along streams. Pennsylvania (*Bartram*), and southward along the Alleghanies: rare. Oct. — Plant usually 6° high.

2. **B. glastifolia**, L'Her. Leaves lanceolate, ascending; achenia obovate, broadly winged; pappus of several short bristles and 2 or 3 short awns. — Rich soil, Pennsylvania to Illinois and southward. Sept., Oct. — Plant 2°–4° high.

* * Heads small, paniced on the slender branches.

3. **B. diffusa**, L'Her. Stem diffusely branched; leaves lance-linear, those on the branchlets very small and awl-shaped; pappus of several very short bristles and 2 short awns. — Prairies near Centralia, Illinois (*Vasey*), and southwestward. Aug. – Oct.

17. BÉLLIS, Tourn. DAISY.

Heads many-flowered, radiate; the rays numerous, pistillate. Scales of the involucre herbaceous, equal, in about 2 rows. Receptacle conical, naked. Achenia obovate, flattened, wingless, and without any pappus. — Low herbs (all but our single species natives of the Old World), either stemless, like the true *Daisy*, *B. perennis*, or leafy-stemmed, as is the following. (The Latin name, from *bellus*, pretty.)

1. **B. integrifolia**, Michx. (WESTERN DAISY.) Annual or biennial, diffusely branched (4'–9' high), smoothish; leaves lanceolate or oblong, the lower spatulate-obovate; heads on slender peduncles; rays pale violet-purple. — Prairies and banks, Kentucky and southwestward. March–June.

18. BRACHYCHÆTA, Torr. & Gr. FALSE GOLDEN-ROD.

Heads and flowers nearly as in *Solidago*, except the pappus, which is a row of minute rather scale-like bristles, shorter than the achenia. — A perennial herb, with rounded or ovate serrate leaves, all the lower ones heart-shaped; the small yellow heads in sessile clusters racemed or spiked on the branches. (Name composed of *βραχύς*, short, and *χαίτη*, bristle, from the pappus.)

1. **B. cordata**, Torr. & Gr. (*Solidago cordata*, *Short*.) Wooded hills, E. Kentucky and southward. Oct. — Plant 2°–4° high, slender, more or less pubescent.

19. SOLIDÀGO, L. GOLDEN-ROD.

Heads few–many-flowered, radiate; the rays 1–16, pistillate. Scales of the oblong involucre appressed, destitute of herbaceous tips (except No. 1). Receptacle small, not chaffy. Achenia many-ribbed, nearly terete. Pappus simple, of equal capillary bristles. — Perennial herbs, with mostly wand-like stems and nearly sessile stem-leaves, never heart-shaped. Heads small, racemed or clustered: flowers both of the disk and ray (except No. 2) yellow. (Name from *solido*, to join, or make whole, in allusion to reputed vulnerary qualities.) Flowering in autumn.

§ 1. **CHRYSÁSTRUM**, Torr. & Gr. Scales of the much imbricated and rigid involucre with abruptly spreading herbaceous tips: heads in clusters or glomerate racemes, disposed in a dense somewhat leafy and interrupted wand-like compound spike.

1. **S. squarrosa**, Muhl. Stem stout (2°–5° high), hairy above; leaves large, oblong, or the lower spatulate-oval and tapering into a margined petiole,

serrate, veiny; disk-flowers 16 - 24, the rays 12 - 16. — Rocky and wooded hills, Maine and W. Vermont to Pennsylvania, and the mountains of Virginia: rather rare.

§ 2. *VIRGAÛREA*, Tourn. *Scales of the involucre destitute of herbaceous tips: rays mostly fewer than the disk-flowers: heads all more or less pedicelled.*

* *Heads clustered in the axils of the feather-veined leaves.*

2. *S. bicolor*, L. *Hoary or grayish with soft hairs; stem mostly simple; leaves oblong or elliptical-lanceolate, acute at both ends, or the lower oval and tapering into a petiole, slightly serrate; clusters or short racemes from the axils of the upper leaves, forming an interrupted spike or crowded panicle; rays small, cream-color or nearly white.* — Var. *CÓNCOLOR* has the rays yellow. — Dry copses and banks: the var. in Pennsylvania and westward.

3. *S. latifolia*, L. *Smooth or nearly so, stem angled, zigzag, simple or paniculate-branched (1° - 3° high); leaves broadly ovate or oval, very strongly and sharply serrate, conspicuously pointed at both ends (thin, 3' - 6' long); heads in very short axillary clusters, or somewhat prolonged at the end of the branches.* — Moist shaded banks: common northward, and along the mountains.

4. *S. cæsia*, L. *Smooth; stem terete, mostly glaucous, at length much branched and diffuse; leaves lanceolate or oblong-lanceolate, serrate, pointed, sessile; heads in very short clusters, or somewhat racemose-panicled on the branches.* — Moist rich woodlands: common.

* * *Racemes terminal, erect, either somewhat simple and wand-like, or compound and panicled, not one-sided: leaves feather-veined. (Not maritime.)*

+ *Heads small: leaves nearly entire, except the lowermost.*

5. *S. virgata*, Michx. *Very smooth throughout; stem strict and simple, wand-like (2° - 4° high), slender, beset with small and entire appressed lanceolate-oblong thickish leaves, which are gradually reduced upwards to mere bracts; the lowest oblong-spatulate; heads crowded in a very narrow compound spicate raceme; rays 5 - 7.* — Damp pine barrens, New Jersey and southward.

6. *S. pubérula*, Nutt. *Stem (1° - 3° high, simple or branched) and panicle minutely hoary; stem-leaves lanceolate, acute, tapering to the base, smoothish; the lower wedge-lanceolate and sparingly toothed; heads very numerous and crowded in compact short racemes forming a prolonged and dense narrow or pyramidal panicle; scales of the involucre linear-awl-shaped, appressed; rays about 10.* — Sandy soil, Maine to Virginia and southward, near the coast.

7. *S. stricta*, Ait. *Very smooth throughout; stem simple, strict (2° - 3° high); leaves lanceolate, pointed, the lower tapering into winged petioles, partly sheathing at the base, minutely serrate above with appressed teeth; racemes much crowded and appressed in a dense wand-like panicle; scales of the involucre linear-oblong, obtuse; rays 5 - 6, small.* — Peat-bogs, Maine to Pennsylvania, Wisconsin, and northward. Root-leaves 6' - 10' long. Flowers earlier than most species, beginning in July.

+ + *Heads rather large, at least for the size of the plant.*

8. *S. speciosa*, Nutt. *Stem stout (3° - 6° high), smooth; leaves thickish, smooth with rough margins, oval or ovate, slightly serrate, the uppermost oblong-lanceolate, the lower contracted into a margined petiole; heads somewhat*

crowded in numerous erect racemes, forming an ample pyramidal or thyrsiform panicle; peduncles and pedicels rough-hairy; scales of the cylindrical involucre oblong, obtuse; rays about 5, large. — Var. *ANGUSTATA* is a dwarf form, with the racemes short and clustered, forming a dense interrupted or compound spike. — Copses, Maine to Wisconsin and southward. — A very handsome species; the lower leaves 4'–6' long and 2'–4' wide in the larger forms.

9. **S. petiolaris**, Ait. *Minutely hoary or downy*; stem strict, simple, (1°–3° high); leaves small ($\frac{1}{2}$ '–2' long), oval or oblong, mucronate, veiny, rough-ciliate; the upper entire and abruptly very short-petioled, the lower often serrate and tapering to the base; heads few, in a wand-like raceme or panicle, on slender bracted pedicels; rays about 10, elongated: scales of the pubescent involucre lanceolate or linear-awl-shaped, the outer ones loose and spreading, more or less foliaceous, — especially in var. *SQUARRULOSA*, Torr. & Gr. — S. W. Illinois (*Dr. Engelmann*), and southward. — The name is misleading, as the leaves are hardly petioled.

10. **S. Virga-aurea**, L. Pubescent or nearly glabrous; stem low (6'–18' high) and simple; leaves lanceolate or oblanceolate, or the lowest spatulate or elliptical-obovate and petioled, serrate with small appressed teeth or nearly entire; racemes thyrsoid or simple, narrow; scales of the involucre lanceolate or linear, acute; rays 8–12. — An extremely variable species in the Old World and in our northern regions; perhaps including several. (Eu.)

Var. **alpina**, Bigel. Dwarf (1'–8' high), with few (1–12) pretty large heads (3''–4'' long, becoming smaller as they increase in number); leaves thickish, mostly smooth; scales of the involucre lanceolate, acute or acutish; rays about 12. — Alpine region of the mountains of Maine, New Hampshire, and New York; and shore of Lake Superior.

Var. **humilis**. Low (6'–12' high) and smooth, bearing several or numerous loosely thyrsoid smaller heads, which, with the peduncles, &c., are mostly somewhat glutinous; scales of the involucre obtuse; rays 6–8, short; leaves varying from narrowly lanceolate and nearly entire to oblanceolate and serrate. (*S. humilis*, Pursh, Torr. & Gr.) — Rocky banks, W. Vermont, and along the Great Lakes northward. Also on islands in the Susquehanna, near Lancaster, Penn., *Prof. Porter*! Great Falls of the Potomac, Virginia, *Dr. Robbins*! — At the base of the White Mountains of New Hampshire, on gravelly banks of streams, occurs a form, with the minutely pubescent stout stem 1°–2° high, the leaves larger and broader, and the heads very numerous in an ample compound raceme; the rays occasionally almost white.

11. **S. thyrsoidea**, E. Meyer. Stem stout (1°–4° high), wand-like, pubescent near the summit, simple; leaves thin, ovate, irregularly and coarsely serrate with sharp salient teeth, large (1'–4' long), all but the uppermost abruptly contracted into long and margined petioles; heads large (5''–6'' long), many-flowered, crowded in an oblong or wand-like raceme or contracted panicle (2'–18' long); scales of the involucre loose and thin, long, lanceolate, taper-pointed; rays 8–10, elongated; achenia smooth. (*S. Virga-aurea*, Pursh. *S. leiocarpa*, DC.) — Wooded sides of mountains, N. Maine to New York (south to the Catskills), shore of Lake Superior, and northward. — Very near an European form of *S. Virga-aurea*.

* * * *Heads in a compound corymb terminating the simple stem, not at all racemose: leaves mostly with a strong midrib.*

12. **S. rigida**, L. *Rough and somewhat hoary with a minute pubescence; stem stout (3°–5° high), very leafy; corymb dense; leaves oval or oblong, copiously feather-veined, thick and rigid; the upper closely sessile by a broad base, slightly serrate, the uppermost entire; heads large, about 34-flowered; the rays 7–10.* — Dry soil, Connecticut to N. Wisconsin and southward.

13. **S. Ohioensis**, Riddell. *Very smooth throughout; stem wand-like, slender, leafy (2°–3° high); stem-leaves oblong-lanceolate, flat, entire, obscurely feather-veined, closely sessile; the lower and radical ones elongated, slightly serrate towards the apex, tapering into long margined petioles; heads numerous, on smooth pedicels, small, 16–20-flowered; the rays 6 or 7.* — Moist meadows or prairies, W. New York to Ohio and Wisconsin. — Root-leaves 1° long; the upper reduced to 1'–2', with rough margins, like the rest.

14. **S. Riddellii**, Frank. *Smooth and stout (2°–4° high), very leafy, the branches of the dense corymb and pedicels rough-pubescent; leaves linear-lanceolate, elongated (4'–6' long), entire, acute, partly clasping or sheathing, conduplicate and mostly recurved, the lowest elongated-lanceolate and tapering into a long keeled petiole, obscurely 3-nerved; heads very numerous, clustered, 20–24-flowered; the rays 7–9.* — Wet grassy prairies, Ohio to Wisconsin and Illinois. — Heads larger than in the last, 2''–3'' long. Stem-leaves upright and partly sheathing at the base, then gradually recurved-spreading.

15. **S. Houghtonii**, Torr. & Gray. *Smooth; stem rather low and slender (1°–2° high); leaves scattered, linear-lanceolate, acutish, flat, entire, tapering into a narrowed slightly clasping base, or the lower into margined petioles; heads few or several, 20–30-flowered; the rays 9 or 10.* — North shore of Lake Michigan; collected in the Michigan State Survey. Swamps at West Bergen, Genesee Co., New York, J. A. Paine, &c. July, Aug. — Leaves rough-margined, 2'–5' long, 2''–4'' wide, 1-nerved, or the lower obscurely 3-nerved above; veins obscure. Heads large, nearly $\frac{1}{2}$ ' long. Scales of the involucre obtuse.

* * * * *Heads in one-sided more or less spreading or recurved racemes: leaves veiny, not 3-ribbed, but sometimes obscurely triple-nerved.*

+ *Leaves thickish, very smooth, entire, elongated, obscurely veiny: heads rather large.*

16. **S. sempervirens**, L. *Smooth and stout (1°–6° high); leaves fleshy, lanceolate, slightly clasping, or the lower ones lanceolate-oblong, obscurely triple-nerved; racemes short, in an open or contracted panicle.* — Varies, in less brackish swamps, with thinner elongated linear-lanceolate leaves, tapering to each end, and more erect racemes in a narrower panicle. — Salt marshes, or rocks on the shore, Maine to Virginia. — Heads showy: the golden rays 8–10.

+ + *Leaves usually ample, serrate, loosely feather-veined, or rarely slightly triple-nerved: heads middle-sized.*

17. **S. elliptica**, Ait. *Smooth; stem stout (1°–3° high), very leafy; leaves elliptical or oblong-lanceolate, acute (2'–3' long), closely sessile, slightly serrate, strongly veined, thick, smooth both sides, shining above; heads in dense spreading racemes which are crowded in a close pyramidal panicle; peduncles and achenia strigose-pubescent.* — Swamps (fresh or brackish) near the coast, Mass. to New Jersey and southward. — Heads showy, 3'' long; the rays 8–12.

18. **S. neglécta**, Torr. & Gray. *Smooth*; stem stout (2° – 3° high); leaves *thickish, smooth both sides, opaque*; the upper *oblong-lanceolate*, mostly acute and nearly entire; the lower *ovate-lanceolate or oblong*, sharply serrate, tapering into a petiole; racemes *short and dense, at length spreading*, disposed in an elongated or pyramidal close panicle; peduncles and achenia nearly glabrous. — Swamps, Maine to Penn. and Wisconsin. — Heads rather large, crowded; the racemes at first erect and scarcely one-sided.

19. **S. pátula**, Muhl. *Stem strongly angled, smooth* (3° – 5° high); leaves ($4'$ – $8'$ long) *ovate*, acute, serrate, pale, *very smooth and veiny underneath*, but the upper surface *very rough*, like shagreen; racemes rather short and numerous on the spreading branches. — Swamps: common.

20. **S. argúta**, Ait. *Smooth throughout* (1° – 4° high); radical and lower stem-leaves *elliptical or lanceolate-oval, sharply serrate* with spreading teeth, *pointed*, tapering into winged and ciliate petioles; the others *lanceolate or oblong*, slightly triple-nerved, tapering to each end, the uppermost entire; racemes *dense, naked, at length elongated and recurved, forming a crowded and flat corymb-like panicle*: rays 8–12, *small*. — Var. 1. **JÚNCEA** has the leaves narrower and less serrate, or all the upper entire. — Var. 2. **SCABRÉLLA** is somewhat roughish-pubescent (Wisconsin, &c.). — Copses and banks: common, especially the first variety. — Well distinguished by its long or drooping racemes, and the closely appressed rigid scales of the involucre, small rays, &c. But the name is a bad one, as even the root-leaves are seldom much toothed.

21. **S. Muhlenbérghii**, Torr. & Gr. *Smooth*; stem angled; leaves (large and thin) *ovate*, and the upper *elliptical-lanceolate, very sharply and strongly serrate, pointed at both ends*, the lowest on margined petioles; racemes *pubescent, spreading, disposed in an elongated open panicle*; rays 6–7, *large*. — Copses and moist woods, New Hampshire to Penn. — Racemes much shorter and looser than in the last; the involucreal scales thin and more slender.

22. **S. linoides**, Solander. *Smooth*; stem slender, simple ($10'$ – $20'$ high); leaves *lanceolate*, serrate with small appressed teeth, narrowed at the base, the lower tapering into margined ciliate petioles, the uppermost oblong; racemes *short, crowded in one or 3–4 small one-sided panicles* ($3'$ – $4'$ long); heads small and few-flowered; rays 1–3. — Bogs, New England (near Boston and Providence), to the pine barrens of New Jersey.

+ + + *Leaves broad, not large, sessile or short-petioled, coarsely and sharply serrate, copiously feather-veined and conspicuously reticulated*: heads small: rays short.

23. **S. altíssima**, L. *Rough-hairy, especially the stem* ($2'$ – $7'$ high); leaves *ovate-lanceolate, elliptical or oblong, often thickish and very rugose*; racemes paniced, spreading; scales of the involucre linear; rays 6–9; the disk-flowers 4–7. — Borders of fields and copses: very common, presenting a great variety of forms: but instead of the tallest, as its name denotes, it is usually one of the lowest of the common *Golden-rods*. It flowers early. Aug. – Sept.

24. **S. ulmifolia**, Muhl. *Stem smooth*, the branches hairy; leaves *thin, elliptical-ovate or oblong-lanceolate, pointed, tapering to the base, loosely veined*, beset with soft hairs beneath; racemes paniced, recurved-spreading; scales of the involucre lanceolate-oblong; rays about 4. — Low copses: common. — Too near the last; distinguished only by its smooth stem and thin larger leaves.

25. **S. Drummóndii**, Torr. & Gr. Stem (1° – 3° high) and lower surface of the broadly ovate or oval somewhat triple-ribbed leaves minutely velvety-pubescent, some of the leaves almost entire; racemes paniced, short; scales of the involucre oblong, obtuse; rays 4 or 5. — Rocks, Illinois opposite St. Louis, and southwestward.

+ + + + Leaves entire or nearly so, thickish, reticulate-veiny, but the veins obscure.

26. **S. pilósa**, Walt. Stem stout, upright (3° – 7° high), clothed with spreading hairs, often paniced at the summit; leaves oblong-lanceolate, roughish, hairy beneath, at least on the midrib, serrulate, the upper ovate-lanceolate or oblong and entire, closely sessile; racemes many, recurved, crowded in a dense pyramidal panicle; rays 7–10, very short. — Low grounds, pine barrens of New Jersey to Virginia, and southward.

27. **S. odóra**, Ait. (SWEET GOLDEN-ROD.) Smooth or nearly so throughout; stem slender (2° – 3° high), often reclined; leaves linear-lanceolate, entire, shining, pellucid-dotted; racemes spreading in a small one-sided panicle; rays 3–4, rather large. — Border of thickets in dry or sandy soil, Vermont and Maine to Kentucky and southward. — The crushed leaves yield a pleasant anisate odor. But an occasional form, var. *INODÓRA*, is nearly scentless.

+ + + + Leaves more or less grayish or hoary, thickish, feather-veined and slightly triple-nerved, obscurely serrate or entire; heads middle-sized.

28. **S. nemoralis**, Ait. Clothed with a minute and close grayish-hoary (soft or roughish) pubescence; stem simple or corymbed at the summit ($\frac{1}{2}^{\circ}$ – $2\frac{1}{2}^{\circ}$ high); leaves oblanceolate or spatulate-oblong, the lower somewhat crenate-toothed and tapering into a petiole; racemes numerous, dense, at length recurved, forming a large and crowded compound raceme or panicle which is usually turned to one side; scales of the involucre linear-oblong, appressed; rays 6–9. — Dry sterile fields: very common. — Flowers very bright yellow, beginning early in Aug.

29. **S. rádula**, Nutt. Stem and oblong or obovate-spatulate leaves rigid and very rough, not hoary, the upper sessile; scales of the involucre oblong, rigid; rays 3–6: otherwise nearly as in *S. nemoralis*, of which it is most probably a greener and rough variety. — Dry Hills, Western Illinois and southwestward.

* * * * Heads in one-sided spreading or recurved racemes, forming an ample panicle; leaves plainly 3-ribbed, or triple-ribbed.

+ Scales of the involucre thickish and rigid, closely imbricated, and with somewhat greenish tips or midrib: leaves rigid, smooth and shining.

30. **S. Shórtii**, Torr. & Gr. Stem slender, simple (1° – 3° high), minutely roughish-pubescent; leaves oblong-lanceolate, acute, the lower sharply serrate above the middle with scattered fine teeth; racemes mostly short in a crowded panicle; achenia silky-pubescent. — Rocks at the Falls of the Ohio, &c. — A handsome species: heads 3'' long, narrow.

31. **S. Missouriénsis**, Nutt. Smooth throughout (1° – 3° high); leaves linear-lanceolate, or the lower broadly lanceolate, tapering to both ends, with very rough margins, the lower very sharply serrate; heads and dense crowded racemes nearly as in No. 20; achenia nearly glabrous. — Dry prairies, from Wisconsin and Illinois southward and westward. — Heads $1\frac{1}{2}$ ''–2'' long.

+ + *Scales of the involucre narrow, thin and membranaceous: racemes mostly elongated and numerous, forming a crowded ample panicle.* (These all present intermediate forms, and should rather be regarded as one polymorphous species.)

32. **S. rupéstris**, Raf. *Stem smooth and slender (2°-3° high); leaves linear-lanceolate, tapering to both ends, smooth and glabrous, entire or nearly so; panicle narrow; heads very small; rays very short.* — Rocky river-banks, Kentucky and Indiana.

33. **S. Canadénsis**, L. *Stem rough-hairy, tall and stout (3°-6° high); leaves lanceolate, pointed, sharply serrate (sometimes almost entire), more or less pubescent beneath and rough above; heads small; rays very short.* — Borders of thickets and fields: very common. — Varies greatly in the roughness and hairiness of the stem and leaves, the latter oblong-lanceolate or elongated linear-lanceolate; — in var. *PRÓCERA*, whitish-woolly underneath; and in var. *SCÁBRA* also very rough above, often entire, and rugose-veined.

34. **S. serótina**, Ait. *Stem very smooth, tall and stout (4°-8° high), often glaucous; leaves lanceolate, pointed, serrate, roughish above, smooth except the veins underneath, which are more or less hairy; rays short.* — Thickets and low grounds: common. — Intermediate in character, and in the size of the heads and rays, between the last and the next.

35. **S. gigánteá**, Ait. *Stem stout (3°-7° high), smooth, often glaucous; leaves quite smooth both sides, lanceolate, taper-pointed, very sharply serrate, except the narrowed base, rough-ciliate; the ample panicle pubescent; rays rather long.* — Copses and fence-rows: common: — presenting many varieties, but with decidedly larger heads and rays than in the preceding. Seldom very tall.

§ 3. **EUTHÀMIA**, Nutt. *Corymbosely much branched: heads small, sessile, in little clusters which are crowded in flat-topped corymbs; the closely appressed scales of the involucre somewhat glutinous: receptacle fimbriate: rays 6-20, short, more numerous than the disk-flowers: leaves narrow, entire, sessile, crowded or fasciated in the axils.*

36. **S. lanceolàta**, L. *Leaves lanceolate-linear, 3-5-nerved; the nerves, margins, and angles of the branches minutely rough-pubescent; heads obovoid-cylindrical, in dense corymbed clusters; rays 15-20.* — River-banks, &c., in moist soil: common. — Stem 2°-4° high: leaves 3'-5' long.

37. **S. tenuifolia**, Pursh. *Smooth, slender; leaves very narrowly linear, mostly 1-nerved, dotted: heads obovoid-club-shaped, in numerous clusters of 2 or 3, disposed in a loose corymb; rays 6-12.* — Sandy fields, Massachusetts to Illinois, and southward: common near the coast.

20. BIGELÓVIA, DC. RAYLESS GOLDEN-ROD.

Heads 3-4-flowered, the flowers all perfect and tubular: rays none. Involucre club-shaped, yellowish; the rigid somewhat glutinous scales linear, closely imbricated and appressed. Receptacle narrow, with an awl-shaped prolongation in the centre. Achenia somewhat obconical, hairy. Pappus a single row of capillary bristles. — A smooth perennial; the slender stem (1°-2° high) simple or branched from the base, naked above, corymbose at the summit, bearing

small heads in a flat-topped corymb. Flowers yellow. Leaves scattered, oblanceolate or linear, 1-3-nerved. (Dedicated by DeCandolle to *Dr. Jacob Bigelow*, author of the *Florula Bostoniensis*, and of the *American Medical Botany*.)

1. **B. nudata**, DC. — Low pine barrens, New Jersey (rare), and southward. Sept.

21. CHRYSÓPSIS, Nutt. GOLDEN ASTER.

Heads many-flowered, radiate; the rays numerous, pistillate. Scales of the involucre linear, imbricated, without herbaceous tips. Receptacle flat. Achenia obovate or linear-oblong, flattened, hairy. Pappus in all the flowers double, the outer of very short and somewhat chaffy bristles, the inner of long capillary bristles. — Chiefly perennial, low herbs, woolly or hairy, with rather large often corymbose heads terminating the branches. Disk and ray-flowers yellow. (Name composed of *χρυσός*, *gold*, and *ὄψις*, *aspect*, from the golden blossoms.)

* *Leaves narrowly lanceolate or linear: achenia linear.*

1. **C. graminifolia**, Nutt. *Silvery-silky*, with long close-pressed hairs; stem slender, often with runners from the base, naked above, bearing few heads; *leaves lanceolate or linear, elongated, grass-like, nerved, shining, entire*. — Dry sandy soil, Delaware to Virginia, and southward. July - Oct.

2. **C. falcata**, Ell. *Stems (4' - 10' high) very woolly; leaves crowded, linear, rigid, about 3-nerved, entire, somewhat recurved or scythe-shaped, hairy*, or smooth when old, sessile; heads (small) corymbed. — Dry sandy soil on the coast, pine barrens of New Jersey to Nantucket and Cape Cod, Mass. Aug.

* * *Leaves oblong or lanceolate, entire or slightly serrate, mostly sessile, veined, not nerved; achenia obovate, flattened.*

3. **C. gossypina**, Nutt. *Densely woolly all over; leaves oblong, obtuse (1' - 2' long)*; heads larger than in the next. — Pine barrens, Virginia and southward. Aug. - Oct.

4. **C. Mariàna**, Nutt. *Silky with long and weak hairs*, or when old smoothish; *leaves oblong*; heads corymbed, on glandular peduncles. — Dry barrens, from S. New York and Penn., southward, near the coast. Aug. - Oct.

5. **C. villosa**, Nutt. *Hirsute and villous-pubescent*; stem corymbosely branched, the branches terminated by single short-peduncled heads; *leaves narrowly oblong, hoary with rough pubescence (as also the involucre), bristly-ciliate toward the base*. — Dry plains and prairies, Wisconsin to Kentucky, and westward. July - Sept.

22. ÍNULA, L. ELECAMPANE.

Outer scales of the involucre sometimes leaf-like. Achenia terete or 4-sided. Pappus simple, of capillary bristles. Anthers with 2 tails at their base. Otherwise much as in the last genus. (The ancient Latin name.)

1. **I. HELÈNIUM**, L. (COMMON ELECAMPANE.) Stout perennial (3° - 5° high); leaves large, woolly beneath; those from the thick root ovate, petioled, the others partly clasping; rays very many, narrow. — Roadsides, escaped from gardens. Aug. — Heads very large. Root mucilaginous. (Nat. from Eu.)

23. PLÛCHEA, Cass. MARSH-FLEABANE.

Heads many-flowered; the flowers all tubular; the central perfect, but sterile, few, with a 5-cleft corolla; all the others with a thread-shaped truncate corolla, pistillate and fertile. Involucre imbricated. Anthers with tails. Achenia grooved. Pappus capillary, in a single row. — Herbs, somewhat glandular, emitting a strong or camphoric odor, the heads in close compound corymbs. Flowers purplish, in summer. (Dedicated to the Abbé *Pluche*.)

1. **P. camphorata**, DC. (SALT-MARSH FLEABANE.) *Annual, minutely viscid, pale* (1° – 2° high); *leaves scarcely petioled, oblong-ovate or lanceolate, thickish, obscurely veiny, serrate; corymb flat; involucre viscid-downy.* (*Coryza camphorata, Bigel. C. Marylandica, Pursh.*) — Salt marshes, Massachusetts to Virginia and southward.

2. **P. foetida**, DC. *Perennial, almost smooth* (2° – 4° high); *leaves distinctly petioled, veiny, oval-lanceolate, pointed at both ends, serrate; corymbs paniced; involucre smooth.* River-banks, Ohio to Illinois, and southward.

3. **P. bifrons**, DC. *Perennial; leaves closely sessile or half-clasping by a somewhat heart-shaped base, lance-oblong, sharply denticulate, veiny* (only 2'–3' long); heads clustered in a corymb; involucre smooth. — Salisbury, Maryland (*W. M. Canby*), and southward.

24. BÁCCHARIS, L. GROUNDSEL-TREE.

Heads many-flowered; the flowers all tubular, dioecious, viz. the pistillate and staminate flowers in separate heads borne by different plants. Involucre imbricated. Corolla of the pistillate flowers very slender and thread-like; of the staminate, larger and 5-lobed. Anthers tailless. Achenia ribbed. Pappus of slender capillary bristles, in the sterile plant scanty and tortuous; in the fertile plant very long and copious. — Shrubs, commonly smooth and resinous or glutinous. Flowers whitish or yellow, autumnal. (The name of some shrub anciently dedicated to *Bacchus*.)

1. **B. halimifolia**, L. Smooth and somewhat scurfy; branches angled; leaves obovate and wedge-form, coarsely toothed, or the upper entire; heads scattered or in leafy panicles; scales of the involucre acutish. — Sea beach, Massachusetts to Virginia, and southward. — Shrub 6° – 12° high; the fertile plant conspicuous in autumn by its very long and white pappus.

2. **B. glomeruliflora**, Pers. Leaves spatulate-oblong; heads larger, sessile in the axils or in clusters; scales of the bell-shaped involucre broader, very obtuse: otherwise like the last. — Pine barrens, E. Virginia and southward.

25. POLÝMNIA, L. LEAF-CUP.

Heads many-flowered, radiate: the rays several (rarely abortive), pistillate; the disk-flowers perfect but sterile. Scales of the involucre in two rows; the outer about 5, leaf-like, large and spreading; the inner small and membranaceous, partly embracing the thick triangular-obovoid achenia. Receptacle flat, with a membranaceous chaff to each flower. Pappus none. — Tall branching

perennial herbs, viscid-hairy, exhaling a heavy odor. Leaves large and thin, opposite, or the uppermost alternate, lobed, and with dilated appendages like stipules at the base. Heads in paniced corymbs. Flowers light yellow; in summer and autumn. (Dedicated to the Muse, *Polyhymnia*, for no obvious reason, as the plants are coarse and inelegant.)

1. **P. Canadensis**, L. *Clammy-hairy*; lower leaves deeply pinnatifid, the uppermost triangular-ovate and 3-5-lobed or angled, petioled; heads small; rays few, obovate or wedge-form, shorter than the involucre, whitish-yellow. — Moist shaded ravines, W. New York (and Weehawken, New Jersey, *Dr. Allen*) to Penn., Wisconsin, and southward along the mountains. — Var. **DISCOIDEA** has the corolla of the ray-flowers reduced to a mere ring around the base of the style. Mt. Carroll, Illinois, *Henry Shimer*.

2. **P. Uvedalia**, L. *Roughish-hairy, stout* (4° – 10° high); leaves broadly ovate, angled and toothed, nearly sessile; the lower palmately lobed, abruptly narrowed into a winged petiole; outer involucre scales very large; rays 10-15, linear-oblong, much longer than the inner scales of the involucre, yellow. — Rich soil, W. New York to Illinois and southward.

26. CHRYSÓGONUM, L. CHRYSOGONUM.

Heads many-flowered, radiate; the rays about 5, pistillate and fertile; the disk-flowers perfect but sterile. Involucre of about 5 exterior leaf-like oblong scales, which exceed the disk, and as many interior shorter and chaff-like concave scales. Receptacle flat, with a linear chaff to each disk-flower. Achenia all in the ray, obovate, obcompressed, 4-angled, each one partly enclosed by the short scale of the involucre behind it; those of the disk-flowers abortive. Pappus a small chaffy crown, 2-3-toothed, and split down the inner side. — A low ($2'$ – $6'$ high), hairy, perennial herb, nearly stemless when it begins to flower, the flowerless shoots forming runners. Leaves opposite, ovate or spatulate, crenate, long-petioled. Heads single, long-peduncled. Flowers yellow. (Name composed of *χρυσός*, *golden*, and *γόνυ*, *knee*.)

1. **C. Virginianum**, L. Dry soil, from Pennsylvania (*Mercersburg, Prof. Porter*) and Illinois southward. May–Aug. — Rays $\frac{1}{2}'$ long.

27. SÍLPHIUM, L. ROSIN-PLANT.

Heads many-flowered, radiate; the rays numerous, pistillate and fertile, their broad flat ovaries imbricated in 2 or 3 rows; the disk-flowers apparently perfect, but with undivided style, and sterile. Scales of the broad and flattish involucre imbricated in several rows, broad and with loose leaf-like summits, except the innermost, which are small and resemble the linear chaff of the flat receptacle. Achenia broad and flat, obcompressed, surrounded by a wing which is notched at the top, destitute of pappus, or with 2 teeth confluent with the winged margin: achenia of the disk sterile and stalk-like. — Coarse and tall rough perennial herbs, with a copious resinous juice, and large corymbose-paniced yellow-flowered heads. (*Σίλφιον*, the ancient name of a plant which produced some gum-resin, perhaps *assafœtida*, was transferred by Linnæus to this American genus.)

* *Stem terete, naked above, alternate-leaved near the base (root very large and thick).*

1. **S. laciniatum**, L. (ROBIN-WEEED. COMPASS-PLANT.) *Rough-bristly throughout; stem stout (3°-6° high), leafy to the top; leaves pinnately parted, petioled but dilated and clasping at the base; their divisions lanceolate or linear, acute, cut-lobed or pinnatifid, rarely entire; heads few (1'-2' broad), somewhat racemed; scales of the involucre ovate, tapering into long and spreading rigid points; achenia broadly winged and deeply notched.* — Prairies, Michigan and Wisconsin, thence southward and westward. July. — Lower and root-leaves vertical, 12'-30' long, ovate in outline; on the wide open prairies disposed to present their edges north and south; hence called *Compass-Plant*.

2. **S. terebinthinaceum**, L. (PRAIRIE DOCK.) *Stem smooth, slender (4°-10° high), paniced at the summit and bearing many (small) heads, leafless except towards the base; leaves ovate and ovate-oblong, somewhat heart-shaped, serrate-toothed, thick, rough, especially beneath (1°-2° long, on slender petioles); scales of the involucre roundish, obtuse, smooth; achenia narrowly winged, slightly notched and 2-toothed.* — Var. **PINNATIFIDUM** has the leaves deeply cut or pinnatifid, but varies into the ordinary form. — Prairies and oak-openings, Ohio and Michigan to Wisconsin and southward. July — Sept.

* * *Stem terete or slightly 4-angled, leafy: leaves undivided (not large).*

3. **S. trifoliatum**, L. *Stem smooth, often glaucous, rather slender (4°-6° high), branched above; stem-leaves lanceolate, pointed, entire or scarcely serrate, rough, short-petioled, in whorls of 3 or 4, the uppermost opposite; heads loosely paniced; achenia rather broadly winged, and sharply 2-toothed at the top.* — Dry plains and banks, W. New York to Wisconsin and southward. Aug.

4. **S. Asteriscus**, L. *Stem hispid (2°-4° high); leaves opposite, or the lower in whorls of 3, the upper alternate, oblong or oval-lanceolate, coarsely toothed, rarely entire, rough-hairy; heads nearly solitary (large); achenia obovate, winged, 2-toothed.* — Dry sandy soil, Virginia and southward.

5. **S. integrifolium**, Michx. *Stem rough, rather stout (2°-4° high), rigid, 4-angular and grooved; leaves all opposite, rigid, lanceolate-ovate, entire, tapering to a sharp point from a roundish heart-shaped and partly clasping base, rough-pubescent or nearly smooth, thick (3'-5' long); heads in a close forking corymb, short-peduncled; achenia broadly winged, deeply notched.* — Var. **LÆVE** has the stem and leaves smooth or nearly so. — Prairies, Michigan to Wisconsin, and southward. Aug.

* * * *Stem square: leaves opposite, connate (thin and large, 6'-15' long).*

6. **S. perfoliatum**, L. (CUP-PLANT.) *Stem stout, often branched above (4°-8° high); leaves ovate, coarsely toothed, the upper united by their bases and forming a cup-shaped disk, the lower abruptly narrowed into winged petioles which are connate by their bases; heads corymbose; achenia winged and variously notched.* — Rich soil along streams, Michigan to Wisconsin, and southward: common. Also escaped from gardens eastward. July.

28. PARTHËNIUM, L. PARTHENIUM.

Heads many-flowered, inconspicuously radiate; the 5 ray-flowers with very short and broad obcordate ligules not projecting beyond the woolly disk, pistil-

late and fertile; the disk-flowers staminate with imperfect styles, sterile. Involucre hemispherical, of 2 ranks of short ovate or roundish scales. Receptacle conical, chaffy. Achenia only in the ray, obcompressed, surrounded by a slender callous margin, crowned with the persistent ray-corolla and a pappus of 2 small chaffy scales. — Leaves alternate. Heads small, corymbed; the flowers whitish. (An ancient name of some plant, from *παρθένος*, *virgin*.)

1. *P. integrifolium*, L. Rough-pubescent perennial (1°–3° high); leaves oblong or ovate, crenate-toothed, or the lower (3'–6' long) cut-lobed below the middle; heads many in a very dense flat corymb. — Dry soil, Maryland to Wisconsin and southward. June–Aug.

29. ÌVA, L. MARSH ELDER. HIGHWATER-SHRUB.

Heads several-flowered, not radiate; the pistillate fertile and the staminate sterile flowers in the same heads, the former few (1–5) and marginal, with a small tubular or no corolla; the latter with a funnel-form 5-toothed corolla. Anthers nearly separate. Scales of the involucre few, roundish. Receptacle small, with narrow chaff among the flowers. Achenia obovoid or lenticular. Pappus none. — Herbaceous or shrubby coarse plants, with thickish leaves, the lower opposite, and small greenish-white heads of flowers; in summer and autumn. (Name of unknown derivation.)

§ 1. *Fertile flowers with a small tubular corolla: involucre simple (heads nodding in the axils of leaf-like bracts, forming spikes or racemes).*

1. *I. frutescens*, L. Shrubby at the base, nearly smooth (3°–8° high); leaves oval or lanceolate, coarsely and sharply toothed, rather fleshy, the upper reduced to linear bracts, in the axils of which the heads are disposed, in leafy paniced racemes; fertile flowers and scales of the involucre 5. — Salt marshes, coast of Massachusetts to Virginia, and southward.

2. *I. ciliata*, Willd. Annual (2°–8° high), rough and hairy; leaves ovate, pointed, coarsely toothed, downy beneath, on slender ciliate petioles; heads in dense spikes, with conspicuous ovate-lanceolate rough-ciliate bracts; scales of the involucre and fertile flowers 3–5. — Moist ground, from Illinois southward.

§ 2. EUPHRÓSYNE, DC. *Fertile flowers 5, in the axils of as many thin membranaceous scales of the involucre, which loosely enwrap the achenia, their corolla a mere rudimentary ring or none.*

3. *I. xanthiifolia*, Nutt. Annual, tall, roughish; leaves nearly all opposite, hoary with minute down, ovate, rhombic, or the lowest heart-shaped, doubly or cut-toothed, or obscurely lobed; heads small, crowded in spikes or clusters disposed in axillary and terminal panicles; bracts inconspicuous. — N. W. Wisconsin (*T. J. Hale*), and northwestward.

30. AMBRÒSIA, Tourn. RAGWEED.

Sterile and fertile flowers occupying different heads on the same plant; the fertile 1–3 together and sessile in the axil of leaves or bracts, at the base of the racemes or spikes of sterile heads. Sterile involucre flattish or top-shaped, composed of 7–12 scales united into a cup, containing 5–20 funnel-form stam-

inate flowers, with slender chaff intermixed, or none. Anthers almost separate. Fertile involucre (fruit) oblong or top-shaped, closed, pointed, resembling an achenium, and usually with 4-8 tubercles or horns near the top in one row, enclosing a single flower which consists of a pistil only; the elongated branches of the style protruding. Achenia ovoid: pappus none. — Homely and coarse weeds, with opposite or alternate lobed or dissected leaves, and inconspicuous greenish or whitish flowers, produced throughout late summer and autumn: our species are all annuals. (*'Αμβροσία*, *the food of the gods*, an ill-chosen name for these miserable weeds.)

§ 1. *Sterile heads sessile, crowded in a dense cylindrical spike, the top-shaped involucre with its truncate margin extended on one side into a large, lanceolate, hooded, recurved, bristly-hairy tooth or appendage; fertile involucre oblong and 4-angled.*

1. **A. bidentata**, Michx. Hairy (1° - 3° high), very leafy; leaves alternate, lanceolate, partly clasping, nearly entire, except a short lobe or tooth on each side near the base. — Prairies of Illinois and southward.

§ 2. *Sterile heads in single or paniced racemes or spikes, the involucre regular.*

* *Leaves opposite, only once lobed: sterile involucre 3-ribbed on one side.*

2. **A. trifida**, L. (GREAT RAGWEED.) Stem stout (4° - 12° high), rough-hairy, as are the large deeply 3-lobed leaves, the lobes oval-lanceolate and serrate; petioles margined; fruit obovate, 5-6-ribbed and tubercled. — Var. INTEGRIFOLIA is only a smaller form, with the upper leaves, or all of them, undivided, ovate or oval. — Moist river-banks: common.

* * *Leaves many of them alternate, all once or twice pinnatifid.*

3. **A. artemisiæfolia**, L. (ROMAN WORMWOOD. HOG-WEED. BITTER-WEED.) Much branched (1° - 3° high), hairy or roughish-pubescent; leaves thin, twice-pinnatifid, smoothish above, paler or hoary beneath; fruit obovoid or globular, armed with about 6 short acute teeth or spines. — Waste places everywhere. — An extremely variable weed, with finely cut leaves; rarely the spikes bear all fertile heads.

4. **A. psilostachya**, DC. Paniculate-branched (2° - 5° high), rough and somewhat hoary with short hispid hairs; leaves once pinnatifid, thickish, the lobes acute, those of the lower leaves often incised; fruit obovoid, without tubercles or with very small ones, pubescent. (*A. coronopifolia*, Torr. & Gr.) — Prairies and plains, Wisconsin, Illinois, and southwestward.

31. XÁNTHIUM, Tourn. COCKLEBUR. CLOTBUR.

Sterile and fertile flowers occupying different heads on the same plant; the latter clustered below, the former in short spikes or racemes above. Sterile involucre and flowers as in Ambrosia, but the scales separate. Fertile involucre closed, coriaceous, ovoid or oblong, clothed with hooked prickles so as to form a rough bur, 2-celled, 2-flowered; the flowers consisting of a pistil with a slender thread-form corolla. Achenia oblong, flat, destitute of pappus. — Coarse and vile weeds, with annual roots, low and branching stout stems, and alternate toothed or lobed petioled leaves; flowering in summer and autumn. (Name from *ξάνθος*, *yellow*, in allusion to the color the plants are said to yield.)

1. **X. strumarium**, L. (COMMON COCKLEBUR.) Rough; stems unarmed; leaves dilated-triangular and more or less heart-shaped, on long petioles, toothed and cut or obscurely lobed; fruit oval or oblong ($\frac{1}{2}$ ' - $\frac{2}{3}$ ' long), pubescent on the lower part of and between the hooked prickles, and with two strong and usually straight beaks at the summit. — Barn-yards, &c. (Nat. from Eu.) — Varies into forms with more spotted stems, and often larger fruit ($\frac{2}{3}$ ' - 1' long), which is either glabrous, glandular, or glandular-hairy, the prickles longer and the beaks often incurved. (X. Canadense, Mill., &c.) — River-banks, &c., common westward; apparently indigenous. And this passes into

Var. **echinatum**. (X. echinatum, Murr., &c.) Fruit turgid (1' long), thickly clothed with long prickles, glandular-hispid, the beaks commonly incurved. — Sandy sea-shore, and along the Great Lakes and rivers. Perhaps an immigrant from farther south. Now widely scattered over the warmer parts of the world.

2. **X. spinosum**, L. (SPINY CLOTBUR.) Hoary-pubescent; stems slender, with slender yellow 3-parted spines at the base of the lanceolate or ovate-lanceolate leaves; these taper into a short petiole, are white-downy beneath, often 2-3-lobed or cut; fruit ($\frac{1}{2}$ ' long) pointed with a single short beak. — Waste places on the sea-board and along rivers, southward. (Nat. from Trop. Amer.?)

32. TETRAGONOTHÈCA, Dill. TETRAGONOTHECA.

Heads many-flowered, radiate; the rays 6 - 9, fertile. Involucre double; the outer of 4 large and leafy ovate scales, which are united below by their margins into a 4-angled or winged cup; the inner of as many small and chaffy scales as there are ray-flowers, and partly clasping their achenia. Receptacle convex or conical, with narrow and membranaceous chaff between the flowers. Achenia roundish and obovoid, flat at the top. Pappus none. — An erect perennial herb, viscidly hairy when young, with opposite and coarsely toothed oval or oblong leaves, their sessile bases sometimes connate, and large single heads of pale yellow flowers, on terminal peduncles. (Name compounded of *τετράγωνος*, four-angled, and *θήκη*, a case, from the shape of the involucre.)

1. **T. helianthoides**, L. — Sandy soil, Virginia and southward. June.

33. ECLÍPTA, L. ECLIPTA.

Heads many-flowered, radiate; the rays short, fertile; the disk-flowers perfect, 4-toothed. Scales of the involucre 10 - 12, in 2 rows, leaf-like, ovate-lanceolate. Receptacle flat, with almost bristle-form chaff between the flowers. Achenia short, 3 - 4-sided, or in the disk laterally flattened, roughened on the sides, hairy at the summit; the pappus none, or an obscure denticulate crown. — Annual or biennial rough herbs, with slender stems and opposite lanceolate or oblong leaves. Heads solitary, small. Flowers whitish: anthers brown. (Name from *ἐκλείπω*, to be deficient, alluding to the absence of pappus.)

1. **E. procumbens**, Michx. Rough with close appressed hairs; stems procumbent, creeping, or ascending; leaves oblong-lanceolate, acute at each end, sessile, slightly serrate; peduncles many times longer than the head. — Var. BRACHÝPODA has the peduncles not more than twice the length of the heads. —

Wet river-banks, Pennsylvania to Illinois, and southward. June - Oct. — All belong to a wide-spread and variable species of the warmer regions, the oldest specific name of which is *E. ALBA*.

34. BORRÍCHIA, Adans. SEA OX-EYE.

Heads many-flowered, radiate; the rays fertile. Scales of the hemispherical involucre imbricated. Receptacle flat, covered with lanceolate rigid and persistent chaff. Achenia somewhat wedge-shaped, 3-4-angled. Pappus a short 4-toothed crown. — Shrubby low maritime plants, coriaceous or fleshy, with opposite nearly entire leaves, and solitary peduncled terminal heads of yellow flowers: anthers blackish. (Named for *Olof Borrich*, a Danish botanist.)

1. *B. frutescens*, DC. Whitened with a minute silky pubescence (6' - 12' high); leaves spatulate-oblong or lanceolate, often toothed near the base; chaff rigidly pointed. — Virginia and southward.

35. HELIÓPSIS, Pers. OX-EYE.

Heads many-flowered, radiate; the rays 10 or more, fertile. Scales of the involucre in 2 or 3 rows; the outer leaf-like and somewhat spreading, the inner shorter than the disk. Receptacle conical: chaff linear. Achenia smooth, 4-angular. Pappus none, or a mere border. — Perennial herbs, like *Helianthus*. Heads showy, peduncled, terminating the stem or branches. Leaves opposite, petioled, triple-ribbed, serrate. Flowers yellow. (Name composed of $\eta\lambda\iota\omicron\varsigma$, *the sun*, and $\delta\psi\iota\varsigma$, *appearance*, from the likeness to the Sunflower.)

1. *H. lævis*, Pers. Nearly smooth (1° - 4° high); leaves ovate-lanceolate or oblong-ovate. — Var. *SCABRA* has roughish foliage, and the involucre somewhat hoary. — Banks and copses: common. Aug.

36. ECHINÀCEA, Mœnch. PURPLE CONE-FLOWER.

Heads many-flowered, radiate; the rays very long, drooping, pistillate but sterile. Scales of the involucre imbricated, lanceolate, spreading. Receptacle conical; the lanceolate spiny-tipped chaff longer than the disk-flowers. Achenia thick and short, 4-sided. Pappus a small toothed border. — Perennial herbs, with the stout and nearly simple stems naked above and terminated by a single large head; the leaves chiefly alternate, 3-5-nerved. Rays rose-purple, rather persistent: disk purplish. (Name formed from $\epsilon\chi\acute{\iota}\nu\omicron\varsigma$, *the Hedgehog*, or *Sea-urchin*, in allusion to the spiny chaff of the disk.)

1. *E. purpurea*, Mœnch. Leaves rough, often serrate; the lowest ovate, 5-nerved, veiny, long-petioled; the others ovate-lanceolate; involucre imbricated in 3-5 rows; stem smooth, or in one variety (*E. serotina*, DC.) rough-bristly, as well as the leaves. — Prairies and banks, from W. Penn. and Ohio southward and westward. July. — Rays 15-20, dull purple (rarely whitish), 1'-2' long. Root thick, black, very pungent to the taste, used in popular medicine under the name of *Black Sampson*. — Probably varies into

2. *E. angustifolia*, DC. Leaves, as well as the slender simple stem, bristly-hairy, lanceolate and linear-lanceolate, 3-nerved, entire; involucre less imbricated; rays 12-15 (2' long), rose-color or red. — Plains from Illinois and Wisconsin southwestward. June - Aug.

37. **RUDBÉCKIA**, L. CONE-FLOWER.

Heads many-flowered, radiate; the rays neutral. Scales of the involucre leaf-like, in about 2 rows, spreading. Receptacle conical or columnar; the short chaff concave, not rigid. Achenia 4-angular, smooth, not margined, flat at the top, with no pappus, or a minute crown-like border. — Chiefly perennial herbs, with alternate leaves, and showy heads terminating the stem or branches; the rays generally long, yellow. (Named in honor of the *Professors Rudbeck*, father and son, predecessors of Linnæus at Upsal.)

* *Disk columnar in fruit, dull greenish-yellow: leaves divided and cut.*

1. **R. laciniata**, L. Stem smooth, branching (3° – 7° high); leaves smooth or roughish, the lowest pinnate, with 5–7 cut or 3-lobed leaflets; upper leaves irregularly 3–5-parted; the lobes ovate-lanceolate, pointed, or the uppermost undivided; heads long-peduncled; chaff truncate and downy at the tip; rays linear ($1'$ – $2'$ long), drooping. — Low thickets: common. July–Sept.

** *Disk globular, pale brownish: lower leaves 3-parted: receptacle sweet-scented.*

2. **R. subtomentosa**, Pursh. Stem branching above (3° – 4° high), downy, as well as the lower side of the ovate or ovate-lanceolate serrate leaves; heads short-peduncled; chaff downy at the blunt apex. — Prairies, Wisconsin, Illinois, and southward.

*** *Disk broadly conical, dark purple or brown: leaves undivided, except No. 3.*

3. **R. triloba**, L. Hairy, *biennial*, much branched (2° – 5° high), the branches slender and spreading; *upper leaves ovate-lanceolate*, sparingly toothed, *the lower 3-lobed*, tapering at the base, *coarsely-serrate* (those from the root pinnately parted or undivided); rays 8, *oval or oblong*; chaff of the black-purple disk smooth, *awned*. — Dry soil, Penn. to Illinois, and southward. Aug. — Heads small, but numerous and showy.

4. **R. speciosa**, Wenderoth. Roughish-hairy (1° – 2° high), branched; the branches upright, elongated and naked above, terminated by single large heads; leaves lanceolate or ovate-lanceolate, pointed at both ends, *petioled*, 3–5-nerved *coarsely and unequally toothed or incised*; involucre much shorter than the numerous elongated ($1'$ – $1\frac{1}{2}'$) rays; chaff of the dark purple disk acutish, smooth. — Dry soil, W. Penn. to Ohio and Virginia. July.

5. **R. fulgida**, Ait. Hairy, the branches naked at the summit and bearing single heads; leaves spatulate-oblong or lanceolate, partly clasping, triple-nerved, *the upper entire, mostly obtuse*; rays about 12, equalling or exceeding the involucre; chaff of the dark purple disk nearly smooth and blunt. — Dry soil, Pennsylvania to Kentucky and southward. — Variable, 1° – 3° high: the rays orange-yellow. Nearly approaches the next.

6. **R. hirta**, L. Very rough and bristly-hairy throughout; stems simple or branched near the base, stout (1° – 2° high), naked above, bearing single large heads; leaves nearly entire; the upper oblong or lanceolate, sessile; the lower spatulate, triple-nerved, petioled; rays (about 14) more or less exceeding the involucre; chaff of the dull brown disk hairy at the tip, acutish. — Dry soil, W. New York to Wisconsin and southward. Now common eastward, as a weed in meadows, of recent introduction, with clover-seed from the West. June–Aug.

38. **LÉPACHYS**, Raf. (OBELISCARIA, DC.)

Heads many-flowered, radiate; the rays few, neutral. Scales of the involucre few and small, spreading. Receptacle oblong or columnar: the chaff truncate, thickened and bearded at the tip, partly embracing the flattened and margined achenia. Pappus none or 2 teeth. — Perennial herbs, with alternate pinnate leaves; the grooved stems or branches naked above, terminated by single showy heads. Rays yellow or party-colored, drooping; the disk grayish. (Name from *λεπίς*, a scale, and *παχύς*, thick, from the thickened tips of the chaff.)

1. **L. pinnata**, Torr. & Gr. Hoary with minute appressed hairs, slender (4° high), branching; leaflets 3 - 7, lanceolate, acute; disk oblong, much shorter than the large and drooping light-yellow rays (which are 2' long). — Dry soil, W. New York (*Dr. Sartwell*), to Wisconsin and southward. July. — The receptacle exhales a pleasant anisate odor when bruised. Achenia slightly margined on the inner edge, obscurely 2-toothed at the top.

39. **HELIÁNTHUS**, L. SUNFLOWER.

Heads many-flowered, radiate; the rays several or many, neutral. Involucre imbricated. Receptacle flattish or convex; the persistent chaff embracing the 4-sided and laterally compressed achenia, which are neither winged nor margined. Pappus very deciduous, of 2 thin chaffy scales on the principal angles of the achenium, and often 2 or more little intermediate scales. — Coarse and stout herbs, with solitary or corymbed heads, and yellow rays; flowering towards autumn. (Named from *ἥλιος*, the sun, and *ἄνθος*, a flower.)

§ 1. *Root annual: leaves alternate: receptacle flat: disk brownish.*

1. **H. ánnuus**, L. (COMMON SUNFLOWER.) Tall, rough; leaves triple-ribbed, ovate or the lower heart-shaped; heads in cult. plant very broad, and rays many. — Escaped from cult. into waste grounds. (Adv. from trop. Amer.)

§ 2. *Root perennial, the creeping rootstocks sometimes tuberous-thickened or tuberiferous.*

* *Disk convex, dark purple: leaves mainly opposite: heads small, except No. 4.*

2. **H. angustifolius**, L. Stem slender (2° - 6° high); leaves long and linear, sessile, entire, with revolute margins, 1-nerved; heads loosely corymbed, long-peduncled; scales of the involucre tapering into narrow and spreading herbaceous tips. — Low pine barrens, New Jersey to Kentucky and southward.

3. **H. atrorubens**, L. *Rough-hairy; stem slender* (2° - 5° high), smooth, and naked and forking above; leaves thin, ovate or oval, or the lowest heart-shaped (3' - 6' long), serrate, abruptly contracted into a margined petiole; heads small, corymbed; scales of the involucre ovate, obtuse, regularly imbricated, appressed, destitute of herbaceous tips; rays 10 - 16; pappus of 2 fringed scales. — Dry soil, Virginia, Illinois, and southward.

4. **H. rigidus**, Desf. *Stem stout* (1° - 3° high), simple or sparingly branched, rough; leaves very thick and rigid, rough both sides, oblong-lanceolate, usually pointed at both ends, nearly sessile, slightly serrate, the lowest oval; heads nearly solitary, pretty large; scales of the involucre ovate or lance-oblong, obtuse, ciliate, appressed, destitute of herbaceous tips; rays 20 - 25; pappus of 2 large and often several small scales. — Dry prairies, Michigan to Illinois, and westward.

* * *Disk convex, yellow: scales of the involucre regularly imbricated and appressed, with somewhat spreading and acute (but not foliaceous) tips: leaves chiefly opposite.*

5. **H. lætiflorus**, Pers. *Stout and rough* (3° – 4° high), branching above; *leaves oval-lanceolate, very rough both sides, narrowed into short petioles, serrate, taper-pointed, the uppermost alternate and nearly entire; heads single or corymbed, on naked peduncles; scales of the involucre ovate-lanceolate, pointed, ciliate.* — Dry open places, Ohio to Illinois, and southward. — Leaves almost as thick as in *H. rigidus*. Rays showy, 1'–2' long.

6. **H. occidentalis**, Riddell. *Somewhat hairy; stem slender, simple, naked above* (1° – 3° high, sending out runners from the base), bearing 1–5 small heads on long peduncles; *lowest leaves oval or lanceolate-ovate, 3-nerved, obscurely serrate, roughish-pubescent beneath, abruptly contracted into long hairy petioles; the upper small and remote, entire; scales of the involucre oval-lanceolate, pointed, ciliate.* — Dry barrens, Ohio to Wisconsin, Kentucky, and southward.

7. **H. cinereus**, var. **Sullivantii**, Torr. & Gr. *Gray with a close roughish pubescence; stem branching above, hairy; leaves ovate-oblong, sessile by a narrowed base, acute, obscurely serrate; the upper small and remote; peduncles slender; scales of the involucre lanceolate, hoary.* — Darby Plains, Ohio, *Sullivant*. Stem 2° – 3° high, bearing few heads, as large as those of the next.

8. **H. mollis**, Lam. *Stem clothed with soft white hairs, simple, leafy, to the top* (2° – 4° high); *leaves ovate, with a broad heart-shaped and clasping base, pointed, nearly entire, hoary above, very soft white-woolly and reticulated underneath; scales of the involucre lanceolate, downy.* — Barrens and Prairies, Ohio to Illinois, and westward.

* * * *Heads very small (about 4" broad): scales of the involucre few, shorter than the yellow disk, irregularly imbricated, appressed, the outer with spreading foliaceous pointed tips: rays 5–8, slender: leaves all but the uppermost opposite.*

9. **H. microcéphalus**, Torr. & Gr. *Stem smooth* (3° – 8° high), with numerous slender branches above; *leaves thin, ovate-lanceolate, taper-pointed, somewhat serrate, veiny, petioled, rough above, downy or hairy underneath; peduncles slender, rough; scales of the involucre ovate and ovate-lanceolate, ciliate.* — Thickets, Pennsylvania to Illinois, and southward.

10. **H. lævigatus**, Torr. & Gr. *Stem slender* (1° – 4° high), simple or sparingly branched, *very smooth and glabrous throughout, as well as the slightly serrate lanceolate leaves.* — Dry soil, Alleghany Mountains, west of the Warm Springs of Virginia, and southward.

* * * * *Heads middle-sized or large: scales of the involucre irregularly imbricated, loose, with spreading foliaceous tips, as long as the yellow disk or longer.*

† *Leaves chiefly alternate or scattered, feather-veined, sometimes obscurely triple-ribbed.*

11. **H. giganteus**, L. *Stem hairy or rough* (3° – 10° high), branched above; *leaves lanceolate, pointed, serrate, very rough above, rough-hairy beneath, narrowed and ciliate at the base, but nearly sessile; scales of the involucre long, linear-lanceolate, pointed, hairy, or strongly ciliate.* — Var. **ambiguus** has most of the leaves opposite and closely sessile by an obtuse base, and approaches No. 14. — Low thickets and swamps: common. Heads somewhat corymbed: the pale yellow rays 15–20.

12. **H. grosse-serratus**, Martens. *Stem smooth and glaucous*, at least below (5° - 10° high); *leaves elongated-lanceolate* or *ovate-lanceolate*, taper-pointed, serrate, rough above, rounded or acute at the base, *petioled, hoary-downy beneath*; scales of the involucre lance-awl-shaped, slightly ciliate. — Dry plains, Ohio to Illinois, and southwestward. — Probably runs into the last.

13. **H. tomentosus**, Michx. *Stem hairy, stout* (4° - 8° high); *leaves oblong-lanceolate, or the lowest ovate*, taper-pointed, obscurely serrate, large ($5'$ - $12'$ long), somewhat petioled, *very rough above, soft-downy beneath*; scales of the involucre with very long and spreading tips, hairy; the chaff and tips of the disk-flowers pubescent. (Disk $1'$ broad; rays 12 - 16 about $1'$ long.) — Rich woods, Illinois? Virginia and southward along the mountains.

+ + *Leaves opposite, or the uppermost alternate, 3-nerved or triple-ribbed.*

14. **H. strumosus**, L. *Stem* (3° - 4° high) smooth below; *leaves ovate-lanceolate, tapering gradually to a point, serrate with small appressed teeth*, abruptly contracted into short margined petioles, rough above, *whitish and naked or minutely downy underneath*; scales of the involucre broadly lanceolate with spreading tips, equalling the disk; rays mostly 10. — Var. **MOLLIS** has the leaves softly downy underneath. — River-banks and low copses. common, especially westward.

15. **H. divaricatus**, L. *Stem simple or forked and corymbed at the top* (1° - 4° high) smooth; *leaves all opposite and divaricate, ovate-lanceolate, 3-nerved from the rounded or truncate sessile base, tapering gradually to a sharp point* ($3'$ - $6'$ long), serrate, *thickish, rough both sides*; scales of the involucre lanceolate from a broad base, pointed, equalling the disk; rays 8 - 12. — Thickets and barrens: common. — Disk $6''$ wide; rays $1'$ long.

16. **H. hirsutus**, Raf. *Stem simple or forked above, stout* (1° - 2° high), *bristly-hairy*; *leaves more or less petioled, ovate-lanceolate, gradually pointed, slightly serrate*, rounded or obtuse at the base, *very rough above, rough-hairy underneath*; scales of the involucre ovate-lanceolate, pointed, equalling the disk; rays about 12. — Dry plains, Ohio to Wisconsin, and southward. — Runs into the last.

17. **H. trachelifolius**, Willd. *Stem loosely branched, tall, hairy*; *leaves thin, ovate-lanceolate or oblong-lanceolate, taper-pointed, sharply serrate, smoothish or roughish-pubescent both sides*, contracted into short petioles; scales of the involucre lanceolate-linear, elongated and very taper-pointed, loose, exceeding the disk; rays 12 - 15. — Copses, Pennsylvania and Ohio to Illinois, and southward. — Probably runs into the next.

18. **H. decapetalus**, L. *Stem branching* (3° - 6° high), smooth below; *leaves thin and green both sides, smooth or roughish, ovate, coarsely serrate*, pointed, abruptly contracted into marginal petioles; scales of the involucre lanceolate-linear, elongated, loosely spreading, the outer longer than the disk; rays about 10. — Var. **FRONDOSUS** has the outer involucral scales foliaceous or changing to leaves. — Copses and low banks of streams: common, especially northward. (*H. multiflorus*, L., may be a cultivated state of this.)

19. **H. doricoides**, Lam. *Stem stout* (5° - 9° high), branching, *rough-hairy above*; *leaves ovate or oblong-lanceolate, pointed, serrate, strongly triple-veined*, rough above, *smoothish or downy underneath*, the lower often heart-shaped and on

margined petioles; scales of the involucre linear-lanceolate, pointed, scarcely exceeding the disk; rays 12-15. — River-bottoms, Pennsylvania to Illinois and southward. — A coarse species, with ample leaves (the lower often 1° long); the upper ones frequently alternate; rootstock thickening into elongated tubers. This is probably the original of

20. **H. TUBEROSUS**, L., the **JERUSALEM ARTICHOKE**, (i. e. *Girasole* of the Italians, meaning the same as sunflower, and corrupted in England into *Jerusalem*), which has all the upper leaves alternate. It has escaped from gardens into fence-rows, &c. in some places.

40. **ACTINÓMERIS**, Nutt. **ACTINOMERIS.**

Heads many-flowered; the rays few or several, neutral, or rarely none. Involucre foliaceous, nearly equal, in 1 to 3 rows. Receptacle convex or conical, chaffy; the chaff embracing the outer margin of the flat (laterally compressed) and winged achenia. Pappus of 2 smooth persistent awns. — Tall and branching perennials, with serrate feather-veined leaves, tapering to the base and mostly decurrent on the stem. Heads corymbed: flowers chiefly yellow. (Name from *ἀκρίς*, a ray, and *μέρις*, a part; alluding to the irregularity of the rays.)

1. **A. squarrosa**, Nutt. *Stem somewhat hairy and winged above* (4°-8° high); leaves alternate or the lower opposite, oblong or ovate-lanceolate, pointed at both ends; heads in an open corymbed panicle; scales of the involucre in 2 rows, the outer linear-spatulate, *reflexed*; rays 4-10, *irregular*; achenia broadly winged; receptacle globular. — Rich soil, Penn. and W. New York (*Sartwell*) to Michigan, Illinois, and southward. Sept.

2. **A. helianthoides**, Nutt. *Stem hairy* (1°-3° high), *widely winged* by the ovate-lanceolate sessile alternate leaves, which are rough above and soft-hairy beneath; heads few; scales of the involucre *not spreading*; rays 8-15, *regular*; achenia oval, slightly winged, tipped with 2 fragile awns; receptacle conical. — Prairies and copses, Ohio to Illinois and southward. July.

41. **COREÓPSIS**, L. **TICKSEED.**

Heads many-flowered, radiate; the rays mostly 8, neutral, rarely wanting. Involucre double; each of about 8 scales, the outer rather foliaceous and somewhat spreading; the inner broader and appressed, nearly membranaceous. Receptacle flat, with membranaceous chaff deciduous with the fruit. Achenia flat, obcompressed (i. e. parallel with the scales of the involucre), often winged, not beaked or narrowed at the top, 2-toothed, 2-awned, or sometimes naked at the summit, the awns not barbed downwardly. — Herbs, generally with opposite leaves, and yellow or party-colored, rarely purple, rays. (Name from *κόψις*, a bug, and *ὄψις*, resemblance; from the form of the fruit.)

§ 2. *Branches of the style truncate: rays rose-color: disk yellow: root perennial.*

1. **C. ròsea**, Nutt. *Stem branching, leafy, smooth* (6'-20' high); leaves opposite, linear, entire; heads small, somewhat corymbed, on short peduncles; outer involucre very short; rays 3-toothed; achenia oblong, wingless; pappus an obscure crown-like border. — Sandy and grassy swamps, Plymouth, Massachusetts, to New Jersey, and southward: rare. Aug.

§ 3. *Corolla of the ray and disk yellow or partly brown: branches of the style tipped with a pointed or acute appendage.*

* *Achenia* nearly orbicular, broadly winged, incurved, furnished with a callous tubercle on the inside at the top and bottom, crowned with 2 small chaff-like denticulate teeth: outer involucre about the length of the inner: rays large, coarsely 3-5-toothed: leaves opposite or the uppermost alternate: heads on long naked peduncles: root in our wild species perennial.

2. **C. auriculata**, Linn. Pubescent or glabrous; stems 1°-4° high, branching, sometimes with runners; leaves mostly petioled, the upper oblong or oval-lanceolate, entire; the lower oval or roundish, some of them variously 3-5-lobed or divided; scales of the outer involucre oblong-linear or lanceolate. — Rich woods and banks, Virginia to Illinois and southward. June - Sept.

3. **C. lanceolata**, L. Smooth or hairy (1°-2° high); stems short, tufted, branched only at the base; leaves all entire (or the lower rarely with a pair of small lateral lobes), lanceolate, sessile, the lowest oblanceolate or spatulate, tapering into petioles; scales of the outer involucre ovate-lanceolate. — Rich or damp soil, Michigan and Illinois to Virginia, and southward. July. Also cultivated in gardens. — Heads showy: rays 1' long.

C. TINCTORIA, Nutt., a native of the plains beyond the Mississippi, with the rays yellow above, and brown-purple towards the base, a common garden biennial or annual, is becoming spontaneous in a few places.

* * *Achenia* oblong, narrowly winged, minutely or obscurely 2-toothed at the summit: scales of the outer involucre narrow, about the length of the inner, all more or less united at the base: rays mostly entire and acute: leaves opposite, sessile, mostly 3-divided, therefore appearing as if whorled: perennial (1°-3° high).

4. **C. senifolia**, Michx. Leaves each divided into 3 sessile ovate-lanceolate entire leaflets, therefore appearing like 6 in a whorl: plant minutely soft-pubescent. — Sandy woods, Virginia and southward. July.

Var. **stellata**, Torr. & Gr. Glabrous, and the leaves narrower. (*C. stellata*, Nutt.) — Virginia, Kentucky, and southward.

5. **C. delphinifolia**, Lam. Glabrous or nearly so; leaves divided into 3 sessile leaflets which are 2-5-parted, their divisions lance-linear (1''-3'' broad), rather rigid; disk brownish. — Pine woods, Virginia and southward. July.

6. **C. verticillata**, L. Glabrous; leaves divided into 3 sessile leaflets which are 1-2-pinnately parted into narrowly linear or filiform divisions. — Damp soil, from Maryland and Michigan southward. Also cultivated in old gardens, but not showy. July - Sept.

7. **C. palmata**, Nutt. Nearly smooth, simple; leaves broadly wedge-shaped, deeply 3-cleft, rigid; the lobes broadly linear, entire, or the middle one 3-lobed. — Prairies, Michigan to Wisconsin, and southwestward. July.

* * * *Achenia* elliptical, narrowly winged, the narrowly notched summit of the wing minutely lacerate-toothed: scales of the outer involucre foliaceous, much smaller than the inner, all united at the base: rays obtuse, entire: leaves opposite, petioled, 3-5-divided: perennial.

8. **C. tripteris**, L. (TALL COREOPSIS.) Smooth; stem simple (4°-9° high), corymbed at the top; leaflets lanceolate, acute, entire. (*Chrysostemma*,

Less.)—Penn. to Michigan, Illinois and southward. Aug. — Sept. — Heads exhaling the odor of anise when bruised: disk turning brownish.

* * * *Achenia* wingless, flat, 2-awned, 2-toothed, or rarely truncate, 1-nerved or ridged on each face: scales of the outer involucre leafy, reflexed or spreading: leaves opposite, petioled, generally pinnately or ternately compound, the leaves or leaflets serrate: biennials or annuals, with the aspect of *Bidens*, intermediate between that genus and *Coreopsis*; — and certain ambiguous (hybridized?) specimens, with the awns barbed some upwards and some downwards, connect the two.

+ Rays conspicuous, golden-yellow.

9. **C. aristosa**, Michx. Somewhat pubescent; leaves 1–2-pinnately 5–7-divided, petioled; leaflets lanceolate, cut-toothed or pinnatifid; heads panicled-corymbose; outer involucre of 10–12 leafy bracts; *achenia* obovate, often obscurely wing-margined, bristly-ciliate, with 2 (rarely 4) long and slender diverging awns as long as the achenium itself. — Swamps, Ohio to Michigan, Wisconsin, and southwestward. Aug. — Oct. — Var. *MUTICA* has two short divergent teeth or points in place of the awns. — W. Illinois and southwestward, where a form (*C. involucrata*, *Nutt.*) occurs with the bracts of the outer involucre more leafy, numerous, and hirsute. Mr. Fritchey sends, from near St. Louis, specimens with short awns and their barbs either spreading or retrorse, and others with long awns retrorsely barbed, — perhaps hybrids with some *Bidens*.

10. **C. trichosperma**, Michx. (TICKSEED SUNFLOWER.) Smooth, branched; leaves short-petioled, 5–7-divided; leaflets lanceolate or linear, cut-toothed, or the upper leaves only 3–5-cleft and almost sessile; heads panicled-corymbose; *achenia* narrowly wedge-oblong or the inner ones wedge-linear, smooth or minutely appressed-hairy, marginless, crowned with 2 erect triangular or awl-shaped stout teeth. — Swamps, Massachusetts to Virginia near the coast. Also Buffalo, New York (*G. W. Clinton*), and Illinois (*Vasey*), where forms with shorter *achenia* approach the Southern *C. aurea*. Aug. — Oct.

+ + Rays none, or rarely small and inconspicuous: outer involucre of few (usually 3–5) loose leafy bracts commonly surpassing the heads: *achenia* minutely appressed-pubescent: plants glabrous, 1°–3° high.

11. **C. discoidea**, Torr. & Gr. Diffusely branched; leaves ternately divided, slender-petioled; leaflets ovate-lanceolate, pointed, coarsely serrate; heads 2''–3'' long; *achenia* linear-wedge-shaped (2''–3'' long), bearing a pair of short and stout upwardly-barbed awns of the length of the corolla. — Wet banks and swamps, Delaware (*W. M. Canby*) to Ohio, Illinois, and southward. July.

Anomalous specimens, from near Philadelphia (*A. H. Smith*, &c.), growing with *Bidens frondosa*, differ from small forms of that only in the less rigid and upturned barbs of the awns.

12. **C. bidentoides**, Nutt. Paniculately branched; leaves undivided, lanceolate, coarsely toothed, taper-pointed, and the base tapering into a slender petiole; heads 6''–10'' long; *achenia* newly subulate tapering downwards, bearing a pair of very slender upwardly roughened awns surpassing the corolla (4'' long), but shorter than the achenium, often also 2 minute teeth alternate with the awns. — Shores of Delaware River, near Philadelphia, and in Delaware, *Nuttall*, *Durand*, *A. H. Smith*, *W. M. Canby*.

42. BÌDENS, L. BUR-MARIGOLD.

Heads many-flowered; the rays when present 3-8, neutral. Involucre double, the outer commonly large and foliaceous. Receptacle flattish; the chaff deciduous with the fruit. Achenia flattened parallel with the scales of the involucre, or slender and 4-sided, crowned with 2 or more rigid and persistent awns which are downwardly barbed. — Annual or perennial herbs, with opposite various leaves, and mostly yellow flowers. (Latin, *bidens*, two-toothed.)

* *Achenis flat, not tapering at the summit.* (All annuals?)

1. **B. frondosa**, L. (COMMON BEGGAR-TICKS.) Smooth or rather hairy, tall (2°-6° high) and branching; leaves 3-5-divided; the leaflets mostly stalked, lanceolate, pointed, coarsely toothed; outer leafy involucre much longer than the head, ciliate below; rays none or few and very small; achenia wedge-obovate, 2-awned, the margins ciliate with upwardly-turned bristles, except near the summit. — Moist waste places: a coarse troublesome weed, the achenia, as in the other species, adhering by their retrorsely barbed awns to the dress, and to the fleece of animals. July - Oct.

2. **B. connata**, Muhl. (SWAMP BEGGAR-TICKS.) Smooth (1°-2° high); leaves lanceolate or oblong-lanceolate, pointed, sharply serrate, tapering into margined petioles which are slightly united at the base; the lower often 3-divided, their lateral divisions united at the base and decurrent on the petiole; scales of the outer involucre longer than the head, few, mostly obtuse; rays none; achenia narrowly wedge-form, 3- (2-4-) awned, the margins minutely retrorsely ciliate. (B. tripartita, Bigel.) — A thin-leaved more petioled form is B. petiolata, Nutt. — E. New England to Illinois, and southward. — Var. COMOSA is stouter, the leaves commonly all simple, upper ones nearly sessile, the heads larger and with more numerous and leafy bracts. — Illinois, Kentucky, and westward. Aug. - Oct.

3. **B. cernua**, L. (SMALLER BUR-MARIGOLD.) Nearly smooth (5'-10' high); leaves all undivided, lanceolate, unequally serrate, scarcely connate; heads nodding, with or without (light yellow) rays: outer involucre longer than the head; achenia wedge-obovate, 4-awned, the margins downwardly barbed. — Wet places, Virginia to Wisconsin, and northward. July - Sept. — Rays, when present, smaller than in the next, the leaves irregularly toothed, and the outer involucre more leaf-like. (Eu.)

4. **B. chrysanthemoides**, Michx. (LARGER BUR-MARIGOLD.) Smooth, erect, or reclining at the base (6'-30' high); leaves lanceolate, tapering at both ends, more or less connate, regularly serrate; outer involucre mostly shorter than the showy golden-yellow (1' long) rays; achenia wedge-shaped, with almost prickly downwardly barbed margins; awns 2, 3, or 4. — Swamps: common. Aug. - Oct. — Northward it runs into No. 3.

** *Achenia linear or awl-shaped, 4-sided, slender, tapering at the summit.*

5. **B. Béckii**, Torr. (WATER MARIGOLD.) Aquatic, perhaps perennial, smooth; stems long and slender, bearing crowded immersed leaves many times dissected into fine capillary divisions; the few emerging leaves lanceolate, slightly connate, toothed; heads single, short-peduncled; involucre much shorter than the showy (golden yellow) rays; achenia linear, thickish, smooth ($\frac{1}{2}$ ' long), bearing 4-6 stout divergent awns which are an inch long and barbed only towards the

apex. — Ponds and slow deep streams, Massachusetts (rare) to Illinois and northward. Aug. — Oct.

6. **B. bipinnata**, L. (SPANISH NEEDLES.) Smooth annual, branched; leaves 1-3-pinnately parted, petioled; leaflets ovate-lanceolate, mostly wedge-shaped at the base; heads small, on slender peduncles; outer involucre of linear scales equalling the short pale yellow rays; achenia slender, 4-grooved and angled, nearly smooth, 3-4-awned. — Dry soil, Connecticut to Illinois and southward.

43. VERBESINA, L. CROWNBEARD.

Heads several-many-flowered; the rays pistillate, few, or sometimes none. Scales of the erect involucre few, imbricated in 2 or more rows. Receptacle rather convex; the chaff concave. Achenia flat (compressed laterally), winged or wingless, 2-awned. — Perennial herbs; the toothed or lobed leaves decurrent on the stem. (“Name metamorphosed from Verbena.”)

1. **V. Siegesbeckia**, Michx. Stem tall, 4-winged; leaves opposite, ovate, triple-nerved, serrate, pointed at both ends, often pubescent beneath (large and thin); heads in compound corymbs; flowers yellow; rays 1-5, lanceolate; achenia wingless. — Rich soil, S. Penn. to Illinois, and southward. July.

2. **V. Virginica**, L. Stem narrowly or interruptedly winged, downy-pubescent, like the lower surface of the ovate-lanceolate feather-veined alternate leaves; heads in compound corymbs; flowers white; rays 3-4, oval; achenia narrowly winged. — Dry soil, Pennsylvania? Illinois, and southward. Aug.

44. DYSODIA, Cav. FETID MARIGOLD.

Heads many-flowered, usually radiate; the rays pistillate. Involucre of one row of scales united into a firm cup, at the base some loose bractlets. Receptacle flat, not chaffy, but beset with short chaffy bristles. Achenia slender, 4-angled. Pappus a row of chaffy scales dissected into numerous rough bristles. — Herbs, mostly annuals or biennials, dotted with large pellucid glands, which give a strong odor (as in TAGÈTES, the FRENCH MARIGOLD of the gardens, which belongs to the same group); the heads terminating the branches: flowers yellow. (Name *δυσωδία*, an ill smell, which the plants exemplify.)

1. **D. chrysanthemoides**, Lag. Nearly smooth, diffusely branched (6'-18' high); leaves opposite, pinnately parted, the narrow lobes bristly-toothed or cut; rays few, scarcely exceeding the involucre. — Roadsides, and banks of rivers, from Illinois southward: a common weed; now migrating eastward, established at Buffalo, N. Y., *G. W. Clinton*. Aug. — Oct.

45. HYMENOPAPPUS, L'Her. HYMENOPAPPUS.

Heads many-flowered; the flowers all tubular and perfect. Scales of the involucre 6-12, loose and broad, thin, the upper part petal-like (usually white). Receptacle small, naked. Corolla with large revolute lobes. Achenia top-shaped, with a slender base, striate. Pappus of 15-20 small and blunt scales in a single row, very thin (whence the name of the genus, from *ὑμῆν*, membrane, and *πάππυς*, pappus). — Biennial or perennial herbs, with alternate mostly dissected leaves, and corymbed small heads of usually whitish flowers.

1. **H. scabiosæus**, L'Her. Somewhat flocculent-woolly when young (1° – 3° high); leaves 1–2-pinnately parted into linear or oblong lobes; scales of the involucre roundish, nearly all whitish. — Sandy barrens, Illinois and southward. May, June.

45½. **ACTINÉLLA**, Pers., Nutt. ACTINELLA.

Heads many-flowered, radiate; the rays several, wedge-oblong, 3-lobed at the apex or 3-toothed, pistillate. Scales of the hemispherical involucre ovate or lanceolate, membranaceous or coriaceous, nearly equal, appressed in 2 or 3 ranks, little shorter than the disk. Receptacle hemispherical or conical, naked. Achenia top-shaped, densely silky-villous. Pappus of 5 or more ovate or lanceolate very thin chaffy scales. — Low herbs, with narrow leaves, dotted or sprinkled with resinous atoms as in the next genus; the solitary heads terminating scapes or slender naked peduncles; flowers yellow. — Natives of the Western plains, &c., and barely entering our borders. (Name a diminutive of *Actinea*, from *ἀκτίς*, ray.)

1. **A. scapōsa**, Nutt., var. **glàbra**. Tufted; leaves crowded on the summit of woolly rootstocks, linear or somewhat spatulate, thickish, sparingly silky-hairy, becoming glabrous; scape ($3'$ – $9'$ high) and involucre more woolly, the scales ovate and obtuse; chaffy scales of the pappus ovate, awnless. — Joliet, Illinois, on an Indian mound (*Dr. Scammon, W. Boott*), and westward.

46. **HELÈNIUM**, L. SNEEZE-WEED.

Heads many-flowered, radiate; the spreading wedge-shaped rays several, 3–5-cleft at the summit, fertile. Involucre small, reflexed, the scales linear or awl-shaped. Receptacle globose or oblong, naked. Achenia top-shaped, ribbed. Pappus of 5–8 thin and 1-nerved chaffy scales, the nerve usually extended into a bristle or point. — Erect, branching herbs, with alternate leaves decurrent on the angled stem and branches, which are terminated by single or corymbed (yellow, rarely purple) heads; often sprinkled with bitter and aromatic resinous globules. (Named after *Helen*, the wife of Menelaus.)

1. **H. autumnàle**, L. (SNEEZE-WEED.) Nearly smooth, perennial; leaves lanceolate, toothed; rays longer than the globular-disk. — Alluvial river-banks: common (except in New England). Sept. — Plant 1° – 3° high, bitter: the corymbed heads showy.

47. **LEPTÓPODA**, Nutt. LEPTOPODA.

Rays neutral. Otherwise nearly as in *Helenium*. — In the true species (of which *L. puberula* and *L. brevifolia* may be found in S. Virginia) the stems are simple, and naked above, like a long peduncle, and bearing a single head (whence the name, from *λεπτός*, slender, and *πούς*, foot); but the following is leafy to the top, and branched; and were better restored to *Helenium*.

1. **L. brachýpoda**, Torr. & Gr. Stem corymbed at the summit (1° – 4° high); leaves oblong-lanceolate, decurrent on the stem; disk globular, brownish; rays ($\frac{1}{2}'$ – $\frac{2}{3}'$ long) yellow, or in one variety brownish-purple, sometimes with an imperfect style. — Damp soil, from Illinois southward. June–Aug.

48. **BALDWINIA**, Nutt. **BALDWINIA.**

Heads globular, many-flowered, radiate; the long and narrowly wedge-shaped rays neutral. Involucre short, of many thickish small scales imbricated in 3 or 4 rows, the outer obovate and obtuse. Receptacle strongly convex, with deep honeycomb-like cells containing the obconical or oblong silky-villous achenia. Pappus of 7-9 lance-oblong erect chaffy scales. — A perennial herb, smoothish, with slender simple stems (2° - 3° high), bearing alternate oblanceolate leaves, and the long naked summit terminated by a showy large head. Rays yellow ($1'$ long); the disk-flowers often turning dark purple. (Named for the late *Dr. William Baldwin.*)

1. **B. uniflora**, Nutt. — Borders of swamps, Virginia and southward. Aug.

49. **MARSHALLIA**, Schreb. **MARSHALLIA.**

Heads many-flowered; the flowers all tubular and perfect. Scales of the involucre linear-lanceolate, foliaceous, erect, in one or two rows, nearly equal. Receptacle convex or conical, with narrowly linear rigid chaff among the flowers. Lobes of the corolla slender, spreading. Achenia top-shaped, 5-angled. Pappus of 5 or 6 membranaceous and pointed chaffy scales. — Smooth and low perennials, with alternate and entire 3-nerved leaves, and solitary heads (resembling those of a Scabious) terminating the naked summit of the simple stem or branches. Flowers purplish; the anthers blue. (Named for *Humphry Marshall*, of Pennsylvania, author of *Arbustum Americanum*, one of the earliest works on the trees and shrubs of this country.)

1. **M. latifolia**, Pursh. Stems leafy; leaves ovate-lanceolate, pointed, sessile. — Dry soil, Virginia and southward. (*M. lanceolata* and *M. angustifolia* may occur in S. Virginia.)

50. **GALINSOGA**, Ruiz & Pav. **GALINSOGA.**

Heads several-flowered, radiate; the rays 4-5, small, roundish, pistillate. Involucre of 4 or 5 ovate thin scales. Receptacle conical, with narrow chaff among the flowers. Achenia angled. Pappus of small oblong cut-fringed chaffy scales (sometimes wanting). — Annual herbs, with opposite triple-nerved thin leaves, and small heads: disk-flowers yellow; rays whitish. (Named for *Galinsoga*, a Spanish botanist.)

1. **G. parviflora**, Cav. Smoothish (1° high); leaves ovate, acute, somewhat toothed; scales of the pappus 8-16. — Waste places, especially eastward; spreading from year to year. (Adv. from S. Amer.)

51. **MARÛTA**, Cass. **MAY-WEED.**

Heads many-flowered, radiate; the rays neutral. Involucre of many small somewhat imbricated scales, shorter than the disk. Receptacle conical, bearing slender chaff, at least near the summit. Achenia obovoid, ribbed, smooth. Pappus none. — Annual acrid herbs, with a strong odor, finely thrice-pinnately divided leaves, and single heads terminating the branches. Rays white, soon

reflexed; the disk yellow. (Derivation unknown.—The genus not distinct enough from the next.)

1. **M. CÔTULA**, DC. (COMMON MAY-WEED.) Scales of the involucre with whitish margins. (*Anthemis Cotula*, L.)—Roadsides: very common. (Nat. from Eu.)

52. ANTHEMIS, L. CHAMOMILE.

Heads and flowers as in *Maruta*, but the rays pistillate. Achenia terete, striate or smooth. Pappus none, or a minute crown.—Herbs with aromatic or strong odor, 1-2-pinnately divided leaves, the branches terminated by single heads. Rays in ours white; the disk yellow. (*Ἀνθεμίς*, the ancient name, given in allusion to the profusion of the flowers.)

1. **A. ARVÉNSIS**, L. (CORN CHAMOMILE.) Pubescent *annual or biennial*, resembling May-weed, but not ill-scented; chaff of the receptacle lanceolate, pointed; pappus a minute border.—Waste places: rare. (Adv. from Eu.)

2. **A. NÓBILIS**, L. (GARDEN CHAMOMILE.) More downy and *perennial*, pleasantly strong-scented; sterile shoots depressed or creeping; leaves very finely dissected; chaff of the receptacle blunt; pappus none.—Established near Lewiston, Delaware, *Nuttall*. (Adv. from Eu.)

53. ACHILLĒA, L. YARROW.

Heads many-flowered, radiate; the rays few, fertile. Involucre imbricated. Receptacle chaffy, flattish. Achenia oblong, flattened, margined. Pappus none.—Perennial herbs, with small corymbose heads. (So named because its virtues are said to have been discovered by *Achilles*.)

1. **A. Millefólium**, L. (COMMON YARROW OR MILFOIL.) Stems simple; *leaves twice-pinnately parted*; the divisions linear, 3-5-cleft, crowded; corymb compound, flat-topped; *involucre oblong*; *rays 4-5, short*, white (sometimes rose-color).—Fields and hills: common northward. Aug. (Eu.)

2. **A. PTÁRMICA**, L. (SNEEZEWORT.) *Leaves simple, lance-linear*, sharply serrate with appressed teeth; corymb loose; *rays 8-12, much longer than the involucre*; flowers white.—Danvers, Massachusetts, &c. (Adv. from Eu.)

54. LEUCÁNTHEMUM, Tourn. OX-EYE DAISY.

Heads many-flowered, radiate; the rays numerous, fertile. Scales of the broad and flat involucre imbricated, with scarious margins. Receptacle flat or convex, naked. Disk-corollas with a flattened tube. Achenia of the disk and ray similar, striate, without pappus.—Perennial herbs, with toothed, pinnatifid, or divided leaves, and single or corymbed heads. Rays white: disk yellow. (Name composed of *λευκός*, *white*, and *ἄνθεμον*, *a flower*, from the white rays.)

1. **L. VULGÁRE**, Lam. (OX-EYE OR WHITE DAISY. WHITE-WEED.) Stem erect, nearly simple, naked above and bearing a single large head; root-leaves spatulate, petioled, the others partly clasping, all cut or pinnatifid-toothed; scales of the involucre with rusty brown margins. (*Chrysánthemum Leucanthemum*, L.—Fields and meadows: too abundant. June, July. A pernicious

weed, with large and showy heads: in Connecticut is a variety with inconspicuous rays. (Nat. from Eu.)

2. **L. PARTHËNIUM**, Godron. (FEVERFEW.) Tall, branched, leafy; leaves twice-pinnately divided; the *divisions ovate, cut*; heads *corymbed*, rather small. (*Matricaria Parthenium*, L. *Pyrethrum Parthenium*, *Smith.*) — Escaped from gardens in some places. (Adv. from Eu.)

55. MATRICÀRIA, Tourn. WILD CHAMOMILE.

Heads many-flowered; the rays pistillate, or wanting. Scales of the involucre imbricated, with scarios margins. Receptacle conical, at least in fruit, naked. Achenia angular, wingless. Pappus a membranaceous crown or border, or none. — Smooth and branching herbs (ours annuals or biennials) with divided leaves and single or corymbed heads. Rays white or none: disk yellow. (Named for reputed medicinal virtues.)

1. **M. INODÒRA**, L. Leaves twice-pinnately divided into fine almost filiform lobes; heads *large*, naked-peduncled, and *with many long rays*; pappus a short crown or border. — (Wild far northward.) Roadsides, Eastport, Maine, *Prof. Verrill*. Aug. (Adv. from Eu.)

2. **M. discoidea**, DC. Low (6'–9' high); leaves 2–3-pinnately parted into short linear lobes; heads *rayless*, short-peduncled; scales of the involucre oval, with broad margins, much shorter than the conical disk; pappus obsolete. — Banks of the Mississippi opposite St. Louis. Probably an immigrant from Oregon, now extending eastward: also established in N. Europe. July–Sept.

56. TANACÈTUM, L. TANSY.

Heads many-flowered, nearly discoid, all fertile; the marginal flowers chiefly pistillate and 3–5-toothed. Scales of the involucre imbricated, dry. Receptacle convex, naked. Achenia angled or ribbed, with a large flat top. Pappus a short crown. — Bitter and acrid strong-scented herbs (ours perennial), with 1–3-pinnately dissected leaves, and corymbed heads. Flowers yellow; in summer. (Name said to be a corruption of *ἀθανασία*, *undying*, from its durable flowers.)

1. **T. VULGÀRE**, L. (COMMON TANSY.) Stem (2°–4° high) smooth; leaflets and the wings of the petiole cut-toothed; corymb dense; pistillate flowers terete; pappus 5-lobed. — Var. *CRISPUM* has the leaves more cut and crisped. — Escaped from gardens to roadsides. (Adv. from Eu.)

2. **T. Huronense**, Nutt. Hairy or woolly when young, stout (1°–3° high); lobes of the leaves oblong; heads large ($\frac{1}{2}$ '– $\frac{3}{4}$ ' wide) and usually few; pistillate flowers flattened, 3–5-cleft; pappus toothed. — St. John's River, Maine (*G. L. Goodale*), shores of the upper Great Lakes, and westward.

57. ARTEMÍSIA, L. WORMWOOD.

Heads discoid, few-many-flowered; the flowers all tubular, the marginal ones pistillate, or sometimes all similar and perfect. Scales of the involucre imbricated, with dry and scarios margins. Receptacle small and flattish, na-

ked. Achenia obovoid, with a small summit and no pappus. — Herbs or shrubby plants, bitter and aromatic, with small heads in paniced spikes or racemes; flowering in summer. Corolla yellow or purplish. (Dedicated to *Artemis*, the Greek Diana.)

§ 1. *Receptacle smooth: marginal flowers pistillate and fertile: disk-flowers perfect but sterile: root perennial, except in No. 4.*

1. **A. dracunculoides**, Pursh. Tall (3° – 5°), somewhat woody at base, slightly hoary or glabrous; leaves linear and entire or the lower 3-cleft; heads small and numerous, paniced. — Sandy banks of streams, S. W. Illinois (*Dr. Vasey, Dr. Mead*) and westward.

2. **A. borealis**, Pallas. Low (3'–6' high), tufted, silky-villous or nearly smooth; lower leaves 3–5-cleft at the apex, or like the others 1–2-pinnately parted, the lobes lanceolate or linear; heads few, hemispherical, pretty large, spiked or racemed. — Shore of Lake Superior, and northward. (Eu.)

3. **A. Canadensis**, Michx. Smooth, or hoary with silky down (1° – 2° high); lower leaves twice-pinnately divided, the upper 3–7-divided; the divisions linear, rather rigid; heads rather large, in paniced racemes. — Shore of all the Great Lakes, &c., and northward. (Eu.)

4. **A. caudata**, Michx. Smooth (2° – 5° high); upper leaves pinnately, the lower 2–3-pinnately divided; the divisions thread-form, diverging; heads small, the racemes in a wand-like elongated panicle; root biennial. — Sandy soil, coast of New Hampshire to Virginia; also Michigan and Illinois.

§ 2. *Receptacle smooth: flowers all fertile, a few pistillate, the others perfect.*

* Tall (1° – 5°) and branching perennials, whitened with fine and close-pressed wool: heads small, ovoid, crowded in leafy panicles.

5. **A. Ludoviciana**, Nutt. (WESTERN MUGWORT.) Whitened woolly throughout; leaves lanceolate, the upper mostly entire, the lower cut-lobed, toothed or pinnatifid; heads larger than in the next, mostly sessile in narrow panicles. — Dry banks, Lakes Huron and Michigan, and southwestward; especially the var. *GNAPHALODES*, which has the elongated nearly entire leaves very woolly both sides.

6. **A. vulgaris**, L. (COMMON MUGWORT.) Leaves mostly glabrous and green above, beneath and the branches white-woolly, all pinnatifid, with the divisions often cut-lobed, linear-lanceolate; heads small in open panicles. — Waste places, near dwellings. (Adv. from Eu.)

** Less branched (1° – 3°) biennial or annual, glabrous: heads densely clustered.

7. **A. biennis**, Willd. (BIENNIAL WORMWOOD.) Lower leaves twice-pinnately parted, the upper pinnatifid; lobes linear, acute, in the lower leaves cut-toothed; heads in short axillary spikes or clusters, crowded in a narrow and glomerate leafy panicle. — Gravelly banks, Ohio to Illinois, and northwestward; rapidly extending eastward by railroad to Buffalo, Philadelphia, &c.

§ 3. *Receptacle hairy; flowers all fertile, the marginal ones pistillate: heads nodding.*

8. **A. absinthium**, L. (COMMON WORMWOOD.) Rather shrubby (2° – 4° high), silky-hoary; leaves 2–3-pinnately parted, lobes lanceolate; heads hemispherical, paniced. — Roadsides, sparingly escaped from gardens. (Adv. from Eu.)

9. **A. frigida**, Willd. Low (6'–20' high), in tufts, slightly woody at the base, white-silky; leaves pinnately parted and 3–5-cleft, the divisions narrow-linear; heads globose, racemose. — Dry hills and rocks, Falls of St. Anthony, Wisconsin (*L. Lesqueureux, T. J. Hale*), Lake Superior, and northwestward.

58. GNAPHALIUM, L. CUDWEED.

Heads many-flowered; the flowers all tubular; the outer pistillate and very slender, the central perfect. Scales of the involucre dry and scarious, white or colored, imbricated in several rows. Receptacle flat, naked. Pappus a single row of capillary rough bristles. — Woolly herbs, with sessile or decurrent leaves, and clustered or corymbed heads; fl. in summer and autumn. Corolla whitish or yellowish. (Name from *γάφαλον*, a *lock of wool*, in allusion to the floccose down of the leaves.)

§ 1. *Achenia terete: pistillate flowers in several rows: bristles of pappus distinct.*

1. **G. decurrens**, Ives. (EVERLASTING.) Stout, erect (2° high) perennial, branched at the top, clammy-pubescent, white-woolly on the branches, bearing numerous heads in dense corymbed clusters; leaves linear-lanceolate, partly clasping, decurrent; scales of the (yellowish-white) involucre oval, acutish. — Hillsides, New Jersey and Penn. to Maine, Michigan, and northward.

2. **G. polycéphalum**, Michx. (COMMON EVERLASTING.) Erect, woolly annual (1'–2' high), fragrant; leaves lanceolate, tapering at the base, with undulate margins, not decurrent, smoothish above; heads clustered at the summit of the paniced-corymbose branches, ovate-conical before expansion, then obovate; scales of the (whitish) involucre ovate and oblong, rather obtuse; perfect flowers few. — Old fields and woods: common.

3. **G. uliginosum**, L. (LOW CUDWEED.) Diffusely branched, woolly annual (3'–6' high); leaves lanceolate or linear, not decurrent; heads (small) in terminal sessile capitata clusters subtended by leaves. — Low grounds by the roadside; common eastward and northward: perhaps introduced. (Eu.)

§ 2. *GAMOCHÆTA*, Weddell. *Achenia and flowers as § 1: bristles of the pappus united at the very base into a ring, so falling off all together.*

4. **G. purpureum**, L. (PURPLISH CUDWEED.) Annual, simple or branched from the base, ascending (6'–20' high), woolly; leaves oblong-spatulate, mostly obtuse, not decurrent, green above, very white with close wool underneath; heads in sessile clusters in the axils of the upper leaves, and spiked at the wand-like summit of the stem; scales of the involucre tawny, the inner often marked with purple. — Sandy or gravelly soil, coast of Maine to Virginia, and southward.

§ 3. *HOMALOTHÈCA*, DC. *Achenia flattened: pistillate flowers in a single marginal row: bristles of the pappus distinct and falling separately, as in § 1.*

5. **G. supinum**, Villars. (MOUNTAIN CUDWEED.) Dwarf and tufted perennial (2' high); leaves linear, woolly; heads solitary or few and spiked on the slender simple flowering stems; scales of the involucre brown, lanceolate, acute. — Alpine summit of Mount Washington, New Hampshire: very rare. (Eu.)

59. ANTENNÀRIA, Gærtn. EVERLASTING.

Heads many-flowered, diœcious or nearly so; the flowers all tubular: pistillate corollas very slender. Scales of the involucre dry and scarious, white or colored, imbricated. Receptacle convex or flat, not chaffy. Pappus a single row of bristles, in the fertile flowers capillary, and in the sterile thickened and club-shaped or barbellate at the summit. — Perennial white-woolly herbs, with entire leaves and corymbed (rarely single) heads. Corolla yellowish. (Name from the resemblance of the sterile pappus to the *antennæ* of certain insects.)

1. **A. margaritacea**, R. Brown. (PEARLY EVERLASTING. *Stem erect* (1° – 2° high), *corymbose* at the summit, with many heads, *leafy*; leaves linear-lanceolate, taper-pointed, sessile; fertile heads often with a few imperfect staminate flowers in the centre; scales of the pearly-white involucre obtuse or rounded. — Dry hills and woods; common northward. Aug.

2. **A. plantaginifolia**, Hook. (PLANTAIN-LEAVED EVERLASTING.) *Spreading by offsets and runners*, low (4'–10' high); leaves silky-woolly when young, at length green above and hoary beneath; those of the *simple and scape-like flowering stems* small, lanceolate, appressed; the radical obovate or oval-spatulate, petioled, ample, 3-nerved; heads in a small crowded corymb; scales of the (mostly white) involucre obtuse in the sterile, and acutish and narrower in the fertile plant. — Var. **MONOCÉPHALA** is an occasional state, with a single larger head. — Sterile knolls and banks: common. March–May.

60. FILÀGO, Tourn. COTTON-ROSE.

Heads many-flowered; the flowers all tubular, the central ones perfect, but often infertile; the others pistillate, very slender and thread-form. Scales of the involucre few and woolly. Receptacle elongated or top-shaped, naked at the summit, but chaffy at the margins or toward the base; the chaff resembling the proper involucral scales, each covering a single pistillate flower. — Pappus of the central flowers capillary, of the outer ones mostly none. — Annual, low, branching woolly herbs, with entire leaves, and small heads in capitate clusters. (Name from *filum*, a thread, in allusion to the cottony hairs of these plants.)

1. **F. GERMÁNICA**, L. (HERBA IMPIA.) *Stem erect*, short, clothed with lanceolate and upright crowded leaves, producing a capitate cluster of woolly heads, from which rise one or more branches, each terminated by a similar head, and so on: — hence the common name applied to it by the old botanists, as if the offspring were undutifully exalting themselves above the parent. — Dry fields, New York to Virginia. July–Oct. (Nat. from Eu.)

61. ERECHTHITES, Raf. FIREWEED.

Heads many-flowered; the flowers all tubular and fertile; the marginal pistillate, with a slender corolla. Scales of the cylindrical involucre in a single row, linear, acute, with a few small bractlets at the base. Receptacle naked. Achenia oblong, tapering at the end. Pappus copious, of very fine and white soft hairs. — Erect and coarse annuals, of rank smell, with alternate simple leaves, and paniculate-corymbed heads of whitish flowers. (The ancient name of some species of Groundsel, probably called after *Erechtheus*.)

1. **E. hieracifolia**, Raf. (FIREWEED.) Often hairy; stem grooved (1° – 5° high); leaves lanceolate or oblong, acute, cut-toothed, sessile; the upper with an auricled clasping base. (*Senecio hieracifolius*, L.) — Moist woods: common, especially northward, and in recent clearings, where the ground has been burned over; whence the popular name. July–Sept.

62. CACÀLIA, L. INDIAN PLANTAIN.

Heads 5–many-flowered; the flowers all tubular and perfect. Scales of the involucre in a single row, with a few bractlets at the base. Receptacle naked. Corolla deeply 5-cleft. Achenia oblong, smooth. Pappus of numerous capillary bristles. — Smooth and tall perennial herbs, with alternate often petioled leaves, and rather large heads, in flat corymbs. Flowers white or whitish. (An ancient name, of uncertain meaning.)

* *Involucre 25–30-flowered, with several bracts at its base: receptacle flat.*

1. **C. suavèolens**, L. Stem grooved (3° – 5° high); leaves triangular-lanceolate, halberd-shaped, pointed, serrate, those of the stem on winged petioles. — Rich woods, Connecticut to Wisconsin and Kentucky: rare. Sept.

* * *Involucre 5-leaved and 5-flowered, its bracts minute or none: receptacle bearing a more or less evident scale-like pointed appendage in the centre.*

2. **C. renifórmis**, Muhl. (GREAT INDIAN PLANTAIN.) Stem (4° – 9° high) grooved and angled; leaves green both sides, dilated fan-shaped, or the lowest kidney-form (1° – 2° broad), repand-toothed and angled, palmately veined, petioled; the teeth pointed; corymbs large. — Rich damp woods, New Jersey to Illinois, and southward along the mountains. Aug.

3. **C. atriplicifolia**, L. (PALE INDIAN P.) Stem terete (3° – 6° high), and with the palmately veined and angulate-lobed leaves glaucous; lower leaves triangular-kidney-form or slightly heart-shaped; the upper rhomboid or wedge-form, toothed. — Rich woodlands, W. New York to Wisconsin, and southward. Aug.

4. **C. tuberósa**, Nutt. (TUBEROUS INDIAN P.) Stem angled and grooved (2° – 6° high), from a thick or tuberous root; leaves green both sides, thick, strongly 5–7-nerved; the lower lance-ovate or oval, nearly entire, tapering into long petioles; the upper on short margined petioles, sometimes toothed at the apex. — Wet prairies, &c., Ohio to Wisconsin, and southward. June.

63. SENÈCIO, L. GROUNDSEL.

Heads many-flowered; the flowers all perfect and tubular, or mostly with radiate marginal ones; the rays pistillate. Scales of the involucre in a single row, or with a few bractlets at the base. Receptacle flat, naked. Pappus of numerous very soft and slender capillary bristles. — Herbs, in the United States, with alternate leaves and solitary or corymbed heads. Flowers chiefly yellow. (Name from *senex*, an old man, alluding to the hoary hairs which cover many species, or to the white hairs of the pappus.)

* *Root annual or in No. 3 biennial: heads several or many in a corymb: herbage labrous or soon becoming so.*

1. **S. vulgāris**, L. (COMMON GROUNDSEL.) Low, corymbosely branched; leaves pinnatifid and toothed, clasping; rays none. — Waste grounds. July–Sept. (Adv. from Eu.)

2. **S. lobatus**, Pers. (BUTTER-WEED.) Rather tall; leaves somewhat fleshy, mostly *lyrate* or *pinnate*, the divisions or leaflets crenate or cut-lobed, variable; heads small in a naked corymb; rays about 12, conspicuous. — Low banks of the Ohio and Mississippi, Illinois and southward. April–July.

3. **S. palustris**, Hook. Biennial, loosely woolly when young; stem stout, 6'–3° high; leaves oblong-lanceolate, *irregularly toothed* or *luciniate*, the upper with a heart-shaped clasping base; rays 20 or more, short, pale yellow; pappus copious and becoming very long. — Wet ground, N. W. Wisconsin (*T. J. Hule*) and northward. June. (Eu.)

* * Root perennial: heads small or middle-sized, in a naked corymb.

4. **S. aureus**, L. (GOLDEN RAGWORT. SQUAW-WEED.) Smooth, or *floccose-woolly* when young (10'–30' high); root-leaves simple and rounded, the larger ones mostly heart-shaped, crenate-toothed, long-petioled; the lower stem-leaves *lyrate*; upper ones lanceolate, cut-pinnatifid, sessile or partly clasping; corymb umbel-like; rays 8–12. — Varies greatly, the leading forms being, — Var. 1. **OBOVATUS**, with the root-leaves round-obovate (growing in drier places). — Var. 2. **BALSAMITÆ**, with root-leaves oblong, spatulate, or lanceolate, sometimes cut-toothed, tapering into the petiole. Rocky places. — Var. 3. **LANCEOLATUS**, *Oakes*, with the leaves all lanceolate-oblong, thin, sharply and unequally toothed, either wedge-shaped or somewhat heart-shaped at the base, the upper merely pinnatifid-cut towards the base. (Cedar swamps, Vermont, *Robbins*.) — Common everywhere: the primary form in swamps. May, June.

5. **S. Elliottii**, Torr. & Gr. Soon smooth, stem simple (1° high), often nearly leafless, bearing a small corymb; root-leaves *thickish*, obovate or roundish, narrowed into a short and winged petiole, or sessile, crenate-toothed, sometimes *lyrate*; stem-leaves small, cut-pinnatifid. — Rich soil, Virginia and southward along the mountains. May.

6. **S. tomentosus**, Michx. (WOOLLY RAGWORT.) Clothed with scarcely deciduous hoary wool (1°–2° high); root-leaves oblong, obtuse, crenate toothed, on slender petioles; the upper sessile; corymb flat-topped; rays 12–15. — Mountains of Pennsylvania (*Pursh*), Maryland, and southward. May. — **S. canus**, Hook., which too closely resembles smaller forms of this, probably occurs within the northern borders of Wisconsin.

* * * Root perennial: heads large and mostly solitary.

7. **S. Pseudo-Árnica**, Less. Loosely white-woolly, sometimes becoming glabrous; stem stout, 6'–12' high, leafy to the top; leaves oblong, repand, tapering into a narrow petiole-like base; heads 1–4, over an inch in diameter; rays 20 or more, large. — Grand Manan Island off Maine (*Prof. Verrill*) and northward.

64. ÁRNICA, L. ARNICA.

Heads many-flowered, radiate; the rays pistillate. Scales of the bell-shaped involucre lanceolate, equal, somewhat in 2 rows. Receptacle flat, fimbriate.

Achenia slender or spindle-shaped. Pappus a single row of rather rigid and strongly roughened-denticulate bristles. — Perennial herbs, chiefly of mountains and cold northern regions, with simple stems, bearing single or corymbed large heads and opposite leaves. Flowers yellow. (Name thought to be a corruption of *Ptarmica*.)

1. **A. móllis**, Hook. Soft-hairy; *stem leafy* (1° - 2° high), bearing 1 to 5 heads; *leaves thin, veiny*, smoothish when old, toothed; the upper *ovate-lanceolate*, closely sessile; the lower narrower, tapering into a margined petiole; scales of the involucre pointed; pappus almost plumose. — Alpine rivulets, mountains of New Hampshire and N. New York, shores of Lake Superior, and northward. July.

2. **A. nudicaulis**, Ell. Hairy and rather glandular (1° - 3° high); *leaves thickish*, 3 - 5-nerved, *ovate or oblong*, all sessile, mostly entire and near the root, those of the *naked stem* small and only one or two pairs; heads several, corymbed, showy. — Damp pine barrens, S. Penn. and southward. April, May.

65. CENTAURĒA, L. STAR-THISTLE.

Heads many-flowered; the flowers all tubular, the marginal often much larger and as it were radiate, sterile. Receptacle bristly. Involucre imbricated, the scales margined or appendaged. Achenia compressed. Pappus wanting, or of a few bristles. — Herbs with alternate leaves and single heads. (Named from the *Centaur*, *Chiron*.)

1. **C. CYANUS**, L. (BLUEBOTTLE.) Scales of the globular involucre fringed-margined; *false rays large*; pappus very short; *leaves linear, entire*, or toothed at the base; root annual. — Roadsides, escaped from gardens. July. — Flowers blue, varying to purplish or white. (Adv. from Eu.)

2. **C. NIGRA**, L. (KNAPWEED.) Scales of the globular involucre appendaged, and with a stiff black fringe; *rays wanting*; pappus very short; *leaves lanceolate*, or the lower lyrate-angled, rough; root perennial. — Waste places, E. New England. Aug. — Flowers purple. (Adv. from Eu.)

3. **C. CALCITRAPA**, L. (STAR THISTLE.) Stem diffusely much branched; *leaves pinnately lobed* or spinulose-toothed; heads sessile, the middle *scales of the ovoid involucre spiny*; pappus none; flowers purple; root annual. — Norfolk, Virginia, and Philadelphia. (Adv. from Eu.)

66. CNĪCUS, Vail. BLESSED THISTLE.

Heads many-flowered; the ray-flowers tubular and sterile, shorter than the rest, which are all tubular and perfect. Scales of the ovoid involucre coriaceous, appressed, extended into a long and rigid pinnately spinose appendage. Receptacle clothed with capillary bristles. Achenia terete, short, strongly striate, crowned with 10 short and horny teeth, and bearing a pappus of 10 elongated rigid bristles, and 10 short bristles alternate with the last in an inner row. — An annual smoothish herb, with clasping scarcely pinnatifid-cut leaves and large bracted heads. Flowers yellow. (Name from *κνίζω*, *to prick*.)

1. **C. BENEDICTUS**, L. — Roadsides, southward: rare, scarcely naturalized. (Adv. from Eu.)

67. **CÍRSIUM**, Tourn. COMMON OR PLUMED THISTLE.

Heads many-flowered; the flowers all tubular, perfect and similar, or rarely imperfectly dioecious. Scales of the ovoid or spherical involucre imbricated in many rows, tipped with a point or prickle. Receptacle thickly clothed with soft bristles or hairs. Achenia oblong, flattish, not ribbed. Pappus of numerous bristles united into a ring at the base, plumose to the middle, deciduous. — Herbs, with sessile alternate leaves, often pinnatifid, and prickly. Heads large, terminal. Flowers reddish-purple, occasionally yellowish, white, or cream-color; in summer. (Name from *κίρσος*, a *swelled vein*, for which the Thistle was a reputed remedy.)

* *Scales of the involucre all tipped with spreading prickles: root biennial.*

1. **C. lanceolatum**, Scop. (COMMON THISTLE.) Leaves decurrent on the stem, forming prickly lobed wings, pinnatifid, rough and bristly above, woolly with deciduous webby hairs beneath, prickly; flowers purple. — Pastures and roadsides, everywhere, at the North. (Nat. from Eu.)

** *Scales of the involucre appressed; the inner ones not prickly: filaments hairy.*

+ *Leaves white-woolly beneath, and sometimes also above: outer scales of the involucre successively shorter, and tipped with short prickles.*

2. **C. Pitcheri**, Torr. & Gr. *White-woolly throughout*, perennial, low; stem stout, very leafy; leaves all pinnately parted into rigid narrowly linear and elongated divisions, with revolute margins; flowers cream-color. — Sandy shores of Lakes Michigan, Huron, and Superior.

3. **C. undulatum**, Spreng. *White-woolly throughout*, biennial, low and stout, leafy; leaves lanceolate-oblong, partly clasping, undulate-pinnatifid, with prickly lobes; flowers reddish-purple. — Islands of L. Huron and Michigan; thence westward.

4. **C. discolor**, Spreng. Biennial; stem grooved, hairy, branched, tall, leafy; leaves all deeply pinnatifid, sparingly hairy and green above, whitened with close wool beneath; the diverging lobes 2-3-cleft, linear-lanceolate, prickly-pointed; flowers pale purple, rarely white. — Meadows and copses.

5. **C. altissimum**, Spreng. Stem downy, branching (3°-10° high), leafy quite to the heads: leaves roughish-hairy above, whitened with close wool beneath, oblong-lanceolate sinuate-toothed, undulate-pinnatifid, or undivided, the lobes or teeth prickly; those from the base pinnatifid; and their lobes short, oblong or triangular; flowers chiefly purple. — Fields and copses, Penn. to Illinois and southward.

6. **C. Virginianum**, Michx. Stem woolly, slender, simple or sparingly branched (1°-3° high); the branches or long peduncles naked: leaves lanceolate, green above, whitened with close wool beneath, ciliate with prickly bristles, entire or sparingly sinuate-lobed, sometimes the lower deeply sinuate-pinnatifid; outer scales of the involucre scarcely prickly; heads small; flowers purple. — Woods and plains, Virginia, Ohio, and southward.

Var. **filipendulum**. Stem stouter, more leafy, corymbosely branched above; the heads on shorter peduncles; leaves pinnatifid; roots tuberous, enlarged below. (*C. filipendulum*, Engelm.) — Prairies of S. Illinois and south-westward.

+ + *Leaves green both sides, or only with loose cobwebby hairs underneath: scales of the involucre scarcely prickly-pointed: heads large.*

7. **C. mūticum**, Michx. (SWAMP THISTLE.) *Stem tall (3°–8° high), angled, smoothish, paniced at the summit, the branches sparingly leafy and bearing single or few rather large naked heads; leaves somewhat hairy above, whitened with loose webby hairs beneath when young, deeply pinnatifid, the divisions lanceolate, acute, cut-lobed, prickly-pointed; scales of the webby and glutinous involucre closely appressed, pointless or barely mucronate; flowers purple; root perennial.* — Swamps and low woods: common.

8. **C. pūmulum**, Spreng. (PASTURE THISTLE.) *Stem low and stout (1°–3° high), hairy, bearing 1–3 very large heads (1½' broad), which are somewhat leafy-bracted at the base; leaves green, lanceolate-oblong, partly clasping, somewhat hairy, pinnatifid, with short and cut very prickly-margined lobes; outer scales of the involucre prickly-pointed, the inner very slender; flowers purple or rarely white (fragrant, 2' long); root biennial.* — Dry fields, Maine to Penn., near the coast, Illinois and westward: common.

9. **C. horridulum**, Michx. (YELLOW THISTLE.) *Stem stout (1°–4° high), webby-haired when young; leaves partly clasping, green, soon smooth, lanceolate, pinnatifid, the short toothed and cut lobes very spiny with yellowish prickles; heads (1'–1½' broad), surrounded at the base by an involucre whorl of leaf-like and very prickly bracts, which equal or exceed the narrow and unarmed scales of the involucre; flowers pale yellow or purple.* — Sandy fields, Massachusetts to Virginia, and southward, near the coast.

* * * *Outer scales of the appressed involucre barely prickly-pointed: filaments nearly smooth: heads imperfectly diacious, small and numerous.*

10. **C. arvénse**, Scop. (CANADA THISTLE.) *Perennial, the roots extensively creeping; leaves oblong or lanceolate, smooth, or slightly woolly beneath, sinuate-pinnatifid, prickly-margined; flowers rose-purple.* — Cultivated fields, pastures, and roadsides: common at the North; a most troublesome weed, extremely difficult to eradicate. (Nat. from Eu.)

68. CARDUUS, Tourn. PLUMELESS THISTLE.

Bristles of the pappus naked (not plumose), merely rough or denticulate. Otherwise as in *Cirsium*. (The ancient Latin name.)

1. **C. nūtans**, L. (MUSK THISTLE.) *Biennial; leaves decurrent, sinuate, spiny; heads solitary, drooping; flowers purple.* — Fields near Harrisburg, Pennsylvania, *Prof. Porter*. (Adv. from Eu.)

69. ONOPÓRDON, Vail. COTTON OR SCOTCH THISTLE.

Heads and flowers nearly as in *Cirsium*. Scales of the involucre coriaceous, tipped with a lanceolate prickly appendage. Receptacle deeply honeycombed. Achenia 4-angled, wrinkled transversely. Bristles of the pappus numerous, slender, not plumose, united at the base into a horny ring. — Coarse, branching annuals, or biennials, with the stems winged by the decurrent base of the lobed and toothed somewhat prickly leaves. Heads large: flowers purple. (The ancient Greek name of the plant.)

1. *O. ACANTHIUM*, L. Stem (2°-4° high) and leaves cotton-woolly; scales linear-awl-shaped. — Roadsides and waste places: rather rare. July-Sept. (Adv. from Eu.)

70. LÁPPA, Tourn. BURDOCK.

Heads many-flowered; the flowers all perfect and similar. Involucre globular; the imbricated scales coriaceous and appressed at the base, tipped with an abrupt and spreading awl-shaped hook-pointed appendage. Receptacle bristly. Achenia oblong, flattened, wrinkled transversely. Pappus short, of numerous rough bristles, not united at the base, deciduous. — Coarse biennial weed, with large unarmed and petioled leaves, and small solitary or clustered heads: flowers purple, rarely white. (Name from *λαβεῖν*, to lay hold, the involucre forming a hooked bur which holds tenaciously to the dress, or the fleece of animals.)

1. *L. OFFICINALIS*, Allioni. — Waste places and around dwellings, in manured soil. — The var. MAJOR (COMMON BURDOCK) has the involucre smoothish; leaves loosely whitish-cottony beneath or becoming smooth, the upper ovate, lower heart-shaped. — Var. TOMENTOSA has the involucre cottony, and is rare. — Var. MINOR has smaller heads with smooth involucre, and smaller smoothish leaves, often tapering at the base; occasionally cut-toothed or laciniate-lobed. (Uxbridge, Mass., *Dr. Robbins.*) July-Oct. (Nat. from Eu.)

SUBORDER II. LIGULIFLORÆ. (CICHORACEÆ.)

71. LÁMPSANA, Tourn. NIPPLE-WORT.

Heads 8-12-flowered. Scales of the cylindrical involucre 8, erect, in one row. Receptacle naked. Achenia oblong. Pappus none. — Slender branching annuals, with angled or toothed leaves, and loosely paniced small heads: flowers yellow. (The *Δαμψάνη* of Dioscorides was evidently a wild Mustard.)

1. *L. COMMUNIS*, L. Nearly smooth; lower leaves ovate, sometimes lyre-shaped. — Roadsides, near Boston, Buffalo, &c. (Adv. from Eu.)

72. CICHORIUM, Tourn. SUCCORY or CICHORY.

Heads several-flowered. Involucre double; the outer of 5 short spreading scales, the inner of 8-10 scales. Achenia striate. Pappus of numerous small chaffy scales, forming a short crown. — Branching perennials, with deep roots; the sessile heads 2 or 3 together, axillary and terminal. Flowers bright blue or varying to purple, showy. (Altered from the Arabian name of the plant.)

1. *C. INTYBUS*, L. Stem-leaves oblong or lanceolate, partly clasping, the lowest runcinate, those of the rigid flowering branches minute. — Roadsides: common near the coast, especially in E. Mass. July-Oct. (Nat. from Eu.)

73. KRÍGIA, Schreber. DWARF DANDELION.

Heads 15-20-flowered. Scales of the involucre several, in about 2 rows. Achenia top-shaped, many-striate or angled. Pappus double; the outer of 5

broad and rounded chaffy scales; the inner of as many alternate slender bristles. — Small annuals or biennials, branched from the base; the leaves chiefly radical, lyrate or toothed; the small heads terminating the naked scapes or branches. Flowers yellow. (Named after *D. Krieg*, an early German botanical collector in this country.)

1. **K. Virgínica**, Willd. Stems or scapes several (1'–10' high); earlier leaves roundish and entire, the others narrower and often pinnatifid. — Var. *DICHÓTOMA* is a branched and leafy summer state. — New England to Illinois and southward. April–Aug.

74. CÝNTHIA, Don. CYNTHIA.

Heads many-flowered. Scales of the involucre several, somewhat in 2 rows. Achenia short, striate. Pappus double; the outer of numerous very small chaffy bristles; the inner of numerous capillary elongated bristles. — Low perennial herbs, nearly smooth and glaucous, with scattered or radical leaves; the scapes or naked peduncles (often bristly at the apex) bearing rather showy single heads. Flowers yellow. (Perhaps named after *Mount Cynthus*.)

1. **C. Virgínica**, Don. *Roots fibrous; stem-leaves* 1–2, oblong or lanceolate-spatulate, clasping, mostly entire; the radical ones on short winged petioles, often toothed, rarely pinnatifid; peduncles 2–5. — Moist banks, New York to Michigan and southward. June. — Stem 1° high, or more.

2. **C. Dándelion**, DC. *Scapes leafless*, from a *tuberous root* (6'–15' high); leaves varying from spatulate-oblong to linear-lanceolate, entire or few-lobed. — Moist ground, Maryland to Kentucky and southward. March–July.

75. LEÓNTODON, L., Juss. HAWKBIT. FALL DANDELION.

Heads many-flowered. Involucre scarcely imbricated, but with several bractlets at the base. Achenia spindle-shaped, striate, all alike. Pappus persistent, composed of plumose bristles which are enlarged and flattened towards the base. — Low and stemless perennials, with toothed or pinnatifid root-leaves, and scapes bearing one or more yellow heads. (Name from *λέων*, a lion, and *ὀδούς*, a tooth, in allusion to the toothed leaves.) — The following belongs to the subgenus *OPORÍNIA*, with a tawny pappus of a single row of equal bristles.

1. **L. AUTUMNÁLE**, L. (FALL DANDELION.) Leaves lacinate-toothed or pinnatifid; scape branched; peduncles thickened at the summit and furnished with small scaly bracts. — Meadows and roadsides: common in E. New England. July–Oct. (Nat. from Eu.)

76. TRÓXIMON, Nutt. TROXIMON.

Head many-flowered. Scales of the bell-shaped involucre ovate or lanceolate, pointed, loosely imbricated in 2 or 3 rows. Achenia smooth, 10-ribbed, not beaked. Pappus longer than the achenium, white, of copious and unequal rigid capillary bristles, some of the larger gradually thickened towards the base. — Perennial herbs, with elongated linear tufted root-leaves, and a simple naked scape. Heads solitary, large: flowers yellow. (Name from *τρώξομαι*, to eat, first applied to some plant with an edible root, like Salsify.)

1. **T. cuspidatum**, Pursh. Leaves lanceolate, elongated, tapering to a sharp point, woolly on the margins; scales of the involucre lanceolate, sharp-pointed. — Prairies, Wisconsin, N. Illinois, and westward. April, May.

77. HIERACIUM, Tourn. HAWKWEED.

Heads 12-many-flowered. Involucre more or less imbricated. Achenia short, oblong or columnar, striate, not beaked. Pappus a single row of tawny and fragile capillary rough bristles. — Perennial herbs, with entire or toothed leaves, and single or paniced heads of mostly yellow flowers; in summer and early autumn. (Name from *ίέραξ*, a hawk.)

* *Heads large and broad: involucre imbricated: achenia tapering towards the base.*

1. **H. Canadense**, Michx. (CANADA HAWKWEED.) Stems simple, leafy, corymbed at the summit (1°–3° high); leaves sessile, lanceolate or ovate-oblong, acute, remotely and very coarsely toothed, somewhat hairy, the uppermost slightly clasping. — Dry woods, northward.

** *Heads small: involucre cylindrical, scarcely imbricated.*

2. **H. scabrum**, Michx. (ROUGH H.) Stem rather stout (1°–3° high), leafy, *rough-hairy*; the stiff panicle at first racemose, at length rather corymbose; the thickish peduncles and the hoary 40–50-flowered involucre densely clothed with dark glandular bristles; *achenia columnar, not tapering at the summit*; leaves obovate or oval, nearly entire, hairy. — Dry open woods; common.

3. **H. longipilum**, Torr. (LONG-BEARDED H.) Stem wand-like, simple, stout (2°–3° high), *very leafy towards the base, naked above*, and bearing a small racemed panicle; the lower portion and both sides of the oblong-lanceolate or spatulate entire leaves thickly clothed with *very long and upright bristles* (often 1' long); peduncles and 20–30-flowered involucre glandular-bristly; *achenia spindle-shaped, narrowed at the apex*. — Prairies, Michigan to Wisconsin and southwestward. — Heads intermediate between the last and the next.

4. **H. Gronovii**, L. (HAIRY H.) Stem wand-like, mostly simple, *leafy and very hairy below, naked above* and forming a long and narrow panicle; leaves oblong or obovate, nearly entire, hairy; the slender peduncles and the 20–30-flowered involucre sparingly glandular-bristly; *achenia spindle-shaped, with a very taper summit*. — Dry sterile soil: common, especially southward. — Varies from 1°–4° high. The small heads and almost beaked fruit distinguish the largest forms from No. 2, and small naked-stemmed states from the next.

5. **H. venosum**, L. (RATTLESNAKE-WEED.) Stem or scape (1°–2° high) *naked or with a single leaf, smooth and slender, forking above into a spreading loose corymb*; root-leaves obovate or oblong, nearly entire, scarcely petioled, thin and pale, purplish and glaucous underneath (often hairy along the midrib), marked above with purple veins; peduncles very slender; involucre 20-flowered; *achenia linear, not tapering upwards*. — Var. *SUBCAULÉSCENS* has the stem more or less leafy next the base. — Dry plains and pine woods: common.

6. **H. paniculatum**, L. (PANICLED H.) Stem slender, *leafy, diffusely branched*, hairy below (2°–3° high); leaves lanceolate, acute at both ends, slightly toothed, smooth; *heads in a loose panicle* (very small), on slender and

diverging peduncles, 12–20-flowered; *achenia short*, not tapering at the summit. — Open woods: rather common.

78. NÁBALUS, Cass. RATTLESNAKE-ROOT.

Heads few-many-flowered. Involucre cylindrical, of 5 to 14 linear scales in a single row, and a few small bractlets at the base. Achenia short, linear-oblong, striate or grooved, not contracted at the apex. Pappus of copious straw-color or brownish and rough capillary bristles.—Perennial herbs, with upright leafy stems arising from spindle-shaped (extremely bitter) tubers, very variable leaves, and racemose-panicled mostly nodding heads. Flowers greenish-white or yellowish, often tinged with purple; in late summer and autumn. (Name probably from *νάβλα*, a *harp*, in allusion to the lyrate leaves which these plants sometimes present.) Species of *Prenánthes*, *L.*

* *Involucre smooth or nearly so, 5–20-flowered.*

1. **N. álbus**, Hook. (WHITE LETUCE. RATTLESNAKE-ROOT.) Smooth and glaucous (2°–4° high); stem *corymbose-panicled at the summit*: leaves angulate or triangular-halberd-form, sinuate-toothed, or 3–5-cleft; the uppermost oblong and undivided; *involucre* (purplish) *of about 8 scales, 8–12-flowered*; *pappus deep cinnamon-color*. — Var. *SERPENTARIA* is a form with deeply divided leaves, their margins often rough-ciliate. — Borders of woods, in rich soil: common, especially northward. — Stouter and more corymbed than the next, with thickish leaves and often purplish branches. Heads 6'' long.

2. **N. altissimus**, Hook. (TALL W.) Smooth; stem tall and slender (3°–6° high); the heads in small axillary and terminal loose clusters forming *a long and wand-like leafy panicle*: leaves membranaceous, all petioled, ovate, heart-shaped, or triangular, and merely toothed or cleft, with naked or winged petioles, or frequently 3–5-parted, with the divisions entire or again cleft; *involucre slender* (greenish), *of 5 scales, 5–6-flowered*; *pappus dirty white*, or pale straw-color. — Rich moist woods: common northward.

3. **N. Fràseri**, DC. (LION'S-FOOT. GALL-OF-THE-EARTH.) Nearly smooth; stem *corymbose-panicled at the summit* (1°–4° high); leaves mostly deltoid, roughish; the lower variously 3–7-lobed, on margined petioles; the upper oblong-lanceolate, mostly undivided, nearly sessile; *involucre* (greenish or purplish, sometimes slightly bristly) *of about 8 scales, 8–12-flowered*; *pappus dull straw-color*. — Varies greatly in foliage: the var. *INTEGRIFOLIUS* has the thickish leaves all undivided and merely toothed. — Dry sandy or sterile soil, S. New England to Virginia and southward.

4. **N. nanus**, DC. Smooth; stem *low and simple* (5'–10' high); the heads in axillary clusters forming a narrow racemed panicle; leaves triangular-halberd-shaped and very variously lobed or cleft, on slender petioles; *involucre* (livid) *10–13-flowered, of about 8 proper scales and several very short bract-like ones, which are triangular-ovate and appressed*; pappus straw-color. — Alpine summits, White Mountains of New Hampshire, and Adirondacks, N. New York.

5. **N. Boóttii**, DC. Stem *simple, dwarf* (5'–6' high), pubescent at the summit; the heads in an almost simple raceme; lowest leaves halberd-shaped or heart-shaped, the middle oblong, the upper lanceolate, nearly entire, taper-

ing into a margined petiole; involucre (livid) 10-18-flowered, of 10-15 very obtuse proper scales, and several linear and loose exterior ones nearly half the length of the former; pappus straw-color. — Higher alpine summits of the mountains of Maine, New Hampshire, and N. New York.

6. **N. virgatus**, DC. (SLENDER RATTLESNAKE-ROOT.) Smooth, slightly glaucous; stem simple (2° - 4° high), prolonged into a naked and slender spiked raceme ($1\frac{1}{2}^{\circ}$ - 2° long); heads clustered and mostly unilateral; leaves lanceolate, acute, closely sessile, the upper reduced to bracts, the lower toothed or pinnatifid; involucre (purplish) of about 8 scales, 8-12-flowered; pappus straw-color. — Sandy pine barrens, New Jersey to Virginia, and southward.

* * Involucre 12-40-flowered, hairy, as well as the peduncles.

7. **N. racemösus**, Hook. Stem wand-like, simple (2° - 5° high), smooth, as well as the oval or oblong-lanceolate denticulate leaves; the lower tapering into winged petioles (rarely cut-pinnatifid,) the upper partly clasping; heads in clusters crowded in a long and narrow interruptedly spiked panicle; involucre about 12-flowered; pappus straw-color. — Plains, Ohio to Illinois and northward. Also Hackensack marshes, New Jersey. — Flowers flesh-color.

8. **N. äsper**, Torr. & Gr. Stem wand-like, simple (2° - 4° high), rough-pubescent, as well as the oval-oblong or broadly lanceolate toothed leaves; heads (mostly erect) in small clusters disposed in a long and narrow compound raceme; involucre 12-14-flowered; pappus straw-color. — Dry prairies and barrens, Ohio to Illinois, and southward. — Flowers larger than No. 7, cream-color.

9. **N. crepidíneus**, DC. Somewhat smooth; stem stout (5° - 8° high), bearing numerous nodding heads in loose clusters on the corymbose-panicled branches; leaves large ($6'$ - $12'$ long), broadly triangular-ovate or halberd-form, strongly toothed, contracted into winged petioles; involucre 20-40-flowered; pappus brown. — Rich soil, Ohio to Illinois and southward. — Involucre blackish; flowers cream-color.

79. LYGODESMIA, Don. LYGODESMIA.

Heads and flowers (5-10) nearly as in Nabalus; the cylindrical involucre more elongated, and the achenia long and slender, tapering at the summit. Pappus whitish. — Smooth, often glaucous, low perennials, with single erect heads of rose-purple flowers terminating almost leafless or rush-like stems or branches. (Name composed of *λύγος*, a twig for wickerwork, and *δεσμός*, a bond, from the twiggy or rush-like stems.)

1. **L. júncea**, Don. Stems (1° high) tufted, branched, striate; lower leaves lance-linear, $1'$ - $2'$ long, rigid, the upper awl-shaped and minute; heads 5-flowered. — St. Croix River, Wisconsin, *T. J. Hale*, and westward, July.

80. CHONDRILLA, Tourn. CHONDRILLA.

Heads few-flowered. Involucre cylindrical, of several narrow linear equal scales, and a row of small bractlets at the base. Achenia terete, several-ribbed, smooth below, roughened at the summit by little scaly projections, from among which springs an abrupt slender beak. Pappus of copious very fine and soft capillary bristles, bright white. — Herbs of the Old World, with wand-like

branching stems, and small heads of yellow flowers. (A name of Dioscorides for some plant which exudes a gum.)

1. **C. JÚNCEA**, L. Biennial, bristly-hairy below, smooth above (1° – 3° high); root-leaves runcinate; stem-leaves few and small, linear; heads scattered on nearly leafless branches, 6"–8" long. — Fields and roadsides, abundant near Alexandria, Virginia, *M. J. Bebb*, *A. H. Curtiss*; perhaps of recent introduction. Aug. (Adv. from Eu.)

81. PYRRHOPÁPPUS, DC. FALSE DANDELION.

Heads, &c. nearly as in *Taraxacum*, but the soft pappus reddish or rusty-color, and with a villous ring at the top of the long beak of the achenium. — Mostly annual or biennial herbs, often branching and leafy below. Heads solitary, terminating the naked summit of the stem or branches. Flowers deep yellow. (Name composed of *πυρρῶς*, *flame-colored*, and *παππός*, *pappus*.)

1. **P. Caroliniánus**, DC. Stem branching (1° – 2° high); leaves oblong or lanceolate, entire, cut, or pinnatifid, the stem-leaves partly clasping. — Sandy fields, from Maryland southward. April–July.

82. TARÁXACUM, Haller. DANDELION.

Head many-flowered. Involucre double, the outer of short scales; the inner of long linear scales, erect in a single row. Achenia terete, oblong, ribbed, and roughened on the ribs, the apex prolonged into a very slender thread-like beak, bearing the pappus of copious soft and white capillary bristles. — Perennials or biennials, producing a tuft of pinnatifid or runcinate radical leaves, and slender naked hollow scapes, bearing a single large head of yellow flowers. (Name from *ταράσσω*, *to disquiet or disorder*, in allusion to medicinal properties.)

1. **T. Dens-leónis**, Desf. (COMMON DANDELION.) Smooth, or at first pubescent; outer involucre reflexed. — Pastures and fields everywhere: probably indigenous in the North. April–Sept. — After blossoming, the inner involucre closes, the slender beak elongates and raises up the pappus while the fruit is forming; the whole involucre is then reflexed, exposing to the wind the naked fruits, with the pappus displayed in an open globular head. (Eu.)

83. LACTŪCA, Tourn. LETTUCE.

Heads several-flowered. Scales of the involucre imbricated in 2 or more sets of unequal lengths. Achenia flat (compressed parallel to the scales of the involucre), abruptly contracted into a long thread-form beak, bearing a copious and fugacious pappus of very soft and white capillary bristles. — Leafy-stemmed herbs, with paniced heads; the flowers of variable color, produced in summer and autumn. (The ancient name of the Lettuce, *L. sativa*; from *lac*, milk, in allusion to the milky juice.)

1. **L. Canadénsis**, L. (WILD LETTUCE.) Biennial, mostly tall; leaves partly clasping, pale beneath; the upper lanceolate and entire; heads about 20-flowered; achenia oval, rather longer than the beak, minutely rugose transversely and roughish, one-ribbed on each face. The typical form (*L. elongata*, *Muhl.*,

and Ed. 2) is tall, with a thick and hollow very leafy stem (4° – 9° high), smooth or nearly so; leaves long, most of them runcinate-pinnatifid; heads very numerous, in a long and narrow naked panicle; flowers mostly pale yellow. — Rich and damp soil, borders of fields or thickets: common, especially northward. — The following are perhaps to be restored as species: —

Var. **integrifolia**, Torr. & Gr. (*L. integrifolia*, *Bigel.*) Stem 3° – 6° high; leaves all undivided, either entire or sharply denticulate; panicle more open; flowers pale yellow, cream-color, or purple. — Open and dry or sterile soil, E. New England near the coast to Illinois and southward.

Var. **sanguinea**, Torr. & Gr. (*L. sanguinea*, *Bigel.*) Lower and less stout (2° – 5° high); leaves all runcinate-pinnatifid, the midrib beneath and lower part of the stem often sparsely bristly-hairy; heads fewer, in a loose open panicle; flowers yellow-purple, reddish with or without a yellow centre, or rarely white. — Open dry ground, Eastern New England to New Jersey, Illinois, and southward.

2. **L. SCARÏOLA**, L. (PRICKLY LETTUCE.) Annual or biennial; stem below sparsely prickly-bristly, as also the midrib on the lower face of the sagittate-clasping oblong or lanceolate spinulose-denticulate vertical leaves; panicle narrow; heads small, few-flowered; achenia striate. — Waste grounds and roadsides, Cambridge, Massachusetts. (Adv. from Eu.)

84. **MULGËDIUM**, Cass. FALSE OR BLUE LETTUCE.

Heads many-flowered. Involucre, &c. as in *Lactuca*. Achenia laterally compressed, striate or ribbed, the summit contracted into a short and thick (or in No. 1 slender) beak or neck of the same texture, expanded at the apex into a ciliate disk, which bears a copious rather deciduous pappus of soft capillary bristles. — Leafy-stemmed herbs, with the general aspect and foliage of *Lactuca*; ours glabrous or nearly so. Heads racemed or paniced; the flowers chiefly blue; in summer. (Name from *mulgeo*, to milk.)

* *Pappus bright white: flowers blue.*

1. **M. pulchellum**, Nutt. *Perennial*, pale or glaucous; stem simple, 1° – 2° high; leaves sessile, oblong- or linear-lanceolate, entire, or the lower runcinate-pinnatifid; heads few and large, racemose, erect; scales of the conical-cylindrical involucre lanceolate, imbricated in 3 or 4 ranks; the peduncles scaly-bracted; *achenia tapering into a slender beak*, almost as in *Lactuca*. — Upper Michigan (*Prof. Porter, &c.*), probably in N. W. Wisconsin: common on the plains westward.

2. **M. acuminatum**, DC. Tall *biennial*, (3° – 6° high), with many small heads in a loose panicle, on diverging peduncles; leaves ovate or ovate-lanceolate, pointed, *barely toothed*, sometimes hairy on the midrib beneath, contracted into a winged petiole, the lowest occasionally sinuate; achenia with a very short beak. — Borders of thickets, New York to Illinois, and southward. — Probably only an entire-leaved state of the next.

3. **M. Floridanum**, DC. *Leaves all lyrate or runcinate*, the upper often with a heart-shaped clasping base; panicle larger: otherwise as No. 2. — Rich soil, Pennsylvania to Illinois and southward.

* * *Pappus tawny: corolla pale blue, or cream-color turning bluish.*

4. **M. leucophæum**, DC. Nearly smooth biennial; stem tall (3°-12° high), very leafy; leaves irregularly pinnatifid, sometimes runcinate, coarsely toothed, the uppermost often undivided and sometimes clasping; heads in a large and dense compound panicle. — Low grounds: rather common.

85. SÓNCHUS, L. SOW-THISTLE.

Heads many-flowered, becoming tumid at the base. Involucre more or less imbricated. Achenia flattened laterally, ribbed or striate, not beaked. Pappus copious, of very white exceedingly soft and fine capillary bristles. — Leafy-stemmed coarse weeds, chiefly smooth and glaucous, with corymbed or umbellate heads of yellow flowers; produced in summer and autumn. (The ancient Greek name.)

* *Annual (1°-5° high): flowers pale yellow.*

1. **S. OLERACEUS**, L. (COMMON SOW-THISTLE.) Stem-leaves runcinate-pinnatifid, or rarely undivided, slightly toothed with soft spiny teeth, clasping by a heart-shaped base, the auricles acute; involucre downy when young; *achenia striate, also wrinkled transversely.* — Waste places in manured soil and around dwellings. (Nat. from Eu.)

2. **S. ASPER**, Vill. (SPINY-LEAVED S.) Stem-leaves less divided and more spiny-toothed, the auricles of the clasping base rounded; *achenia margined, 3-nerved on each side, smooth.* — With and like the last. (Nat. from Eu.)

* * *Perennial, with creeping rootstocks: flowers bright yellow, in large heads.*

3. **S. ARVÉNSIS**, L. (FIELD S.) Leaves runcinate-pinnatifid, spiny-toothed, clasping by a heart-shaped base; peduncles and involucre bristly; *achenia transversely wrinkled on the ribs.* — Roadsides, &c., New England and New York: becoming more common. (Nat. from Eu.)

ORDER 54. LOBELIACEÆ. (LOBELIA FAMILY.)

Herbs with milky juice, alternate leaves, and scattered flowers, an irregular monopetalous 5-lobed corolla; the 5 stamens free from the corolla, and united into a tube commonly by their filaments and always by their anthers. — Calyx-tube adherent to the many-seeded pod. Style 1: stigma often fringed. Seeds anatropous, with a small straight embryo, in copious albumen. — Acrid poisonous plants (rather to be regarded as a part of the next order), represented only by the genus

1. LOBÈLIA, L. LOBELIA.

Calyx 5-cleft, with a short tube. Corolla with a straight tube, split down on the (apparently) upper side, somewhat 2-lipped; the upper lip of 2 rather erect lobes, the lower lip spreading and 3-cleft. Two of the anthers in our species bearded at the top. Pod 2-celled, many-seeded, opening at the top. — Flowers axillary or chiefly in bracted racemes; in summer and early autumn. (Dedicated to *Matthias De l'Obel*, an early Flemish herbalist.)

* *Flowers deep red, large: stem simple.*

1. **L. cardinalis**, L. (CARDINAL-FLOWER.) Tall (2° – 4° high), smoothish; leaves oblong-lanceolate, slightly toothed; raceme elongated, rather 1-sided; the pedicels much shorter than the leaf-like bracts. — Low grounds: common. — Perennial by offsets, with large and very showy intensely red flowers, — rarely varying to rose-color! (Plymouth, *Mr. Gilbert*), or even to white!

** *Flowers blue, or blue variegated with white.*

+ *Stems leafy to the top, simple (1° – 3° high) from a perennial root; leaves oblong or ovate-lanceolate: sinuses of the calyx with conspicuous deflexed auricles: flowers crowded in a long spike or dense raceme.*

2. **L. syphilitica**, L. (GREAT LOBELIA.) Somewhat hairy; leaves thin, acute at both ends ($2'$ – $6'$ long), irregularly serrate; flowers (nearly $1'$ long) pedicelled, longer than the leafy bracts; calyx hirsute, the lobes half the length of the corolla, the short tube hemispherical. — Low grounds: common. — Flowers light blue, rarely white.

3. **L. puberula**, Michx. *Finely soft-pubescent; leaves thickish, obtuse ($1'$ – $2'$ long), with small glandular teeth; spike rather 1-sided; calyx-lobes (and ovate bracts) little shorter than the corolla, the hairy tube top-shaped.* — Moist grounds, New Jersey to Illinois and southward. — Corolla bright blue, $\frac{1}{2}'$ long.

4. **L. leptostachys**, A. DC. *Smooth above; leaves obtuse, denticulate, oblong-lanceolate, the upper gradually reduced to awl-shaped bracts; raceme spike-like, long and dense; lobes of the calyx nearly equalling the corolla, the auricles in the form of 10 awl-shaped appendages as long as the hemispherical tube.* — Sandy soil, Illinois and southward. — Corolla $3''$ – $4''$ long.

+ + *Stems leafy, mostly simple ($1'$ – $2\frac{1}{2}'$ high) from a perennial root: leaves lanceolate or oblong-lanceolate: calyx-tube hemispherical, the sinuses destitute of auricles: flowers pretty large ($\frac{2}{3}'$ – $1'$ long) and showy, in a loose nearly 1-sided raceme: anthers sometimes bearded on the back.*

5. **L. glandulosa**, Walt. Sparingly pubescent: leaves, bracts, and usually the lobes of the calyx strongly glandular-toothed; calyx-tube densely hispid, rarely sparsely so, or smoothish. — Moist places, Virginia and southward.

6. **L. amœna**, Michx. *Glabrous (rarely minutely pubescent); leaves and bracts usually glandular-toothed; calyx-lobes entire and slender.* — Shady moist places, Virginia and southward.

+ + + *Stems leafy: calyx with no auricles or appendages at the sinuses: flowers small $\frac{1}{4}'$ – $\frac{1}{2}'$ long, racemed: roots slender, annual or biennial, or perhaps sometimes perennial.*

+ + *More or less pubescent, at least below: leaves oblong or ovate: stems angled or striate: racemes spike-like: corolla pale blue.*

7. **L. inflata**, L. (INDIAN TOBACCO.) Stems paniculately much branched from an annual root, pubescent with spreading hairs ($9'$ – $18'$ high); leaves ovate or oblong, toothed, gradually diminishing into leaf-like bracts, which exceed the lower short-pedicelled flowers, calyx-tube ovoid, the pod inflated. — Dry open fields: common. — Corolla only $1\frac{1}{2}''$ – $2''$ long. Plant poisonous and a noted quack medicine.

8. **L. spicàta**, Lam. Stem slender, *strict and simple* (1° – 3° high) from a biennial or perhaps perennial root, below and the barely denticulate leaves minutely pubescent; lower and root-leaves obovate or spatulate, the upper reduced to linear or club-shaped bracts; *raceme long and naked*, mostly dense and many-flowered; calyx-tube short, obconical or becoming almost hemispherical. (*L. Claytoniana*, Michx. *L. pallida*, Muhl.) — Moist or dry, mostly gravelly or sandy soil: rather common, at least southward and westward. — A slender and smaller flowered variety (beginning to blossom in June) grows in swamps at Lancaster, Penn., *Prof. Porter*. — Corolla ordinarily 4" long.

++ ++ *Glabrous or nearly so: leaves small, linear or lanceolate, only those from the root obovate or spatulate, the uppermost reduced to setaceous bracts, all entire or barely denticulate: stems very slender, simple or becoming paniculately branched above: racemes loosely several-flowered.*

9. **L. Nuttállii**, Rœm. & Sch. Stem very slender (1° – 2° high), terete; *pedicels mostly longer than the bract and shorter than the flower*, usually with very minute bractlets near the base; *calyx-tube very short, depressed-hemispherical in fruit*, the globular pod half free; corolla pale blue, barely 3" long. — Sandy swamps, from Long Island, New Jersey, and the adjacent lower borders of Pennsylvania, southward.

10. **L. Kálmii**, L. Stem mostly low (4'–18' high) minutely angled; *pedicels filiform, not exceeding the linear or setaceous bracts but as long as the flower, minutely 2-bracteolate or 2-glandular above the middle; calyx-tube top-shaped or obovoid* with an acute base, fully half the length of the lobes, in fruit rather longer than they, smooth, covering the whole pod; corolla bright light blue, 4"–5" long. — Wet limestone rocks and banks, Northern New England to Wisconsin and northward along the St. Lawrence and Great Lakes, and through New York southward to Lancaster, Penn. (*Prof. Porter*).

11. **L. Cánbyi**, n. sp. Stem strict (1° – 2° high), minutely angled; *pedicels shorter than the bracts and mostly shorter than the flower*, minutely roughened under a lens; *bractlets none; calyx-tube top-shaped, acute at the base, and only half the length of the lobes* (which, with the linear leaves, are sparsely glandular-denticulate along the margins), in fruit becoming oblong, covering the whole pod; corolla deep blue (fully 5" long), more or less *bearded in the throat*. — Wet places, pine barrens of New Jersey, especially at Quaker Bridge, *Wm. M. Canby, C. E. Smith, &c.* (Also South Carolina, *M. A. Curtis.*) Aug., Sept. — Leaves 1', rarely $1\frac{1}{2}'$ long, numerous, gradually diminishing in size up to the raceme, the largest 2" wide. Pod nearly 3" long.

+ + + + *Stem simple from a perennial root, and nearly leafless, except at or near the base: flowers in a simple loose raceme, light blue: leaves fleshy: calyx-tube acute at the base, top-shaped: auricles none.*

12. **L. paludòsa**, Nutt. Nearly smooth; stem slender (1° – $2\frac{1}{2}^{\circ}$ high); *leaves thickish but flat, scattered near the base, linear-spatulate or oblong-linear, glandular-denticulate, mostly tapering into a petiole; lower lip of the corolla bearded in the middle; calyx-tube about half the length of the short lobes, becoming hemispherical in fruit*. — Wet bogs, Delaware (*Nuttall*) and southward. — Corolla 5"–6" long.

13. **L. Dortmanna**, L. (WATER LOBELIA.) Very smooth; *scape thickish* (5'–12' high), *few-flowered*; *leaves all tufted at the root, linear, terete, hollow*, with a partition lengthwise, sessile; lower lip of the corolla slightly hairy; calyx-tube about as long as the lobes, in fruit much longer. — In the gravelly borders of ponds, New York, New England, and northward. — Corolla 6''–8'' long. (Eu.)

ORDER 55. **CAMPANULACEÆ.** (CAMPANULA FAMILY.)

Herbs, with milky juice, alternate leaves, and scattered flowers; the calyx adherent to the ovary; the regular 5-lobed corolla bell-shaped, valvate in the bud; the 5 stamens free from the corolla and usually distinct. — Style 1, beset with collecting hairs above: stigmas 2 or more. Pod 2–several-celled, many-seeded. Seed small, anatropous, with a straight embryo in fleshy albumen. — Flowers generally blue and showy. — Sparingly represented in America, and in the Northern States by only two genera.

1. **CAMPANULA**, Tourn. BELLFLOWER.

Calyx 5-cleft. Corolla generally bell-shaped, 5-lobed. Stamens 5, separate; the filaments broad and membranaceous at the base. Stigmas and cells of the pod 3 in our species, the short pod opening on the sides by as many valves or holes. — Herbs, with terminal or axillary flowers; in summer. (A diminutive of the Italian *campana*, a bell, from the shape of the corolla.)

* *Indigenous species, perennials, except perhaps No. 2.*

+ *Flowers loosely panicled (or rarely solitary), long-peduncled: pods nodding.*

1. **C. rotundifolia**, L. (HAREBELL.) Slender, branching (5'–12' high), 1–10-flowered; *root-leaves round-heart-shaped* or ovate, mostly toothed or crenate, long-petioled, early withering away; *stem-leaves numerous, linear or narrowly lanceolate, entire, smooth*; *calyx-lobes awl-shaped*, varying from $\frac{1}{3}$ to $\frac{2}{3}$ the length of the bright-blue corolla (which is 6''–9'' long). — Rocky shaded banks: common northward, and along the mountains. — A delicate and pretty, but variable species, with a most inappropriate name, since the round root-leaves are rarely obvious. (Eu.)

Var. **linifolia**. Stems more upright and rather rigid; the lowest leaves varying from heart-shaped to ovate-lanceolate; corolla $\frac{2}{3}$ '–1' long. (*C. linifolia*, Lam.) — Shores of the Great Lakes, and northwestward. (Eu.)

2. **C. aparinoides**, Pursh. (MARSH BELLFLOWER.) *Stem simple and slender, weak* (8'–20' high), few-flowered, somewhat 3-angled, *rough backwards on the angles, as are the slightly toothed edges and midrib of the linear-lanceolate leaves; peduncles diverging, slender; lobes of the calyx triangular, half the length of the bell-shaped nearly white corolla.* (*C. erinoides*, Muhl.) — Bogs and wet meadows, among high grass. — Plant with somewhat the habit of a Galium; the corolla barely 4'' long.

3. **C. divaricata**, Michx. Very smooth; stem loosely branched (1°–3° high); *leaves oblong-lanceolate, pointed at both ends, coarsely and sharply toothed*;

flowers numerous on the branches of the large compound panicle, *calyx-lobes awl-shaped*, about half the length of the pale-blue small corolla (of 3" long); *style protruded*. — Dry woods and rocks, mountains of Virginia, Kentucky, and southward.

+ + *Flowers numerous and nearly sessile, crowded in a long more or less leafy spike: corolla almost wheel-shaped, deeply 5-lobed: pods erect.*

4. **C. Americana**, L. (TALL BELLFLOWER.) Stem mostly simple (3°-6° high); leaves ovate and ovate-lanceolate, taper-pointed, serrate, mostly on margined petioles, thin, somewhat hairy (2½'-6' long); the slender style protruded and curved. — Moist rich soil, New York to Wisconsin, and southward. — Spike 1°-2° long. Corolla light blue, 1' broad.

* *European species escaped from gardens into roadsides; both perennials.*

5. **C. GLOMERATA**, L. (CLUSTERED B.) Somewhat hairy, stout and erect, 1° high; stem-leaves oblong or lanceolate, cordate-clasping; flowers sessile, clustered in the upper axils, forming a leafy head; corolla open-bell-shaped, 1' long. — Danvers, Mass., &c. (Adv. from Eu.)

6. **C. RAPUNCULOIDES**, L. Smoothish, slender, erect; stem-leaves ovate-lanceolate, pointed, the lower long-petioled and heart-shaped; flowers nodding, single in the axil of bracts, forming racemes; corolla oblong, 1' long. — E. Massachusetts; and Richfield Springs, New York, *G. W. Clinton*. (Adv. from Eu.)

2. SPECULARIA, Heister. VENUS'S LOOKING-GLASS.

Calyx 5- (or 3-4-) lobed. Corolla wheel-shaped, 5-lobed. Stamens 5, separate; the membranaceous hairy filaments shorter than the anthers. Stigmas 3. Pod prismatic or elongated-oblong, 3-celled, opening by 3 small lateral valves. — Low annuals; the earlier flowers in the American species (§ TRIODALLUS, Raf.) minute and fruiting precociously in the bud, without expanding their imperfect corolla. (Name from *Speculum Veneris*, the early name of the common European species.)

1. **S. perfoliata**, A. DC. Somewhat hairy (3'-20' high); leaves roundish or ovate, clasping by the heart-shaped base, toothed; flowers sessile, solitary or 2-3 together in the axils; the upper or later ones only with a conspicuous and expanding (purple-blue) corolla; pod oblong, opening rather below the middle. — Sterile open ground: common. May-Aug.

ORDER 56. ERICACEÆ. (HEATH FAMILY.)

Shrubs, sometimes herbs, with the flowers regular or nearly so: the stamens as many or twice as many as the 4-5-lobed or 4-5-petalled corolla, free from but inserted with it: anthers 2-celled, commonly appendaged or opening by terminal chinks or pores, introrse (except in Suborder 3): style 1: ovary 3-10-celled. Pollen compound, of 4 united grains, except in Suborder 4. Seeds small, anatropous. Embryo small, or sometimes minute, in fleshy albumen. — A large family, very various in many of the characters, comprising four well-marked suborders, as follows: —

SUBORDER I. **VACCINIEÆ.** WHORTLEBERRY FAMILY.

Calyx-tube adherent to the ovary, which forms an edible berry or berry-like fruit, crowned with the short calyx-teeth. Anthers 2-parted. — Shrubs or somewhat woody plants, with scaly buds.

1. **Gaylussacia.** Ovary 8–10-celled, with a single ovule in each cell. Fruit a berried drupe with 8 or 10 small seed-like nutlets.
2. **Vaccinium.** Berry 4–6-celled (or imperfectly 8–10-celled by false partitions), many-seeded. Anther-cells tapering upward into a tube.
3. **Chiogenes.** Berry 4-celled, many-seeded, its summit free. Anther-cells not prolonged into a tube, but each 2-pointed.

SUBORDER II. **ERICINEÆ.** PROPER HEATH FAMILY.

Calyx free from the ovary. Corolla monopetalous, or sometimes polypetalous, hypogynous. — Shrubs or small trees.

Tribe I. ARBUTEÆ. Fruit indehiscent, a berry or drupe. Corolla deciduous.

4. **Arctostaphylos.** Corolla urn-shaped. Drupe berry-like, 5–10-seeded.

Tribe II. ANDROMEDEÆ. Fruit a loculicidal pod. Corolla deciduous.

* Anthers upright in the bud. Corolla monopetalous.

+ Anther-cells opening through their whole length; not appendaged.

5. **Epigæa.** Corolla salver-shaped. Calyx of 5 separate dry and pointed sepals.

+ + Anther-cells opening only at the top. Corolla not salver-shaped.

+ + Calyx becoming enlarged and berry-like in fruit.

6. **Gaultheria.** Calyx 5-cleft, in fruit enclosing the small pod. Anthers 4-awned at the top.

+ + + Calyx dry, not becoming fleshy after flowering.

7. **Leucothoe.** Calyx imbricated in the bud. Corolla cylindraceous, 5-toothed. Pod depressed, 5-lobed, the valves entire.

8. **Cassandra.** Calyx imbricated in the bud. Corolla cylindraceous, 5-toothed. Pod splitting when ripe into an outer and inner layer, the inner of 10 valves.

9. **Cassiope.** Calyx imbricated. Corolla broadly campanulate, deeply 4–5-cleft. Pod globular-ovoid, 4–5-valved, the valves 2-cleft.

10. **Andromeda.** Calyx valvate and very early open in the bud. Pod globular. Seeds mostly hanging on the central placenta.

11. **Oxydendrum.** Calyx valvate and opening early in the bud. Pod oblong-pyramidal. Seeds all ascending.

** Anthers turned over outwardly in the bud. Corolla of 5 separate petals.

12. **Clethra.** Sepals 5. Stamens 10: anther-cells opening by a terminal hole or chink. Style 3-cleft at the apex. Pod 3-valved.

Tribe III. ERICEÆ. Corolla persistent, becoming scarious after flowering.

13. **Calluna.** Corolla bell-shaped, 4-parted. Pod opening septicidally.

Tribe IV. RHODOREÆ. Fruit a septicidal pod. Corolla deciduous.

* Anther-cells opening by a hole or chink at the top.

+ Flowers not from scaly buds; the bracts leaf-like or coriaceous.

14. **Phyllodoce.** Corolla ovate or urn-shaped. Leaves narrow and heath-like.

15. **Kalmia.** Corolla broadly bell-shaped or wheel-shaped, with 10 pouches receiving as many anthers. Leaves oblong or linear.

+ + Flowers developed from large scaly buds, the scales or bracts caducous.

16. **Menziesia.** Corolla globular-bell-shaped, 4-toothed. Stamens 8. Leaves deciduous.

17. **Azalea.** Corolla open funnel-form, 5-lobed. Stamens 5. Leaves deciduous.

18. **Rhododendron**. Corolla bell-shaped or funnel-form. Stamens 10. Leaves evergreen.
 19. **Rhodora**. Corolla irregular, two petals nearly separate. Stamens 10. Leaves deciduous.
 20. **Ledum**. Corolla regular, all 5 petals nearly separate. Stamens 5-10. Leaves evergreen.
 * * Anther-cells opening lengthwise. Leaves evergreen. Bud-scales firm and persistent.
 21. **Loiseleuria**. Corolla deeply 5-cleft. Stamens 5, included.
 22. **Leiophyllum**. Corolla of 5 separate petals. Stamens 10, exserted.

SUBORDER III. **PYROLEÆ**. PYROLA FAMILY.

Calyx free from the ovary. Corolla polypetalous. Anthers extrorse in the bud. Seeds with a loose and translucent cellular coat much larger than the nucleus. — Nearly herbaceous and broad-leaved evergreens.

23. **Pyrola**. Flowers in a raceme. Petals not widely spreading. Filaments awl-shaped. Style long. Valves of the pod cobwebby on the edges.
 24. **Moneses**. Flower single. Petals widely spreading. Filaments not dilated in the middle: anthers conspicuously 2-horned. Style straight, exserted: stigma 5-rayed. Valves of the pod smooth on the edges.
 25. **Chimaphila**. Flowers corymb or umbelled. Petals widely spreading. Filaments dilated in the middle: anthers 2-horned. Style very short and top-shaped, covered by a broad and orbicular stigma. Valves of the pod smooth on the edges.

SUBORDER IV. **MONOTROPEÆ**. INDIAN-PIPE FAMILY.

Flowers nearly as in Suborders 2 or 3, but the plants herbaceous, root-parasitic, entirely destitute of green foliage, and with the aspect of Beechdrops. Seeds as in Suborder 3.

* Corolla monopetalous: anthers 2-celled.

26. **Pterospora**. Corolla ovate, 5-toothed: anthers 2-awned on the back, opening lengthwise.
 27. **Schweinitzia**. Corolla broadly bell-shaped, 5-lobed: anthers opening at the top.

* * Corolla of 4 or 5 separate petals: calyx imperfect or bract-like.

28. **Monotropa**. Petals narrow. Anthers kidney-shaped, opening across the top.

I. **GAYLUSSÁCIA**, H. B. K. HUCKLEBERRY.

Corolla tubular, ovoid, or bell-shaped; the border 5-cleft. Stamens 10: anthers awnless; the cells tapering upward into more or less of a tube, opening by a chink at the end. Fruit a berry-like drupe, containing 10 seed-like nutlets. — Branching shrubs, with the aspect of *Vaccinium*, commonly sprinkled with resinous dots; the flowers (white tinged with purple or red) in lateral and bracted racemes. (Named for the distinguished chemist, *Gay-Lussac*.)

* *Leaves thick and evergreen, not resinous-dotted.*

1. **G. brachýcera**, Gray. (BOX-HUCKLEBERRY.) Very smooth (1° high); leaves oval, finely crenate-toothed; racemes short and nearly sessile; pedicels very short; corolla cylindrical-bell-shaped. — Dry Woods, Perry Co., Pennsylvania, near Bloomfield (*Prof. Baird*), and mountains of Virginia. May. — Leaves in shape and aspect like those of the Box.

* * *Leaves deciduous, entire, sprinkled more or less with resinous or waxy atoms.*

2. **G. dumosa**, Torr. & Gr. (DWARF HUCKLEBERRY.) Somewhat hairy and glandular, low (1°-5° high from a creeping base), bushy; leaves obovate-

oblong, mucronate, green both sides, rather thick and shining when old; racemes elongated; bracts leaf-like, oval, persistent, as long as the pedicels; ovary bristly or glandular; corolla bell-shaped; fruit black (insipid). — Var. *HIRTÉLLA* has the young branchlets, racemes, and often the leaves hairy. — Sandy-low soil, Maine to Penn. and Virginia, near the coast, and southward. June.

3. *G. frondosa*, Torr. & Gr. (BLUE TANGLE. DANGLEBERRY.) Smooth (3°–6° high); branches slender and divergent; leaves obovate-oblong, blunt, pale, glaucous beneath; racemes slender, loose; bracts oblong or linear, deciduous, shorter than the slender drooping pedicels; corolla globular-bell-shaped; fruit dark blue with a white bloom (sweet and edible). — Low copses, coast of New England to Kentucky, and southward. May, June.

4. *G. resinosa*, Torr. & Gr. (BLACK HUCKLEBERRY.) Much branched, rigid, slightly pubescent when young (1°–3° high); leaves oval, oblong-ovate, or oblong, thickly clothed and at first clammy, as well as the flowers, with shining resinous globules; racemes short, clustered, one-sided; pedicels about the length of the flowers; bracts and bractlets (reddish) small and deciduous; corolla ovoid-conical, or at length cylindrical with an open mouth; fruit black, without bloom (pleasant, very rarely white). — Woodlands and swamps: common (except southwestward towards the Mississippi). May, June. — The common *Huckleberry* of the North.

2. VACCÍNIUM, L. CRANBERRY. BLUEBERRY. BILBERRY.

Corolla various in shape; the limb 4–5-cleft, revolute. Stamens 8 or 10: anthers sometimes 2-awned on the back; the cells separate and prolonged upwards into a tube, opening by a hole at the apex. Berry 4–5-celled, many-seeded, or sometimes 8–10-celled by a false partition stretching from the back of each cell to the placenta. — Shrubs with solitary, clustered, or racemed flowers: the corolla white or reddish. (Ancient Latin name, of obscure derivation.)

§ 1. *OXYCÓCCUS*, Tourn. Ovary 4-celled: corolla 4-parted, the long narrow divisions revolute: anthers 8, awnless, tapering above into very long tubes: pedicels slender.

* Stems very slender, creeping or trailing: leaves small, entire, whitened beneath, evergreen: pedicels erect, with the pale rose-colored flower nodding on their summit: corolla deeply 4-parted: berries red, acid.

1. *V. Oxycoccus*, L. (SMALL CRANBERRY.) Stems very slender (4'–9' long); leaves ovate, acute, with strongly revolute margins (2''–3'' long); pedicels 1–4, terminal; filaments more than half the length of the anthers. (*Oxycoccus vulgaris*, Pursh.) — Peat-bogs, New England and Pennsylvania to Wisconsin, and northward. June. — Berry 3''–4'' broad, often speckled with white when young; seldom gathered for the market. (Eu.)

2. *V. macrocárpum*, Ait. (LARGE OR AMERICAN CRANBERRY.) Stems elongated (1°–3° long), the flowering branches ascending; leaves oblong, obtuse, glaucous underneath, less revolute (4''–6'' long); pedicels several, becoming lateral; filaments scarcely one third the length of the anthers. (*Oxycoccus macrocárpum*, Pers.) — Peat-bogs, Virginia to Wisconsin, and everywhere northward, but scarcely westward. June. — Berry $\frac{1}{2}$ '–1' long.

* * *Stem upright and leaves deciduous, as in common Blueberries: flowers axillary and solitary: corolla deeply 4-cleft: berries turning purple, insipid.*

3. **V. erythrocarpon**, Michx. Smooth, divergently branched (1° – 4° high); leaves oblong-lanceolate, taper-pointed, bristly serrate, thin. — Wooded hills, mountains of Virginia and southward. July.

§ 2. **VITIS-IDÆA**, Tourn. Ovary 4–5-celled: corolla bell-shaped, 4–5-lobed: anthers 8–10, awnless: filaments hairy: flowers in short and bracted nodding racemes: leaves evergreen: berries red or purple.

4. **V. Vitis-Idæa**, L. (COWBERRY.) Low (6'–10' high); branches erect from tufted creeping stems; leaves obovate, with revolute margins, dark green, smooth and shining above, dotted with blackish bristly points underneath; corolla bell-shaped, 4-cleft. — Higher mountains of New England, also on the coast of Maine, and at Danvers, Massachusetts (*Oakes*), and northward. June. — Berries dark red, acid and rather bitter, mealy, barely edible. (Eu.)

§ 3. **PICROCÓCCUS**, Nutt. Ovary more or less 10-celled by false partitions: berries greenish, hardly edible, ripening few seeds: corolla open-bell-shaped, 5-lobed: anthers 10, extended into very long much exerted tubes, 2-awned on the back: flowers on slender pedicels, singly in the axils of the upper leaves or leaf-like bracts, forming leafy racemes, not articulated: leaves thin, deciduous.

5. **V. stamineum**, L. (DEERBERRY. SQUAW HUCKLEBERRY.) Diffusely branched (2° – 3° high), somewhat pubescent; leaves ovate or oval, pale, glaucous or whitish underneath; corolla greenish-white or purplish; berries globular or pear-shaped, large, greenish, mawkish. — Dry woods, Maine to Michigan and southward, mainly eastward. May, June.

§ 4. **BATODÉNDRON**, Nutt. Ovary more or less 10-celled by false partitions: berries black: corolla short-bell-shaped, 5-toothed: anthers 10, included, conspicuously 2-awned on the back, and extended into slender tubes: filaments hairy: flowers on slender pedicels singly in the axils of coriaceous shining leaves, or racemed at the end of the branches, articulated just below the ovary!

6. **V. arboreum**, Marshall. (FARKLE-BERRY.) Tall (8° – 15° high), smoothish; leaves oval or obovate, entire or denticulate, mucronate, bright green and shining above, at the South evergreen; corolla white; berries mealy, insipid, ripening late. — Dry ground, Makanda, S. Illinois (*Dr. Vasey*), probably also in Virginia, and southward. June.

§ 5. **EUVACCÍNIUM**. Ovary 4–5-celled, with no trace of false partitions: corolla urn-shaped or globular, 4–5-toothed: anthers 2-awned on the back: filaments smooth: flowers axillary, solitary, or 2 or 3 together: berries blue or black, edible: northern or alpine plants, with deciduous leaves.

* *Parts of the flower mostly in fours: stamens 8.*

7. **V. uliginosum**, L. (BOG BILBERRY.) Low and spreading (4'–18' high), tufted; leaves entire, dull, obovate or oblong, pale and slightly pubescent underneath; flowers single or 2–3 together from a scaly bud, almost sessile; corolla short, urn-shaped; berries black with a bloom, sweet. — Alpine tops of the high mountains of New England and New York, shore of Lake Superior, and northward. (Eu.)

* * *Parts of the flower in fives: stamens 10: leaves membranaceous: flowers solitary on short axillary peduncles, nodding.*

8. **V. cæspitosum**, Michx. Dwarf (3'–5' high), tufted; leaves obovate, narrowed at the base, smooth and shining, serrate; corolla oblong, slightly urn-shaped; berries blue. — Alpine region of the White Mountains of New Hampshire; and high northward.

9. **V. ovalifolium**, Smith. Straggling, 3°–10° high; leaves elliptical, obtuse, nearly entire, pale, mostly glaucous beneath, smooth; corolla ovoid; berries blue. — Peat-bogs, Keweenaw Co., Lake Superior, *Dr. Robbins* (and far westward). May.

10. **V. myrtilloides**, Hook. More erect, 1°–4° high; branchlets somewhat angled; leaves mostly ovate and acute or pointed, sharply and closely serrulate, bright green, nearly smooth; border of the calyx almost entire; corolla depressed-globular, rather large; berries large, black, rather acid. — Woods and bluffs, Keweenaw Co., Lake Superior, *Dr. Robbins*. (Lake Huron, *Dr. Todd*; and northwestward.) May, June. — Pedicels 3"–6" long, drooping in flower, erect in fruit.

§ 6. **CYANOCÓCCUS**. *Ovary more or less completely 10-celled by false partitions: corolla oblong-cylindrical or slightly urn-shaped, 5-toothed: anthers 10, awnless: filaments hairy: berries blue or black with a bloom (sweet): flowers in clusters or very short racemes from scaly buds separate from and rather preceding the leaves, on short pedicels, appearing in early spring. (Leaves deciduous in the Northern species or proper Blueberries.)*

11. **V. Pennsylvanicum**, Lam. (DWARF BLUEBERRY.) Dwarf (6'–15' high), smooth; leaves lanceolate or oblong, distinctly serrulate with bristle-pointed teeth, smooth and shining both sides (or sometimes downy on the midrib underneath); corolla short, cylindrical-bell-shaped. — Var **ANGUSTIFOLIUM** is a high mountain or boreal form, 3'–6' high, with narrower lanceolate leaves. (*V. angustifolium*, *At.*) — Dry hills and woods: common from Pennsylvania and N. Illinois far northward. — Branches green, angled, warty. Berries abundant, large and sweet, ripening early in July: the earliest blueberry or blue huckleberry in the market.

12. **V. Canadense**, Kalm. (CANADA BLUEBERRY.) Low (1°–2° high); leaves oblong-lanceolate or elliptical, entire, downy both sides, as well as the crowded branchlets; corolla shorter: otherwise as the last, into which it seems to pass. — Swamps or moist woods, Maine to Wisconsin, and northward.

13. **V. vacillans**, Solander. (LOW BLUEBERRY.) Low (1°–2½° high), glabrous; leaves obovate or oval, very pale or dull, glaucous, at least underneath, minutely ciliolate-serrulate or entire; corolla between bell-shaped and cylindrical, the mouth somewhat contracted. — Dry woodlands, especially in sandy soil, New England to Virginia and N. Illinois. — Branches yellowish-green. Berries ripening later than those of No. 11.

14. **V. corymbosum**, L. (COMMON or SWAMP-BLUEBERRY.) Tall (5°–10° high); leaves ovate, oval, oblong, or elliptical-lanceolate; corolla varying from turgid-ovate and cylindrical-urn-shaped to oblong-cylindrical. — Swamps and low thickets: everywhere common, except southwestward. — This yields

the common *blueberry* or *blue huckleberry* of the latter part of the season. The typical form has the leaves entire and more or less pubescent, at least when young, as also the branchlets. The species exhibits the greatest variety of forms; the last of those here mentioned is the most remarkable, and the only one which has any claims to be regarded as a species.

Var. **glábrum**, is wholly or nearly glabrous throughout; the leaves entire.

Var. **amœnum**, has the leaves bristly-ciliate, shining above, green both sides, beneath somewhat pubescent on the veins. (V. amœnum, *Ait.*, &c.)

Var. **pállidum**, has the leaves mostly glabrous, pale or whitish, glaucous especially underneath, serrulate with bristly teeth. (V. pallidum, *Ait.*)

Var. **atrocóccum**, has the leaves entire, downy or woolly underneath even when old, as also the branchlets; berries smaller, black, without bloom. (V. fuscátum, *Ait.*? & Ed. 1.)

3. CHIÓGENES, Salisb. CREEPING SNOWBERRY.

Calyx-tube adherent to the lower part of the ovary; the limb 4-parted. Corolla bell-shaped, deeply 4-cleft. Stamens 8, included, inserted on an 8-toothed epigynous disk: filaments very short and broad: anther-cells ovate-oblong, quite separate, not awned on the back, but each minutely 2-pointed at the apex, and opening by a large chink down to the middle. Berry white, globular, crowned with the 4-toothed calyx, rather dry, 4-celled, many-seeded. — A trailing and creeping evergreen, with very slender and scarcely woody stems, and small Thyme-like, ovate and pointed leaves on short petioles, with revolute margins, smooth above, the lower surface and the branches beset with rigid rusty bristles. Flowers very small, solitary in the axils, on short nodding peduncles, with 2 large bractlets under the calyx. (Name from *χιών*, *snow*, and *γένος*, *offspring*, in allusion to the snow-white berries.)

1. **C. hispídula**, Torr. & Gr. — Peat-bogs, and mossy mountain woods, in the shade of evergreens; common northward, extending southward in the Alleghanies. May. — Plant with the aromatic flavor of Gaultheria or Birch. Leaves 3''–4'' long. Berries 3'' broad, bright white.

4. ARCTOSTÁPHYLOS, Adans. BEARBERRY.

Corolla ovate and urn-shaped, with a short revolute 5-toothed limb. Stamens 10, included: anthers with 2 reflexed awns on the back near the apex, opening by terminal pores. Drupe berry-like, with 5–10 seed-like nutlets. — Shrubs, with alternate leaves, and scaly-bracted nearly white flowers in terminal racemes or clusters. Fruit austere. (Name composed of *ἄρκτος*, *a bear*, and *σταφυλή*, *a grape or berry*, the Greek of the popular name.)

1. **A. Uva-úrsi**, Spreng. (BEARBERRY.) Trailing; leaves thick and evergreen, obovate or spatulate, entire, smooth; fruit red. (*Árbutus Uva-ursi*, *L.*) — Rocks and bare hills, New Jersey to Wisconsin and northward. May. (Eu.)

2. **A. alpína**, Spreng. (ALPINE BEARBERRY.) Dwarf, tufted and depressed; leaves deciduous, serrate, wrinkled with strong netted veins, obovate; fruit black. — Alpine region of the White Mountains, New Hampshire, Mount Katahdin, Maine, and high northward. (Eu.)

5. **EPIGÆA**, L. GROUND LAUREL. TRAILING ÁRBUTUS.

Corolla salver-form; the tube hairy inside, as long as the ovate-lanceolate pointed and scale-like nearly distinct sepals. Stamens 10, with slender filaments: anthers oblong, awnless, opening lengthwise. Style slender, its apex (as in *Pyrola*) forming a sort of ring or collar around and partly adnate to the 5 little lobes of the stigma. Pod depressed-globular, 5-lobed, 5-celled, many-seeded. — A prostrate or trailing scarcely shrubby plant, bristly with rusty hairs, with evergreen and reticulated rounded and heart-shaped alternate leaves, on slender petioles, and with rose-colored flowers in small axillary clusters, from scaly bracts. (Name composed of *ἐπί*, upon, and *γη*, the earth, from the trailing growth.)

1. **E. rèpens**, L. — Sandy woods, or sometimes in rocky soil, especially in the shade of pines: common in many places, especially eastward. — Flowers appearing in early spring, exhaling a rich spicy fragrance. In New England called **MAYFLOWER**.

6. **GAULTHÈRIA**, Kalm. AROMATIC WINTERGREEN.

Corolla cylindrical-ovoid or a little urn-shaped, 5-toothed. Stamens 10, included: anther-cells each 2-awned at the summit, opening by a terminal pore. Pod depressed, 5-lobed, 5-celled, 5-valved, many-seeded, enclosed when ripe by the calyx, which thickens and turns fleshy, so as to appear as a globular red berry! — Shrubs, or almost herbaceous plants, with alternate evergreen leaves and axillary (nearly white) flowers: pedicels with 2 bractlets. (Dedicated by Kalm to "*Dr. Gauthier*," of Quebec; *Linn. Amœn. Acad.* 3, p. 15. The true orthography, as ascertained by Prof. Brunet from the old records in Quebec, is *Gaultier*; so that the orthography of the genus, if changed at all, should be **GAULTIERA**.)

1. **G. procumbens**, L. (CREEPING WINTERGREEN.) Stems slender and extensively creeping on or below the surface; the flowering branches ascending, leafy at the summit (3' - 5' high); leaves obovate or oval, obscurely serrate; flowers few, mostly single in the axils, nodding. — Cool damp woods, mostly in the shade of evergreens especially northward, and southward along the Alleghanies. July. — The bright red berries (formed of the calyx) and the foliage have the well-known spicy-aromatic flavor of the Sweet Birch. In the interior of the country it is called *Wintergreen*, or sometimes *Tea-berry*. Eastward it is called *Checkerberry* or *Partridge-berry* (names also applied to *Mitchella*, the latter especially so), also *Boxberry*.

7. **LEUCÓTHOË**, Don. LEUCOTHOË.

Calyx of 5 nearly distinct sepals, imbricated in the bud, not enlarged nor fleshy in fruit. Corolla ovate or cylindraceous, 5-toothed. Stamens 10: anthers naked, or the cells with 1 or 2 erect awns at the apex, opening by a pore. Pod depressed, more or less 5-lobed, 5-celled, 5-valved, the sutures not thickened; valves entire: the many-seeded placenta borne on the summit of the short columella, mostly pendulous. — Shrubs with petioled and serrulate leaves, and white scaly-bracted flowers crowded in axillary or terminal spiked racemes. (A mythological name.)

§ 1. **LEUCOTHÖË** proper. *Anthers awnless; the cells sometimes obscurely 2-pointed: stigma depressed-capitate, 5-rayed: racemes sessile (dense), produced at the time of flowering from scaly buds in the axils of the coriaceous and shining persistent leaves of the preceding year, shorter than they: bracts persistent: bractlets at the base of the short pedicels. (Seed-coat loose and cellular, wing-like.)*

1. **L. axillàris**, Don. *Leaves lanceolate-oblong or oval, abruptly pointed or acute, somewhat spinulose-serrulate, on very short petioles; sepals broadly ovate. (Andromeda axillaris, Lam.)* — Banks of streams, Virginia, in the low country, and southward. Feb. — April. — Shrub 2° — 4° high.

2. **L. Catesbæi**. *Leaves ovate-lanceolate, taper-pointed, serrulate with ciliate-spinulose appressed teeth, conspicuously petioled (3' — 6' long); sepals ovate-oblong, often acute. (Andr. Catesbæi, Walt. A. axillaris, Michx. A. spinulosa, Pursh. L. spinulosa, Don.)* — Moist banks of streams, Virginia along the mountains, and southward. May. — Shrub 2° — 4° high, with long spreading or recurved branches. Flowers exhaling the unpleasant scent of Chestnut-blossoms.

§ 2. **EÛBOTRYS**, Nutt. *Anthers awned: stigma simple: bractlets close to the calyx, and, like the sepals, of a rigid texture, ovate or lanceolate, pointed: placentæ not pendulous: flowers very short-pedicelled, in long one-sided racemes, which mostly terminate the branches, formed with them in the summer, but the flower-buds not expanding till the following spring: bracts awl-shaped, deciduous: leaves membranaceous and deciduous, serrulate, the midrib and veins beneath pubescent.*

3. **L. recúrva**, Buckley. *Branches and racemes recurved-spreading; leaves lanceolate or ovate, taper-pointed; sepals ovate; anther-cells 1-awned; pod 5-lobed; seeds flat and cellular-winged.* — Dry hills, Alleghanies of Virginia and southward. April. — Lower and more straggling than the next.

4. **L. racemòsa**. *Branches and racemes mostly erect; leaves oblong or oval-lanceolate, acute; sepals ovate-lanceolate; anther-cells each 2-awned; pod not lobed; seeds angled and wingless. (Andromeda racemosa & A. paniculata, L.)* — Moist thickets, Massachusetts to Virginia, near the coast, and southward. May, June. — Shrub 4° — 6° high. Corolla cylindrical.

8. CASSÁNDRA, Don. LEATHER-LEAF.

Calyx of 5 distinct rigid ovate and acute sepals, imbricated in the bud, and with a pair of similar bractlets. Corolla cylindrical-oblong, 5-toothed. Stamens 10: anther-cells tapering into a tubular beak, and opening by a pore at the apex, awnless. Pod depressed, 5-celled, many-seeded; the pericarp of 2 layers, the outer 5-valved, the cartilaginous inner layer at length 10-valved. Seeds flattened, wingless. — Low and much branched shrubs, with nearly ever-green and coriaceous leaves, which are scurfy, especially underneath. Flowers white, in the axils of the upper small leaves, forming small 1-sided leafy racemes; the flower-buds formed in the summer and expanding early the next spring. (*Cassandra*, a daughter of Priam and Hecuba.)

1. **C. calyculàta**, Don. *Leaves oblong, obtuse, flat. (Andromeda calyculata, L.)* — Wet bogs: common northward: rare westward to North Wisconsin. (Eu.)

9. CASSIOPE, Don. CASSIOPE.

Calyx without bractlets, of 4 or 5 nearly distinct ovate sepals, imbricated in the bud. Corolla broadly campanulate, deeply 4-5-cleft. Stamens 8 or 10: anthers fixed by their apex; the ovoid cells each opening by a large terminal pore, and bearing a long recurved awn behind. Pod ovoid or globular, 4-5-celled, 4-5-valved; the valves 2-cleft: placentæ many-seeded, pendulous from the summit of the columella. Seeds smooth and wingless. — Small, arctic or alpine evergreen plants, resembling Club-Mosses or Heaths. Flowers solitary, nodding on slender erect peduncles, white or rose-color. (*Cassiope* was the mother of *Andromeda*.)

1. **C. hypnoides**, Don. Tufted and procumbent, moss-like (1' - 4' high); leaves needle-shaped, imbricated; corolla 5-cleft; style short and conical. (*Andromeda hypnoides*, L.) — Alpine summits of the Adirondack Mountains, New York (*Dr. Parry*), White Mountains of New Hampshire, Mount Katahdin, Maine (*Mr. Young*), and high northward. (Eu.)

10. ANDRÓMEDA, L. (in part). ANDROMEDA.

Calyx without bractlets, of 5 nearly or partly distinct sepals, valvate in the bud, but very soon separate or open. Corolla 5-toothed. Stamens 10: anthers fixed near the middle, the cells opening by a terminal pore. Pod globular, 5-celled, 5-valved; the many-seeded placentæ borne on the summit or middle of the columella. — Shrubs, with umbelled, clustered, or paniced and racemed (mostly white) flowers. (Fancifully named by Linnæus for *A. polifolia*, in allusion to the fable of *Andromeda*.)

§ 1. ANDRÓMEDA proper. *Corolla globular-urn-shaped: filaments bearded, not appendaged: anthers short, the cells each surmounted by a slender ascending awn: seeds turned in all directions, oval, with a close and hard smooth coat: flowers in a terminal umbel: pedicels from the axils of ovate persistent scaly bracts; leaves evergreen.*

1. **A. polifolia**, L. Smooth and glaucous (6' - 18' high; leaves thick, lanceolate or oblong-linear, with strongly revolute margins, white beneath. — Cold bogs, from Pennsylvania northward. May. (Eu.)

§ 2. PORTŪNA, Nutt. *Corolla ovoid-urn-shaped and 5-angled: filaments not appendaged: anthers oblong, the cells each bearing a long reflexed awn near the insertion: seeds mostly pendulous, and with a loose cellular coat: flowers in axillary and terminal naked racemes, formed in summer, but the blossoms expanding the following spring: pedicels 1-sided, bracted and with minute bractlets, recurved: leaves thick and evergreen.*

2. **A. floribunda**, Pursh. Branches bristly when young; leaves lance-oblong, acute or pointed (2' long), petioled, serrulate and bristly-ciliate; racemes dense, crowded in panicles. — Moist hills, in the Alleghanies from Virginia southward. April. — A very leafy shrub, 2° - 10° high, bearing abundance of handsome flowers.

§ 3. PIÉRIS, Don. *Corolla ovoid-oblong or cylindraceous: filaments slender and awl-shaped, usually appendaged with a spreading or recurved bristle on each side*

at or below the apex: anthers oblong, awnless: sutures of the 5-angular pod with a more or less thickened line or ridge, which often falls away separately when the pod opens: seeds turned in all directions, oblong, with a thin and rather loose reticulated coat: flowers in umbel-like clusters variously arranged.

3. **A. Mariàna**, L. (STAGGER-BUSH.) Nearly glabrous; leaves deciduous, but rather coriaceous, oval or oblong, veiny; flowers large and nodding, in clusters from axillary scaly buds, which are crowded on naked branches of the preceding year; sepals leaf-like, deciduous with the leaves. — Sandy low places, Rhode Island to Virginia near the coast, and southward. May, June. — Shrub 2°–4° high: foliage said to poison lambs and calves.

§ 4. **LYÒNIA**, Nutt. *Calyx 5-cleft: corolla globular, pubescent: filaments and anthers destitute of awns or appendages, or the former sometimes 2-setose near the apex: pods prominently ribbed at the sutures, the ribs at length separating or separable: seeds slender, all pendulous, with a loose and thin cellular coat: flowers small, mostly in clusters which are racemosed-panicled: bracts minute and deciduous: leaves pubescent or scurfy beneath.*

4. **A. ligustrina**, Muhl. Leaves deciduous, not scurfy, smoothish when old, obovate-oblong varying to oblong-lanceolate; flowers racemose-panicled on branchlets of the preceding year. — Swamps and low thickets, New England to Penn., Virginia, and southward. June, July. — Shrub 4°–10° high.

11. OXYDÉNDRUM, DC. SORREL-TREE. SOUR-WOOD.

Calyx without bractlets, of 5 almost distinct sepals, valvate in the bud. Corolla ovate, 5-toothed, puberulent. Stamens 10: anthers fixed near the base, linear, awnless; the cells tapering upwards, and opening by a long chink. Pod oblong-pyramidal, 5-celled, 5-valved; the many-seeded placenta at the base of the cells. Seeds *all ascending*, slender, the thin and loose reticulated coat extended at both ends into awl-shaped appendages. — A tree with deciduous, oblong-lanceolate, pointed, soon smooth, serrulate leaves, on slender petioles, and white flowers in long one-sided racemes clustered in an open panicle, terminating the branches of the season. Bracts and bractlets minute, deciduous. Foliage sour to the taste (whence the name, from *ὄξύς*, *sour*, and *δένδρον*, *tree*).

1. **O. arboreum**, DC. (*Andromeda arborea*, L.) — Rich woods, from Penn. and Ohio southward, mostly along the Alleghanies. June, July. — Tree 40°–60° high. Leaves in size and shape like those of the Peach.

12. CLÈTHRA, L. WHITE ALDER. SWEET PEPPERBUSH.

Calyx of 5 sepals, imbricated in the bud. Corolla of 5 distinct obovate-oblong petals. Stamens 10, often exerted: anthers inversely arrow-shaped, inverted and reflexed in the bud, opening by terminal pores or short slits. Style slender, 3-cleft at the apex. Pod 3-valved, 3-celled, many-seeded, enclosed in the calyx. Shrubs or trees, with alternate and serrate deciduous leaves, and white flowers in terminal hoary racemes. Bracts deciduous. (*Κλήθρα*, the ancient Greek name of the Alder, which this genus somewhat resembles in foliage.)

1. **C. alnifolia**, L. *Leaves wedge-obovate, sharply serrate, entire towards the base, prominently straight-veined, smooth, green both sides; racemes upright, paniced; bracts shorter than the flowers; filaments smooth.* — Wet copses, Maine to Virginia near the coast, and southward. — Shrub 3°–10° high, covered in July and August with handsome fragrant blossoms. — In the South are varieties with the leaves rather scabrous, and pubescent or white-downy beneath.

2. **C. acuminata**, Michx. *Leaves oval or oblong, pointed, thin, finely serrate (5'–7' long), pale beneath; racemes solitary, drooping; bracts longer than the flowers; filaments and pods hairy.* — Woods in the Alleghanies, Virginia and southward. July. — A tall shrub or small tree.

13. CALLUNA, Salisb. HEATHER.

Calyx of 4 colored sepals. Corolla bell-shaped, 4-parted, much shorter and less conspicuous than the calyx, both becoming scarious and persistent. Stamens 8, distinct: anthers with a pair of deflexed appendages on the back, the cells opening each by a long chink. Pod 4-celled, septicidally 4-valved. — Evergreen undershrub, with no scaly buds, opposite and minute leaves (mostly extended at base into 2 sharp auricles), crowded and imbricated on the branches. Flowers axillary, or terminating very short shoots and crowded on the branches, forming close mostly one-sided spikes or spike-like racemes, rose-colored or sometimes white, small, bracted by 2 or 3 pairs of leaves, the innermost of which are more or less scarious. (Named from *καλλύνω*, to brush or sweep, brooms being made of its twigs.)

1. **C. vulgaris**, Salisb. (*C. Atlantica*, Seemann, *Jour. Bot.* 4, p. 305, t. 53. *Erica vulgaris*, L.) — Low grounds, Tewksbury, Massachusetts (*Jackson Dawson, &c.*, a small patch); border of forest on Cape Elizabeth, Maine (*Mr. Pickard*, from *Dr. Wood*); also Nova Scotia, Cape Breton, and Newfoundland; mostly local. (See various articles in *Amer. Jour. Sci.*) July, Aug. (Eu.)

14. PHYLLODOCE, Salisb. PHYLLODOCE.

Corolla urn-shaped or bell-shaped, 5-toothed, deciduous. Stamens 10: anthers pointless, shorter than the filaments, opening by terminal pores. Pod 5-celled, 5-valved, septicidal (as are all the succeeding), many-seeded. — Low alpine Heath-like evergreen undershrubs, clothed with scattered linear and obtuse rough-margined leaves. Flowers usually nodding on solitary or umbelled peduncles at the summit of the branches. (“A mythological name.”)

1. **P. taxifolia**, Salisb. Corolla oblong-urn-shaped, purplish, smooth; style included. (*Menziesia cærulea*, *Smith.*) — Alpine summits of the mountains of New Hampshire and Maine, and northward. July. (Eu.)

15. KÁLMIA, L. AMERICAN LAUREL.

Calyx 5-parted. Corolla between wheel-shaped and bell-shaped, 5-lobed, furnished with 10 depressions in which the 10 anthers are severally lodged; filaments long and thread-form. Pod globose, 5-celled, many-seeded. — Evergreen mostly smooth shrubs, with alternate or opposite entire coriaceous leaves,

naked buds, and showy flowers. (Dedicated to *Peter Kalm*, a pupil of Linnæus who travelled in this country about the middle of the last century, afterwards Professor at Abo.)

§ 1. *Flowers in simple or clustered naked umbel-like corymbs: pedicels from the axils of small and firm foliaceous persistent bracts: calyx smaller than the pod, persistent: leaves glabrous.*

1. **K. latifolia**, L. (CALICO-BUSH. MOUNTAIN LAUREL. SPOONWOOD.) *Leaves mostly alternate, bright green both sides, ovate-lanceolate or elliptical, tapering to each end, petioled; corymbs terminal, many-flowered, clammy-pubescent; pod depressed, glandular.*— Rocky hills and damp soil, rather common from Maine to Ohio and Kentucky, as a shrub 4°–8° high; but in the mountains from Penn. southward forming dense thickets, and often tree-like (10°–20° high). May, June.—Flowers profuse, large and very showy, varying from deep rose-color to nearly white, clammy.

2. **K. angustifolia**, L. (SHEEP LAUREL. LAMBKILL.) *Leaves commonly opposite or in threes, pale or whitish underneath, light green above, narrowly oblong, obtuse, petioled; corymbs lateral (appearing later than the shoots of the season), slightly glandular, many-flowered; pod depressed, nearly smooth; pedicels recurved in fruit.*—Hillsides: common. May, June.—Shrub 2°–3° high: the flowers more crimson and two thirds smaller than in the last.

3. **K. glauca**, Ait. (PALE LAUREL.) *Branchlets 2-edged: leaves opposite, nearly sessile, oblong, white-glaucous beneath, with revolute margins; corymbs terminal, few-flowered, smooth; bracts large; pod ovoid, smooth.*—Var. **ROSMARINIFOLIA** has linear and strongly revolute leaves.—Cold peat-bogs and mountains, from Pennsylvania northward. May, June.—Straggling, about 1° high. Flowers $\frac{1}{2}$ broad, lilac-purple.

§ 2. *Flowers scattered, solitary in the axils of the leaves of the season: calyx leafy, larger than the pod, nearly equalling the corolla, at length deciduous: leaves (alternate and opposite) and branches bristly-hairy.*

4. **K. hirsuta**, Walt. *Branches terete; leaves oblong or lanceolate (4'' long), becoming glabrous.*—Sandy pine-barren swamps, E. Virginia and southward. May–Sept.—Shrub 1° high. Corolla rose-color.

16. MENZIESIA, Smith. MENZIESIA.

Calyx very small and flattish, 4-toothed or 4-lobed. Corolla cylindraceous-urn-shaped and soon bell-shaped, obtusely 4-lobed. Stamens 8, included: anther-cells opening at the top by an oblique pore. Pod ovoid, woody, 4-celled, 4-valved, many-seeded. Seeds narrow, with a loose coat.—A low shrub; the straggling branches and the oblong-obovate alternate deciduous leaves (like those of *Azalea*) hairy and ciliate with rusty rather chaff-like bristles. Flowers small, developed with the leaves, in terminal clusters from scaly buds, greenish-white and purplish, nodding. (Named for *Archibald Menzies*, who in Vancouver's voyage brought the species from the Northwest Coast.)

1. **M. ferruginea**, Smith, var. **globularis**. Corolla rather shorter and broader than in the Oregon plant.—Alleghany Mountains, S. Pennsylvania to Virginia, &c. (Also beyond Lake Superior.) June.

17. AZALEA, L. FALSE HONEYSUCKLE. AZALEA.

Calyx 5-parted, often minute. Corolla funnel-form, 5-lobed, slightly irregular; the lobes spreading. Stamens 5, with long exserted filaments, usually declined, as well as the similar style: anthers short, opening by terminal pores, pointless. Pod 5-celled, 5-valved, many-seeded. Seeds scale-like. — Upright shrubs, with alternate and obovate or oblong deciduous leaves, which are entire, ciliate, and mucronate with a glandular point. Flowers large and showy, often glandular and glutinous outside, in umbelled clusters from large scaly-imbriated terminal buds. (Name from ἀζαλέος, *arid*, — most inappropriate as applied to our species, which grow in swamps.)

* *Flowers appearing after the leaves.*

1. **A. arboréscens**, Pursh. (SMOOTH AZALEA.) *Branchlets smooth; leaves obovate, obtuse, very smooth both sides, shining above, glaucous beneath, the margins bristly-ciliate; calyx-lobes long and conspicuous; corolla slightly clammy; stamens and style very much exserted.* — Mountains of Penn. to Virginia, and southward. June. — Shrub 3°–10° high, with thickish leaves, and very fragrant rose-colored blossoms larger than in No. 3.

2. **A. viscosa**, L. (CLAMMY A. WHITE SWAMP-HONEYSUCKLE.) *Branchlets bristly, as well as the margins and midrib of the oblong-obovate otherwise smooth leaves; calyx-lobes minute; corolla clammy, the tube much longer than the lobes; stamens and especially the style exserted.* — Var. *GLAUCA* has the leaves paler, often white-glaucous underneath or both sides, sometimes rough-hairy. — Var. *NŪTIDA* is dwarf, with oblanceolate leaves green both sides. — Swamps, Maine to Kentucky, mostly near the coast. June, July. — Shrub 4°–10° high, with clammy fragrant flowers, white or tinged with rose-color.

** *Flowers appearing before or with the leaves.*

3. **A. nudiflora**, L. (PURPLE A. PINKSTER-FLOWER.) *Branchlets rather hairy; leaves obovate or oblong, downy underneath; calyx very short; tube of the corolla scarcely longer than the ample lobes, slightly glandular; stamens and style much exserted.* — Swamps, Massachusetts and New York to Virginia, and southward. April, May. — Shrub 2°–6° high; the showy flowers varying from flesh-color to pink and purple. There are numberless varieties, some of them with 10 stamens.

4. **A. calendulæcea**, Michx. (FLAME-COLORED AZALEA.) *Branchlets and obovate or oblong leaves hairy; calyx-lobes oblong, rather conspicuous; tube of the corolla shorter than the lobes, hairy; stamens and style much exserted.* — Woods, mountains of Penn. to Virginia, Kentucky, and southward. May. — Shrub 3°–10° high, covered just when the leaves appear with a profusion of large orange blossoms, usually turning to flame-color, not fragrant.

18. RHODODÉNDRON, L. ROSE-BAY.

Calyx 5-parted, minute in our species. Corolla bell-shaped or partly funnel-form, sometimes slightly irregular, 5-lobed. Stamens 10 (rarely fewer), commonly declined: anthers, pods, &c. as in Azalea. — Shrubs or low trees, with evergreen entire alternate leaves, and ample showy flowers, in compact terminal

corymbs or clusters, from large scaly-bracted buds. (*Ῥοδόδενδρον*, *rose-tree*; the ancient name.)

1. **R. máximum**, L. (GREAT LAUREL.) *Leaves elliptical-oblong or lance-oblong, acute, narrowed towards the base, very smooth, with somewhat revolute margins; pedicels viscid; corolla bell-shaped.*—Damp deep woods, sparingly from Maine to Ohio, but very common along shaded water-courses through the mountainous parts of Pennsylvania and southward. July. — Shrub or tree 6°–20° high. Leaves 4'–10' long, very thick. Corolla an inch broad, pale rose-color or nearly white, greenish in the throat on the upper side, and spotted with yellow or reddish.

2. **R. Catawbiense**, Michx. *Leaves oval or oblong, rounded at both ends, smooth, pale beneath (3'–5' long); corolla broadly bell-shaped, lilac-purple; pedicels rusty-downy.*—High Alleghanies, Virginia and southward. June. — Shrub 3°–6° high.

3. **R. Lappónicum**, Wahl. (LAPLAND ROSE-BAY.) *Dwarf, prostrate; leaves elliptical, obtuse, dotted (like the branches) with rusty scales; umbels few-flowered; corolla open bell-shaped, dotted; stamens 5–10.*—Alpine summits of the mountains of Maine, New Hampshire, and New York. July. — Shrub 6' high, in broad tufts: leaves $\frac{1}{2}$ ' long. Corolla violet-purple. (Eu.)

19. RHODÒRA, Duhamel. RHODORA.

Calyx minute, 5-toothed. Corolla irregular and 2-lipped; the upper lip usually 3-lobed or 3-cleft, and the lower two-parted or of 2 distinct spreading petals. Stamens 10, and with the slender style declined. Otherwise as in *Azalea*. (Name from *Ῥόδον*, *a rose*, from the color of the showy flowers.)

1. **R. Canadensis**, L. —Damp cold woods and swamps, New England to Penn. and northward, or on mountains. May. —A handsome low shrub, with the oblong deciduous leaves whitish and downy underneath; the showy rose-purple (rarely white) flowers in umbel-like clusters, on short peduncles, appearing rather earlier than the leaves.

20. LÈDUM, L. LABRADOR TEA.

Calyx 5-toothed, very small. Corolla of 5 obovate and spreading distinct petals. Stamens 5–10: anthers opening by terminal pores. Pod 5-celled, splitting from the base upwards, many-seeded: placenta borne on the summit of the columella. —Low shrubs, with the alternate entire leaves clothed with rusty wool underneath, persistent, the margins revolute: herbage slightly fragrant when bruised. Flowers white, small, in terminal umbel-like clusters from large scaly buds: bracts or scales thin and caducous. (*Λήδον*, the ancient Greek name of the *Cistus*.)

1. **L. latifolium**, Ait. Leaves elliptical or oblong; stamens 5, sometimes 6 or 7; pod oblong. —Cold bogs and damp mountain woods, New England to Pennsylvania, Wisconsin, and northward. June. —Shrub 2°–5° high. (*L. PALÚSTRE*, L., grows in British America. It is distinguished by its linear leaves, uniformly 10 stamens, and *oval pods*.) (Eu.)

21. LOISELEÛRIA, Desv. ALPINE AZALEA.

Calyx 5-parted, nearly as long as the rather bell-shaped and deeply 5-cleft regular corolla. Stamens 5, not declined, included: anthers opening lengthwise. Style short. Pod ovoid, 2-3-celled, many-seeded, 2-3-valved; the valves 2-cleft from the apex: placenta borne on the middle of the columella. — A small depressed evergreen shrubby plant, much branched and tufted, smooth, with coriaceous opposite elliptical leaves, on short petioles, with revolute margins. Flowers small, white or rose-color, 2-5 in a cluster, from a terminal scaly bud; the scales or bracts thick and persistent. (Named for *Loiseleur Delongchamps*, a French botanist.)

1. *L. procumbens*, Desv. (*Azalea procumbens*, L.) — Alpine summits of the White Mountains, New Hampshire, on rocks. June. (Eu.)

22. LEIOPHYLLUM, Pers. SAND MYRTLE.

Calyx 5-parted. Corolla of 5 distinct obovate-oblong petals, spreading. Stamens 10, exerted: anthers opening lengthwise. Pod 2-3-celled, splitting from the apex downward, many-seeded. — A low much-branched evergreen, with the aspect, foliage, &c. of the preceding genus, but the crowded leaves sometimes alternate, scarcely petioled. Flowers small, white, in terminal umbel-like clusters. (Name formed of *λείος*, smooth, and *φύλλον*, foliage, from the leaves.)

1. *L. buxifolium*, Ell. — Sandy pine barrens of New Jersey, and mountain-tops in Virginia? and southward. May. — Shrub 6' - 10' high: leaves oval or oblong, smooth and shining, 3" - 6" long.

23. PYROLA, Tourn. WINTERGREEN. SHIN-LEAF.

Calyx 5-parted, persistent. Petals 5, concave and more or less converging, deciduous. Stamens 10: filaments awl-shaped, naked: anthers extrorse in the bud, but in the flower inverted by the inflexion of the apex of the filament, more or less 4-celled, opening by a pair of pores at the blunt or somewhat 2-horned base which by the inversion becomes the apparent apex! Style generally long: stigma 5-lobed or 5-rayed. Pod depressed-globose, 5-lobed, 5-celled, 5-valved from the base upwards (loculicidal); the valves cobwebby on the edges. Seeds minute, innumerable, resembling saw-dust, with a very loose cellular-reticulated coat. — Low and smooth perennial herbs, with running subterranean shoots, bearing a cluster of rounded and petioled evergreen root-leaves, and a simple raceme of nodding flowers, on an upright more or less scaly-bracted scape. (Name a diminutive of *Pyrus*, the Pear-tree, from some fancied resemblance in the foliage, which is not obvious.)

§ 1. *Stamens ascending: style turned down and towards the apex usually more or less curved upwards, longer than the campanulate-connivent or somewhat expanding petals: stigma much narrower than the truncate and somewhat excavated apex of the style which forms a sort of ring or collar, the 5 lobes at first very short and included, at length usually protruded.* (Leaves denticulate or entire.)

1. *P. rotundifolia*, L. Leaves orbicular, thick, shining, usually shorter than the petiole; raceme elongated, many-flowered; calyx-lobes lanceolate or oblong-

lanceolate, acutish, with somewhat spreading tips, *one half or one third the length of the roundish-obovate rather spreading (chiefly white) petals; anther-cells nearly blunt.* — Damp or sandy woods: common, especially northward. June, July. — Scape 6'–12' high, many-bracted: flowers $\frac{3}{4}$ ' broad. — Exhibits many varieties, such as: Var. *INCARNATA*, with flesh-colored flowers; calyx-lobes triangular-lanceolate. — Var. *ASARIFOLIA*, with oblate or round-reniform leaves, and triangular-ovate calyx-lobes of about half the length of the white or flesh-colored petals. (*P. asarifolia*, *Michx.*) Common northward. — Var. *ULIGINOSA*, with roundish-oval or somewhat kidney-shaped smaller leaves (1'–1½' wide), and ovate acute calyx-lobes, about one quarter of the length of the reddish or purple petals; flowers rather smaller, few or several. (*P. uliginosa*, *Torr. & Gr.*) Cold bogs, N. New England to Wisconsin, and northward. (Eu.)

2. **P. elliptica**, Nutt. (SHIN-LEAF.) *Leaves thin and dull, elliptical or obovate-oval, usually longer than the margined petiole; raceme many-flowered; calyx-lobes ovate, acute, not one fourth the length of the obovate rather spreading (greenish-white) petals; anther-cells blunt.* — Rich woods, New England to Pennsylvania, Wisconsin, and northward: common. June. — Scape and flowers nearly as large as in the preceding.

3. **P. chlorantha**, Swartz. *Leaves small (1' long), roundish, thick, dull, shorter than the petiole; scape few-flowered, naked (5'–8' high); calyx-lobes roundish-ovate, very short; the elliptical petals converging (greenish-white); anther-cells contracted below the orifice into a distinct neck or horn: style little exerted.* — Open woods, New England to Penn., Wisconsin, and northward. June, July.

4. **P. oxypétala**, C. F. Austin, n. sp. *Leaves ovate, small (8''–12'' long), shorter than the slender petiole; scape (7'–8' high) several-flowered; flowers on ascending pedicels, not nodding; calyx-lobes triangular-ovate, acute, short; petals lanceolate-oblong, acuminate, greenish; anthers conspicuously mucronate at the apex, obtusely 2-horned (as in No. 3) at the base, not inverted; style straightish, scarcely exerted.* — Wooded hill near Deposit, Delaware Co., New York, June 1, 1860, *C. F. Austin*. — Stigma as in No. 3; the calyx-lobes, &c., different, so that it can hardly be a monstrosity of that species.

§ 2. *Stamens regular: style straight, much narrower than the expanded depressed 5-rayed stigma: petals erect and connivent.*

5. **P. secúnda**, L. Subcaulescent; *leaves ovate, thin, longer than the petiole, scattered, finely serrate; racemes dense and spike-like, the numerous small (greenish-white) flowers all turned to one side, scarcely nodding; calyx-lobes ovate, very much shorter than the oblong oval petals; style long, exerted.* — Rich woods: common eastward and northward. July. — Scape 3'–6' high. (Eu.)

Var. **pùmila** (Paine, *Cat. Pl. Oneida Co.*) is a remarkable form, growing in high peat-bogs from Otsego Co., New York, Lake Superior, and northward (indicated by Chamisso in Russian America); *leaves orbicular or broadly oval, very thin (5''–12' long); scape 2'–4' high, 3–8-flowered.* July, Aug.

6. **P. minor**, L. *Leaves roundish, slightly crenulate, thickish, mostly longer than the margined petiole; raceme spiked; calyx-lobes triangular-ovate, very much shorter than the nearly globose corolla; style short and included.* — Woods at the base of the White Mountains, New Hampshire, Lake Superior (*L. Foots*),

and northward. July, Aug. — Scape 5'–10' high. Flowers small, crowded, white or rose-color. (Eu.)

24. MONÈSES, Salisb. ONE-FLOWERED PYROLA.

Petals 5, widely spreading, orbicular. Filaments awl-shaped, naked: anthers as in *Pyrola*, but conspicuously 2-horned. Style straight, exserted: stigma large, peltate, with 5 narrow and conspicuous radiating lobes. Valves of the pod naked, as in the next genus. (Parts of the flower occasionally in fours.) Scape 1-flowered. Otherwise as in *Pyrola*: intermediate between it and *Chimaphila*. (Name formed of *μόνος*, *single*, and *ἡσυχία*, *delight*, from the pretty and solitary flower.)

1. *M. uniflora*. (*Pyrola uniflora*, L.) — Deep cold woods, from Penn. and New England northward. June. — A small perennial, with the rounded and veiny serrate thin leaves (6''–9'' long), clustered at the ascending apex of creeping subterranean shoots; the 1–2-bracted scape (2'–4' high) bearing a white or rose-colored terminal flower 6'' wide. (Eu.)

25. CHIMÁPHILA, Pursh. PIPSISSEWA.

Petals 5, concave, orbicular, widely spreading. Stamens 10: filaments enlarged and hairy in the middle: anthers as in *Pyrola*, but more or less conspicuously 2-horned. Style very short, inversely conical, nearly immersed in the depressed summit of the globular ovary: stigma broad and orbicular, disk-shaped, the border 5-crenate. Pod, &c. as in *Pyrola*, but splitting from the apex downwards, the edges of the valves not woolly. — Low, nearly herbaceous plants, with long running underground shoots, and evergreen thick and shining leaves, somewhat whorled or scattered along the short ascending stems: the fragrant (white or purplish) flowers corymbed or umbelled on a terminal peduncle. (Name from *χειμα*, *winter*, and *φιλέω*, *to love*, in allusion to one of the popular names, viz. *Wintergreen*.)

1. *C. umbellata*, Nutt. (PRINCE'S PINE. PIPSISSEWA.) *Leaves wedge-lanceolate, acute at the base, sharply serrate, not spotted; peduncles 4–7-flowered.* — Dry woods: common. June. — Plant 4'–10' high, leafy: petals flesh-color: anthers violet. (Eu.)

2. *C. maculata*, Pursh. (SPOTTED WINTERGREEN.) *Leaves ovate-lanceolate, obtuse at the base, remotely toothed, the upper surface variegated with white; peduncles 1–5-flowered.* — Dry woods: most common in the Middle States. June, July. — Plant 3'–6' high.

26. PTERÓSPORA, Nutt. PINE-DROPS.

Calyx 5-parted. Corolla ovate, urn-shaped, 5-toothed, persistent. Stamens 10: anthers 2-celled, awned on the back, opening lengthwise. Style short: stigma 5-lobed. Pod globose, depressed, 5-lobed, 5-celled, loculicidal, but the valves cohering with the columella. Seeds very numerous, ovoid, tapering to each end, the apex expanded into a broad reticulated wing many times larger than the body of the seed. — A stout and simple purplish-brown clammy-pubes-

cent root-parasitic herb (1°-2° high); the wand-like stem furnished towards the base with scattered lanceolate scales in place of leaves, above bearing many nodding (white) flowers, resembling those of *Andromeda*, in a long bracted raceme. (Name from *πτερόν*, a wing, and *σπορά*, seed, alluding to the singular wing borne by the seeds.)

1. *P. Andromedæa*, Nutt. — Hard clay soil, parasitic apparently on the roots of pines, from Vermont, Peekskill and Albany, N. Y., and N. Pennsylvania northward and westward: rare.

27. SCHWEINÍTZIA, Ell. SWEET PINE-SAP.

Calyx of 5 oblong-lanceolate acute scale-like sepals, erect, persistent. Corolla persistent, bell-shaped, rather fleshy, 5-lobed, slightly 5-gibbous at the base. Stamens 10: anthers much shorter than the filaments, fixed near the summit, awnless; the two sac-shaped cells opening at the top. Pod ovoid. 5-celled, with a short and thick style, and a large 5-angular stigma. Seeds innumerable. — A low and smooth brownish plant, 3'-4' high, with the aspect of *Monotropa*, scaly-bracted, the flowers several in a terminal spike, at first nodding, flesh-color, exhaling the fragrance of violets. (Named for the late *L. D. von Schweinitz*.)

1. *S. odorata*, Ell. — Woods, parasitic on the roots of herbs, Maryland and southward: rare. April.

28. MONÓTROPÁ, L. INDIAN PIPE. PINE-SAP.

Calyx of 2-5 lanceolate bract-like scales, deciduous. Corolla of 4 or 5 separate erect spatulate or wedge-shaped scale-like petals, which are gibbous or saccate at the base, and tardily deciduous. Stamens 8 or 10: filaments awl-shaped: anthers kidney-shaped, becoming 1-celled, opening across the top. Style columnar: stigma disk-like, 4-5-rayed. Pod ovoid, 8-10-grooved, 4-5-celled, loculicidal: the very thick placentæ covered with innumerable minute seeds, which have a very loose coat. — Low and fleshy herbs, tawny, reddish, or white, parasitic on roots, or growing on decomposing vegetable matter like a Fungus; the clustered stems springing from a ball of matted fibrous rootlets, furnished with scales or bracts in place of leaves, 1-several-flowered; the flowering summit at first nodding, in fruit erect. (Name composed of *μόνος*, one, and *τρόπος*, turn, from the summit of the stem turned to one side.)

§ 1. *MONOTROPA*, Nutt. *Plant inodorous, with a single 5-petalled and 10-androus flower at the summit; the calyx of 2-4 irregular scales or bracts: anthers transverse, opening by 2 chinks: style short and thick: stigma naked.*

1. *M. uniflora*, L. (INDIAN PIPE. CORPSE-PLANT.) Smooth, waxy-white (turning blackish in drying, 3'-8' high); stigma naked. — Dark and rich woods: common. June-Aug.

§ 2. *HYPÓPITYS*, Dill. *Plant commonly fragrant: flowers several in a scaly raceme; the terminal one usually 5-petalled and 10-androus, while the rest are 4-petalled and 8-androus; the bract-like sepals mostly as many as the petals: anthers opening by a continuous line into 2 very unequal valves, the smaller one erect and appearing like a continuation of the filament: style longer than the ovary, hollow.*

2. **M. Hypópitys**, L. (PINE-SAP. FALSE BEECH-DROPS.) Somewhat pubescent or downy, tawny, whitish, or reddish (4'–12' high); pod globular or oval; stigma ciliate. — Occurs in various forms: the more pubescent is *M. lanuginosa*, *Michx.* — Oak and pine woods: common. June–Aug. (Eu.)

ORDER 57. GALACÍNEÆ. (GALAX FAMILY.)

Character that of the following genus; which is kept as a distinct order until the true relationship is ascertained.

1. GÀLAX, L. GALAX.

Calyx of 5 small and separate sepals, persistent. Petals 5, hypogynous, obovate-spatulate, rather erect, deciduous. Stamens hypogynous: filaments united in a 10-toothed tube, slightly cohering with the base of the petals, the 5 teeth opposite the petals naked, the 5 alternate ones shorter and bearing each a roundish 1-celled anther, which opens across the top. Pollen simple. Style short: stigma 3-lobed. Pod ovoid, 3-celled, loculicidally 3-valved: columella none. Seeds numerous, the cellular loose coat tapering to each end. Embryo straight in fleshy albumen, more than half its length. — A smooth herb, with a thick matted tuft of scaly creeping rootstocks, beset with fibrous red roots, sending up round-heart-shaped crenate-toothed and veiny shining leaves (about 2' wide) on slender petioles, and a slender naked scape, 1°–2° high, bearing a wand-like spike or raceme of small and minutely-bracted white flowers. (Name from *γάλα*, *milk*, — of no conceivable application to this plant.)

1. **G. aphýlla**, L. — Open woods, Virginia and southward. June.

ORDER 58. AQUIFOLIACÆ. (HOLLY FAMILY.)

Trees or shrubs, with small axillary 4–8-merous flowers, a minute calyx free from the 4–8-celled ovary and the 4–8-seeded berry-like drupe; the stamens as many as the divisions of the almost or quite 4–8-petalled corolla and alternate with them, attached to their very base. — Corolla imbricated in the bud. Anthers opening lengthwise. Stigmas 4–8, or united into one, nearly sessile. Seeds suspended and solitary in each cell, anatropous, with a minute embryo in fleshy albumen. Leaves simple, mostly alternate. Flowers white or greenish. — A small family, here represented by only two genera (*Prinos* being placed under *Ilex*); related to the Celastraceæ, &c., among the Polypetalous orders as much as to the following order.

1. ÌLEX, L. (*Ilex* & *Prinos*, L.) HOLLY.

Flowers more or less dioeciously polygamous. Calyx 4–6-toothed. Petals 4–6, separate, or united only at the base, oval or obovate, obtuse, spreading. Stamens 4–6. The berry-like drupe containing 4–6 little nutlets. — Leaves alternate. Fertile flowers inclined to be solitary, and the sterile or partly sterile flowers to be clustered in the axils. (The ancient Latin name of the Holly-Oak rather than of the Holly.)

§ 1. **AQUIFOLIUM**, Tourn. *Parts of the flower commonly in fours, sometimes in fives or sixes: drupe red, its nutlets ribbed, veiny, or one-grooved on the back: leaves (mostly smooth) coriaceous and evergreen.*

* *Leaves armed with spiny teeth: trees.*

1. **I. opaca**, Ait. (AMERICAN HOLLY.) *Leaves oval, flat, the wavy margins with scattered spiny teeth; flowers in loose clusters along the base of the young branches and in the axils; calyx-teeth acute.*—Moist woodlands, Maine to Pennsylvania, near the coast, and more common from Virginia southward. June.—Tree 20°–40° high; the deep green foliage less glossy, the berries not so bright red, and their nutlets not so veiny, as in the European Holly (*I. AQUIFOLIUM*, L.).

** *Leaves serrate or entire, not spiny: shrubs.*

2. **I. Cassine**, L. (CASSENA. YAUPON.) *Leaves lance-ovate or elliptical, crenate (1'–1½' long); flower-clusters nearly sessile, smooth; calyx-teeth obtuse.*—Virginia and southward along the coast. May.—Leaves used for tea by the people along the coast, as they were to make the celebrated *black drink* of the North Carolina Indians.

3. **I. myrtifolia**, Walt. *Leaves linear-lanceolate or linear-oblong, sparingly and sharply serrate or entire (1' long); peduncles slender and 3–9-flowered, or the more fertile shorter and 1-flowered, smooth; calyx-teeth acute.*—Coast of Virginia and southward. May.—Probably a var. or the next.

4. **I. Dahoon**, Walt. (DAHOON HOLLY.) *Leaves oblanceolate or oblong, entire, or sharply serrate towards the apex, with revolute margins (2'–3' long), the midrib and peduncles pubescent; calyx-teeth acute.*—Swamps, coast of Virginia and southward. May, June.

§ 2. **PRINOIDES**. *Parts of the (polygamous or dioecious) flowers in fours or fives (rarely in sixes): drupe red or purple, the nutlets striate-many-ribbed on the back: leaves deciduous: shrubs.*

5. **I. decidua**, Walt. *Leaves wedge-oblong or lance-obovate, obtusely serrate, downy on the midrib beneath, shining above, becoming thickish; peduncles of the sterile flowers longer than the petioles, of the fertile short; calyx-teeth smooth, acute.*—Wet grounds, Virginia, Illinois, and southward. May.

6. **I. monticola**, Gray. *Leaves ovate or lance-oblong, ample (3'–5' long), taper-pointed, thin-membranaceous, smooth, sharply serrate; fertile flowers very short-peduncled; calyx ciliate.* (*I. ambigua*, Torr. *I. montana*, Ed. 1, not *Prinos montanus*, Sw.)—Damp woods, Taconic and Catskill Mountains, and Cattaraugus Co., New York (*G. W. Clinton*), through Pennsylvania (east to Northampton Co. *Mr. Wolle*, *Prof. T. Green*), and southward along the Alleghanies. May.

7. **I. mollis**, Gray. *Leaves soft downy beneath, oval, ovate, or oblong, taper-pointed at both ends, especially at the apex, thin-membranaceous, sharply serrulate; sterile flowers very numerous in umbel-like clusters, the pedicels shorter than the petiole and (with the calyx) soft-downy, the fertile peduncles very short.* (*Prinos pubescens*, Michx. herb. *P. ambiguus*, Pursh, not Michx.)—Burgoon's Gap, Alleghanies of Pennsylvania (*J. R. Lowrie*, in fruit), and along the mountains in the Southern States.—Resembles the last.

§ 3. **PRINOS**, L. *Parts of the sterile flowers in fours, fives, or sixes, those of the fertile flowers commonly in sixes (rarely in fives, sevens, or eights) : nutlets smooth and even : shrubs.*

* *Leaves deciduous : flowers in sessile clusters, or the fertile solitary : fruit bright red.*

8. **I. verticillata**, Gray. (BLACK ALDER. WINTERBERRY.) *Leaves obovate, oval, or wedge-lanceolate, pointed, acute at the base, serrate, downy on the veins beneath ; flowers all very short-peduncled. (Prinos verticillatus, L.) — Low grounds : common. May, June.*

9. **I. lævigata**, Gray. (SMOOTH WINTERBERRY.) *Leaves lanceolate or oblong-lanceolate, pointed at both ends, appressed-serrulate, shining above, beneath mostly glabrous ; sterile flowers long-peduncled. (Prinos lævigatus, Pursh.) — Wet grounds, Maine to the mountains of Virginia. June. — Fruit larger than in the last, ripening earlier in the autumn.*

** *Leaves coriaceous, evergreen, shining above, often black-dotted beneath : fruit black.*

10. **I. glabra**, Gray. (INKBERRY.) *Leaves wedge-lanceolate or oblong, sparingly toothed towards the apex, smooth ; peduncles ($\frac{1}{2}$ long) of the sterile flowers 3-6-flowered, of the fertile 1-flowered ; calyx-teeth rather blunt. (Prinos glaber, L.) — Sandy grounds, Cape Ann, Massachusetts, to Virginia and southward near the coast. June. — Shrub 2°-3° high.*

2. NEMOPÁNTHES, Raf. MOUNTAIN HOLLY.

Flowers polygamo-dicecious. Calyx in the sterile flowers of 4-5 minute deciduous teeth ; in the fertile ones obsolete. Petals 4-5, oblong-linear, spreading, distinct. Stamens 4-5 : filaments slender. Drupe with 4-5 bony nutlets, light red. — A much-branched shrub, with ash-gray bark, alternate and oblong deciduous leaves on slender petioles, entire, or slightly toothed, smooth. Flowers on long and slender axillary peduncles, solitary, or sparingly clustered. (Name said by the author to mean "flower with a filiform peduncle," therefore probably composed of *νήμα*, a thread, *πούς*, *q. foot*, and *ἄθος*, a flower.)

1. **N. Canadensis**, DC. (*Ilex Canadensis*, Michx.) — Damp cold woods, from the mountains of Virginia to Maine, Wisconsin, and northward : common at the north. May.

ORDER 59. EBENACEÆ. (EBONY FAMILY.)

Trees or shrubs, with alternate entire leaves, and polygamous regular flowers which have a calyx free from the 3-12-celled ovary ; the stamens 2-4 times as many as the lobes of the corolla, often in pairs before them, their anthers turned inwards, and the fruit a several-celled berry. Ovules 1 or 2, suspended from the summit of each cell. Seeds anatropous, mostly single in each cell, large and flat, with a smooth coriaceous integument : the embryo shorter than the hard albumen, with a long radicle and flat cotyledons. Styles wholly or partly separate. — Wood hard and dark-colored. No milky juice. — A small family, chiefly tropical, represented here only by the Persimmon.

1. **DIOSPÝROS**, L. DATE-PLUM. PERSIMMON.

Calyx 4-6-lobed. Corolla 4-6-lobed, convolute in the bud. Stamens commonly 16 in the sterile flowers, and 8 in the fertile, in the latter imperfect. Berry large, globular, surrounded at the base by the thickish calyx, 4-8-celled, 4-8-seeded. — Flowers diœciously polygamous, the fertile axillary and solitary, the sterile smaller and often clustered. (Name, Διός, of Jove, and πύρος, grain.)

1. **D. Virginiana**, L. (COMMON PERSIMMON.) Leaves ovate-oblong, smooth or nearly so; peduncles very short; calyx 4-parted; corolla between bell-shaped and urn-shaped; styles 4, two-lobed at the apex; ovary 8-celled. — Woods and old fields, Rhode Island and New York to Illinois, and southward. June. — Tree 20°-60° high, with very hard blackish wood, thickish leaves, a pale yellow corolla, and a plum-like fruit, 1' in diameter, which is exceedingly astringent when green, yellow when ripe, and sweet and edible after exposure to frost.

ORDER 60. **SAPOTACEÆ**. (SAPPODILLA FAMILY.)

Trees or shrubs, mostly with a milky juice, simple and entire alternate leaves (often rusty-downy beneath), small and perfect regular flowers usually in axillary clusters; the calyx free and persistent; the fertile stamens commonly as many as the lobes of the hypogynous short corolla and opposite them, inserted on its tube, along with one or more rows of appendages and scales, or sterile stamens; anthers turned outwards; ovary 4-12-celled, with a single anatropous ovule in each cell; seeds large. — Albumen mostly none; but the large embryo with thickened cotyledons. Style single, pointed. — A small, mostly tropical order, producing the Sappodilla or Star-apple, and some other edible fruits, represented in our district only by the genus

1. **BUMELIA**, Swartz. BUMELIA.

Calyx 5-parted. Corolla 5-cleft, with a pair of internal appendages at each sinus. Fertile stamens 5: anthers arrow-shaped. Sterile stamens 5, petal-like, alternate with the lobes of the corolla. Ovary 5-celled. Fruit small, resembling a cherry, black, containing a large ovoid and erect seed, with a roundish scar at its base. — Flowers small, white, in fascicles from the axil of the leaves. Branches often spiny. Leaves often fascicled on short spurs. Wood very hard. (The ancient name of a kind of Ash.)

1. **B. lycioides**, Gærtn. (SOUTHERN BUCKTHORN.) Spiny (10°-25° high); leaves wedge-oblong varying to oval-lanceolate, with a tapering base, often acute, reticulated, nearly glabrous (2'-4' long); clusters densely many-flowered; fruit ovoid. — Moist ground, Illinois, Kentucky, and southward. May, June.

2. **B. lanuginosa**, Pers. Spiny (10°-40° high); leaves oblong-obovate or wedge-obovate, rusty-woolly beneath, obtuse (1½'-3' long); clusters 6-12-flowered; fruit globular. (*B. lanuginosa* & *tomentosa*, A. DC.) — Woods, Illinois, opposite St. Louis, and southward, — a variety with the leaves less woolly and rusty beneath (*B. oblongifolia*, Nutt.), passing towards No. 1. July.

ORDER 61. **STYRACACEÆ.** (STORAX FAMILY.)

Shrubs or trees, with alternate simple leaves destitute of stipules, and perfect regular flowers; the calyx either free or adherent to the 2-5-celled ovary; the corolla of 4-8 petals, commonly more or less united at the base; the stamens twice as many as the petals or more numerous, monadelphous or polyadelphous at the base; style 1; fruit dry or drupe-like, 1-5-celled, the cells commonly 1-seeded.—Seeds anatropous. Embryo nearly the length of the albumen: radicle slender, as long as or longer than the flat cotyledons. Corolla hypogynous when the calyx is free: the stamens adherent to its base. Ovules 2 or more in each cell.—A small family, mostly of warm countries, comprising two very distinct tribes, which are sometimes separated as suborders or orders.

Tribe I. STYRACEÆ. Calyx 4-8-toothed or entire. Stamens 2-4 times as many as the petals: anthers linear or oblong, adnate, introrse. Ovules or part of them ascending, — Flowers white, handsome. Pubescence soft and stellate.

1. **Styrax.** Calyx coherent only with the base of the 3-celled ovary. Corolla mostly 5, parted. Fruit 1-celled, 1-seeded.
2. **Halesia.** Calyx coherent with the whole surface of the 2-4-celled ovary, which is 2-4, winged and 2-4-celled in fruit. Corolla 4-lobed.

Tribe II. SYMPLICINEÆ. Calyx 5-cleft. Stamens usually very numerous: anthers short, innate. Ovules pendulous.— Flowers yellow. Pubescence simple.

3. **Symplocos.** Calyx coherent. Petals 5, united merely at the base.

1. **STÏRAX,** Tourn. STORAX.

Calyx truncate, somewhat 5-toothed, the base (in our species) coherent with the base of the 3-celled many-ovuled ovary. Corolla 5-parted (rarely 4-8-parted), large; the lobes mostly soft-downy. Stamens twice as many as the lobes of the corolla: filaments flat, united at the base into a short tube: anthers linear, adnate. Fruit globular, its base surrounded by the persistent calyx, 1-celled, mostly 1-seeded, dry, often 3-valved. Seed globular, erect, with a hard coat.— Shrubs or small trees, with commonly deciduous leaves, and axillary or leafy-racemed white and showy flowers, on drooping peduncles; produced in spring. Pubescence scurfy or stellate. (ἡ Στῦραξ, the ancient Greek name of the tree which produces storax.)

1. **S. grandifolia,** Ait. Leaves obovate, acute or pointed, *white-tomentose beneath* (3'-6' long); *flowers mostly in elongated racemes*; corolla ($\frac{1}{2}$ ' long) convolute-imbriated in the bud.— Woods, Virginia and southward.

2. **S. pulverulenta,** Michx. Leaves oval or obovate (about 1' long), *above sparingly puberulent, and scurfy-tomentose beneath*; *flowers* ($\frac{1}{2}$ ' long) 1-3 together in the axils and at the tips of the branches, fragrant.— Low pine barrens, Virginia (*Pursh*) and southward.— Shrub 1°-4° high.

3. **S. Americana,** Lam. Leaves oblong, acute at both ends (1'-3' long), *smooth, or barely pulverulent beneath*; *flowers axillary or in 3-4-flowered racemes* ($\frac{1}{2}$ ' long); corolla valvate in the bud. (*S. glabrum* and *S. læve, Ell.*)— Margin of swamps, Virginia and southward.— Shrub 4°-8° high.

2. **HALËSIA**, Ellis. SNOWDROP OR SILVER-BELL-TREE.

Calyx inversely conical, 4-toothed; the tube 4-ribbed, coherent with the 2-4-celled ovary. Petals 4, united at the base, or oftener to the middle, into an open bell-shaped corolla, convolute or imbricated in the bud. Stamens 8-16: filaments united into a ring at the base, and usually a little coherent with the base of the corolla: anthers linear-oblong. Ovules 4 in each cell. Fruit large and dry, 2-4-winged, within bony and 1-4-celled. Seeds single, cylindrical.—Shrubs or small trees, with large and veiny pointed deciduous leaves, and showy white flowers, drooping on slender pedicels, in clusters or short racemes, from axillary buds of the preceding year. Pubescence partly stellate. (Named for *Stephen Hales*, author of *Vegetable Statics*, &c.)

1. **H. tetráptera**, L. Leaves oblong-ovate; fruit 4-winged. — Banks of streams, upper part of Virginia, also on the Ohio River at Evansville (*Short*), and southward. Fruit $1\frac{1}{2}$ ' long.

3. **SÝMPLOCOS**, Jacq. (HÓPEA, L.) SWEET-LEAF.

Calyx 5-cleft, the tube coherent with the lower part of the 3-celled ovary. Petals 5, imbricated in the bud, lightly united at the base. Stamens very numerous, in 5 clusters, one cohering with the base of each petal: filaments slender: anthers very short. Fruit drupe-like or dry, mostly 1-celled and 1-seeded. — Shrubs or small trees, the leaves commonly turning yellowish in drying, and furnishing a yellow dye. Flowers in axillary clusters or racemes, yellow. (Name *σύμπλοκος*, *connected*, from the union of the stamens.)

1. **S. tinctoria**, L'Her. (HORSE-SUGAR, &c.) Leaves elongated-oblong, acute, obscurely toothed, thickish, almost persistent, minutely pubescent and pale beneath (3'-5' long); flowers 6-14, in close and bracted clusters, odorous. — Rich ground, Virginia and southward. April.—Leaves sweet, greedily eaten by cattle.

ORDER 62. **PLANTAGINÀCÆÆ**. (PLANTAIN FAMILY.)

Chiefly stemless herbs, with regular 4-merous spiked flowers, the stamens inserted on the tube of the dry and membranaceous veinless monopetalous corolla, alternate with its lobes; — chiefly represented by the genus

1. **PLANTÀGO**, L. PLANTAIN. RIBGRASS.

Calyx of 4 imbricated persistent sepals, mostly with dry membranaceous margins. Corolla salver-form, withering on the pod, the border 4-parted. Stamens 4, or rarely 2, in all or some flowers with long and weak exerted filaments, and fugacious 2-celled anthers. Ovary 2- (or in No. 5 falsely 3-4-) celled, with 1-several ovules in each cell. Style and long hairy stigma single, filiform. Pod 2-celled, 2-several-seeded, opening all round by a transverse line, so that the top falls off like a lid, and the loose partition (which bears the peltate seeds) falls away. Embryo straight, in fleshy albumen. — Leaves ribbed. Flowers whitish, small, in a bracted spike or head, raised on a naked scape. (The Latin name.)

§ 1. *Flowers all perfect and alike, and with the 4 stamens and filiform stigma much exerted, but dichogamous, i. e. with the stigma exerted from the tip of the corolla a day or so before it expands and the anthers are hung out (an arrangement for cross-fertilization): lobes of the corolla spreading or reflexed after flowering.*

* *Leaves 5-7-ribbed, mostly broad: spike long and slender, smooth: seeds not concave on the inner face: root perennial, or perhaps annual in No. 2 and 3.*

1. **P. MAJOR**, L. (COMMON PLANTAIN.) Smooth or rather hairy, rarely roughish; leaves ovate, oblong, oval, or slightly heart-shaped, often toothed, abruptly narrowed into a channelled petiole; spike dense; *pod* 7-16-seeded. — Moist-grounds, everywhere near dwellings. June-Sept. — A small and rougher form in salt marshes. (Nat. from Eu., but probably indigenous high north.)

2. **P. KAMTSCHÁTICA**, Cham. Much resembles small forms of the preceding; but sepals and bract narrower, and *pod* 4-seeded. (P. Rugelii, Decaisne.) — Buffalo, N. Y., Mr. Day, and sparingly in the south. (Apparently adv.)

3. **P. sparsiflora**, Michx. Slender (3'-18' high), smoothish or hairy; leaves lanceolate or oblong, 3-5-nerved, tapering to both ends, denticulate or entire; spike sparsely-flowered, very slender; lobes of the corolla acute; *pod* 2-seeded. — Mound City, Illinois (Dr. Vasey), and southward. July-Sept.

4. **P. cordata**, Lam. Tall, glabrous; leaves heart-shaped or round-ovate (3'-8' long), long-petioled, the ribs rising from the midrib; spike at length loosely flowered; bracts round-ovate, fleshy; *pod* 2-4-seeded. — Along rivulets, New York to Wisconsin (rare), and southward. April-June.

* * *Leaves linear, thick and fleshy, without ribs, or when dry obscurely 3-nerved: spike slender: tube of the corolla hairy below: seeds not hollowed.*

5. **P. maritima**, L., var. **juncoides**. Smooth, or the scape slightly pubescent; leaves flat or flattish and channelled, erect, nearly as long as the scape (5'-12'), mostly entire; *pod* 2-celled or incompletely 3-4-celled, 2-4-seeded; root on our coast annual or biennial. (P. juncoides, Lam.) — Salt marshes, from New Jersey northward. Near Boston a depauperate form, 2'-5' high, little fleshy, grows in sand beyond the influence of salt water (D. Murray). The perennial P. maritima occurs in New Brunswick, &c., perhaps in Maine.

* * * *Leaves 3-5-ribbed, narrow: spike thick and dense, at first or throughout very short: two of the scarious sepals generally united into one: seeds only 2, hollowed on the inner face.*

6. **P. lanceolata**, L. (RIBGRASS. RIPLEGRASS. ENGLISH PLANTAIN.) Mostly hairy; scape grooved-angled, at length much longer than the lanceolate or lance-oblong leaves, slender (9'-2° high); root perennial. — Dry fields: common eastward. (Nat. from Eu.)

§ 2. *Flowers of two sorts on distinct plants, apparently polygamo-dioecious; the mostly sterile with the usual large anthers on long capillary filaments, and the lobes of the corolla reflexed or spreading; the truly fertile with minute anthers on short included filaments, and the corolla usually closing permanently over the apex of the fruit: seeds not hollowed on the face: small annuals or biennials.*

* *Stamens 4: spike dense.*

7. **P. Virginica**, L. Hairy or hoary-pubescent (2'-9' high); leaves oblong, varying to obovate and spatulate-lanceolate, 3-5-nerved, slightly or

coarsely and sparingly toothed; seeds usually 2.—Sandy grounds, Rhode Island to Illinois and southward. May—Sept.

* * *Stamens only 2: leaves narrowly linear or thread-shaped, barely 1-ribbed: spike mostly slender, of few or many crowded or scattered small flowers.*

8. **P. pusilla**, Nutt. Minutely pubescent (1'—4' high); leaves entire; *pod* short-ovoid, 4-seeded, little exceeding the calyx and bract.—Dry hills, New York to Illinois, and southward. April—Aug.

9. **P. heterophylla**, Nutt. Leaves rather fleshy, acute, entire, or some of them below 2—4-lobed or toothed; *pod* oblong-conoidal, 10—28-seeded, nearly twice the length of the calyx and bract. (*P. pusilla*, DeCaisne, in DC.)—Low or sandy grounds, from Philadelphia southward. April—June.—Plant 2'—8' high.

§ 3. *Flowers all commonly fertile, but of 2 sorts on different plants; the more common with very small anthers on short filaments (perhaps early and close-fertilized); others with large anthers on long-exserted filaments: corolla with broad round lobes permanently widely spreading: seeds 2 (one in each cell), boat-shaped, deeply hollowed on the face: mostly annuals, with narrow and woolly or hairy leaves.*

10. **P. Patagónica**, Jacq. Silky-woolly, or becoming naked; leaves 1—3-nerved; spike cylindrical or oblong, dense; sepals very obtuse, scarious, with a thick centre. (Found through almost the whole length of America.)

Var. **gnaphalioides**, Gray. White with silky wool; leaves varying from oblong-linear to filiform; spike very dense ($\frac{1}{4}$ '—4' long), woolly; bracts not exceeding the calyx. (*P. Lagopus*, Pursh. *P. gnaphalioides*, Nutt.)—Dry plains, W. Wisconsin and southwestward.—Runs through var. SPINULOSA and var. NUDA into

Var. **aristata**, Gray. Loosely hairy and green, or becoming glabrous; bracts awned, 2—3 times the length of the flowers. (*P. aristata*, Michx., &c.)—Illinois and southwestward.

ORDER 63. PLUMBAGINACEÆ. (LEADWORT FAMILY.)

Maritime herbs, with regular 5-merous flowers, a plaited calyx, the 5 stamens opposite the separate petals or the lobes of the corolla, and the free ovary one-celled, with a solitary ovule hanging from a long cord which rises from the base of the cell.—The STATICEÆ or MARSH-ROSEMARY TRIBE only is represented in our region, in gardens by the THRIFT (*ARMERIA VULGARIS*), on the coast by a single species of

1. STÁTICE, Tourn. SEA-LAVENDER. MARSH-ROSEMARY.

Flowers scattered or loosely spiked and 1-sided on the branches, 2—3-bracted. Calyx funnel-form, dry and membranaceous, persistent. Corolla of 5 nearly or quite distinct petals, with long claws, the 5 stamens severally attached to their bases. Styles 5, rarely 3, separate. Fruit membranous and indehiscent, 1-seeded, in the bottom of the calyx. Embryo straight, in mealy albumen.—Sea-side perennials, with thick and stalked radical leaves; the naked flowering stems or

scapes branched into panicles. (*Σταυρίκη*, an ancient name given to this or some other herb, on account of its astringency.)

1. **S. Limonium**, L. Leaves oblong, spatulate, or obovate-lanceolate, 1-ribbed, tipped with a deciduous bristly point, petioled; scape much-branched, corymbose-panicled (1°-2° high); spikelets 1-3-flowered; calyx-tube hairy on the angles, the lobes ovate-triangular, with as many teeth in the sinuses. — Root thick and woody, very astringent. Flowers lavender-color. (Eu.)

Var. **Caroliniana** (S. Caroliniana, *Walt.*, &c.), the plant of the Northern States, has a hollow scape, more erect branches, at length scattered flowers, and sharper calyx-lobes. — Salt marshes along the coast, extending northward (where it passes into *S. Bahusiensis*, *Fries*). Aug., Sept. (Eu.)

ORDER 64. **PRIMULACEÆ.** (PRIMROSE FAMILY.)

Herbs, with simple leaves, and regular perfect flowers, the stamens as many as the lobes of the monopetalous (rarely polypetalous) corolla and inserted opposite them, a 1-celled ovary with a central free placenta rising from the base, bearing several or many seeds. — Calyx free from the ovary, or in *Samolus* partly coherent. (Corolla none in *Glaux*.) Stamens 4 or 5, rarely 6 or 8. Style and stigma one. Seeds with a small embryo in fleshy albumen, amphitropous and fixed by the middle, except in Tribe 4.

Tribe I. PRIMULÆ. Pod free from the calyx, opening by valves or teeth.

* Stemless: leaves all in a cluster from the root.

1. **Primula.** Corolla funnel-form or salver-shaped, open at the throat. Stamens included.
2. **Androsace.** Corolla short, very small, constricted at the throat. Stamens included.
3. **Dodecatheon.** Corolla reflexed, 5-parted. Stamens exserted: filaments united.

* * Stems leafy: corolla wheel-shaped (or in *Glaux* none).

4. **Trientalis.** Corolla mostly 7-parted. Stem leafy only at the summit.
5. **Lysimachia.** Corolla 5-6-parted or 5-6-petalled. Stems leafy throughout.
6. **Glaux.** Corolla none: the calyx petal-like.

Tribe II. ANAGALLIDÆ. Pod circumcissile. Otherwise as in Tribe I.

7. **Anagallis.** Corolla longer than the calyx, 5-parted. Leaves opposite.
8. **Centunculus.** Corolla shorter than the calyx, 4-5-cleft. Leaves alternate.

Tribe III. SAMOLEÆ. Pod partly adherent to the calyx, opening by valves.

9. **Samolus.** Corolla bell-shaped and with 5 sterile filaments in the sinuses.

Tribe IV. HOTTONIÆ. Pod as in Tribe I. Seeds fixed by the base, anatropous.

10. **Hottonia.** Corolla salver-shaped. Immersed leaves pectinately dissected.

1. **PRÍMULA**, L. PRIMROSE. COWSLIP.

Calyx tubular, angled, 5-cleft. Corolla salver-shaped, enlarging above the insertion of the stamens; the 5 lobes often notched or inversely heart-shaped. Stamens 5, included. Pod many-seeded, splitting at the top into 5 valves or 10 teeth. — Low perennial herbs, producing a tuft of veiny leaves at the root, and simple scapes, bearing the flowers in an umbel. (Name a diminutive of *primus*, from the flowering of the true Primrose in early spring.)

1. **P. farinosa**, L. (BIRD'S-EYE PRIMROSE.) Leaves elliptical or obovate-lanceolate, the lower surface and the 3-20-flowered involucre, &c. covered with

a white mealiness: corolla pale lilac with a yellow eye. — Shores of Lakes St. Clair and Huron; also Mount Kineo, Maine (*A. H. Smith*), and northward. June, July. — Scape 3' - 10' high. (Eu.)

2. **P. Mistassinica**, Michx. Leaves spatulate or wedge-oblong, thin and veiny, *not mealy*; involucre 1-8-flowered; lobes of the flesh-colored corolla broadly and deeply obcordate. — Shores of the Upper Lakes: also Crooked Lake (*Sartwell*) and Annsville, Oneida County, New York (*Knieskern* and *Vasey*), Willoughby Mountain, Vermont (*Wood*, &c.), and northward. May. — A pretty species, 2' - 6' high. (Eu.)

2. ANDRÓSACE, Tourn. ANDROSACE.

Calyx 5-cleft; the tube short. Corolla salver-shaped or funnel-form; the tube shorter than the calyx, contracted at the throat; the limb 5-parted. Stamens and style included. Pod 5-valved. — Small herbs, with clustered root-leaves, and very small solitary or umbelled flowers. (An ancient and obscure name, thought by Linnaeus to be formed of *ἀνδρός*, of man, and *σάκος*, a shield.)

1. **A. occidentalis**, Pursh. Smoothish annual; scapes diffuse (2' - 4' high), many-flowered; leaves and leaflets of the involucre oblong-ovate, entire, sessile; calyx-lobes leafy, triangular-lanceolate, longer than the (white) corolla. — Bare hills on the Mississippi, Illinois, and northwestward.

3. DODECÀTHEON, L. AMERICAN COWSLIP.

Calyx deeply 5-cleft; the divisions lanceolate, reflexed. Corolla with a very short tube, a thickened throat, and a 5-parted reflexed limb; the divisions long and narrow. Filaments short, monadelphous at the base: anthers long and linear, approximate in a slender cone. — Perennial smooth herb, with fibrous roots, a cluster of oblong or spatulate leaves, and a simple naked scape, involucre with small bracts at the summit, bearing an ample umbel of showy flowers, nodding on slender peduncles. Corolla rose-color, or sometimes white. (Name fancifully assumed from *δώδεκα*, twelve, and *θεοί*, gods.)

1. **D. Meadia**, L. — Rich woods, Pennsylvania and Maryland to Wisconsin, and southwestward. May, June. — Very handsome in cultivation. In the West called SHOOTING-STAR.

4. TRIENTÀLIS, L. CHICKWEED-WINTERGREEN.

Calyx mostly 7-parted; the divisions linear-lanceolate, pointed. Corolla mostly 7-parted, spreading, flat, without any tube. Filaments slender, united in a ring at the base: anthers oblong, revolute after flowering. Pod few-seeded. — Low and smooth perennials, with simple erect stems, bearing a few alternate usually minute and scale-like leaves below, and a whorl of thin veiny leaves at the summit. Peduncles one or more, very slender, bearing a delicate white and star-shaped flower. (A Latin name, meaning the third part of a foot, alluding to the height of the plant.)

1. **T. Americana**, Pursh. (STAR-FLOWER.) Leaves elongated-lanceolate, tapering to both ends; petals finely pointed. — Damp cold woods: common northward, and southward in the mountains. May.

5. **LYSIMACHIA**, Tourn. LOOSESTRIFE.

Calyx 5- (rarely 6-7-) parted. Corolla wheel-shaped, 5- (or 6-7-) parted, or even of as many separate petals. Stamens as many. Pod globose, 5-10-valved, few-many-seeded. — Leafy-stemmed perennials, with mostly yellow flowers, either axillary or in a terminal raceme. (Named in honor of King *Lysimachus*, or from *λύσις*, a release from, *μάχη*, strife.)

§ 1. **NAUMBURGIA**, Moench. *Leaves opposite, sessile, minutely glandular-dotted: parts of the flower not rarely 6, sometimes 7; the narrow petals almost or quite distinct, and with an interposed small tooth, sprinkled with purplish dots: filaments slender, distinct, equal: anthers short: pod few-seeded.*

1. **L. thyrsoiflora**, L. (TUFTED LOOSESTRIFE.) Smooth; stem simple (1°-2° high); lower leaves reduced to scales, the rest lanceolate, the axils of one or two pairs of the middle ones bearing a short-peduncled head-like or spike-like cluster of light yellow small flowers; divisions of the corolla lance-linear. (*Naumburgia thyrsoiflora*, *Reichenb.*, and *Ed. 2.*) — Cold wet swamps, from Penn. northward. June, July. (Eu.)

§ 2. **TRIDÝNIA**, Raf. *Leaves opposite or whorled, sessile, dotted: calyx and golden-yellow corolla streaked with dark lines: filaments mostly unequal, plainly monadelphous at the base, with no interposed sterile ones: anthers short: pod 5-valved, ripening only 2-5 seeds.*

2. **L. stricta**, Ait. Smooth, at length branched, very leafy; leaves opposite or rarely alternate, lanceolate, acute at each end; flowers on slender pedicels in a long raceme (5'-12'), which is leafy at the base; or, in var. **PRODUCTA**, leafy for fully half its length: lobes of the corolla lance-oblong. — Low grounds: common. June-Aug. — Stems 1°-2° high, often bearing oblong or moniliform bulblets in the axils.

3. **L. quadrifolia**, L. Somewhat hairy; stem simple (1°-2° high); leaves whorled in fours or fives (rarely in threes or sixes) ovate-lanceolate; flowers on long capillary peduncles from the axils of the leaves; lobes of the corolla ovate-oblong. — Moist or sandy soil: common. June. — A variety has the leaves varying to opposite and partly alternate, some of the upper reduced to bracts shorter than the peduncles. Near New York, Washington, &c.

§ 3. **STEIRONEMA**, Raf. *Leaves opposite, not dotted, glabrous, mostly ciliate at the base: flowers nodding on slender peduncles from the axils of the upper leaves: corolla light yellow, not streaked nor dotted; the lobes broadly ovate, pointed, with undulate or denticulate margins, little exceeding the sepals: filaments nearly equal, scarcely monadelphous, with the rudiments of a sterile set interposed at the base in the form of slender teeth or processes: anthers linear, at length curved: pod 5-10-valved, or bursting irregularly, 10-20-seeded.*

4. **L. ciliata**, L. Stem erect (2°-3° high); leaves lanceolate-ovate (3'-6' long), tapering to an acute point, rounded or heart-shaped at the base, all on long and fringed petioles; corolla longer than the calyx. — Low ground and thickets: common. July.

5. **L. radicans**, Hook. Stem slender, soon reclined, the elongated branches often rooting in the mud; leaves ovate-lanceolate, mostly rounded at the base, on slen-

der petioles: corolla about the length of the calyx. — Swampy river-banks, West Virginia (*Aikin*) and southward. — Leaves and flowers nearly one half smaller than in the last.

6. **L. lanceolata**, Walt. *Stem erect* (10'–20' high); *leaves lanceolate, varying to oblong and to linear, narrowed into a short margined petiole* or tapering base, or the lowest short and broad on long petioles. — Var. **HÝBRIDA** is merely the broader-leaved form. Var. **ANGUSTIFÓLIA** (*L. angustifolia*, *Lam.*) is a slender branching form, with the upper leaves narrowly lanceolate or linear, and acute at both ends. — Low grounds: not uncommon, especially westward and southward. June–Aug.

7. **L. longifolia**, Pursh. *Stem erect*, 4-angled, slender (1°–3° high), often branched below; *stem-leaves sessile, narrowly linear, elongated* (2'–4' long, 2''–3'' wide), smooth and shining, rather rigid, obtuse, the margins often a little revolute, the veins obscure; the lowest oblong or spatulate; corolla (8''–9'' broad) longer than the calyx, the lobes conspicuously pointed. (*L. revoluta*, *Nutt.*) — Moist soil, Western New York and Pennsylvania to Wisconsin and Illinois. July–Sept.

§ 4. *Introduced European species of true Lysimachia.*

8. **L. NUMMULARIA**, L. (MONEYWORT.) Smooth; stems trailing and creeping; leaves roundish, small, short-petioled; peduncles axillary, 1-flowered; divisions of the corolla broadly ovate, obtuse, longer than the lance-ovate calyx-lobes and the stamens; filaments slightly monadelphous at the base. — Escaped from gardens into damp ground in some places. July–Sept.

6. **GLAÛX**, L. SEA-MILKWORT.

Calyx bell-shaped, 5-cleft; the lobes ovate, petal-like. Corolla wanting. Stamens 5, on the base of the calyx, alternate with its lobes. Pod 5-valved, few-seeded. — A low and leafy fleshy perennial, with opposite oblong and entire sessile leaves, and solitary nearly sessile (purplish and white) flowers in their axils. (An ancient Greek name, from *γλαυκός*, *sea-green*.)

1. **G. maritima**, L. — Sea-shore of New England from Cape Cod northward. Also beyond the Mississippi northwestward. June. (Eu.)

7. **ANAGÁLLIS**, Tourn. PIMPERNEL.

Calyx 5-parted. Corolla wheel-shaped, with almost no tube, 5-parted, longer than the calyx; the divisions broad. Stamens 5: filaments bearded. Pod membranaceous, circumscissile, the top falling off like a lid, many-seeded. — Low, spreading or procumbent herbs, mostly annuals, with opposite or whorled entire leaves, and solitary flowers on axillary peduncles. (The ancient Greek name, probably from *ἀνά*, *again*, and *αγάλλω*, *to delight in*.)

1. **A. ARVÉNSIS**, L. (COMMON PIMPERNEL.) Leaves ovate, sessile, shorter than the peduncles; petals obovate, obtuse, fringed with minute teeth or stalked glands. — Waste sandy fields. June–Aug. — Flowers variable in size, scarlet, sometimes purple, blue, or white, quickly closing at the approach of bad weather; whence the English popular name of "*Poor Man's Weather-glass*." (Nat. from Eu.)

8. CENTÚNCULUS, L. CHAFFWEED.

Calyx 4-5-parted. Corolla shorter than the calyx, 4-5-cleft, wheel-shaped, with an urn-shaped short tube, usually withering on the summit of the pod (which is like that of *Anagallis*). Stamens 4 or 5: filaments beardless.— Small annuals, with alternate entire leaves, and solitary inconspicuous flowers in their axils. (Derivation obscure.)

1. *C. mínimus*, L. Stems ascending (2'-5' long); leaves ovate, obovate, or spatulate-oblong; flowers nearly sessile, the parts mostly in fours. (*C. lanceolatus*, *Michx.*)— Low grounds, Illinois and southward. (Eu.)

9. SÁMOLUS, L. WATER PIMPERNEL. BROOK-WEED.

Calyx 5-cleft; the tube adherent to the base of the ovary. Corolla somewhat bell-shaped, 5-cleft, commonly with 5 sterile filaments in the sinuses. True stamens 5, on the tube of the corolla, included. Pod 5-valved at the summit, many-seeded.— Smooth herbs, with alternate entire leaves, and small white flowers in racemes. ("According to Pliny, an ancient Druidical name.")

1. *S. Valerándi*, L. Stem erect (6'-12' high), leafy; leaves obovate; bracts none; bractlets on the middle of the slender ascending pedicels; calyxlobes ovate, shorter than the corolla. (Eu.)

Var. *Americánus*, Gray. More slender, becoming diffusely branched; racemes often paniced, the pedicels longer and spreading; bractlets, flowers, and pods smaller. (*S. floribúndus*, *H. B. K.*)— Wet places: common. June-Sept.

10. HOTTÓNIA, L. FEATHERFOIL. WATER VIOLET.

Calyx 5-parted, the divisions linear. Corolla salver-shaped, with a short tube; the limb 5-parted. Stamens 5, included. Pod many-seeded, 5-valved; the valves cohering at the base and summit. Seeds attached by their base, anatropous.— Aquatic perennials, with the immersed leaves pectinate, and the erect hollow flower-stems almost leafless. Flowers white or whitish, whorled at the joints, forming a sort of interrupted raceme. (Named for *Prof. Hotton*, a botanist of Leyden, in the 17th century.)

1. *H. infláta*, Ell. Leaves dissected into thread-like divisions, scattered on the floating and rooting stems, and crowded at the base of the cluster of peduncles, which are strongly inflated between the joints (often as thick as one's finger); pedicels short.— Pools and ditches, New England to Kentucky, and southward. June-Aug.

ORDER 65. LENTIBULACEÆ. (BLADDERWORT FAMILY.)

Small herbs (growing in water or wet places), with a 2-lipped calyx, and a 2-lipped personate corolla, 2 stamens with (confluently) one-celled anthers, and a one-celled ovary with a free central placenta, bearing several anatropous seeds, with a thick straight embryo, and no albumen.— Corolla deeply 2-lipped, spurred at the base in front; the palate usually bearded. Ovary

free: style very short or none: stigma 1-2-lipped, the lower lip larger and with a prominent palate. Pod often bursting irregularly. Scapes 1-few-flowered. — Consists mostly of the two following genera:—

1. UTRICULÀRIA, L. BLADDERWORT.

Lips of the 2-parted calyx entire, or nearly so. Corolla personate, the palate on the lower lip projecting, often closing the throat. Anthers convergent. — Aquatic and immersed, with capillary dissected leaves bearing little bladders, which are filled with air and float the plant at the time of flowering; or rooting in the mud, and sometimes with few or no leaves or bladders. Scapes 1-few-flowered; usually flowering all summer. (Name from *utriculus*, a little bladder.)

* *Upper leaves in a whorl on the otherwise naked scape, floating by means of large bladders formed of the inflated petioles; the lower leaves dissected and capillary, bearing little bladders: rootlets few or none.*

1. **U. inflata**, Walt. (INFLATED BLADDERWORT.) Swimming free; bladder-like petioles oblong, pointed at the ends, and branched near the apex, bearing fine thread-like divisions; flowers 5-10 (large, yellow); the appressed spur half the length of the corolla; style distinct. — Ponds, Maine to Virginia, and southward, near the coast.

* * *Scapes naked (except some small scaly bracts), from immersed branching stems, which commonly swim free, and bear capillary dissected leaves furnished with small air-bladders on their lobes: roots few and not-affixed, or none. (Mostly perennial, propagated from year to year by a sort of buds.)*

— *Flowers all alike, yellow, several in a raceme: pedicels nodding in fruit.*

2. **U. vulgaris**, L. (GREATER BLADDERWORT.) Immersed stems (1°-3° long) crowded with 2-3-pinnately many-parted capillary leaves, bearing many bladders; scapes 5-12-flowered (6'-12' long); corolla closed (6"-9" broad, the sides reflexed; spur conical, rather shorter than the lower lip, thick and blunt in the European and the high northern plant; in the common Var. AMERICANA (*U. macrorhiza*, *LeConte*), less thick and rather acute. — Common in ponds and slow streams. (Eu.)

3. **U. minor**, L. (SMALLER B.) Leaves scattered on the thread-like immersed stems, 2-4 times forked, short; scapes weak, 2-8-flowered (3'-7' high); upper lip of the gaping corolla not longer than the depressed palate; spur very short and blunt, or almost none. — Shallow water, Connecticut to N. Illinois and northward. — Corolla 2"-3" broad. (Eu.)

— — *Flowers of 2 sorts; viz. the usual sort (3-7) in a raceme, their pedicels ascending, the corolla yellow; and more fertile ones solitary and scattered along the leafy stems, on short soon reflexed peduncles, fruiting in the bud, the corolla minute and never expanding.*

4. **U. clandestina**, Nutt. Leaves numerous on the slender immersed stems, several times forked, capillary, copiously bladder-bearing; scapes slender (3'-5' high); lips of the corolla nearly equal in length, the lower broader and 3-lobed, somewhat longer than the approximate thick and blunt spur. — Ponds, E. New England, W. New York, and New Jersey.

+ + + *Flowers all alike, few (1-5): pedicels erect in fruit.*

+ + *Corolla yellow: scape and pedicels filiform: spur ascending or horizontal.*

5. **U. intermedia**, Hayne. *Leaves crowded on the immersed stems, 2-ranked, 4-5 times forked, rigid; the divisions linear-awl-shaped, minutely bristle-toothed along the margins, not bladder-bearing, the bladders being on separate leafless branches; upper lip of the corolla much longer than the palate; spur conical-oblong, acute, appressed to the very broad (6''-8'') lower lip and nearly as long as it.*—Shallow pools, New England and New Jersey to Ohio, Wisconsin, and northward.—Leafy stems 3'-6' long. Scapes 3'-7' high. (Eu.)

6. **U. striata**, LeConte. *Leaves crowded or whorled on the small immersed stems, several times forked, capillary, bladder-bearing; flowers 2-5 (6'' broad), on long pedicels; lips of the corolla nearly equal, broad and expanded; the upper undulate, concave, plaited-striate in the middle; spur nearly linear, obtuse, approaching and almost equalling the lower lip.*—Shallow pools in pine barrens, Long Island, New Jersey, and southward.—Scape 8'-12' high.

7. **U. biflora**, Lam. *Scape (2'-5' high) 1-3-flowered, at the base bearing somewhat elongated submersed branches with capillary root-like leaves and numerous bladders; spur oblong, equalling the lower lip; seeds scale-shaped; otherwise resembles the next.*—Shallow water, Illinois and southward.

8. **U. gibba**, L. *Scape (1'-3' high), 1-2-flowered, at the base furnished with very slender short branches, bearing sparingly dissected capillary root-like leaves and scattered bladders; lips of the corolla broad and rounded, nearly equal; the lower with the sides reflexed (4''-5'' long), exceeding the very thick and blunt conical gibbous spur.*—Shallow water, Virginia to Massachusetts, N. New York and N. Illinois.

+ + + *Corolla violet-purple.*

9. **U. purpurea**, Walt. ? *Leaves whorled along the long immersed free floating stems, petioled, decomposed, capillary, bearing many bladders; flowers 2-4 (6'' wide); spur appressed to the 3-lobed 2-saccate lower lip of the corolla and about half its length.* (*U. saccata*, LeConte.)—Ponds, Maine to Virginia, and southward.—Scape 3'-6' high, not scaly below.

* * * *Scape solitary, slender and naked, or with a few small scales; the base rooting in the mud or soil: leaves small, awl-shaped or grass-like, often raised out of the water, commonly few or fugacious: air-bladders few on the leaves or rootlets, or commonly none.*

+ *Flower purple, solitary: leaves bearing a few delicate lobes.*

10. **U. resupinata**, Greene. *Scape (2'-8' high) 2-bracted above; leaves thread-like, on delicate creeping branches; corolla (4''-5'' long) deeply 2-parted; spur oblong-conical, very obtuse, shorter than the dilated lower lip and remote from it, both ascending, the flower resting transversely on the summit of the scape.*—Sandy margins of ponds, E. Maine to Rhode Island.

+ + *Flowers 2-10, (chiefly) yellow: leaves entire, rarely seen.*

11. **U. cornuta**, Michx. *Stem strict (3'-1° high), 2-10-flowered; pedicels not longer than the calyx; lower lip of the corolla large and helmet-shaped, its centre very convex and projecting, while the sides are strongly reflexed; upper lip obovate and much smaller; spur awl-shaped, turned downward and outward,*

about as long as the lower lip. — Peat-bogs, or sandy swamps: common both northward and southward. — Flowers close together, large.

12. **U. subulata**, L. Stem capillary (3' - 5' high); *pedicels capillary*; *lower lip of the corolla flat* or with its margins recurved, *equally 3-lobed*, much larger than the ovate upper one; *spur oblong*, acute, straight, *appressed* to the lower lip, which it nearly equals in length. — Sandy swamps, pine-barrens of New Jersey, Virginia, and southward. June. — Corolla 2" - 4" broad.

U. — Walter characterizes his *U. purpurea* as with "*floribus parvis*." Elliott mentions that he once saw, near Savannah, a small terrestrial species, like *U. subulata*, but purple-flowered, which he took for Walter's plant. Mr. J. A. Paine, Jr. found in the pine barrens of New Jersey, in Sept., 1866, a few minute specimens of this sort, with "faint pink-purple corolla, not larger than a pin's head." It is left for further investigation.

2. PINGUÍCULA, L. BUTTERWORT.

Upper lip of the calyx 3-cleft, the lower 2-cleft. Corolla with an open hairy or spotted palate. — Small and stemless perennials, growing on damp rocks, with 1-flowered scapes, and broad and entire leaves, all clustered at the root, soft-fleshy, mostly greasy to the touch (whence the name, from *pinguis*, fat).

1. **P. vulgaris**, L. Leaves ovate or elliptical; scape and calyx a little pubescent; lips of the violet corolla very unequal, the tube funnel-form; spur straightish. — Wet rocks, W. New York to Lake Superior, and northward. July. (Eu.)

ORDER 66. BIGNONIACEÆ. (BIGNONIA FAMILY.)

Woody or rarely herbaceous plants, monopetalous, didynamous or diandrous, with the ovary commonly 2-celled by the meeting of the two parietal placentæ or of a projection from them, many-seeded: the large seeds with a flat embryo and no albumen. — Calyx 2-lipped, 5-cleft, or entire. Corolla tubular or bell-shaped, 5-lobed, somewhat irregular or 2-lipped, deciduous; the lower lobé largest. Stamens inserted on the corolla; the fifth or posterior one, and sometimes the shorter pair also, sterile or rudimentary: anthers of 2 diverging cells. Ovary free, bearing a long style, with a 2-lipped stigma. — Leaves compound or simple, opposite, rarely alternate. Flowers large and showy. — Chiefly a tropical family: to which are appended several outlying groups, such as the PEDALINEÆ, represented by *Martynia*, &c.

* Woody plants, with dry pods. Seeds transverse, very flat, winged; the broad and leaf-like cotyledons notched at both ends.

1. **Bignonia**. Pod flattened parallel with the partition. Leaves compound, tendril-bearing.

2. **Tecoma**. Pod flattish contrary to the partition. Leaves compound, without tendrils.

3. **Catalpa**. Pod as in No. 2. Fertile stamens only 2. Trees: leaves simple.

** Herbs. Fruit fleshy outside, woody within. Seeds attached by one end, not winged; the cotyledons thick and entire.

4. **Martynia**. Fertile stamens 2 or 4. Corolla bell-shaped, 5-lobed. Fruit beaked.

1. **BIGNONIA**, Tourn. BIGNONIA.

Calyx truncate, or slightly 5-toothed. Corolla somewhat bell-shaped, 5-lobed and rather 2-lipped. Stamens 4, often showing a rudiment of the fifth. Pod 2-celled, flattened parallel with the valves and partition. Seeds transversely winged. — Woody climbers, with chiefly compound leaves, terminating in a tendril. (Named for the *Abbé Bignon*.)

1. **B. capreolata**, L. Smooth; leaves of 2 ovate or oblong leaflets and a branched tendril, often with a pair of accessory leaves in the axil resembling stipules; peduncles few and clustered, 1-flowered. — Rich soil, Virginia to S. Illinois and southward. April. — Stems climbing tall trees; π transverse section of the wood showing a cross. Corolla orange, 2' long. Pod 6' long. Seeds with the wing $1\frac{1}{2}$ ' long.

2. **TÉCOMA**, Juss. TRUMPET-FLOWER.

Calyx bell-shaped, 5-toothed. Corolla funnel-form, 5-lobed, a little irregular. Stamens 4. Pod 2-celled, the partition contrary to the convex valves. Seeds transversely winged. — Woody climbers, with compound leaves. (Abridged from the Mexican name.)

1. **T. radicans**, Juss. (TRUMPET CREEPER.) Climbing by rootlets; leaves pinnate; leaflets 5-11, ovate, pointed, toothed; flowers corymbed; stamens not protruded beyond the tubular-funnel-form corolla. (*Bignonia radicans*, L.) — Rich soil, Pennsylvania to Illinois and southward; but cultivated farther north. July-Sept. — Corolla 2' - 3' long, orange and scarlet, showy.

3. **CATÁLPA**, Scop., Walt. CATALPA. INDIAN BEAN.

Calyx deeply 2-lipped. Corolla bell-shaped, swelling; the undulate 5-lobed spreading border irregular and 2-lipped. Fertile stamens 2, or sometimes 4; the 1 or 3 others sterile and rudimentary. Pod very long and slender, nearly cylindrical, 2-celled; the partition contrary to the valves. Seeds winged on each side, the wings cut into a fringe. (The aboriginal name.)

1. **C. bignonioides**, Walt. Leaves heart-shaped, pointed, downy beneath; flowers in open compound panicles. — S. Illinois? and southward. Cultivated in the Northern States: a well-known ornamental tree, with large leaves, and showy flowers, which are white, slightly tinged with violet, and dotted with purple and yellow in the throat, appearing in July. Pods hanging till the next spring, often 1° long.

4. **MARTÝNIA**, L. UNICORN-PLANT.

Calyx 5-cleft, mostly unequal. Corolla gibbous, bell-shaped, 5-lobed and somewhat 2-lipped. Fertile stamens 4, or only 2. Pod fleshy, the flesh at length falling away in 2 valves; the inner part woody, terminated by a beak, which at length splits into 2 hooked horns, and opens at the apex between the horns, imperfectly 5-celled, owing to the divergence of the two plates of each of the two partitions or placentæ, leaving a space in the centre, while by reaching and cohering with the walls of the fruit they form 4 other cells. Seeds several, wing-

less, with a thickened and roughened coat. — Low branching annuals, clammy-pubescent, exhaling a heavy odor: stems thickish: leaves simple, rounded. Flowers racemed, large. (Dedicated to *Prof. John Martyn*, of Cambridge.)

1. **M. proboscidea**, Glox. Leaves heart-shaped, oblique, entire, or undulate, the upper alternate; corolla dull white or purplish, or spotted with yellow and purple; endocarp of the fruit crested on one side, long-beaked. — Banks of the Mississippi in S. Illinois (probably indigenous) and southwestward. Also common in gardens. July – Oct.

ORDER 67. OROBANCHACEÆ. (BROOM-RAPE FAMILY.)

Herbs destitute of green foliage (root-parasites) monopetalous, didynamous, the ovary one-celled with 2 or 4 parietal placentæ; pod very many-seeded: seeds minute, with albumen, and a very minute embryo. — Calyx persistent, 4 – 5-toothed or parted. Corolla tubular, more or less 2-lipped, ringent, persistent and withering; the upper lip entire or 2-lobed, the lower 3-lobed. Stamens 4, didynamous, inserted on the tube of the corolla: anthers 2-celled, persistent. Ovary free, ovoid, pointed with a long style which is curved at the apex: stigma large. Pod 1-celled, 2-valved; the valves each bearing on their face one placenta or a pair. Seeds very numerous, minute, anatropous, the minute embryo at the base of transparent albumen. — Low, thick or fleshy herbs, bearing scales in place of leaves, lurid yellowish or brownish throughout. Flowers solitary or spiked.

* Flowers of two sorts: stems branching.

1. **Epiphegus**. Upper flowers sterile, with a tubular corolla; the lower fertile, with the corolla minute and not expanding. Bracts inconspicuous.

* * Flowers all alike and perfect: stems mostly simple.

2. **Conopholis**. Flowers spiked. Calyx with 2 bractlets, and split on the lower side. Stamens protruded. Corolla 2-lipped.

3. **Phelipæa**. Flowers spiked or paniced. Calyx with 2 bractlets, and regularly 5-cleft. Corolla 2-lipped. Stamens included.

4. **Aphyllon**. Flowers solitary, without bractlets. Calyx regularly 5-cleft. Corolla with the border almost equally 5-lobed. Stamens included.

1. EPIPHÈGUS, Nutt. BEECH-DROPS. CANCER-ROOT.

Flowers racemose or spiked, scattered on the branches; the upper sterile, with a long tubular corolla and long filaments and style; the lower fertile, with a very short corolla which seldom opens, but is forced off from the base by the growth of the pod: the stamens and style very short. Calyx 5-toothed. Stigma capitate, a little 2-lobed. Pod 2-valved at the apex, with 2 approximate placentæ on each valve. — Herbs slender, purplish or yellowish-brown, much branched, with small and scattered scales, 6' – 12' high. (Name composed of *ἐπί*, upon, and *φηγός*, the *Beech*, because it grows on the roots of that tree.)

1. **E. Virginiána**, Bart. (*E. Americanus*, Nutt.) — Common under *Beech*-trees, parasitic on their roots. Aug. – Oct. — Corolla of the upper (sterile) flowers whitish and purple, 6'' – 8'' long, curved, 4-toothed.

2. **CONÓPHOLIS**, Wallroth. SQUAW-ROOT. CANCER-ROOT.

Flowers in a thick scaly spike, perfect, with 2 bractlets at the base of the irregularly 4-5-toothed calyx; its tube split down on the lower side. Corolla tubular, swollen at the base, strongly 2-lipped; the upper lip arched, notched at the summit; the lower shorter, 3-parted, spreading. Stamens protruded. Stigma depressed. Pod with 4 placentæ, a pair on the middle of each valve. — Upper scales forming bracts to the flowers; the lower covering each other in regular order, not unlike those of a fir-cone (whence the name, from *κῶνος*, a cone, and *φολίς*, a scale.)

1. **C. Americana**, Wallroth. (*Orobánche Americana*, L.) — Oak woods: not rare, growing in clusters among fallen leaves. May, June. — A singular plant, chestnut-colored or yellowish throughout, as thick as a man's thumb, 3'-6' long, covered with scales, which are at first fleshy, then dry and hard.

3. **PHELIPÆA**, Tourn. BROOM-RAPE.

Flowers perfect, crowded in a spike, raceme, or clustered panicle, with a pair of bractlets at the base of the regular 4-5-cleft calyx. Corolla 2-lipped; the upper lip 2-lobed or notched; the lower 3-parted. Stamens included. Ovary with a gland at the base on the upper side. Pod with 4 placentæ, two on the middle of each valve. — Stems rather thick, scaly. (Named for *L. & J. Phelipeaux*, patrons of science in the time of Tournefort.)

1. **P. Ludoviciana**, Don. Glandular-pubescent, branched (3'-12' high); the flowers spiked in close clusters; corolla somewhat curved, twice the length of the narrow lanceolate calyx-lobes; the lips equal in length. — Illinois (*E. Hall*) and westward. Oct.

4. **APHÝLLON**, Mitchell. NAKED BROOM-RAPE.

Flowers perfect, solitary on long naked scapes or peduncles, without bractlets. Calyx 5-cleft, regular. Corolla with a long curved tube and a spreading border, somewhat 2-lipped; the upper lip deeply 2-cleft, its lobes similar to the 3 of the lower lip. Stamens included. Stigma broadly 2-lipped. Capsule with 4 equidistant placentæ, 2 borne on each valve half-way between the midrib and the margin. Plants brownish or yellowish. Flowers (purplish) and scapes minutely glandular-pubescent. (Name from a privative and *φύλλον*, foliage, alluding to the naked stalks.) — Perhaps rather a section of *Phelipæa*.

1. **A. uniflorum**, Torr. & Gr. (ONE-FLOWERED CANCER-ROOT.) *Stem subterranean or nearly so, very short, scaly, often branched, each branch sending up 1-3 slender one-flowered scapes (3'-5' high); divisions of the calyx lanceawl-shaped, half the length of the corolla.* (*Orobánche uniflora*, L.) — Woods: not rare. April, May. — Corolla 1' long, with 2 yellow bearded folds in the throat, the lobes obovate.

2. **A. fasciculatum**, Torr. & Gr. *Scaly stem erect and rising 3'-4' out of the ground, mostly longer than the crowded peduncles; divisions of the calyx triangular, very much shorter than the corolla, which has rounded short lobes.* (*Orobánche fasciculata*, Nutt.) — Islands in Lake Michigan (*Engelmann*), N. Illinois (*Vasey*), and northwestward. May.

ORDER 68. SCROPHULARIACEÆ. (FIGWORT FAMILY.)

Chiefly herbs (rarely trees), with didynamous or diandrous (or very rarely 5 perfect) stamens inserted on the tube of the 2-lipped or more or less irregular corolla, the lobes of which are imbricated in the bud: fruit a 2-celled and usually many-seeded pod, with the placenta in the axis: seeds anatropous, with a small embryo in copious albumen. — Style single: stigma entire or 2-lobed. Leaves and inflorescence various; but the flowers not terminal in any genuine representatives of the order. — A large order of bitterish, some of them narcotic-poisonous plants: the two principal groups generally distinguishable by the æstivation of the corolla.

I. ANTIRRHINIDÆ. Upper lip of the corolla covering the lower in the bud (with occasional exceptions in *Mimulus*, &c.). Pod usually septicial.

Tribe I. VERBASCEÆ. Corolla nearly wheel-shaped. Flowers in a simple spike or raceme. Leaves all alternate.

1. **Verbascum.** Stamens 5, all with anthers, and 3 or all of them with bearded filaments.

Tribe II. ANTIRRHINÆÆ. Corolla tubular, with a spur or sac at the base below, the throat usually with a palate. Pod opening by chinks or holes. Flowers in simple racemes or axillary. Lower leaves usually opposite or whorled.

2. **Linaria.** Corolla spurred at the base; the palate seldom closing the throat.

3. **Antirrhinum.** Corolla merely saccate at the base; the palate closing the throat.

Tribe III. CHELONEÆ. Corolla tubular, or 2-lipped, not spurred nor saccate below. Pod 2-4-valved. Leaves opposite. Inflorescence usually compound; the flowers in small clusters or cymes in the axils of the leaves or bracts, the clusters spiked or racemed; or when reduced to a single flower the peduncle 2-bracteate. Stamens 4, with mostly a rudiment of the fifth.

4. **Scrophularia.** Corolla inflated, globular or oblong, with four erect lobes and one spreading one. Rudiment of the sterile stamen a scale on the upper lip.

5. **Collinsia.** Corolla 2-cleft, the short tube saccate on the upper side; the middle lobe of the lower lip sac-like and enclosing the declined stamens.

6. **Chelone.** Corolla tubular, inflated above. Sterile stamen shorter than the others. Anthers very woolly. Seeds winged.

7. **Pentstemon.** Corolla tubular. Sterile stamen about as long as the rest. Seeds wingless.

Tribe IV. GRATIOLEÆ. Corolla tubular, not saccate nor spurred. Pod 2-valved. Inflorescence simple; the flowers being single in the axil of the bracts or leaves, the peduncles bractless. Leaves all or the lower ones opposite. No rudiment of a fifth stamen.

* Stamens 4, all anther-bearing and similar.

8. **Mimulus.** Calyx prismatic, 5-angled, 5-toothed. Corolla elongated.

9. **Conoclea.** Calyx 5-parted, the divisions equal. Corolla short.

10. **Herpestis.** Calyx 5-parted, unequal, the upper division largest. Corolla short.

** Anther-bearing stamens 2: usually also a pair of sterile filaments.

11. **Gratiola.** Calyx 5-parted. Stamens included; the sterile pair short or none.

12. **Ilysanthes.** Calyx 5-parted. Stamens included; the sterile filaments protruded.

II. RHINANTHIDÆ. Under lip or the lateral lobes of the corolla covering the upper in the bud. Pod commonly loculicidal.

Tribe V. SIBTHORPIÆ, VERONICEÆ, &c. Corolla wheel-shaped, salver-shaped, or bell-shaped. Stamens 2 or 4, not approaching each other in pairs nor strongly didynamous: anthers 2-celled.

13. **Micranthemum.** Calyx 4-toothed or cleft. Upper lip of corolla short or none. Stamens 2, anterior : filaments with an appendage. Leaves opposite. Flowers axillary.
14. **Limosella.** Calyx 5-toothed. Corolla open bell-shaped, 5-cleft, nearly regular. Stamens 4. Leaves alternate or fascicled, fleshy. Flowers axillary.
15. **Synthyris.** Calyx 4-parted. Corolla bell-shaped, 2-4-lobed, irregular. Stamens 2 or 4. Leaves alternate. Flowers racemed.
16. **Veronica.** Calyx 4- (rarely 3-5-) parted. Corolla wheel-shaped or salver-shaped, almost regular. Stamens 2. Leaves chiefly opposite or whorled. Flowers racemed.
- Tribe VI. BUCHNEREÆ.** Corolla salver-shaped. Stamens 4, approximate in pairs : anthers 1-celled. Upper leaves alternate. Flowers in a spike.
17. **Buchnera.** Calyx tubular, 5-toothed. Limb of the elongated corolla 5-cleft.
- Tribe VII. GERARDIÆ.** Corolla inflated or tubular, with a spreading and slightly unequal 5-lobed limb. Stamens 4, approximate in pairs : anthers 2-celled. Leaves opposite, or the uppermost alternate.
18. **Seymeria.** Stamens nearly equal. Tube of the corolla broad, not longer than the lobes.
19. **Gerardia.** Stamens strongly unequal, included.
- Tribe VIII. EUPHRASIEÆ.** Corolla tubular, 2-lipped ; the upper lip narrow, erect or arched, enclosing the 4 usually strongly didynamous stamens.
- * Anther-cells unequal and separated. Pod many-seeded.
20. **Castilleja.** Calyx tubular, cleft down the lower, and often also on the upper, side.
- ** Anther-cells equal. Pod many-several-seeded.
21. **Schwalbea.** Calyx 5-toothed, very oblique, the upper tooth much the smallest.
22. **Euphrasia.** Calyx 4-cleft. Upper lip of the corolla 2-lobed. Pod oblong.
23. **Rhinanthus.** Calyx inflated, ovate. Pod orbicular : seeds winged.
24. **Pedicularis.** Calyx not inflated. Pod ovate or sword-shaped : seeds wingless.
- *** Anther-cells equal. Pod 1-4-seeded.
25. **Melampyrum.** Calyx 4-cleft. Ovary 2-celled, 4-ovuled. Pod flat, oblique.

1. VERBÁSCUM, L. MULLEIN.

Calyx 5-parted. Corolla 5-lobed, open or concave, wheel-shaped ; the lobes broad and rounded, a little unequal. Stamens 5 ; all the filaments, or the 3 upper, woolly. Style flattened at the apex. Pod globular, many-seeded. — Tall and usually woolly biennial herbs, with alternate leaves, those of the stem sessile or decurrent. Flowers in large terminal racemes, ephemeral ; in summer. (The ancient Latin name, altered from *Barbascum*.)

1. **V. THÁPSUS, L.** (COMMON MULLEIN.) *Densely woolly throughout ; stem tall and stout, simple, winged by the decurrent bases of the oblong acute leaves ; flowers (yellow, very rarely white) in a prolonged and very dense cylindrical spike ; lower stamens usually beardless.* — Fields, &c. : common. (Nat. from Eu.)

2. **V. BLATTÁRIA, L.** (MOTH M.) *Green and smoothish, slender ; lower leaves petioled, oblong, doubly serrate, sometimes lyre-shaped, the upper partly clasping ; raceme loose ; filaments all bearded with violet wool.* — Roadsides : not rare eastward. Corolla either yellow, or white with a tinge of purple. (Nat. from Eu.)

3. **V. LYCHNÍTIS, L.** (WHITE M.) *Clothed with a thin powdery wooliness ; stem and branches angled above ; leaves ovate, acute, not decurrent, greenish above ; flowers (yellow, rarely white) in a pyramidal panicle ; filaments with whitish wool.* — Waste places, Penn. to New York : rare : hybridizes spontaneously with the common Mullein. (Adv. from Eu.)

2. LINÀRIA, Tourn. TOAD-FLAX.

Calyx 5-parted. Corolla personate, with the prominent palate often nearly closing the throat, spurred at the base on the lower side. Stamens 4. Pod thin, opening below the summit by one or two pores or chinks. Seeds many. — Herbs, with at least all the upper leaves alternate: fl. in summer. (Name from *Linum*, the Flax, which the leaves of some species resemble.)

* *Leaves sessile, narrow: plant glabrous, erect, leafy.*

1. **L. Canadensis**, Spreng. (WILD TOAD-FLAX.) *Slender annual or biennial, mostly simple, with scattered linear leaves; those from prostrate shoots oblong, crowded, and mostly opposite or whorled; flowers blue (very small), in a slender raceme, short-pedicelled; spur thread-shaped (occasionally wanting).* — Sandy soil: common.

2. **L. vulgàris**, Mill. (TOAD-FLAX. BUTTER-AND-EGGS. RAMSTED.) *Perennial, pale (1°-3° high); leaves alternate, crowded, linear or lanceolate, acutish; flowers crowded in a dense raceme, yellow (1' long); spur awl-shaped; seeds flattened and margined.* — Old fields and roadsides: common eastward, extending westward: a showy but pernicious weed. — The *Peloria* state, with a regular 5-cleft border to the corolla, 5 spurs, and 5 stamens, has been observed in Pennsylvania by *Dr. Darlington*. (Nat. from Eu.)

3. **L. GENISTIFÒLIA**, Mill. *Glaucous perennial, paniculate-branched; leaves lanceolate, acute, often partly clasping; flowers scattered, yellow (smaller than in No. 2); seeds angled and wrinkled.* — Roadsides, New York, near the city (*Prof. H. J. Clark, Lesquereux*). (Adv. from Eu.)

** *Leaves petioled, broad, veiny, hairy: stems procumbent.*

4. **L. ELÁTINE**, Mill. Branching annual; leaves alternate, ovate and halberd-shaped, mostly shorter than the slender axillary peduncles; flowers small, yellow and purplish; sepals lanceolate, very acute. — Fields and banks, eastward: scarce. (Adv. from Eu.)

3. ANTIRRHÌNUM, L. SNAPDRAGON.

Corolla saccate at the base, the throat closed by the large bearded palate. Seeds oblong-truncate. Otherwise nearly as *Linaria*. Corolla commonly showy, resembling the face of an animal or a mask; whence the name (from *ἀντι*, in comparison with, and *ῥίον*, a snout.) Fl. summer and autumn.

1. **A. ORÓNTIUM**, L. A small flowered annual or biennial, low, erect; leaves lance-linear; spike loose, leafy; sepals longer than the purplish or white corolla. — About gardens, and old fields in Virginia. (Adv. from Eu.)

2. **A. MÀJUS**, L. (LARGE SNAPDRAGON.) A large-flowered perennial, with oblong smooth leaves and a glandular-downy raceme; sepals short; corolla 1½' - 2' long, purple or white. — Eastward escaping from gardens. (Adv. from Eu.)

4. SCROPHULÀRIA, Tourn. FIGWORT.

Calyx deeply 5-cleft. Corolla with a somewhat globular tube; the 4 upper lobes of the short border erect (the two upper longer), the lower spreading. Stamens 4, declined, with the anther-cells transverse and confluent into one;

the vestige of the fifth stamen forms a scale-like rudiment at the summit of the tube of the corolla. Pod many-seeded. — Rank herbs, with mostly opposite leaves, and small greenish-purple or lurid flowers in loose cymes, forming a terminal narrow panicle. (So called because a reputed remedy for *scrofula*.)

1. **S. nodosa**, L. Smooth perennial (3°–4° high); stem 4-sided; leaves ovate, oblong, or the upper lanceolate, cut-serrate, rounded or heart-shaped at the base. (*S. Marilandica*, L.) — Damp copses and banks. June–Aug. (Eu.)

5. COLLÍNSIA, Nutt. COLLINSIA.

Calyx deeply 5-cleft. Corolla declined, with the tube saccate or bulging at the base on the upper side, deeply 2-lipped; the upper lip 2-cleft, its lobes partly folded backwards; the lower 3-cleft, its middle lobe keeled and sac-like, enclosing the 4 declined stamens and style. Fifth stamen a slender rudiment. Pod 4-many-seeded. — Slender branching annuals or biennials, with opposite leaves, and handsome party-colored flowers in umbel-like clusters, appearing whorled in the axils of the upper leaves. (Dedicated to the late *Zaccheus Collins*, of Philadelphia, an accurate botanist.)

1. **C. vérna**, Nutt. Slender (6'–20' high); lower leaves ovate; the upper ovate-lanceolate, clasping by the heart-shaped base, toothed; *whorls about 6-flowered; flowers long-peduncled; corolla (blue and white) twice the length of the calyx.* — Moist soil, W. New York to Wisconsin and Kentucky. May, June.

2. **C. parviflora**, Dougl. Small; lower leaves ovate or rounded; the upper oblong-lanceolate, mostly entire; *whorls 2–6-flowered; flowers short-peduncled; the small (blue) corolla scarcely exceeding the calyx.* — Shore of Lake Superior and westward.

6. CHELÒNE, Tourn. TURTLE-HEAD. SNAKE-HEAD.

Calyx of 5 distinct imbricated sepals. Corolla inflated-tubular, with the mouth a little open; the upper lip broad and arched, keeled in the middle, notched at the apex; the lower woolly-bearded in the throat, 3-lobed at the apex, the middle lobe smallest. Stamens 4, with woolly filaments and very woolly heart-shaped anthers; and a fifth sterile filament smaller than the others. Seeds many, wing-margined. — Smooth perennials, with upright branching stems, opposite serrate leaves, and large white or purple flowers, which are nearly sessile in spikes or clusters, and closely imbricated with round-ovate concave bracts and bractlets. (Name from *χελώνη*, a *tortoise*, the corolla resembling in shape the head of a reptile.)

1. **C. glabra**, L. Leaves very short-petioled, lanceolate or lance-oblong, pointed, variable in width, &c.: the flowers white, rose-color, or purple. (Also *C. obliqua*, L., &c.) — Wet places: common. July–Sept. — Called also SHELL-FLOWER, BALMONY, &c.

7. PENTSTÈMON, Mitchell. BEARD-TONGUE. PENTSTEMON.

Calyx 5-parted. Corolla tubular and more or less inflated, or bell-shaped, either decidedly or slightly 2-lipped; the upper lip 2-lobed, and the lower 3-cleft. Stamens 4, declined at the base, ascending above; and a fifth sterile filament

usually as long as the others, either naked or bearded. Seeds numerous, wingless.—Perennials, branched from the base, simple above, with opposite leaves, the upper sessile and mostly clasping. Flowers mostly showy, thyrsoïd or racemose-panicled. (Name from *πέντε*, *five*, and *στήμων*, *stamen*; the fifth stamen being present and conspicuous, although sterile.)

* *Sterile filament bearded down one side: flowers numerous in a loose and somewhat clammy panicle, white or purplish.*

1. **P. pubescens**, Solander. More or less pubescent (1°–3° high); stem-leaves lanceolate from a clasping base, serrate or sometimes entire; *corolla* 2-lipped, gradually widened upwards, somewhat flattened and one-ridged on the upper side, and with 2 infolded lines on the lower which are bearded inside; the *throat* almost closed; lower lip rather longer than the upper; sterile filament densely yellow-bearded. —Varies greatly in the foliage, sometimes nearly glabrous, when it is *P. lævigatus*, Solander, &c.—Dry banks, Connecticut to Wisconsin, and southward. June–Sept.

2. **P. Digitalis**, Nutt. Nearly glabrous (2°–4° high); stem-leaves oblong- or ovate-lanceolate, clasping, serrulate or entire; *corolla* abruptly inflated and almost bell-shaped from a narrow base, beardless, with the *throat* widely open, the spreading lobes nearly equal; sterile filament sparingly bearded. (Flowers more showy than in the preceding, mostly white, over 1' long). — Rich soil, Alexandria, Virginia (*A. H. Curtiss*), to Illinois, and southward. June–Aug.

** *Sterile filament nearly beardless, dilated and hooked at the end (rarely wanting).*

3. **P. grandiflorus**, Fraser. Very smooth and glaucous; stems simple (1°–3° high); leaves thickish, ovate or rounded, the upper clasping; flowers (showy, 2' long) on short pedicels, in a long and narrow raceme rather than panicle; *corolla* oblong-bell-shaped, almost regular, bluish or lilac-purple. —Prairies, W. Wisconsin (Falls of St. Anthony, *Lapham*), and westward. June.

8. MÍMULUS, L. MONKEY-FLOWER.

Calyx prismatic, 5-angled, 5-toothed, the upper tooth largest. *Corolla* tubular; the upper lip erect or reflexed-spreading, 2-lobed; the lower spreading, 3-lobed. Stamens 4. Stigma 2-lipped, the lips ovate. Seeds numerous. —Herbs, with opposite leaves, and mostly handsome flowers on solitary axillary and bractless peduncles. (Name from *μῦώ*, *an ape*, on account of the gaping *corolla*.)

* *Erect from a perennial root, glabrous: leaves feather-veined: corolla violet-purple.*

1. **M. ringens**, L. Stem square (1°–2° high); leaves oblong or lanceolate, pointed, clasping by a heart-shaped base, serrate; peduncles longer than the flower; calyx-teeth taper-pointed, nearly equal. —Wet places: common. July–Sept. — Flower 1'–1½' long, rarely white.

2. **M. alatus**, Ait. Stem somewhat winged at the angles; leaves oblong-ovate, tapering into a petiole; peduncles shorter than the calyx, which has very short abruptly pointed teeth: otherwise like the last. — Low grounds, Connecticut to Illinois, and southward.

** *Diffusely spreading: leaves several-nerved and veiny: corolla yellow.*

3. **M. Jamésii**, Torr. Smooth or smoothish; stems creeping at the base; stem-leaves roundish or kidney-shaped, nearly sessile, equalling the peduncles;

calyx ovate, inflated in fruit, the upper tooth much the largest. — In cool springs, N. Michigan, Wisconsin, Illinois, and westward. — Flowers small, little larger than in the cultivated *M. MOSCHATUS* or MUSK-PLANT.

9. CONÒBEA, Aublet. (CAPRÀRIA, Michx.)

Calyx 5-parted, equal. Upper lip of the corolla 3-lobed, the lower 3-parted. Stamens 4, fertile: anthers approximate. Style 2-lobed at the apex, the lobes wedge-form. Seeds numerous. — Low branching herbs, with opposite leaves, and small solitary flowers on axillary 2-bractleted peduncles. (Name unexplained.)

1. *C. multifida*, Benth. Diffusely spreading, much branched, minutely pubescent, annual; leaves petioled, pinnately parted, divisions linear-wedge-shaped; corolla (greenish-white) scarcely longer than the calyx. — River-banks, Ohio to Illinois, and southward; also adventive below Philadelphia. July–Sept.

10. HERPÉSTIS, Gærtn. HERPESTIS.

Calyx 5-parted; the upper division broadest, the innermost often very narrow. Upper lip of the corolla entire, notched or 2-cleft; the lower 3-lobed. Stamens 4, all fertile. Style dilated or 2-lobed at the apex. Seeds numerous. — Low herbs, with opposite leaves, and solitary axillary flowers; in summer: ours rather succulent perennials. (Name from *ἔρπησις*, a creeping thing, the species being chiefly procumbent.)

* *Upper lip of the blue corolla merely notched: leaves many-nerved.*

1. *H. rotundifolia*, Pursh. Nearly smooth, creeping; leaves round-obovate, half clasping ($\frac{1}{2}$ '–1' long); peduncles twice or thrice the length of the calyx, the upper sepal ovate. — Wet places, Illinois and southward.

2. *H. amplexicaulis*, Pursh. Stems hairy, creeping at the base; leaves ovate, clasping; peduncles shorter than the calyx; upper sepal heart-shaped. — Wet places, New Jersey and southward. — Aromatic when bruised.

** *Corolla (bluish) almost equally 5-cleft, the upper lip being 2-parted: calyx 2-bracted: stamens almost equal: leaves nearly nerveless.*

3. *H. Monnièra*, H. B. K. Smooth, somewhat creeping; leaves obovate or wedge-shaped. — Maryland and southward along the coast.

11. GRATIOLA, L. HEDGE-HYSSOP.

Calyx 5-parted, the divisions narrow and nearly equal. Upper lip of the corolla entire or 2-cleft, the lower 3-cleft. Fertile stamens 2, included, posterior; the anterior mere sterile filaments, or wanting. Style dilated or 2-lipped at the apex. Pod 4-valved, many-seeded. — Low herbs, mostly perennials, some apparently annuals, with opposite sessile leaves, and axillary 1-flowered peduncles, usually with 2 bractlets at the base of the calyx. Flowering all summer; all inhabiting wet or damp places. (Name from *gratia*, grace or favor, on account of supposed excellent medicinal properties.)

§ 1. *Antthers with a broad connective: the cells transverse: stems mostly diffusely branched, or creeping at base, soft viscid-pubescent or smooth.*

* *Sterile filaments minute or none: corolla whitish, with the tube yellowish.*

1. **G. Virginiana**, L. Stem clammy-puberulent above (4'–6' high); leaves lanceolate, narrowed at the base, sparingly toothed; peduncles almost equaling the leaves ($\frac{1}{2}$ '–1' long); pod ovoid (2" long). — Very common.

2. **G. sphærocárpa**, Ell. Smooth, rather stout (5'–10' high); leaves lance-ovate or oblong, toothed; peduncles scarcely longer than the calyx and the large (3") globular pod. — Maryland to Illinois, and southward.

** *Sterile filaments slender, tipped with a little head: leaves short ($\frac{1}{2}$ '–1' long).*

3. **G. viscòsa**, Schweinitz. Clammy-pubescent or glandular; leaves ovate-lanceolate or oblong, acute, toothed, mostly shorter than the peduncles; corolla whitish, yellow within. — Kentucky and southward.

4. **G. àurea**, Muhl. Nearly glabrous; leaves lanceolate or oblong-linear, entire, equalling the peduncles; corolla golden-yellow ($\frac{1}{2}$ ' long). — Sandy swamps, N. Vermont and New Hampshire to Virginia, and southward.

§ 2. *Anthers with no broad connective; the cells vertical: hairy plants, with erect rigid and more simple stems, from an apparently annual root: sterile filaments tipped with a head.*

5. **G. pilòsa**, Michx. Leaves ovate or oblong, sparingly toothed, sessile ($\frac{1}{2}$ ' long); flowers nearly sessile; corolla white, scarcely exceeding the calyx. — Low ground, Camden Co., New Jersey (C. E. Smith, C. F. Parker), Maryland, and southward.

12. ILYSÁNTHES, Raf. (LINDÉRNIA, Muhl.)

Calyx 5-parted, nearly equal. Upper lip of the corolla short, erect, 2-lobed; the lower larger and spreading, 3-cleft. Fertile stamens 2, included, posterior; the anterior pair sterile, inserted in the throat of the corolla, 2-lobed, without anthers; one of the lobes glandular; the other smooth, usually short and tooth-like. Style 2-lipped at the apex. Pod ovate or oblong, many-seeded. — Small and smooth annuals, with opposite leaves, and small axillary (purplish) flowers, or the upper racemed, produced all summer. (Name from ἰλύς, mud, or mire, and ἄνθος, flower.)

1. **I. gratioloides**, Benth. (FALSE PIMPERNEL.) Much branched, diffusely spreading (4'–8' high); leaves ovate, rounded, or oblong, sparingly toothed or entire, the upper partly clasping. (Capraria gratioloides, L. Lindernia dilatata, & L. attenuata, Muhl.) — Wet places: common.

13. MICRÁNTHEMUM, Michx. (Including HEMIANTHUS, Nutt.)

Calyx 4-lobed or 4- (rarely 5-) parted. Corolla short, 2-lipped, with the upper lip considerably shorter than the lower, or 1-lipped, the upper lip obsolete; lower lip 3-cleft, the middle lobe longest. Stamens 2, anterior, the short filament with a glandular (mostly basal) appendage: anthers 2-celled, didymous. No sterile filaments. Style short, its apex or the stigma 2-lobed. Pod globular, thin, with a very delicate or evanescent partition, several-many-seeded. — Small, smooth, depressed and tufted or creeping annuals, in mud or shallow water, with opposite and entire rounded or spatulate sessile leaves, and minute

white or purplish flowers solitary in the axils of some of the middle leaves (usually one axil floriferous, that of the other leaf sterile). (Name formed of *μικρός*, *small*, and *ἄνθεμον*, *flower*.)—The section *HEMIÁNTHUS* (from *ἡμι*, *half*, and *ἄνθος*, *flower*) includes the species like ours, of which there are several discovered by C. Wright in Cuba, having the upper lip of the corolla very short or obsolete, and mostly slender or subulate stigmas.

1. *M. Nuttállii*. (*Hemianthus micranthemoides*, *Nutt.*) Branches ascending, 1'-2' high; leaves obovate-spatulate or oval; peduncles at length recurved, about the length of the calyx, which is bell-shaped, 4-toothed and usually split down on one side, in fruit becoming pear-shaped; middle lobe of the corolla linear-oblong, nearly twice the length of the lateral ones; appendage of the stamens nearly as long as the filament itself; stigmas subulate.—Tidal muddy bands of the Delaware River, and southward. Aug. - Oct.

14. LIMOSÉLLA, L. MUDWORT.

Calyx bell-shaped, 5-toothed. Corolla short, widely bell-shaped, 5-cleft, nearly regular. Stamens 4: anthers confluent 1-celled. Style short, club-shaped. Pod globular, many-seeded; the partition thin and vanishing.—Small annuals, growing in mud, usually near the sea-shore, creeping by slender runners, without ascending stems; the entire fleshy leaves in dense clusters around the simple 1-flowered peduncles. Flowers small, white or purplish. (Name a diminutive of *limus*, mud, in which these little plants delight to grow.)

1. *L. aquática*, L.: var. *tenuifolia*, Hoffm. Leaves (with no blade distinct from the petiole) awl-shaped or thread-form. (*L. tenuifolia*, *Nutt.* *L. subulata*, *Ives.*)—In brackish tidal mud, from New Jersey northward. Aug., Sept.—Plant 1'-2' high. (Eu.)

15. SÝNTHYRIS, Benth. SYNTHYRIS.

Calyx 4-parted. Corolla somewhat bell-shaped, variously 2-4-lobed or cleft. Stamens 2, inserted just below the sinuses on each side of the upper lobe of the corolla, occasionally with another pair from the other sinuses, exserted: anther-cells not confluent into one. Style slender: stigma simple. Pod flattened, rounded, obtuse or notched, 2-grooved, 2-celled (rarely 3-lobed and 3-celled), many-seeded, loculicidal; the valves cohering below with the columella.—Perennial herbs, with the simple scape-like stems beset with partly-clasping bract-like alternate leaves, the root-leaves rounded and petioled, crenate. Flowers in a raceme or spike, with bracted pedicels. (Name composed of *σύν*, *together*, and *θύρίς*, *a little door*; evidently in allusion to the closed valves of the pod.)

1. *S. Houghtoniána*, Benth. Hairy; root-leaves round-ovate, heart-shaped; raceme spiked, dense (5'-12'); corolla not longer than the calyx, usually 2-3-parted.—High prairies and hills, Wisconsin, *Houghton*, *Lapham*. Michigan, *Wright*. Illinois, *Mead*. May.—Corolla greenish-white, for the most part deeply 2-parted, with the upper lip entire, a little longer and narrower than the lower, which is 3-toothed; but occasionally 3-parted, with the upper lip notched or 2-lobed. When there are 4 stamens the lower are later than the others.

16. VERÓNICA, L. SPEEDWELL.

Calyx 4- (rarely 3-5-) parted. Corolla wheel-shaped or salver-shaped, the border 4-parted (rarely 5-parted); the lateral lobes or the lower one commonly narrower than the others. Stamens 2, one each side of the upper lobe of the corolla, exserted: anther-cells confluent at the apex. Style entire: stigma single. Pod flattened, obtuse or notched at the apex, 2-celled, few-many-seeded. — Chiefly herbs: leaves mostly opposite or whorled: flowers blue, flesh-color, or white. (Derivation doubtful; perhaps the flower of *St. Veronica*.)

§ 1. *Tall perennials, with mostly whorled leaves: racemes terminal, dense, spiked: bracts very small: tube of the corolla longer than its limb and much longer than the calyx; both sometimes 5-cleft.* (*Leptándra, Nutt.*)

1. **V. Virginica**, L. (CULVER'S-ROOT. CULVER'S PHYSIC.) Smooth or rather downy; stem simple, straight (2°-6° high); leaves whorled in fours to sevens, short-petioled, lanceolate, pointed, finely serrate; spikes paniced; stamens much exserted. — Rich woods, Vermont to Wisconsin and southward: often cultivated. July, Aug. — Corolla small, nearly white. Pod oblong-ovate, not notched, opening by 4 teeth at the apex, many-seeded.

§ 2. *Perennials with opposite usually serrate leaves: flowers in axillary mostly opposite racemes: corolla wheel-shaped (pale blue): pod notched, many-seeded.*

2. **V. Anagállis**, L. (WATER SPEEDWELL.) Smooth, creeping and rooting at the base, then erect; leaves sessile, most of them clasping by a heart-shaped base, ovate-lanceolate, acute, serrate or entire (2' - 3' long); pedicels spreading; pod slightly notched. — Brooks and ditches, especially northward: rare eastward. June - Aug. — Corolla pale blue with purple stripes. (Eu.)

3. **V. Americana**, Schweinitz. (AMERICAN BROOKLIME.) Smooth, decumbent at the base, then erect (8' - 15' high); leaves mostly petioled, ovate or oblong, acutish, serrate, thickish, truncate or slightly heart-shaped at the base; the slender pedicels spreading; pod turgid. (*V. Beccabúnga, Amer. authors.*) — Brooks and ditches: common. June - Aug.

§ 3. *Perennials, with diffuse or ascending branches from a decumbent base: leaves opposite: racemes axillary, from alternate or sometimes opposite axils: corolla wheel-shaped: pod strongly flattened, several-seeded.*

4. **V. scutelláta**, L. (MARSH SPEEDWELL.) Smooth, slender and weak (6' - 12' high); leaves sessile, linear, acute, remotely denticulate; racemes 1 or 2, very slender and zigzag; flowers few and scattered, on elongated spreading or reflexed pedicels; pod very flat, much broader than long, notched at both ends or didymous. — Bogs: common northward. June - Aug. (Eu.)

5. **V. officinális**, L. (COMMON SPEEDWELL.) Pubescent; stem prostrate, rooting at the base; leaves short-petioled, obovate-elliptical or wedge-oblong, obtuse, serrate; racemes densely many-flowered; pedicels shorter than the calyx; pod obovate-triangular, broadly notched. — Dry hills and open woods. July. (Eu.)

§ 4. *Leaves opposite: flowers in a terminal raceme, the lower bracts resembling the stem-leaves: corolla wheel-shaped: pods flat, several-seeded.*

* *Perennials (mostly turning blackish in drying).*

6. **V. alpina**, L. Stem branched from the base, erect, simple (2' - 6' high); leaves elliptical, or the lowest rounded, entire or toothed; nearly sessile;

raceme hairy, few-flowered, crowded; pod obovate, notched. — Alpine summits of the White Mountains of New Hampshire. (Eu.)

7. *V. serpyllifolia*, L. (THYME-LEAVED SPEEDWELL.) Much branched at the creeping base, *nearly smooth*; branches ascending and simple (2' - 4' high); *leaves ovate or oblong, obscurely crenate, the lowest petioled and rounded, the upper passing into lanceolate bracts*; *raceme loose*; pod rounded, broader than long, obtusely notched. — Roadsides and fields: common: introduced and indigenous. May - July. — Corolla whitish, or pale blue, with deeper stripes. (Eu.)

* * *Annuals or biennials: floral leaves like those of the stem, so that the flowers appear to be axillary and solitary: corolla shorter than the calyx.*

8. *V. peregrina*, L. (NECKWEED. PURSLANE SPEEDWELL.) *Nearly smooth, erect (4' - 9' high), branched; lowest leaves petioled, oval-oblong, toothed, thickish; the others sessile, obtuse*; the upper oblong-linear and entire, longer than the almost sessile (whitish) flowers; *pod orbicular, slightly notched, many-seeded*. — Waste and cultivated grounds: common: appearing like an introduced weed. April - June.

9. *V. arvensis*, L. (CORN SPEEDWELL.) Simple or diffusely branched (3' - 8' high), *hairy; lower leaves petioled, ovate, crenate*; the uppermost sessile, lanceolate, entire; peduncles shorter than the calyx; *pod inversely heart-shaped, the lobes rounded*. — Cultivated grounds: rather common. (Nat. from Eu.)

§ 5. *Annuals or biennials (prostrate-spreading, hairy): stem-leaves opposite (all petioled), the upper alternate and bearing solitary peduncled flowers in their axils: corolla wheel-shaped: pod flat: seeds cup-shaped. Rare or local immigrants, and only near the coast, from Massachusetts to Virginia.*

10. *V. agræstis*, L. (FIELD SPEEDWELL.) *Leaves round or ovate, crenate-toothed*; the floral somewhat similar, *about the length of the recurved peduncles*; calyx-lobes oblong; flower small; *ovary many-ovuled, but the nearly orbicular and sharply notched pod 1 - 2-seeded*. — Sandy fields. (Adv. from Eu.)

11. *V. buxbaumii*, Tenore. *Leaves round or heart-ovate, crenately cut-toothed ($\frac{2}{3}$ ' - 1' long), shorter than the peduncles*; flower large (nearly $\frac{1}{2}$ ' wide, blue); calyx-lobes lanceolate, widely spreading in fruit; *pod obcordate-triangular, broadly notched, 16 - 24-seeded*. — Waste grounds. (Adv. from Eu.)

12. *V. hederæfolia*, L. (IVY-LEAVED SPEEDWELL.) *Leaves rounded or heart-shaped, 3 - 7-toothed or lobed, shorter than the peduncles*; calyx-lobes somewhat heart-shaped; flowers small; *pod turgid, 2-lobed, 2 - 4-seeded*. — Shaded places. April - June. (Adv. from Eu.)

17. BUCHNERA, L. BLUE-HEARTS.

Calyx tubular, obscurely nerved, 5-toothed. Corolla salver-form, with a straight or curved tube, and an almost equally 5-cleft limb: the lobes oblong or wedge-obovate, flat. Stamens 4, included, approximate in pairs: anthers one-celled (the other cell wanting). Style club-shaped and entire at the apex. Pod 2-valved, many-seeded. — Perennial rough-hairy herbs (doubtless root-parasitic), turning blackish in drying, with opposite leaves, or the uppermost alternate; the flowers opposite in a terminal spike, bracted and with 2 bractlets. (Named in honor of J. G. Buchner, an early German botanist.)

1. **B. Americana**, L. Rough-hairy; stem wand-like (1° – 2° high); lower leaves obovate-oblong, the others oblong and lanceolate, sparingly and coarsely toothed, veiny; the uppermost linear-lanceolate, entire; spike interrupted; calyx longer than the bracts, one third the length of the deep-purple corolla. — Plains, W. New York to Wisconsin and southward. June–Aug.

18. SEYMÈRIA, Pursh. SEYMERIA.

Calyx bell-shaped, deeply 5-cleft. Corolla with a short and broad tube, not longer than the 5 ovate or oblong nearly equal and spreading lobes. Stamens 4, somewhat equal: anthers approximate by pairs, oblong, 2-celled; the cells equal and pointless. Pod many-seeded. — Erect branching herbs, with the general aspect and character of *Gerardia*: leaves mostly opposite and dissected or pinnatifid, the uppermost alternate and bract-like. Flowers yellow, interruptedly racemed or spiked. (Named for *Henry Seymer*, an English naturalist.)

1. **S. macrophýlla**, Nutt. (MULLEIN-FOXGLOVE.) Rather pubescent (4° – 5° high); leaves large, the lower pinnately divided, with the broadly lanceolate divisions pinnatifid and incised, the upper lanceolate; tube of the corolla incurved, very woolly inside, as are the filaments except their apex; style short, dilated and notched at the point; pod ovate, pointed. — Shady river-banks, Ohio to Illinois, and southwestward. July.

19. GERÁRDIA, L. GERARDIA.

Calyx bell-shaped, 5-toothed or 5-cleft. Corolla campanulate-funnel-form, or somewhat tubular, swelling above, with 5 more or less unequal spreading lobes, the 2 upper usually rather smaller and more united. Stamens 4, strongly didynamous, included, hairy: anthers approaching by pairs, 2-celled; the cells parallel, often pointed at the base. Style elongated, mostly enlarged and flattened at the apex. Pod globular or ovate, pointed, many-seeded. — Erect branching herbs (more or less root-parasitic): stem-leaves opposite, or the upper alternate, the uppermost reduced to bracts and subtending 1-flowered peduncles, which often form a raceme or spike. Flowers showy, purple or yellow; in late summer and autumn. (Dedicated to the celebrated herbalist, *John Gerarde*.)

§ 1. GERARDIA proper. *Calyx-teeth short: corolla purple or rose-color: anthers all alike, nearly pointless: leaves linear, entire.* (Our species are all branching annuals. They still need revision, in connection with those of the Southern States.)

* *Peduncles shorter or moderately longer than the calyx: stem erect.*

1. **G. purpúrea**, L. (PURPLE GERARDIA.) Stem (8'–20' high) with long and rigid widely spreading branches; leaves linear, acute, rough-margined; flowers large (1' long), bright purple, often downy; calyx-teeth sharp-pointed, shorter than the tube. — Low grounds, Maine to Wisconsin and southward, mostly towards the coast.

2. **G. marítima**, Raf. (SEA-SIDE G.) Low (4'–12' high), with shorter branches; leaves rather fleshy and obtuse, as are the short calyx-teeth; corolla $\frac{1}{2}$ ' long. — Salt marshes along the coast.

3. **G. áspera**, Dougl. Sparingly branched (1° – 2° high); leaves long and narrowly linear, rough; pedicels once or twice the length of the calyx, which has

lanceolate acute teeth nearly as long as the tube; corolla larger than in No. 1.—Damp grounds, Illinois, Wisconsin, and westward.

* * *Peduncles long and filiform, commonly exceeding the leaves: stems diffusely branched, slender (8'–20' high): corolla light purple, 5''–7'' long.*

4. **G. tenuifolia**, Vahl. (SLENDER G.) *Leaves narrowly linear, acute, the floral ones mostly like the others; calyx-teeth very short, acute; pod globular, not exceeding the calyx.*—Dry woods: common.

5. **G. setacea**, Walt. *Leaves bristle-shaped, as are the branchlets, or the lower linear; pod ovate, mostly longer than the calyx, which has short setaceous teeth.* (G. Skinneriana, Wood.)—Dry grounds, Penn. to Wisconsin, and southward.

§ 2. **DASYSTOMA**, Raf. *Calyx 5-cleft, the lobes often toothed: corolla yellow; the tube elongated, woolly inside, as well as the anthers and filaments: anthers all alike, scarcely included, the cells awn-pointed at the base: leaves rather large, all of them or only the lower pinnatifid or toothed.* (Perennials.)

6. **G. flava**, L. partly. (DOWNY FALSE FOXGLOVE.) *Pubescent with a fine close down; stem (3°–4° high) mostly simple; leaves ovate-lanceolate or oblong, obtuse, entire, or the lower usually sinuate-toothed or pinnatifid; peduncles very short; calyx-lobes oblong, obtuse, rather shorter than the tube.*—Open woods, especially in the Middle States.—Corolla 1½' long.

7. **G. quercifolia**, Pursh. (SMOOTH FALSE FOXGLOVE.) *Smooth and glaucous (3°–6° high), usually branching; lower leaves commonly twice-pinnatifid; the upper oblong-lanceolate, pinnatifid or entire; peduncles nearly as long as the calyx, the lance-linear acute lobes of which are as long as the at length inflated tube.*—Rich woods, especially southward.—Corolla 2' long.

8. **G. integrifolia**, Gray. *Smooth, not glaucous; stem (1°–2° high) mostly simple; leaves lanceolate, acute, entire, or the lowest obscurely toothed; peduncles shorter than the calyx.* (*Dasytoma quercifolia*, var. ? *integrifolia*, Benth.)—Woods and barrens, Pennsylvania to Illinois, and southward along the mountains.—Corolla 1' long.

9. **G. grandiflora**, Benth. *Minutely downy; stem much branched (3°–4° high); leaves ovate-lanceolate, coarsely toothed or cut, the lower pinnatifid; peduncles rather shorter than the calyx; corolla (2' long) 4 times the length of the broadly lanceolate entire or toothed calyx-lobes.* (*Dasytoma Drummondii*, Benth.)—Oak openings, Wisconsin (*Lapham*), Illinois (*Vasey*) and southward.—Intermediate between *G. flava* and the next.

10. **G. pedicularia**, L. *Smoothish or pubescent, much branched (2°–3° high, very leafy); leaves ovate-lanceolate, pinnatifid, and the lobes cut and toothed; peduncles longer than the hairy mostly serrate calyx-lobes.*—Dry copses: common.

§ 3. **OTOPHYLLA**, Benth. *Calyx deeply 5-cleft, the lobes unequal: corolla purple (rarely white), sparingly hairy inside, as well as the very unequal stamens: anthers pointless, those of the shorter pair much smaller.* (Annuals?)

11. **G. auriculata**, Michx. *Rough-hairy; stem erect, nearly simple (9'–20' high); leaves lanceolate or ovate-lanceolate, sessile, the lower entire, the others with an oblong-lanceolate lobe on each side at the base; flowers nearly sessile in the axils (1' long).*—Low grounds, Penn. to Michigan, Illinois, and southward.

20. CASTILLEIA, Mutis. PAINTED-CUP.

Calyx tubular, flattened, cleft at the summit on the anterior, and usually on the posterior side also; the divisions entire or 2-lobed. Tube of the corolla included in the calyx; its upper lip long and narrow, arched and keeled, flattened laterally, enclosing the 4 unequal stamens; the lower lip short, 3-lobed. Anther-cells oblong-linear, unequal, the outer fixed by the middle, the inner pendulous. Pod many-seeded. — Herbs (root-parasitic), with alternate entire or cut-lobed leaves; the floral ones usually dilated, colored, and more showy than the yellow or purplish spiked flowers. (Dedicated to *Castillejo*, a Spanish botanist.)

1. *C. coccinea*, Spreng. (SCARLET PAINTED-CUP.) Hairy biennial or annual; stem simple; root-leaves clustered; those of the stem incised; the floral 3-cleft, *bright scarlet* towards the summit (rarely yellow); *calyx almost equally 2-cleft, the lobes nearly entire*, about the length of the pale yellow corolla. (*Euchroma coccinea*, Nutt.) — Sandy low grounds: not uncommon. May – Aug.

2. *C. pallida*, Kunth. Smooth or sparingly hairy perennial; at the summit woolly; leaves often incised; the floral oblong or obovate, *whitish*, rarely tinged with purple; *calyx cleft more deeply in front, the divisions 2-cleft*, the ovate-oblong lobes mostly shorter than the whitish corolla. (*C. septentrionalis*, Lindl.) — Alpine region of the White Mountains, New Hampshire, and Green Mountains, Vermont, Lake Superior and northward. Aug. (Eu.)

3. *C. sessiliflora*, Pursh. Hairy and low perennial (6' – 9' high); leaves mostly 3-cleft, with narrow diverging lobes; the floral broader, *not colored*: spike many-flowered, crowded; *calyx deeper cleft in front, the divisions deeply 2-cleft*, shorter than the tube of the long and narrow greenish-yellow corolla; which has the *lobes of the lower lip slender*, pointed, about half the length of the upper. — Prairies, Wisconsin (*Lapham*), Illinois, and westward. — Corolla 2' long.

21. SCHWÁLBEA, Gronov. CHAFF-SEED.

Calyx oblique, tubular, 10 – 12-ribbed, 5-toothed; the posterior tooth much smallest, the 2 anterior united much higher than the others. Upper lip of the corolla arched, oblong, entire; the lower rather shorter, erect, 2-plaited, with 3 very short and broad obtuse lobes. Stamens 4, included in the upper lip: anther-cells equal and parallel, obscurely pointed at the base. Pod ovate, many-seeded. Seeds linear, with a loose chaff-like coat. — A perennial minutely pubescent upright herb (1° – 2° high), with leafy simple stems, terminated by a loose spike of rather large dull purplish-yellow flowers; the leaves alternate, sessile, 3-nerved, entire, ovate or oblong, the upper gradually reduced into narrow bracts. Pedicels very short, with 2 bractlets under the calyx. (Dedicated to *C. G. Schwalbe*, an obscure German botanist.)

1. *S. Americana*, L. — Wet sandy soil, Cape Cod at Sandwich (*B. D. Greene*), near Albany, New York, pine barrens of New Jersey, and southward near the coast. May – July.

22. EUPHRASIA, Tourn. EYEBRIGHT.

Calyx tubular or bell-shaped, 4-cleft. Upper lip of the corolla erect, scarcely arched, 2-lobed, the lobes broad and spreading; lower lip spreading, 3-cleft, the

lobes obtuse or notched. Stamens 4, under the upper lip: anther-cells equal, pointed at the base. Pod oblong, flattened. Seeds numerous. — Herbs with branching stems, and opposite toothed or cut leaves. Flowers small, spiked. (Name *εὐφρασία*, *cheerfulness*, in allusion to its reputed medicinal properties.)

1. **E. officinalis**, L. Low annual; leaves ovate or lanceolate, the lowest crenate, the floral bristly-toothed; lobes of the lower lip of the (whitish, yellowish, or bluish) corolla notched. — Alpine summits of the White Mountains, New Hampshire (*Oakes*), Lake Superior, and northward. A dwarf variety, 1' - 5' high, with very small flowers. (*E. pusilla*, *Godet, mss.*) (Eu.)

23. RHINÁNTHUS, L. YELLOW-RATTLE.

Calyx membranaceous, flattened, much inflated in fruit, 4-toothed. Upper lip of the corolla arched, ovate, obtuse, flattened, entire at the summit, but furnished with a minute tooth on each side below the apex; lower lip 3-lobed. Stamens 4, under the upper lip: anthers approximate, hairy, transverse; the cells equal, pointless. Pod orbicular, flattened. Seeds many, orbicular, winged. — Annual upright herbs, with opposite leaves; the lower oblong or linear; the upper lanceolate, toothed; the floral rounded and cut-serrate with bristly teeth; the solitary yellow flowers nearly sessile in their axils, and crowded in a one-sided spike. (Name composed of *ῥίη*, a snout, and *ἄνθος*, a flower, from the beaked upper lip of the corolla in some species formerly of this genus.)

1. **R. Crista-gállii**, L. (COMMON YELLOW-RATTLE.) Leaves oblong or lanceolate; seeds broadly winged (when ripe they rattle in the inflated calyx, whence the popular name). — Plymouth, Mass. (probably introduced), White Mountains, N. Hampshire (*Tuckerman*), Lake Superior, and northward. (Eu.)

24. PEDICULÀRIS, Tourn. LOUSEWORT.

Calyx various. Corolla strongly 2-lipped; the upper lip arched, flattened, often beaked at the apex; the lower erect at the base, 2-crested above, 3-lobed; the lobes commonly spreading, the lateral ones rounded and larger. Stamens 4, under the upper lip: anthers transverse; the cells equal, pointless. Pod ovate or lanceolate, mostly oblique, several-seeded. — Perennial herbs, with chiefly pinnatifid leaves, the floral bract-like, and rather large flowers in a spike. (Name from *pediculus*, a louse; of no obvious application.)

1. **P. Canadénsis**, L. (COMMON LOUSEWORT. WOOD BETONY.) Hairy; stems simple, clustered (5' - 12' high); leaves scattered; the lowest pinnately parted; the others half-pinnatifid; spike short and dense; calyx split in front, otherwise almost entire, oblique; upper lip of the (dull greenish-yellow and purplish) corolla hooded, incurved, 2-toothed under the apex; pod flat, somewhat sword-shaped. — Copses and banks: common. May - July.

2. **P. lanceolata**, Michx. Stem upright (1° - 3° high), nearly simple, mostly smooth; leaves partly opposite, oblong-lanceolate, doubly cut-toothed; spike crowded; calyx 2-lobed, leafy-crested; upper lip of the (pale yellow) corolla incurved and bearing a short truncate beak at the apex; the lower erect, so as nearly to close the throat; pod ovate, scarcely longer than the calyx. (*P. pallida*, *Pursh.*) — Swamps, Connecticut to Virginia and Wisconsin. Aug., Sept.

25. MELAMPYRUM, Tourn. COW-WHEAT.

Calyx bell-shaped, 4-cleft; the taper lobes sharp-pointed. Tube of the corolla cylindrical, enlarging above; upper lip arched, compressed, straight in front; the lower erect-spreading, biconvex, 3-lobed at the apex. Stamens 4, under the upper lip: anthers approximate, oblong, nearly vertical, hairy; the equal cells minutely pointed at the base. Ovary with 2 ovules in each cell. Pod flattened, oblique, 1-4-seeded. — Erect branching annuals, with opposite leaves, the lower entire, the upper mostly larger and fringed with bristly teeth at the base. Flowers scattered and solitary in the axils of the upper leaves in our species. (Name composed of μέλας, *black*, and πῦρος, *wheat*; from the color of the seeds of field species in Europe, as they appear mixed with grain.)

1. **M. Americanum**, Michx. Leaves lanceolate, short-petioled, the lower entire; the floral ones similar, or abrupt at the base and beset with a few bristly teeth; calyx-teeth linear-awl-shaped, not half the length of the slender tube of the pale greenish-yellow corolla. (M. pratense, var. Americanum, *Benth.*) — Open woods: common. June-Sept. — Plant 6'-12' high. Corolla 5'' long, more slender than in M. pratense, sometimes tinged with purple.

ORDER 69. ACANTHACEÆ. (ACANTHUS FAMILY.)

Chiefly herbs, with opposite simple leaves, didynamous or diandrous stamens, inserted on the tube of the more or less 2-lipped corolla, the lobes of which are convolute or imbricated in the bud; fruit a 2-celled and few- (4-12-) seeded pod; seeds anatropous, without albumen, usually flat and supported by hooked projections of the placenta. — Flowers commonly much bracted. Calyx 5-cleft. Style thread-form: stigma simple or 2-cleft. Pod loculicidal, usually flattened contrary to the valves and partition. Seed with albumen in Elytraria of the Southern States, according to Dr. Feay. Cotyledons broad and flat. — Mucilaginous and slightly bitter, not noxious. A large family in the warmer parts of the world: represented in gardens by THUNBERGIA, which differs from the rest by the globular pod and seeds, the latter not on hooks (*retinacula*); in the Northern States by only two indigenous genera.

1. DIANTHÈRA, Gronov. WATER-WILLOW.

Calyx 5-parted. Corolla deeply 2-lipped; the upper lip erect, notched; the lower spreading, 3-parted, external in the bud. Stamens 2: anthers 2-celled, the cells separated and somewhat unequal. Pod obovate, flattened, contracted at the base into a short stalk, 4-seeded. — Perennial herbs, growing in water or wet places, with entire leaves, and purplish flowers in axillary peduncled spikes or heads. (Name formed of δῖς, *double*, and ἀνθηρά, *anther*; the separated cells giving the appearance of two anthers on each filament.)

1. **D. Americana**, L. Leaves linear-lanceolate, elongated; spikes oblong, dense, long-peduncled. (Justicia pedunculosa, *Michx.* Rhytiglossa, *Nees.*) — Borders of streams and ponds, N. W. Vermont to Wisconsin, Virginia, and southward. July-Sept.

2. RUÉLLIA, L. (DIPTERACÁNTHUS, Nees, & Ed. 2.)

Calyx 5-parted. Corolla funnel-form, the spreading ample border almost equally and regularly 5-cleft, convolute in the bud. Stamens 4, included, didynamous: cells of the somewhat arrow-shaped anthers parallel and nearly equal. Pod narrow, in our species (of the section DIPTERACÁNTHUS) somewhat flattened, contracted and seedless at the base, above 8-12-seeded. Seeds with a mucilaginous coat, when wet under the microscope exhibiting innumerable tapering short bristles, their walls marked with rings or spirals. — Perennials, with rather large and showy blue or purple flowers, mostly in axillary clusters, sometimes also with small flowers precociously close-fertilized in the bud. Calyx often 2-bracteolate. (Named for the early herbalist, *John Ruelle*.)

1. **R. ciliósa**, Pursh. *Hirsute* with soft whitish hairs (1° – 3° high); *leaves nearly sessile, oval or ovate-oblong* ($1'$ – $2'$ long); *flowers 1-3 and almost sessile in the axils; tube of the corolla* ($1'$ – $1\frac{1}{2}'$ long) *fully twice the length of the setaceous calyx-lobes*; the throat short. (*Dipteracanthus ciliosus*, Nees.) — Dry soil, Michigan to Illinois, and southward. June – Sept.

2. **R. strépens**, L. *Glabrous or sparingly pubescent* (1° – 4° high); *leaves narrowed at the base into a petiole, ovate, obovate, or mostly oblong* ($2\frac{1}{2}'$ – $5'$ long); *tube of the corolla* (about $1'$ long) *little longer than the dilated portion, slightly exceeding the lanceolate or linear calyx-lobes*. (*Dipteracanthus strepens*, Nees.) — Flowers 1-5 in each axil, rarely on a slender peduncle, usually almost sessile; sometimes many and closely crowded, then mostly fruiting in the bud, (when it is *D. micránthus*, Engelm. & Gr.). — Rich soil, Pennsylvania to Wisconsin, and southward. July – Sept.

ORDER 70. VERBENÀCÆÆ. (VERVAIN FAMILY.)

Herbs or shrubs, with opposite leaves, more or less 2-lipped or irregular corolla, and didynamous stamens, the 2-4-celled (in Phryma 1-celled) fruit dry or drupaceous, usually splitting when ripe into as many 1-seeded indehiscent nutlets; differing from the following order in the ovary not being 4-lobed, the style therefore terminal, and the plants seldom aromatic or furnishing a volatile oil. — Seeds with a straight embryo and little or no albumen. — A large order in the warmer parts of the world, sparingly represented in cool regions.

Tribe I. VERBENÆÆ. Ovary 2-4-celled, and with an erect anatropous ovule in each cell: radicle inferior.

1. **Verbena**. Flowers in spikes or heads. Calyx tubular. Fruit splitting into 4 nutlets.
2. **Lippia**. Flowers in spikes or heads. Calyx short, 2-cleft. Fruit splitting into 2 nutlets.
3. **Callicarpa**. Flowers in axillary cymes. Calyx short. Fruit berry-like, with 4 nutlets.

Tribe II. PHRYMÆÆ. Ovary 1-celled: ovule erect, orthotropous: radicle superior.

4. **Phryma**. Flowers in slender spikes. Calyx cylindrical, 2-lipped. Fruit an achenium.

1. VERBÈNA, L. VERVAIN.

Calyx tubular, 5-toothed, one of the teeth often shorter than the others. Corolla tubular, often curved, salver-form; the border somewhat unequally 5-cleft.

Stamens included; the upper pair occasionally without anthers. Style slender: stigma capitate. Fruit splitting into 4 seed-like nutlets. — Flowers sessile, in single or often paniced spikes, bracted; produced all summer. (The Latin name for any sacred herb: derivation obscure.) — The species present numerous spontaneous hybrids.

§ 1. *Anthers not appendaged: erect herbs, with slender spikes.*

* *Leaves undivided: root perennial.*

1. **V. angustifolia**, Michx. Low (6'–18' high), often simple; leaves narrowly lanceolate, tapering to the base, sessile, roughish, slightly toothed; spikes few or single; the purple flowers crowded, larger than in the next. — Dry soil, Amherst, Mass., to Wisconsin and southward: rare northward.

2. **V. hastata**, L. (BLUE VERVAIN.) Tall (4°–6° high); *leaves lanceolate or oblong-lanceolate*, taper-pointed, cut-serrate, *petioled*, the lower often lobed and sometimes *hulberd-shaped* at the base; *spikes linear, erect, densely flowered*, corymbed or paniced. (*V. paniculata*, Lam., when the leaves are not lobed.) — Low and waste grounds: common. At the north probably immigrant from the south.

3. **V. urticifolia**, L. (NETTLE-LEAVED or WHITE V.) Rather tall; *leaves oval or oblong-ovate, acute, coarsely serrate, petioled*; *spikes very slender, at length much elongated, with the flowers remote*, loosely paniced, very small, white. — Old fields and roadsides: apparently immigrant.

4. **V. stricta**, Vent. (HOARY V.) *Downy with soft whitish hairs*; stem nearly simple (1° ± 2° high); *leaves sessile, obovate or oblong, serrate*; *spikes thick and very densely flowered*, somewhat clustered, hairy. — Barrens, Ohio to Wisconsin, and southward. — Flowers blue, pretty large.

* * *Leaves cleft or pinnatifid, narrowed at the base: root perennial?*

5. **V. officinalis**, L. (EUROPEAN V.) *Erect, loosely branched* (1°–3° high); *leaves pinnatifid or 3-cleft, oblong-lanceolate*, sessile, smooth above, the lobes cut and toothed; *spikes paniced, very slender*; *bracts small*, much shorter than the very small purplish flowers. (*V. spuria*, L.) — Roadsides; chiefly southward: scarce. (Nat. from Eu.)

6. **V. bracteosa**, Michx. *Widely spreading or procumbent, hairy*; *leaves wedge-lanceolate, cut-pinnatifid or 3-cleft*, short-petioled; *spikes single, remotely flowered*; *bracts large and leafy*, the lower pinnatifid, longer than the small purple flowers. — Waste places, Wisconsin to Kentucky and southward.

§ 2. *Anthers of the longer stamens tipped with a glandular appendage.*

7. **V. Aubletia**, L. *Annual, rather hairy, spreading or ascending*; *leaves obovate-oblong with a wedge-shaped base, 3-cleft and cut or pinnatifid*; *spikes peduncled, flat-topped in flower*; *bracts shorter than the calyx*; *flowers showy, light purple*. — Prairies and rocks, from Illinois southward. Also cultivated.

2. LÍPPIA, L. (ZAPÁNIA, Juss.)

Calyx short, often flattened, 2–4-toothed, or 2-lipped. Corolla 2-lipped: upper lip notched; the lower much larger, 3-lobed. Stamens included. Style slender: stigma obliquely capitate. Fruit 2-celled, 2-seeded. (Dedicated to *Augustus Lippi*, an Italian naturalist and traveller.)

1. **L. lanceolata**, Michx. (FOG-FRUIT.) Procumbent or creeping, roughish, green; leaves oblanceolate or wedge-spatulate, serrate above; peduncles axillary, slender, bearing solitary closely bracted heads of bluish-white flowers; calyx 2-cleft, the divisions sharply keeled, — River-banks, Pennsylvania to Illinois and southward. July — Sept.

3. CALLICÁRPA, L. CALLICARPA.

Calyx 4-5-toothed, short. Corolla tubular-bell-shaped, 4-5-lobed, nearly regular. Stamens 4, nearly equal, exserted: anthers opening at the apex. Style slender, thickened upwards. Fruit a small berry-like drupe, with 4 nutlets. — Shrubs, with scurfy pubescence, and small flowers in axillary cymes. (Name formed of *καλλος*, *beauty*, and *καρπός*, *fruit*.)

1. **C. Americana**, L. (FRENCH MULBERRY.) Leaves ovate-oblong with a tapering base, toothed, whitish beneath; calyx obscurely 4-toothed; fruits violet-color. — Rich soil, Virginia and southward. May — July.

4. PHRÛMA, L. LOPSEED.

Calyx cylindrical, 2-lipped; the upper lip of 3 bristle-awl-shaped teeth; the lower shorter, 2-toothed. Corolla 2-lipped; upper lip notched; the lower much larger, 3-lobed. Stamens included. Style slender: stigma 2-lobed. Fruit dry, in the bottom of the calyx, oblong, 1-celled and 1-seeded! Seed orthotropous. Radicle pointing upwards; cotyledons convolute round their axis. — A perennial herb, with slender branching stems, and coarsely toothed ovate leaves, the lower long-petioled; the small opposite flowers in elongated and slender terminal spikes, reflexed in fruit, and bent close against the axis. Corolla purplish or pale rose-color. (Derivation of the name unknown.)

1. **P. Leptostachya**, L. — Woods and copses: common. July. — Plant (2°-3° high); leaves 3' - 5' long, thin.

ORDER 71. LABIATÆ. (MINT FAMILY.)

Chiefly herbs, with square stems, opposite aromatic leaves, more or less 2-lipped corolla, didynamous or diandrous stamens, and a deeply 4-lobed ovary, which forms in fruit 4 little seed-like nutlets or achenia, surrounding the base of the single style in the bottom of the persistent calyx, each filled with a single erect seed. — Nutlets smooth or barely roughish and fixed by their base, except in the first tribe. Albumen mostly none. Embryo straight (except in *Scutellaria*): radicle at the base of the fruit. Upper lip of the corolla 2-lobed or sometimes entire; the lower 3-lobed. Stamens inserted on the tube of the corolla. Style 2-lobed at the apex. Flowers axillary, chiefly in cymose clusters, these often aggregated in terminal spikes or racemes. Foliage mostly dotted with small glands containing a volatile oil, upon which depends the warmth and aroma of the plants of this large and well-known family. (More abundant in the Old World than the New. One third of our genera and many of the species are merely introduced plants.)

Tribe I. AJUGOIDEÆ. Stamens 4, ascending (curved upwards) and parallel, usually projecting from the notch of the upper side of the (not evidently 2-lipped) 5-lobed corolla. Nutlets reticulated and pitted, obliquely attached by the inside near the base.

* Lobes of the corolla all declined (turned forwards): stamens exerted.

1. **Teucrium.** Lower lobe of the corolla much larger than the others. Calyx 5-toothed.
2. **Trichostema.** Lobes of the corolla scarcely unequal. Calyx 5-cleft, oblique.

** Lobes of the corolla almost equally spreading: stamens nearly included.

3. **Isanthus.** Calyx bell-shaped, 5-cleft, almost equalling the small corolla.

Tribe II. SATUREIÆ. Stamens 4, the inferior pair longer, or only 2, distant, straight, diverging, or converging under the upper lip: anthers 2-celled. Lobes of the corolla flat and spreading.

* Corolla not evidently 2-lipped, but almost equally 4-lobed, small. Stamens erect, distant.

4. **Mentha.** Fertile stamens 4, nearly equal.
5. **Lycopus.** Fertile stamens 2; and often 2 sterile filaments without anthers.

** Corolla more or less 2-lipped; the tube naked (not bearded) within.

+ Stamens only 2, distant: no rudiments of the upper pair.

6. **Cunila.** Calyx very hairy in the throat, equally 5-toothed. Corolla small.
+ + Stamens 4, all with anthers.
7. **Hyssopus.** Calyx tubular, 15-nerved, naked in the throat, equally 5-toothed. Stamens exerted, diverging.
8. **Pycnanthemum.** Calyx ovate or short-tubular, 10-13-nerved, naked in the throat, equally 5-toothed or somewhat 2-lipped. Flowers in dense heads or clusters.
9. **Origanum.** Calyx ovate-bell-shaped, hairy in the throat, 13-nerved, 5-toothed. Stamens diverging. Flowers spiked, and with large colored bracts.
10. **Thymus.** Calyx ovate, nodding in fruit, hairy in the throat, 10-13-nerved, 2-lipped. Stamens distant. Bracts minute. Leaves very small.
11. **Satureia.** Calyx bell-shaped, naked in the throat, 10-nerved, equally 5-toothed. Stamens somewhat ascending.
12. **Calamintha.** Calyx tubular, often hairy in the throat, 13-nerved, 2-lipped. Tube of the corolla straight. Stamens connivent at the summit in pairs under the upper lip.
13. **Melissa.** Calyx tubular-bell-shaped, 2-lipped, flattish on the upper side. Tube of the corolla curved upwards. Stamens curved above, connivent under the erect upper lip.
+ + + Stamens only 2 with anthers, ascending, and a pair of small sterile filaments.
14. **Hedeoma.** Calyx gibbous on the lower side, hairy in the throat. Flowers loose.
* * * Corolla 2-lipped and with a bearded ring inside at the bottom of the enlarged throat.
15. **Collinsonia.** Calyx enlarged and declined in fruit, 2-lipped. Lower lobe of the corolla much larger than the other four. Stamens 2 or 4, long, diverging.

Tribe III. MONARDEÆ. Stamens 2 (sometimes also with mere rudiments of the upper pair), ascending and parallel: anthers either apparently or really 1-celled. Corolla 2-lipped.

16. **Salvia.** Calyx 2-lipped. Anthers with a long connective astride the filament, bearing a linear cell at the upper end, and none or an imperfect cell on the lower.
17. **Monarda.** Calyx tubular and elongated, equally 5-toothed. Anthers of 2 cells confluent into one; the connective inconspicuous.
18. **Blephilia.** Calyx ovate-tubular, 2-lipped. Anthers as in the preceding.

Tribe IV. NEPETEÆ. Stamens 4, the superior (inner) pair longer than the inferior! ascending or diverging. Corolla 2-lipped; the upper lip concave or arched, the lower spreading. Calyx mostly 15-nerved.

19. **Lophanthus.** Stamens divergent; the upper pair curved downwards; the lower ascending: anther-cells nearly parallel.
20. **Nepeta.** Stamens all ascending; the anthers approximate in pairs; the cells at length widely diverging. Calyx more or less curved.

21. **Dracocephalum.** Stamens nearly as in the preceding. Calyx straight, the upper lip or upper tooth commonly larger.
22. **Cedronella.** Stamens all ascending. Anther-cells parallel.
- Tribe V. STACHYDEÆ.** Stamens 4, ascending and parallel; the inferior (outer) pair longer than the superior, except in No. 33. Anthers usually approximate in pairs. Corolla 2-lipped; the upper lip concave or arched. Calyx 5-10-nerved.
- * Calyx not 2-lipped, thin and membranaceous, inflated-bell-shaped in fruit.
23. **Synandra.** Calyx almost equally 4-lobed! Anther-cells widely divergent.
24. **Physostegia.** Calyx 5-toothed or 5-lobed. Anther-cells parallel.
- ** Calyx 2-lipped, closed in fruit.
25. **Brunella.** Calyx nerved and veiny; upper lip flat, 3-toothed, the lower 2-cleft.
26. **Scutellaria.** Calyx with a helmet-like projection on the upper side; the lips entire.
- *** Calyx not 2-lipped, nor the tube inflated, 5-10-toothed.
- + Stamens included in the tube of the corolla.
27. **Marrubium.** Calyx tubular, 5-10-nerved, and with 5 or 10 awl-shaped teeth.
- + + Stamens projecting beyond the tube of the corolla.
- ++ Anthers opening transversely by two unequal valves; the smaller valve ciliate.
28. **Galeopsis.** Calyx tubular-bell-shaped; the 5 teeth spiny-pointed.
- ++ + Anthers opening lengthwise.
29. **Stachys.** Calyx tubular-bell-shaped. Nutlets rounded at the top. Stamens after shedding the pollen often turned downward.
30. **Leonurus.** Calyx top-shaped, the rigid and spiny-pointed teeth soon spreading. Nutlets truncate and acutely 3-angled at the top.
31. **Lamium.** Calyx-teeth not spiny-pointed. Nutlets sharply 3-angled, truncate at the top.
32. **Ballota.** Calyx somewhat funnel-form, the 5-10 teeth united at the base into a spreading border. Nutlets roundish at the top. Upper lip of the corolla erect.
33. **Phlomis.** Calyx tubular, the 5 teeth abruptly awned. Upper lip of the corolla arched.

1. TEUCRIUM, L. GERMANDER.

Calyx 5-toothed. Corolla with the 4 upper lobes nearly equal, oblong, turned forward, so that there seems to be no upper lip; the lower one much larger. Stamens 4, exerted from the deep cleft between the 2 upper lobes of the corolla: anther-cells confluent. (Named for *Teucer*, king of Troy.)

1. **T. Canadense**, L. (AMERICAN GERMANDER. WOOD SAGE.) Herbaceous perennial, downy; stem erect (1°-3° high); leaves ovate-lanceolate, serrate, rounded at the base, short-petioled, hoary underneath; the floral scarcely longer than the oblique unequally-toothed calyx; whorls about 6-flowered, crowded in a long and simple wand-like spike.—Low grounds: not rare. July-Sept.—Corolla pale purple, rarely white.

2. TRICHOSTEMA, L. BLUE CURLS.

Calyx bell-shaped, oblique, deeply 5-cleft; the 3 upper teeth elongated and partly united, the 2 lower very short. Corolla 5-lobed; the lobes narrowly oblong, declined, nearly equal in length; the 3 lower more or less united. Stamens 4, with very long capillary filaments, exerted much beyond the corolla, curved: anther-cells divergent and at length confluent.—Low annuals, somewhat clammy-glandular and balsamic, branched, with entire leaves, and mostly solitary 1-flowered pedicels terminating the branches, becoming lateral by the production of axillary branchlets, and the flower appearing to be reversed,

namely, the short teeth of the calyx upward, &c. Corolla blue, varying to purple, rarely white, small; in summer and autumn. (Name composed of *θρίξ*, hair, and *στῆμα*, stamen, from the capillary filaments.)

1. **T. dichótomum**, L. (BASTARD PENNYROYAL.) *Leaves lance-oblong or rhombic-lanceolate*, rarely lance-linear, short-petioled. — Sandy fields, New England to Kentucky, and southward, chiefly eastward. — Stamens $\frac{1}{2}$ ' long. Corolla blue: a pink variety near Hartford, Conn., *C. H. Olmstead*.

2. **T. lineàre**, Nutt. *Leaves linear*, nearly smooth. — Sandy pine barrens of New Jersey, and southward. — More slender and less forked than the last.

3. ISÁNTHUS, Michx. FALSE PENNYROYAL.

Calyx bell-shaped, 5-lobed, equal, enlarged in fruit. Corolla little longer than the calyx; the border bell-shaped, with 5 nearly equal and obovate spreading lobes. Stamens 4, slightly didynamous, incurved-ascending, scarcely exceeding the corolla. — A low, much branched annual, clammy-pubescent, with nearly entire lance-oblong 3-nerved leaves, and small pale blue flowers on axillary 1-3-flowered peduncles. (Name from *ἴσος*, equal, and *ἄνθος*, flower, referring to the almost regular corolla.)

1. **I. cæruleus**, Michx. — Gravelly banks and fields, Maine to Illinois, and southward. July, Aug. — Corolla 2" long.

4. MÉNTHA, L. MINT.

Calyx bell-shaped or tubular, 5-toothed, equal or nearly so. Corolla with a short included tube; the bell-shaped border somewhat equally 4-cleft; the upper lobe broadest, entire or notched. Stamens 4, equal, erect, distant. — Odeorous perennial herbs, with the small flowers mostly in close clusters, forming axillary capitate whorls, sometimes approximated in interrupted spikes; produced in summer; of two sorts as to the length of the stamens in most species. Corolla pale purple or whitish. (*Μίνθη* of Theophrastus, from a Nymph of that name, fabled to have been changed into Mint by the jealous Proserpine.)

* *Inflorescence terminal, forming narrow spikes: leaves sessile or nearly so.*

1. **M. ROTUNDIFÓLIA**, L. (ROUND-LEAVED MINT.) *Soft-hairy or downy; leaves round-ovate* and somewhat heart-shaped, *rugose*, crenate-toothed. — Wet places, Cumberland, Maine, *J. Blake*. (Adv. from Eu.)

2. **M. VÍRIDIS**, L. (SPEARMINT.) *Nearly smooth; leaves oblong- or ovate-lanceolate*, unequally serrate. — Wet places: common. (Nat. from Eu.)

* * *Inflorescence mostly terminal, forming interrupted spikes or heads: leaves petioled.*

3. **M. PIPERÍTA**, L. (PEPPERMINT.) *Smooth; leaves ovate-oblong, acute; spikes loose*. — Var. *SUBHIRSUTA*, Benth., has the petioles, veins of the leaves, &c. rather hairy. — Low grounds, and along brooks: less naturalized than the last; and like it multiplying rapidly by running under-ground shoots. (Nat. from Eu.)

4. **M. AQUÁTICA**, L. (WATER MINT.) *Pubescent or smoothish; leaves ovate or round-ovate; flowers in a terminal globular or interrupted and oblong head*, often with one or more clusters in the axils of the upper leaves; calyx and

usually the pedicels hairy. The common form has the stems *hairy downwards*. — Muddy shores, Nanticoke River, Delaware, *W. M. Canby*. — Var. *GLABRATA*, Benth., a nearly smooth form, differing from Peppermint in the rounder leaves and spike of 2 or 3 rounded heads (*M. citræta*, *Ehrh.*). — Litchfield, Connecticut, *Dr. T. F. Allen*. (Nat. from Eu.)

* * * *Inflorescence axillary, the globular whorls or clusters all in the axils of the leaves, the uppermost axils not flower-bearing: leaves more or less petioled, toothed. (The species apparently run together.)*

5. *M. SATIVA*, L. (WHORLED MINT.) *Stem hairy downwards; leaves ovate; calyx oblong-cylindrical with very slender teeth. Intermediate between the last and the next, apparently runs into both. — River-banks, Lancaster, Penn., Prof. Porter. (Adv. from Eu.)*

6. *M. ARVENSIS*, L. (CORN MINT.) *Lower and smaller-leaved than the last; calyx bell-shaped, the teeth short and broader. — Moist fields: rare. (Adv. from Eu.)*

7. *M. Canadensis*, L. (WILD MINT.) *Leaves varying from ovate-oblong to lanceolate, tapering to both ends; calyx oblong-bell-shaped, the teeth rather short; hairs on the stem when present not conspicuously reflexed. The commoner form is more or less hairy, and has nearly the odor of Pennyroyal. — Var. GLABRATA, Benth. (M. borealis, Michx.) is smoothish, "the scent pleasanter, more like that of Monarda." (Prof. Porter.) — Shady wet places, New England to Kentucky and northward.*

5. LÝCOPUS, L. WATER HOREHOUND.

Calyx bell-shaped, 4-5-toothed, naked in the throat. Corolla bell-shaped, scarcely longer than the calyx, nearly equally 4-lobed. Stamens 2, distant; the upper pair either sterile rudiments or wanting. Nutlets with thickened margins. — Perennial low herbs, resembling Mints, with sharply toothed or pinnatifid leaves, the floral ones similar and much longer than the dense axillary whorls of small mostly white flowers; in summer. (Name compounded of *λύκος*, a wolf, and *πούς*, foot, from some fancied likeness in the leaves.)

1. *L. Virginicus*, L. (BUGLE-WEED.) *Stem obtusely 4-angled (6' - 18' high), producing long and slender runners from the base; leaves oblong or ovate-lanceolate, toothed, entire towards the base, short-petioled; calyx-teeth 4, ovate, bluntish and pointless. — Shady moist places: common, especially northward. — Smooth, often purplish, with small capitate clusters of very small flowers. — The depauperate, few-flowered form, often tuberiferous at base (L. uniflorus, Michx., and L. pumilus, Vahl), Lake Superior and northward.*

2. *L. Europæus*, L. *Stem sharply 4-angled (1° - 3° high); leaves ovate-oblong or oblong-lanceolate, sinuate-toothed or pinnatifid, usually more or less petioled; whorls many-flowered; calyx-teeth 5, triangular-lanceolate, tapering to a rigid very sharp point; nutlets (smooth or glandular-roughened at the top) equaling or exceeding the calyx-tube. (Eu.) — Includes several nominal species (the sterile filaments variable); among them in our district is*

Var. *sessilifolius*. *Nearly smooth, producing slender leafy runners from the decumbent base; leaves oblong-ovate, closely sessile or almost clasping, re-*

motely toothed; sterile filaments minute and slender. — Atsion Creek, New Jersey, *W. M. Canby*. — Nearest the var. *parvifolius*, *Miquel*, from Japan: probably *L. rubellus*, *Mœnch*, is a closely related form.

Var. **integrifolius**. Stems often producing slender runners; leaves oblong-lanceolate, varying to narrowly lanceolate (*L. angustifolius*, *Nutt.*), much acuminate at both ends (2'–4' long), slender-petioled, sharply serrate. — Common westward.

Var. **sinuatus**. (*L. sinuatus*, *Benth.* *L. exaltatus* & *L. sinuatus*, *Ell.*) Much branched, smooth or smoothish; runners short or none; leaves mostly more tapering to both ends than in the European form, varying from cut-toothed to pinnatifid; sterile filaments mostly with a globular or spatulate tip. — Common in wet grounds.

6. CUNILA, L. DITTANY.

Calyx ovate-tubular, equally 5-toothed, very hairy in the throat. Corolla 2-lipped; upper lip erect, flattish, mostly notched; the lower spreading, 3-cleft. Stamens 2, erect, exserted, distant: no sterile filaments. — Perennials, with small white or purplish flowers, in corymbed cymes or clusters. (An ancient Latin name, of unknown origin.)

1. **C. Mariàna**, L. (COMMON DITTANY.) Stems tufted, corymbosely much branched (1° high); leaves smooth, ovate, serrate, rounded or heart-shaped at the base, nearly sessile, dotted (1' long); cymes peduncled; calyx striate. — Dry hills, S. New York to Ohio, Illinois, and southward. July–Sept.

7. HYSSÖPUS, L. HYSSOP.

Calyx tubular, 15-nerved, equally 5-toothed, naked in the throat. Corolla short, 2-lipped; upper lip erect, flat, obscurely notched; the lower 3-cleft, with the middle lobe larger and 2-cleft. Stamens 4, exserted, diverging. — A perennial herb, with wand-like simple branches, lanceolate or linear entire leaves, and blue-purple flowers in small clusters, crowded in a spike. (The ancient name.)

1. **H. OFFICINALIS**, L. — Roadsides, &c., sparingly escaped from gardens. (Adv. from Eu.)

8. PYCNÁNTHUM, Michx. MOUNTAIN MINT. BASIL.

Calyx ovate-oblong or tubular, about 13-nerved, equally 5-toothed, or the three upper teeth more or less united, naked in the throat. Corolla short, more or less 2-lipped; the upper lip straight, nearly flat, entire or slightly notched; the lower 3-cleft, its lobes all ovate and obtuse. Stamens 4, distant, the lower pair rather longer: anther-cells parallel. — Perennial upright herbs, with a pungent mint-like flavor, corymbosely branched above; the floral leaves often whitened; the many-flowered whorls dense, crowded with bracts, and usually forming terminal heads or close cymes. Corolla whitish or purplish, the lips mostly dotted with purple. Fl. summer and early autumn. — Varies, like the Mints, with the stamens exserted or included in different flowers. (Name composed of *πυκνός*, *dense*, and *ἄνθεμον*, *a blossom*, from the dense inflorescence.)

* *Calyx scarcely at all 2-lipped, the teeth and bracts awl-shaped and awn-pointed, rigid, naked, as long as the corolla: flowers in rather dense mostly terminal heads: leaves rigid, slightly petioled.*

1. **P. aristatum**, Michx. Minutely hoary-puberulent (1° – 2° high); leaves ovate-oblong and oblong-lanceolate, acute, sparingly denticulate-serrate ($1'$ – $2'$ long), roundish at the base. — Pine barrens, from New Jersey southward.

Var. **hyssopifolium**, Gray. Leaves narrowly oblong or broadly linear, nearly entire and obtuse. (*P. hyssopifolium*, Benth.) — Virginia and southward.

* * *Calyx 2-lipped from the greater union more or less of the 3 upper teeth, which, with the bracts, are subulate and bearded with some spreading hairs: flowers in dense and compound flattened cymes, which become considerably expanded in fruit: leaves membranaceous, petioled. (Species 2–5 incline to run together.)*

2. **P. Tullia**, Benth. Leaves greener and loosely soft-downy, only the floral ones whitened, otherwise resembling those of the next; cymes dense; bracts much surpassing the flowers, their long awn-like points and the awn-pointed calyx-teeth bearded with long loose hairs. — Dry ground, mountains of Virginia and southward.

3. **P. incanum**, Michx. Leaves ovate-oblong, acute, remotely toothed, downy above and mostly hoary with whitish wool underneath, the uppermost whitened both sides; cymes open; bracts linear-awl-shaped and, with the calyx-teeth, more or less awn-pointed. — Rocky woods and hills, New England to Michigan, and southward. — Plant 2° – 4° high, the taste intermediate between that of Pennyroyal and Spearmint, as in most of the following species. Very variable.

4. **P. clinopodioides**, Torr. & Gr. Leaves oblong-lanceolate, scarcely toothed, short-petioled, not whitened; the upper surface often smooth, the lower as well as the stem downy; cymes contracted; bracts and calyx-teeth short-subulate, the latter nearly one half shorter than the tube. — Dry copses, S. New York to Pennsylvania. Connects No. 3 with No. 5.

* * * *Calyx usually almost equally 5-toothed: flowers crowded in loose heads or dense clusters at the end of the branches and in the uppermost axils; the bracts shorter than the 2-lipped corollas: leaves almost sessile.*

5. **P. Torrèyi**, Benth. Somewhat pubescent; stem strict and nearly simple (2° – 3° high); leaves thin, linear-lanceolate, tapering to both ends (mostly $2'$ long and $2''$ – $3''$ wide), nearly entire; the awl-shaped calyx-teeth and bracts canescent. — Dry soil, S. New York to Pennsylvania. — Intermediate in aspect between No. 4 and No. 8.

6. **P. pilosum**, Nutt. More or less downy with long and soft whitish hairs, much branched above; leaves lanceolate, acute at both ends, or the lower ovate-lanceolate, nearly entire, the floral not whitened; calyx-teeth ovate-lanceolate, acute, and with the bracts hoary-hairy. — Dry hills and plains, W. Pennsylvania to Illinois, and southward.

7. **P. muticum**, Pers. Minutely hoary throughout, or becoming almost smooth, corymbosely much branched (1° – $2\frac{1}{2}^{\circ}$ high); leaves ovate or broadly ovate-lanceolate, varying to lanceolate, rather rigid, acute, rounded or slightly heart-shaped at the base, mostly sessile and minutely sharp-toothed, prominently veined, green when old; the floral ones, bracts, and triangular or ovate calyx-teeth, hoary with a

fine close down. (*Brachystemum verticillatum*, Michx.) — Dry hills, Maine to Ohio, Kentucky, and southward. — Flowers in very dense clusters; the outer bracts ovate-lanceolate and pointed, the others pointless,

* * * * *Calyx equally 5-toothed: flowers collected in dense and globular, often fasciated, small and numerous heads, which are crowded in terminal corymbs: bracts rigid, closely appressed, shorter than the flowers: lips of the corolla very short: leaves narrow, sessile, entire, rigid, crowded and clustered in the axils.*

8. **P. lanceolatum**, Pursh. *Smoothish or minutely pubescent* (2° high); *leaves lanceolate or lance-linear, obtuse at the base; heads downy; calyx-teeth short and triangular.* — Dry thickets: not rare.

9. **P. linifolium**, Pursh. *Smoother and leaves narrower and heads less downy than in the last; the narrower bracts and lance-awl-shaped calyx-teeth pungently pointed.* — S. New England to Illinois, and southward.

* * * * * *Calyx equally 5-toothed: flowers collected in few and solitary large and globular heads (terminal, and in the upper axils of the membranaceous petioled leaves); the bracts loose, ciliate-bearded.*

10. **P. montanum**, Michx. *Stem* (1°–3° high) *and ovate- or oblong-lanceolate serrate leaves glabrous; bracts very acute or awl-pointed, the outermost ovate and leaf-like, the inner linear; teeth of the tubular calyx short and acute.* — Alleghanies, from S. Virginia southward. — Flavor warm and pleasant. Foliage and heads resembling *Monarda*.

9. ORÍGANUM, L. WILD MARJORAM.

Calyx ovate-bell-shaped, hairy in the throat, striate, 5-toothed. Tube of the corolla about the length of the calyx, 2-lipped; the upper lip rather erect and slightly notched; the lower longer, of 3 nearly equal spreading lobes. Stamens 4, exserted, diverging. — Perennials, with nearly entire leaves, and purplish flowers crowded in cylindrical or oblong spikes, imbricated with colored bracts. (An ancient Greek name, said to be composed of *ὄρος*, a mountain, and *γάμος*, delight.)

1. **O. vulgare**, L. Upright, hairy, corymbose at the summit; leaves petioled, round-ovate; bracts ovate, obtuse, purplish. — Dry banks: scarce. June–Oct. (Nat. from Eu.)

10. THÏMUS, L. THYME.

Calyx ovate, 2-lipped, 13-nerved, hairy in the throat; the upper lip 3-toothed, spreading; the lower 2-cleft, with the awl-shaped divisions ciliate. Corolla short, slightly 2-lipped; the upper lip straight and flattish, notched at the apex; the lower 3-cleft. Stamens 4, straight and distant, usually exserted. — Low perennials, with small and entire strongly-veined leaves, and purplish or whitish flowers. (The ancient Greek name of the Thyme, probably from *θύω*, to burn perfume, because it was used for incense.)

1. **T. serpyllum**, L. (CREEPING THYME.) Prostrate; leaves green, flat, ovate, entire, short-petioled, flowers crowded at the end of the branches. — Old fields, Eastern New England and Pennsylvania: rare. (Adv. from Eu.) The GARDEN THYME is **T. vulgaris**, L.

11. SATURĒIA, L. SAVORY.

Calyx bell-shaped, 10-nerved, equally 5-toothed, naked in the throat. Corolla 2-lipped; the upper lip erect, flat, nearly entire, the lower 3-cleft. Stamens 4, somewhat ascending. — Aromatic plants, with narrow entire leaves, often clustered, and somewhat spiked purplish flowers. (The ancient Latin name.)

1. **S. HORTÉNSIS**, L. (SUMMER SAVORY.) Pubescent annual; clusters few-flowered; bracts small or none. — Prairies of Illinois, and rocky islands at the Falls of the Ohio, *Short*: escaped from gardens. (Adv. from Eu.)

12. CALAMÍNTHA, Mœnch. CALAMINTH.

Calyx tubular, 13-nerved, mostly hairy in the throat, 2-lipped; the upper lip 3-cleft, the lower 2-cleft. Corolla with a straight tube and an inflated throat, distinctly 2-lipped; the upper lip erect, flattish, entire or notched; the lower spreading, 3-parted, the middle lobe usually largest. Stamens 4, mostly ascending; the anthers usually approximate in pairs. — Perennials, with mostly purplish or whitish flowers; produced all summer: inflorescence various. (Name composed of *καλός*, beautiful, and *μίνθα*, Mint.)

§ 1. **CALAMÍNTHA**, Mœnch. *Calyx striate, scarcely gibbous at the base: clusters of flowers loose and peduncled in the axils of the leaves, and forming a raceme at the summit: bracts minute.*

1. **C. NÉPETA**, Link. (BASIL-THYME.) Soft hairy; stem ascending (1°–3° high); leaves petioled, broadly ovate, obtuse, crenate; corolla (3" long) about twice the length of the calyx. — Dry hills, Virginia, &c. (Nat. from Eu.)

§ 2. **CALOMELÍSSA**, Benth. *Calyx nearly as § 1: whorls few—several-flowered, sessile; flowers on slender naked pedicels; the bracts at their base linear or oblong, leaf-like.*

2. **C. glabélla**, Benth. Smooth; stems diffuse or spreading (1°–2° long); leaves slightly petioled, oblong or oblong-linear, narrowed at the base (8"–2' long), sparingly toothed, or nearly entire; clusters 6–10-flowered; corolla (purplish, 5"–6" long) fully twice the length of the calyx, the teeth of the latter awl-pointed. (*Cunila glabella*, Michx. *Micromeria*, Benth.) — Limestone banks, near Frankfort, Kentucky (*Short*), and southward.

Var. **Nuttállii**, Gray. Smaller; the flowering stems more upright (5'–9 high), with narrower mostly entire leaves and fewer-flowered clusters; while sterile runners from the base bear ovate thickish leaves only 2"–5" long. (*C. Nuttallii*, Benth. *Micromeria glabella*, var. *angustifolia*, Torr.) — Wet limestone rocks, Niagara Falls to Wisconsin, Central Ohio (*Sullivan*), and southwestward. — Appearing very distinct, but united by southwestern specimens.

§ 3. **CLINOPODIUM**, L. *Calyx more or less gibbous below: clusters sessile and many-flowered, dense, crowded with awl-shaped bracts.*

3. **C. Clinopodium**, Benth. (BASIL.) Hairy, erect (1°–2° high); leaves ovate, petioled, nearly entire; flowers (pale purple) in globular clusters; hairy bracts as long as the calyx. (*Clinopodium vulgare*, L.) — Borders of thickets and fields. Naturalized extensively: but apparently also indigenous about the upper Great Lakes. (Eu.)

13. MELÍSSA, L. BALM.

Calyx with the upper lip flattened and 3-toothed, the lower 2-cleft. Corolla with a recurved-ascending tube. Stamens 4, curved and conniving under the upper lip. Otherwise nearly as Calamintha. — Clusters few-flowered, loose, one-sided, with few and mostly ovate bracts resembling the leaves. (Name from *μήλισσα*, a bee; the flowers yielding abundance of honey.)

1. *M. officinalis*, L. (COMMON BALM.) Upright, branching; leaves broadly ovate, crenate-toothed, lemon-scented; corolla nearly white. — Sparingly escaped from gardens. (Nat. from Eu.)

14. HEDEÒMA, Pers. MOCK PENNYROYAL.

Calyx ovoid or tubular, gibbous on the lower side near the base, 13-nerved, bearded in the throat, 2-lipped; the upper lip 3-toothed, the lower 2-cleft. Corolla 2-lipped; the upper lip erect, flat, notched at the apex; the lower spreading, 3-cleft. Fertile stamens 2; the upper pair reduced to sterile filaments or wanting. — Low, odorous annuals, with small leaves, and loose axillary clusters of flowers (in summer), often forming terminal leafy racemes. (Altered from *Ἡδυόσμον*, an ancient name of Mint, from its sweet scent.)

1. *H. pulegioides*, Pers. (AMERICAN PENNYROYAL.) Erect, branching, hairy; leaves petioled, oblong-ovate, obscurely serrate, the floral similar; whorls few-flowered; corolla (bluish, pubescent) scarcely exceeding the calyx; sterile filaments tipped with a little head. — Open barren woods and fields. — The taste and odor nearly of the true Pennyroyal (*Mentha Pulegium*) of Europe.

2. *H. hispida*, Pursh. Erect, hairy (2'–5' high); leaves sessile, linear, entire, the floral similar and exceeding the flowers; corolla scarcely longer than the ciliate hispid calyx. — Dry hills, W. Illinois and westward.

15. COLLINSÒNIA, L. HORSE-BALM.

Calyx ovate, enlarged and declined in fruit, 2-lipped; upper lip truncate and flattened, 3-toothed, the lower 2-cleft. Corolla elongated, expanded at the throat, somewhat 2-lipped; the 4 upper lobes nearly equal, but the lower much larger and longer, pendent, toothed or lacerate-fringed. Stamens 2 (sometimes 4, the upper pair shorter), much exserted, diverging; anther-cells divergent. — Strong-scented perennials, with large ovate leaves, and yellowish flowers on slender pedicels, in loose and paniced terminal racemes. (Named in honor of *Peter Collinson*, a well-known patron of science and correspondent of Linnæus, who introduced it into England.)

1. *C. Canadensis*, L. (RICH-WEED. STONE-ROOT.) Nearly smooth (1°–3° high); leaves serrate, pointed, petioled (3'–6' long); panicle loose; stamens 2. — Rich moist woods: common. July–Sept. — Corolla 8''–9'' long, lemon-scented.

16. SÁLVIA, L. SAGE.

Calyx naked in the throat, 2-lipped; the upper lip 3-toothed or entire, the lower 2-cleft. Corolla deeply 2-lipped, ringent; the upper lip straight or scythe-shaped, entire or barely notched; the lower spreading or pendent, 3-lobed, the

middle lobe larger. Stamens 2, on short filaments, jointed with the elongated transverse connective, one end of which ascending under the upper lip bears a linear 1-celled (half-) anther, the other usually descending bears an imperfect or deformed (half-) anther. — Flowers mostly large and showy, in spiked, racemed, or paniced whorls, produced in summer. (Name from *salvo*, to save, in allusion to the reputed healing qualities of Sage.)

1. **S. lyrata**, L. (LYRE-LEAVED SAGE.) Low perennial (10'–20' high), somewhat hairy; stem nearly simple and naked; root-leaves obovate, lyre-shaped or sinuate-pinnatifid, sometimes almost entire; those of the stem mostly a single pair, smaller and narrower; the floral oblong-linear, not longer than the calyx; whorls loose and distant, forming an interrupted raceme; upper lip of the blue-purple pubescent corolla short, straight, not vaulted. — Woodlands and meadows, New Jersey to Ohio, Illinois, and southward.

2. **S. urticifolia**, L. (NETTLE-LEAVED SAGE.) Downy with clammy hairs, leafy; leaves rhombic-ovate, pointed, crenate, rounded or slightly heart-shaped at the base, narrowed into a short petiole, the floral nearly similar; whorls remote, many-flowered; upper lip of the blue corolla erect, one third the length of the lower; style bearded. — Woodlands, from Maryland southward. — Corolla 4'' long; the lateral lobes deflexed, the middle notched.

17. MONÁRDA, L. HORSE-MINT.

Calyx tubular, elongated, 15-nerved, nearly equally 5-toothed, usually hairy in the throat. Corolla elongated, with a slightly expanded throat, and a strongly 2-lipped limb; the lips linear or oblong, somewhat equal; the upper erect, entire or slightly notched; the lower spreading, 3-lobed at the apex, the lateral lobes ovate and obtuse, the middle one narrower and slightly notched. Stamens 2, elongated, ascending, inserted in the throat of the corolla: anthers linear (the divaricate cells confluent at the junction). — Odorous erect herbs, with entire or toothed leaves, and pretty large flowers in a few whorled heads, closely surrounded with bracts. (Dedicated to *Nicolas Monardes*, author of many tracts upon medicinal and other useful plants, especially those of the New World, in the latter half of the 16th century.)

* *Stamens and style exerted beyond the narrow acute upper lip of the corolla: root perennial: leaves lance-ovate or oblong, with a rounded or slightly heart-shaped base.*

1. **M. didyma**, L. (OSWEGO TEA.) Somewhat hairy (2° high); leaves petioled, pointed; the floral ones and the large outer bracts tinged with red; calyx smooth, incurved, nearly naked in the throat; corolla smooth (2' long), bright red, showy. — Moist woods by streams, New England to Wisconsin northward, and southward in the Alleghanies: often cultivated (under the name of *Balm* or *Bee-Balm*). July, Aug.

2. **M. fistulosa**, L. (WILD BERGAMOT.) Smoothish or downy; leaves petioled; the uppermost and outer bracts somewhat colored (whitish or purplish); calyx slightly curved, very hairy in the throat; corolla purplish, rose-color, or almost white, smooth or hairy. — Woods and rocky banks, Vermont to Wisconsin, and southward, especially westward. July–Sept. — Very variable in appearance, 2°–5° high; the pale corolla smaller than in the last.

3. **M. Bradburiàna**, Beck. *Leaves nearly sessile, clothed with long soft hairs, especially underneath*; the floral and the outer bracts somewhat heart-shaped, purplish; *calyx smoothish, contracted above, very hairy in the throat, with awl-shaped awned teeth*; corolla smoothish, bearded at the tip of the upper lip, scarcely twice the length of the calyx, pale purplish, the lower lip dotted with purple. — Oak-openings and woods, Ohio to Illinois, and westward. May — July.

* * *Stamens not exceeding the upper lip of the short corolla: annuals or biennials.*

4. **M. punctàta**, L. (HORSE-MINT.) *Minutely downy (2°–3° high)*; leaves petioled, lanceolate, narrowed at the base; bracts lanceolate, obtuse at the base, sessile, yellowish and purple; teeth of the downy calyx short and rigid, awnless; corolla nearly smooth, yellowish, the upper lip spotted with purple, notched at the apex, the tube scarcely exceeding the calyx. — Sandy fields and dry banks, New York to Illinois and southward. July — Sept. — Very odorous and pungent.

18. BLEPHÍLIA, Raf. BLEPHILIA.

Calyx ovoid-tubular, 13-nerved, 2-lipped, naked in the throat; upper lip with 3 awned teeth, the lower with 2 nearly awnless teeth. Corolla inflated in the throat, strongly and nearly equally 2-lipped; the upper lip erect, entire; the lower spreading, 3-cleft, with the lateral lobes ovate and rounded, larger than the oblong and notched middle one. Stamens 2, ascending, exerted (the rudiments of the upper pair minute or none): anthers, &c. as in *Monarda*. — Perennial herbs, with nearly the foliage, &c. of *Monarda*; the small pale bluish purple flowers crowded in axillary and terminal globose capitate whorls; in summer. (Name from *βλεφάρις*, the *eyelash*, in reference to the hairy-fringed bracts and calyx-teeth.)

1. **B. ciliàta**, Raf. *Somewhat downy (1°–2° high)*; *leaves almost sessile, oblong-ovate, narrowed at the base, whitish-downy underneath*; outer bracts ovate, acute, colored, ciliate, as long as the calyx; corolla hairy. (*Monarda ciliata*, L.) — Dry open places, Penn. to Kentucky and Wisconsin.

2. **B. hirsùta**, Benth. *Hairy throughout*; *leaves long-petioled, ovate, pointed, rounded or heart-shaped at the base*; the lower floral ones similar, the uppermost and the bracts *linear-awl-shaped*, shorter than the long-haired calyx; corolla smoothish, pale, with darker purple spots. (*B. nepetoides*, Raf. *Monarda hirsuta*, Pursh.) — Damp rich woods, E. Vermont to Wisconsin and Kentucky. — Plant 2°–3° high, with spreading branches, and numerous close whorls, the lower remote.

19. LOPHÁNTHUS, Benth. GIANT HYSSOP.

Calyx tubular-bell-shaped, 15-nerved, oblique, 5-toothed, the upper teeth rather longer than the others. Corolla 2-lipped; the upper lip nearly erect, 2-lobed; the lower somewhat spreading, 3-cleft, with the middle lobe crenate. Stamens 4, exerted; the upper pair declined; the lower and shorter pair ascending, so that the pairs cross. Anther-cells nearly parallel. — Perennial tall herbs, with petioled serrate leaves, and small flowers crowded in interrupted terminal spikes; in summer. (Name from *λόφος*, a *crest*, and *ἄνθος*, a *flower*.)

1. **L. nepetoides**, Benth. *Smooth*, or nearly so; leaves ovate, somewhat pointed, coarsely crenate-toothed (2' - 4' long); *calyx-teeth ovate, rather obtuse, little shorter than the pale greenish-yellow corolla.* — Borders of woods, W. Vermont to Wisconsin, and southward. — Stem stout, 4° - 6° high, sharply 4-angled. Spikes 2' - 6' long, crowded with the ovate pointed bracts.

2. **L. scrophulariæfolius**, Benth. Stem (obtusely 4-angled) and lower surface of the ovate or somewhat heart-shaped acute leaves more or less *pubescent*; *calyx-teeth lanceolate, acute, shorter than the purplish corolla* (spikes 4' - 15' long): otherwise like the last. — Same geographical range.

3. **L. anisatus**, Benth. (ANISE HYSSOP.) *Smooth*, but the ovate acute leaves *glaucous-white underneath* with minute down; *calyx-teeth lanceolate, acute.* — Plains, Wisconsin and northwestward. — Foliage with the scent of anise.

20. NÉPETA, L. CAT-MINT.

Calyx tubular, often incurved, obliquely 5-toothed. Corolla dilated in the throat, 2-lipped; the upper lip erect, rather concave, notched or 2-cleft; the lower spreading, 3-cleft, the middle lobe largest, either 2-lobed or entire. Stamens 4, ascending under the upper lip, the lower pair shorter. Anthers approximate in pairs; the cells divergent. — Perennial herbs. (The Latin name, thought to be derived from *Nepete*, an Etrurian city.)

§ 1. *Cymose clusters rather dense and many-flowered, forming interrupted spikes or racemes: upper floral leaves small and bract-like.*

1. **N. CATARIA**, L. (CATNIP.) Downy, erect, branched; leaves heart-shaped, oblong, deeply crenate, whitish-downy underneath; corolla whitish, dotted with purple. — Near dwellings: a very common weed. July - Sept. (Adv. from Eu.)

§ 2. **GLECHÔMA**, L. *Leaves all alike: the axillary clusters loosely few-flowered.*

2. **N. GLECHÔMA**, Benth. (GROUND IVY. GILL.) Creeping and trailing; leaves petioled, round kidney-shaped, crenate, green both sides; corolla thrice the length of the calyx, light blue. (*Glechoma hederacea*, L.) — Damp waste grounds near dwellings. May - Aug. — Anther-cells diverging at a right angle, each pair approximate and forming a cross. (Adv. from Eu.)

21. DRACOCÉPHALUM, L. DRAGON-HEAD.

Calyx tubular, 13 - 15-nerved, straight, 5-toothed; the upper tooth usually much largest. Corolla 2-lipped; the upper lip slightly arched and notched; the lower spreading, 3-cleft, with its middle lobe largest and 2-cleft or notched at the end. Stamens 4, ascending under the upper lip; the lower pair shorter. Anthers approximate by pairs, the cells divergent. — Whorls many-flowered, mostly spiked or capitate, and with awn-toothed or fringed leafy bracts. (Name from *δράκων*, a dragon, and *κεφαλή*, head, alluding to the form of the corolla in the original species.)

1. **D. parviflorum**, Nutt. Annual or biennial; stem erect, leafy (8' - 20' high); leaves ovate-lanceolate, sharply cut-toothed, petioled; whorls crowded in a terminal head or spike; upper tooth of the calyx ovate, nearly equalling

the bluish small slender corolla. — Rocky places, from St. Lawrence Co., New York, to Wisconsin along the Great Lakes, northward. May — Aug.

22. CEDRONÉLLA, Moench. CEDRONELLA.

Calyx rather obliquely 5-toothed, many-nerved. Corolla ample, expanded at the throat, 2-lipped; the upper lip flattish or concave, 2-lobed; the lower 3-cleft, spreading, the middle lobe largest. Stamens 4, ascending; the lower pair shorter. Anther-cells parallel. — Sweet-scented perennials, with pale purplish flowers. (Name a diminutive of *κέδριον*, oil of Cedar, from the aromatic leaves of the original species, *C. triphylla*, the *Balm-of-Gilead* of English gardens.)

1. *C. cordata*, Benth. Low, with slender runners, hairy; leaves broadly heart-shaped, crenate, petioled, the floral shorter than the calyx; whorls few-flowered, at the summit of short ascending stems; corolla hairy inside ($1\frac{1}{2}'$ long); stamens shorter than the upper lip. (*Dracocephalum cordatum*, Nutt.) — Low shady banks, W. Penn. to Kentucky, and southward along the mountains. June.

23. SYNÁNDRA, Nutt. SYNANDRA.

Calyx bell-shaped, inflated, membranaceous, irregularly veiny, almost equally 4-toothed! Corolla with a long tube, much expanded above and at the throat; the upper lip slightly arched, entire; the lower spreading and 3-cleft, with ovate lobes, the middle one broadest and notched at the end. Stamens 4, ascending: filaments hairy: anthers approximate in pairs under the upper lip; the two upper each with one fertile and one smaller sterile cell, the latter cohering with each other (whence the name; from *σύν*, together, and *άνθη*, for anther).

1. *S. grandiflora*, Nutt. — Shaded banks, Ohio, Kentucky, and southward. June. — A perennial? hairy herb, 1° high. Lower leaves long-petioled, broadly ovate, heart-shaped, crenate, thin; the floral sessile, gradually reduced to bracts, each with a single sessile flower. Corolla $1\frac{1}{2}'$ long, yellowish-white.

24. PHYSOSTÈGIA, Benth. FALSE DRAGON-HEAD.

Calyx nearly equally 5-toothed, obscurely 10-nerved, short-tubular or bell-shaped, more or less enlarged and slightly inflated in fruit. Corolla funnel-form with a much inflated throat, 2-lipped; the upper lip erect, nearly entire; the lower 3-parted, spreading, small: its middle lobe larger, broad and rounded, notched. Stamens 4, ascending under the upper lip: anthers approximate; the cells parallel. — Smooth perennials, with upright wand-like stems, and sessile lanceolate or oblong mostly serrate leaves. Flowers large and showy, rose or flesh-color variegated with purple, opposite, crowded in simple or paniced terminal leafless spikes. (Name from *φύσα*, a bladder, and *στέγη*, a covering.)

1. *P. Virginiana*, Benth. Corolla 1' or more long, gaping, the upper lip slightly arching; calyx-tube oblong- or obconical-bell-shaped. (*Dracocephalum Virginianum*, L.) — Wet banks, &c., W. New York to Wisconsin and southward. July — Sept. — Varies from 1° — 4° high, stout or slender; the leaves from oblong-obovate (the lower) to narrowly lanceolate, and from very sharply toothed to nearly entire; the flowers either crowded or scattered.

25. **BRUNÉLLA**, Tourn. (*Prunella*, L.) SELF-HEAL.

Calyx tubular-bell-shaped, somewhat 10-nerved and reticulated-veiny, flattened on the upper side, naked in the throat, closed in fruit, 2-lipped; the upper lip broad and flat, truncate, with 3 short teeth; the lower 2-cleft. Corolla ascending, slightly contracted at the throat, and dilated at the lower side just beneath it, 2-lipped; the upper lip erect, arched, entire; the lower reflexed-spreading, 3-cleft; its lateral lobes oblong; the middle one rounded, concave, crenulate. Stamens 4, ascending under the upper lip: filaments 2-toothed at the apex, the lower tooth bearing the anther. Anthers approximate in pairs, their cells diverging. — Low perennials, with nearly simple stems, and 3-flowered clusters of flowers sessile in the axils of round and bract-like membranaceous floral leaves, imbricated in a close spike or head. (Name said to be taken from the German *braune*, a disease of the throat, for which this plant was a reputed remedy.)

2. **B. vulgaris**, L. (COMMON SELF-HEAL OR HEAL-ALL.) Leaves ovate-oblong, entire or toothed, petioled, hairy or smoothish; corolla (violet or flesh-color, rarely white) not twice the length of the purplish calyx. — Woods and fields: common. June – Sept. (Eu.)

26. **SCUTELLÀRIA**, L. SKULLCAP.

Calyx bell-shaped in flower, 2-lipped; the lips entire, closed in fruit, the upper with a helmet-like at length concave and enlarged appendage on the back (the upper sepal); calyx splitting to the base at maturity, the upper lip usually falling away. Corolla with an elongated curved ascending tube, dilated at the throat, 2-lipped; the upper lip arched, entire or barely notched; the lateral lobes mostly connected with the upper rather than the lower lip; the lower lobe or lip spreading and convex, notched at the apex. Stamens 4, ascending under the upper lip: anthers approximate in pairs, ciliate or bearded; those of the lower stamens 1-celled (halved), of the upper 2-celled and heart-shaped. — Bitter perennial herbs, not aromatic, with axillary or else spiked or racemed flowers; in summer: the short peduncles or pedicels chiefly opposite, 1-flowered, often 1-sided. (Name from *scutella*, a dish, in allusion to the form of the appendage to the fruiting calyx.)

* *Flowers (blue) in terminal (single or paniced) racemes; the floral leaves, except the lower ones, being small, and reduced to bracts.*

+ *Lips short, nearly equal in length, the lateral lobes rather distinct, and almost as long as the straightish or scarcely incurved upper lip: leaves on slender petioles.*

1. **S. versicolor**, Nutt. *Soft hairy*, the hairs of the inflorescence, &c. partly viscid-glandular; stem mostly erect (1°–3° high); *leaves ovate or round-ovate, chiefly heart-shaped*, crenate-toothed, very veiny, *rugose*, the floral reduced to broadly ovate entire bracts about equalling the glandular-hairy calyx; racemes mostly simple. — River-banks, &c., Penn. to Wisconsin and southward. — Corolla 9" long, with a slender tube, below whitish, the lower lip purple-spotted; the upper deep blue; the lateral lobes belonging as much to the lower as to the upper lip. — *S. saxatilis*, var. ? *pilosior*, *Benth.*, is probably a smaller

form of this, as is *S. rugosa*, Wood. (The latter from Harper's Ferry, Dr. Aikin, according to Wood.)

2. *S. saxatilis*, Riddell. *Smoothish or slightly hairy*; stem weak, ascending (6'–18' long), often producing runners, branched; *leaves ovate or ovate-oblong and mostly heart-shaped*, coarsely crenate-toothed (1'–2' long), *thin, obtuse*; upper bracts oblong or ovate, small; racemes loose.—Moist shaded banks, S. Ohio, Virginia, and Kentucky, and southward in the mountains.—Corolla 8" long, the lateral lobes connected with the straightish upper lip.

+ + *Lateral lobes of the corolla small, much shorter than the decidedly arched or incurved upper lip, and connected with it: stem erect: leaves moderately petioled, except in No. 6.*

3. *S. canescens*, Nutt. Stem branched (2°–4° high) above, with the *panicled many-flowered racemes, flowers, and the lower surface of the ovate or lance-ovate acute* (at the base acute, obtuse, or cordate) *crenate leaves whitish with fine soft down*, often becoming rather glabrous; bracts oblong or lanceolate; upper lip of the corolla shorter than the lower.—Rich ground, Penn. to Illinois and southward.—Corolla 8" long.

4. *S. serrata*, Andrews. *Green and nearly glabrous*; stem rather simple (1°–3° high), with single loosely-flowered racemes; *leaves serrate, acuminate at both ends, ovate or ovate-oblong*; calyx, &c. somewhat hairy; *lips of the corolla equal in length* (corolla 1' long, the tube more tapering below than in the last, which this resembles).—Woods, Maryland, Illinois, and southward.

5. *S. pilosa*, Michx. *Pubescent with spreading hairs*; stem nearly simple (1°–3° high); *leaves rather distant, crenate, oblong-ovate, obtuse*, varying to roundish-ovate, the lower abrupt or heart-shaped at the base and long-petioled, the upper on short margined petioles, veiny; bracts oblong-spatulate; racemes short, often branched; *corolla* (6"–8" long) *rather narrow*, the lower lip a little shorter. (*S. hirsuta*, Short, is a large form.)—Dry ground, S. New York to Michigan and southward.

6. *S. integrifolia*, L. *Downy all over with a minute hoariness*; stem commonly simple (1°–2° high); *leaves oblong-lanceolate or linear, mostly entire*, obtuse, very short-petioled; raceme often branched; *corolla* (1' long) *much enlarged above, the ample lips equal in length*.—Borders of thickets, from Bridgewater, Mass. (Mr. Howard), to Pennsylvania and southward.

* * *Flowers (blue or violet, short-peduncled) solitary in the axils of the upper mostly sessile leaves, which are similar to the lower ones.*

+ *Corolla* (2"–3" long) *seldom thrice the length of the calyx; the short lips nearly equal in length, the upper lip concave.*

7. *S. nervosa*, Pursh. *Smooth, simple or branched, slender* (10'–20' high); *lower leaves roundish; the middle ones ovate, toothed, somewhat heart-shaped* (1' long); the floral ovate-lanceolate, entire; nerve-like veins prominent beneath. (*S. gracilis*, Nutt.)—Moist thickets, New York to Illinois and southward.

8. *S. parvula*, Michx. *Minutely downy, dwarf* (3'–6' high), branched and spreading; *lowest leaves round-ovate; the others ovate or lance-ovate, obtuse, all entire or nearly so, slightly heart-shaped* (6"–8" long). (*S. ambigua*, Nutt.)—Dry banks, W. New England to Wisconsin and southward. May, June.

+ + Corolla (8''–9'' long), with a slender tube: lower lip large and rather longer than the somewhat arched upper lip: stem simple.

9. *S. galericulata*, L. Smooth or a little downy, erect (1°–2° high); leaves ovate-lanceolate, acute, serrate, roundish and slightly heart-shaped at the base (1'–2' long).—Wet shady places: common northward. (Eu.)

* * * Flowers small (blue, 3'' long), in axillary and often also in terminal one-sided racemes; the lower floral leaves like the others, the upper small and bract-like.

10. *S. lateriflora*, L. Smooth; stem upright, much branched (1°–2° high); leaves lanceolate-ovate or ovate-oblong, pointed, coarsely serrate, rounded at the base, petioled (2'–3' long).—Wet shaded places: common.—A quack having formerly vaunted its virtues as a remedy for hydrophobia, this species bears the name of *Mad-dog Skullicap*.

27. MARRÛBIUM, L. HOREHOUND.

Calyx tubular, 5–10-nerved, nearly equally 5–10-toothed; the teeth more or less spiny-pointed and spreading at maturity. Upper lip of the corolla erect, notched; the lower spreading, 3-cleft, its middle lobe broadest. Stamens 4, included in the tube of the corolla. Nutlets not truncate.—Whitish-woolly bitter-aromatic perennials, branched at the base, with rugose and crenate or cut leaves, and many-flowered axillary whorls. (A name of Pliny, said to be derived from the Hebrew *marrob*, a bitter juice.)

1. *M. vulgare*, L. (COMMON HOREHOUND.) Stems ascending; leaves round-ovate, petioled, crenate-toothed; whorls capitate; calyx with 10 recurved teeth, the alternate ones shorter; corolla small, white.—Escaped from gardens into waste places. (Nat. from Eu.)

28. GALEÓPSIS, L. HEMP-NETTLE.

Calyx tubular-bell-shaped, about 5-nerved, with 5 somewhat equal and spiny-tipped teeth. Corolla dilated at the throat; the upper lip ovate, arched, entire; the lower 3-cleft, spreading; the lateral lobes ovate, the middle one inversely heart-shaped; palate with 2 teeth at the sinuses. Stamens 4, ascending under the upper lip: *anther-cells transversely 2-valved*; the inner valve of each cell bristly-fringed, the outer one larger and naked.—Annuals, with spreading branches, and several—many-flowered whorls in the axils of floral leaves which are nearly like the lower ones. (Name composed of *γαλέη*, a weasel, and *ὄψις*, resemblance, from some fancied likeness of the corolla to the head of a weasel.)

1. *G. tetrahit*, L. (COMMON HEMP-NETTLE.) Stem swollen below the joints, bristly-hairy; leaves ovate, coarsely serrate; corolla purplish, or variegated, about twice the length of the calyx; or, in var. *grandiflora*, 3–4 times the length of the calyx, often yellowish with a purple spot on the lower lip.—Waste places: rather common. Aug., Sept. (Nat. from Eu.)

2. *G. ladanum*, L. (RED H.) Stem smooth or pubescent; leaves oblong-lanceolate, more or less downy; corolla red or rose-color (the throat often spotted with yellow), much exceeding the calyx.—Eastern New England: rare. Aug. (Adv. from Eu.)

29. STACHYS, L. HEDGE-NETTLE.

Calyx tubular-bell-shaped, 5-10-nerved, equally 5-toothed, or the upper teeth united to form an upper lip. Corolla not dilated at the throat; the upper lip erect or rather spreading, often arched, entire or nearly so; the lower usually longer and spreading, 3-lobed, with the middle lobe largest and nearly entire. Stamens 4, ascending under the upper lip (often reflexed on the throat after flowering): anthers approximate in pairs. Nutlets obtuse, not truncate. — Whorls 2-many-flowered, approximate in a terminal raceme or spike (whence the name, from *σπάχης*, a spike). Flowering in summer.

* *Root annual: stems decumbent, low.*

1. **S. ARVÉNSIS**, L. (WOUNDWORT.) Hairy; leaves petioled, ovate, obtuse, crenate, heart-shaped at the base; axillary whorls 4-6-flowered, distant; corolla (purplish) scarcely longer than the soon declined unarmed calyx. — Waste places, E. Massachusetts: scarce. (Adv. from Eu.)

** *Root perennial: stem erect.*

2. **S. PALÚSTRIS**, L. Stem 4-angled (2° - 3° high), leafy, hirsute with spreading or reflexed hairs, especially on the angles; leaves sessile, or the lower short-petioled, oblong- or ovate-lanceolate, crenately serrate, rounded or heart-shaped at the base, downy or hairy-pubescent, obtusish ($2'$ - $4'$ long), the upper floral ones shorter than the nearly sessile calyx; whorls 6-10-flowered, the upper crowded into an interrupted spike; calyx hispid; the lance-subulate teeth somewhat spiny, half the length of the purple corolla, diverging in fruit. — Wet banks of streams, &c., mostly northward. (Eu.) — To this, for the present, we must refer all the following as varieties, different as some of them are: —

Var. **áspera**. (*S. aspera*, Michx.) Stem more commonly smooth on the sides, the angles beset with stiff reflexed bristles; leaves hairy or smoothish, pointed, the lower petioled, the lower floral as long as the flowers; spike often slender and more interrupted; calyx-tube rather narrower and the teeth more awl-shaped and spiny. — Common in wet grounds. — This passes into

Var. **glàbra**. (*S. glabra*, Riddell, *suppl. cat. Ohio pl.* 1836.) More slender, smooth and glabrous throughout, or with few bristly hairs; leaves oblong- or ovate-lanceolate, taper-pointed, more sharply toothed, mostly rounded or truncate at the base, all petioled. — W. New York to Michigan and southwestward.

Var. **cordàta**. (*S. cordata*, Riddell, *l. c.* *S. Nuttallii*, Shuttlew.) Stem beset with spreading or reflexed bristly hairs; leaves hairy or smoothish, oblong, heart-shaped at the narrowed base, all more or less petioled; calyx-teeth sometimes shorter. — Common westward and southward.

3. **S. hyssopifòlia**, Michx. Smooth and glabrous, or nearly so; stems slender (1° high), the angles sometimes reflexed-bristly; leaves linear-oblong, or narrowly linear, sessile, obscurely toothed towards the apex; whorls 4-6-flowered, rather distant; corolla (light purple) twice or thrice the length of the triangular-awl-shaped spreading calyx-teeth. — Wet sandy places, Massachusetts to Virginia; also Michigan, Illinois and southward.

BETÓNICA OFFICINÁLIS, the WOOD BETONY of Europe, — of a genus hardly distinct from *Stachys*, — was found by C. J. Sprague in a thicket at Newton, Massachusetts.

30. LEONÛRUS, L. MOTHERWORT.

Calyx top-shaped, 5-nerved, with 5 nearly equal teeth which are awl-shaped, and when old rather spiny-pointed and spreading. Upper lip of the corolla oblong and entire, somewhat arched; the lower spreading, 3-lobed; its middle lobe larger, broad and inversely heart-shaped, the lateral ones oblong. Stamens 4, ascending under the upper lip: anthers approximate in pairs, the valves naked. Nutlets truncate and sharply 3-angled. — Upright herbs, with cut-lobed leaves, and close whorls of flowers in their axils; in summer. (Name from λέων, a lion, and οὐρά, tail, i. e. *Lion's-tail*.)

1. **L. CARDIACA**, L. (COMMON MOTHERWORT.) Tall perennial; leaves long-petioled; the lower rounded, palmately lobed; the floral wedge-shaped at the base, 3-cleft, the lobes lanceolate; upper lip of the pale purple corolla bearded. — Waste places, around dwellings. (Nat. from Eu.)

2. **L. MARRUBIÁSTRUM**, L. Tall biennial, with elongated branches; stem-leaves oblong-ovate, coarsely toothed; corolla (whitish) shorter than the calyx-teeth; the tube naked within; lower lip rather erect. — Roadsides, Pennsylvania: rare. (Adv. from Eu.)

31. LÁMIUM, L. DEAD-NETTLE.

Calyx tubular-bell-shaped, about 5-nerved, with 5 nearly equal awl-pointed teeth. Corolla dilated at the throat; the upper lip ovate or oblong, arched, narrowed at the base; the middle lobe of the spreading lower lip broad, notched at the apex, contracted as if stalked at the base; the lateral ones small, at the margin of the throat. Stamens 4, ascending under the upper lip: anthers approximate in pairs, 2-celled, the cells divergent. Nutlets truncate at the apex. — Herbs, decumbent at the base, the lowest leaves small and long-petioled, the middle ones heart-shaped and doubly toothed, the floral subtending the whorled clusters of flowers; produced from spring to autumn. (Name from λαιμος, the throat, in allusion to the ringent corolla.)

* *Annuals or biennials, low: flowers small, purple, in few whorls or heads.*

1. **L. AMPLEXICAÛLE**, L. Leaves rounded, deeply crenate-toothed or cut, the upper ones clasping; corolla elongated, upper lip bearded, the lower spotted; lateral lobes truncate. — Cultivated grounds. (Adv. from Eu.)

2. **L. PURPÛREUM**, L. Leaves roundish or oblong, heart-shaped, crenate-toothed, all petioled. — Cult. grounds, Pennsylvania. (Adv. from Eu.)

* * *Perennial, taller: flowers larger, in several axillary whorls.*

3. **L. ÁLBUM**, L. Hairy; leaves ovate, heart-shaped, petioled; calyx-teeth very slender, spreading; corolla white, the tube curved upwards, obliquely contracted near the base, where there is a ring of hairs inside; lateral lobes of lower lip bearing a long slender tooth. — Waste ground near Boston, *D. Murray*. (Adv. from Eu.)

32. BALLÛTA, L. FETID HOREHOUND.

Calyx nearly funnel-form; the 10-ribbed tube expanded above into a spreading regular border, with 5-10 teeth. Anthers exerted beyond the tube of

the corolla, approximate in pairs. Otherwise much as in *Marrubium*. (The Greek name, of uncertain origin.)

1. **B. NIGRA**, L. (BLACK HOREHOUND.) More or less hairy, but green, erect; the root perennial; leaves ovate, toothed; whorls many-flowered, dense; calyx-teeth 5, longer than the tube of the purplish corolla. — Waste places, Massachusetts and Connecticut: scarce. (Adv. from Eu.)

33. PHLÒMIS, L. JERUSALEM SAGE.

Calyx tubular, 5-10-ribbed, truncate or equally 5-toothed. Upper lip of the corolla arched; the lower spreading, 3-cleft. Stamens 4, ascending and approximate in pairs under the upper lip; the filaments of the upper pair with an awl-shaped appendage at the base, *longer than the others* in *P. tuberosa*, &c.: anther-cells divergent and confluent. — Leaves rugose. Whorls dense and many-flowered, axillary, remote, bracted. (An old Greek name of a woolly species, of obscure derivation.)

1. **P. TUBEROSA**, L. Tall perennial (3°-5° high), nearly smooth; leaves ovate-heart-shaped, crenate, petioled; the floral oblong-lanceolate; bracts awl-shaped, hairy; upper lip of the purple corolla densely bearded with white hairs on the inside. — Shore of Lake Ontario near Rochester. June, July. (Adv. from Eu.)

ORDER 72. BORRAGINACEÆ. (BORAGE FAMILY.)

Chiefly rough-hairy herbs (not aromatic), with alternate entire leaves, and symmetrical flowers with a 5-parted calyx, a regular 5-lobed corolla (except in No. 1), 5 stamens inserted on its tube, a single style and a deeply 4-lobed ovary (as in Labiatae), which forms in fruit 4 seed-like nutlets, each with a single seed. — Albumen none. Cotyledons plano-convex: radicle pointing to the apex of the fruit. Stigmas 1 or 2. Calyx valvate, the corolla imbricated (in *Myosotis convolute*) in the bud. Flowers mostly on one side of the branches of a reduced cyme, imitating a spike or raceme, which is rolled up from the end, and straightens as the blossoms expand (circinate or scorpioid), often bractless. (A rather large family of innocent, mucilaginous, and slightly bitter plants; the roots of some species yielding a red dye.)

Tribe I. BORRAGEÆ. Ovary deeply 4-parted, forming as many separate 1-seeded nutlets in fruit; the style rising from the centre between them. (Root frequently red.)

* Corolla naked and open (without scales) in the throat, somewhat irregular! Nutlets erect

1. **Echium**. Corolla funnel-form, unequally 5-lobed. Stamens protruded.

** Corolla with 5 scales closing the throat. Nutlets not prickly, erect; the scar broad.

2. **Lycopsis**. Corolla funnel-form, slightly curved and oblique: scales blunt and hairy.

3. **Symphytum**. Corolla tubular, and enlarged at the summit: scales awl-shaped.

*** Corolla open, with folds rather than scales in the throat. Nutlets smooth, erect; scar small.
— Lobes of the tubular corolla imbricated in the bud.

4. **Onosmodium**. Nutlets stony, smooth. Lobes of the corolla acute and erect.

5. **Lithospermum**. Nutlets stony, smooth. Lobes of the corolla spreading, rounded.

6. **Mertensia.** Nutlets fleshy, fixed by the inner angle. Lobes of the corolla rounded.
 — Lobes of the short salver-shaped corolla convolute in the bud.
7. **Myosotis.** Nutlets hard and smooth. Flowers all or most of them bractless.
- * * * * Corolla with 5 scales closing the throat. Nutlets prickly, laterally fixed to the central column or the base of the style, often recumbent.
8. **Echinosperrnum.** Corolla salver-shaped. Nutlets erect, prickly on the margin.
9. **Cynoglossum.** Corolla funnel-form. Nutlets oblique or depressed, prickly all over.
- Tribe II. HELIOTROPEÆ.** Ovary not lobed, tipped with the simple style: the fruit separating when ripe into 2 or 4 nutlets.
10. **Heliotropium.** Throat of the short salver-shaped corolla open. Nutlets 1-celled.
11. **Heliohytum.** Throat of the corolla contracted. Nutlets 2, each 2-celled, i. e. 4 in 2 pairs and sometimes a pair of empty false cells.

1. ÈCHIU M, Tourn. VIPER'S BUGLOSS.

Corolla with a cylindraceous or funnel-form tube, and a more or less unequal spreading 5-lobed border; the lobes rounded, the expanded throat naked. Stamens mostly exerted, unequal. Style thread-form. Nutlets roughened or wrinkled, fixed by a flat base. (A name of Dioscorides, from ἔχλις, a viper.)

1. **E. VULGARE, L.** (BLUE-WEED.) Rough-bristly biennial; stem erect (2° high), mostly simple; stem-leaves linear-lanceolate, sessile; flowers showy, in short lateral clusters, disposed in a long and narrow raceme; corolla reddish-purple changing to brilliant blue (rarely pale). — Roadsides and meadows: rather rare northward; but a troublesome weed in cultivated fields in Virginia. June. (Nat. from Eu.)

2. LYCÓPSIS, L. BUGLOSS.

Corolla funnel-shaped, with a curved tube and a slightly unequal limb; the throat closed with 5 convex obtuse bristly scales placed opposite the lobes. Stamens and style included. Nutlets rough-wrinkled, erect, fixed by a hollowed-out base. — Annuals. (Name from λύκος, a wolf, and ὄψις, face.)

1. **L. ARVENSIS, L.** (SMALL BUGLOSS.) Very rough-bristly (1° high); leaves lanceolate; flowers in leafy raceme-like clusters; calyx as long as the tube of the small blue corolla. — Dry or sandy fields, New England to Virginia: scarce. (Adv. from Eu.)

3. SÝMPHYTUM, Tourn. COMFREY.

Corolla oblong-tubular, inflated above, 5-toothed; the short teeth spreading; the throat closed with 5 converging linear-awl-shaped scales. Stamens included: anthers elongated. Style thread-form. Nutlets smooth, ovate, erect, fixed by the large hollowed base, which is finely toothed on its margin. — Coarse perennial herbs, with thickened bitterish mucilaginous roots; the nodding raceme-like clusters either single or in pairs. (Name from συμφείν, to grow together, probably in allusion to its reputed healing virtues.)

1. **S. OFFICINALE, L.** (COMMON COMFREY.) Hairy, branched, winged above by the decurrent leaves; the lower leaves ovate-lanceolate, tapering into a petiole, the upper narrower; corolla yellowish-white, rarely purplish. — Moist places; escaped from gardens. June. (Adv. from Eu.)

4. **ONOSMÒDIUM**, Michx. FALSE GROMWELL.

Calyx 5-parted; the divisions linear and erect. Corolla tubular, or tubular-funnel-form, naked in the throat (the sinuses minutely hooded-inflexed); the 5 acute lobes converging or barely spreading. Anthers oblong-linear or arrow-shaped, mucronate, inserted in the throat of the corolla. Style thread-form, much exerted. Nutlets bony, ovoid, smooth, erect, fixed by the base; the scar minute, not hollowed out. — Chiefly perennial herbs, coarse and hispid, with oblong and sessile ribbed-veined leaves, and white, greenish, or yellowish flowers, in at length elongated and erect leafy raceme-like clusters; in summer. — Our species all belong to true *ONOSMODIUM*, having the anthers all included, smooth, and on very short filaments; the corolla only once or twice the length of the calyx. (Named from the resemblance to the genus *Onosma*, which means *ass-smell*.)

1. **O. Virginiànum**, DC. *Clothed all over with harsh and rigid appressed short bristles*; stems rather slender (1° – 2° high); leaves narrowly oblong, or oblong-lanceolate ($1'$ – $2\frac{1}{2}'$ long), the lower narrowed at the base; lobes of the narrow corolla lance-awl-shaped, sparingly bearded outside with long bristles. (*O. hispidum*, Michx. *Lithospermum Virginianum*, L.!) — Banks and hillsides, S. New England to Virginia and southward.

2. **O. Carolinànum**, DC. (excl. syn. Michx.) *Shaggy all over with long and spreading bristly hairs*; stem stout, upright (3° – 4° high); leaves ovate-lanceolate or oblong-lanceolate, acute; lobes of the rather broad corolla ovate-triangular or triangular-lanceolate, thickly hirsute outside. (*O. mólle*, Beck, &c. *Lithospermum Carolinianum*, Lam.) — River-banks, W. New York to Wisconsin and southward. — Perhaps passes into the next.

3. **O. mólle**, Michx. *Hoary with finer and soft mostly appressed hairs*; leaves oblong-ovate, obtusish, strongly ribbed, lobes of the rather narrow corolla triangular and sharp-pointed, thickly hirsute outside. — Dry grounds, Ohio to Illinois and southward.

5. **LITHOSPÉRMUM**, Tourn. GROMWELL. PUCCOON.

Corolla funnel-form, or sometimes salver-shaped; the open throat naked, or with a more or less evident transverse fold or scale-like appendage opposite each lobe; the spreading limb 5-cleft; its lobes rounded. Anthers oblong, almost sessile, included. Nutlets ovate, smooth or roughened, mostly bony or stony, fixed by the base; the scar nearly flat. — Herbs, with thickish and commonly red roots and sessile leaves; the flowers solitary and as if axillary, or spiked and leafy-bracted: sometimes dimorphous as to insertion of stamens and length of style. (Name formed of *λίθος*, stone, and *σπέρμα*, seed, from the hard nutlets.)

§ 1. *Nutlets tubercled or rough-wrinkled and pitted, gray and dull: throat of the (nearly white) corolla destitute of any evident folds or appendages.*

1. **L. ARVÉNSE**, L. (CORN GROMWELL.) *Minutely rough-hoary annual or biennial*; stems erect ($6'$ – $12'$ high); leaves lanceolate or linear, veinless; corolla scarcely longer than the calyx. — Sandy banks and roadsides. May–Aug. (Nat. from Eu.)

§ 2. *Nutlets smooth and shining, mostly white like ivory, occasionally dotted with pores: corolla in our species greenish-white or cream-color, small, with 5 small but distinct pubescent scales in the throat. (Root perennial.)*

2. **L. angustifolium**, Michx. Minutely and slightly hoary, roughish, much branched, erect or spreading (6'–15' high); leaves linear, rigid, 1-nerved; corolla not longer than the calyx; the short peduncles in fruit mostly recurved; nutlets more or less pitted when young, rarely bright white, but smooth and shining. — River-banks, from Illinois southward and westward. May.

3. **L. OFFICINALE**, L. (COMMON GROMWELL.) Much branched above, erect (1°–2° high); leaves thinnish, broadly lanceolate, acute, with a few distinct veins, rough above, soft-pubescent beneath; corolla exceeding the calyx; nutlets very smooth and even. — Roadsides, &c.: rather rare. (Nat. from Eu.)

4. **L. latifolium**, Michx. Stem loosely branched, erect (2°–3° high), rough; leaves ovate and ovate-lanceolate, mostly taper-pointed (even the floral ones 2'–4' long), ribbed-veined, roughish above, finely soft-pubescent beneath, the root-leaves large and rounded; corolla shorter than the calyx; nutlets very smooth or sparingly impressed-punctate, shining, turgid (2" long). — Borders of woods, W. New York and Penn. to Wisconsin and southwestward. June.

§ 3. **BÁTSCHIA**, Gmelin. *Nutlets smooth and shining: corolla large, salver-shaped or nearly so, deep orange-yellow, somewhat pubescent outside, the tube 2–4 times longer than the calyx, the throat more or less appendaged. (Roots perennial, long and deep, yielding a red dye.)*

* *Tube of the corolla from one half to twice longer than the calyx, not much longer than its ample limb, the lobes entire; the appendages glandular and adherent (especially when the stamens are at the base of the tube), or slightly arched.*

5. **L. hirtum**, Lehm. (HAIRY PUCCOON.) *Hispid* with bristly hairs (1°–2° high); stem-leaves lanceolate or linear, those of the flowering branches ovate-oblong, bristly-ciliate; corolla woolly-bearded at the base inside; flowers distinctly peduncled; fruiting calyx ($\frac{1}{2}$ ' long) 3–4 times longer than the nutlets. (Also *L. sericeum*, Lehm. *Batschia Caroliniensis*, Gmel. *B. Gmelini*, Mich.) — Dry woods, New York to Wisconsin, Virginia, and southward and westward. April–June. — Flowers crowded, showy: limb of the corolla $\frac{3}{4}$ '–1' broad.

6. **L. canescens**, Lehm. (HOARY PUCCOON OR ALKANET.) *Softly hairy* and more or less hoary (6'–15' high); leaves obtuse, linear-oblong, or the upper ovate-oblong, more or less downy beneath and roughish with close appressed hairs above; corolla naked at the base within; flowers sessile; fruiting calyx (3" long) barely twice the length of the nutlets. (*Batschia canescens*, Michx.) — Open woods and plains, New York to Kentucky and northwestward. May. — Limb of the showy corolla smaller and the calyx shorter than in the last.

** *Tube of the corolla 2–4 times the length of the calyx and of its erose-toothed or crenulate lobes, the appendages more projecting. (Pentalophus, A. DC.)*

7. **L. longiflorum**, Spreng. Minutely strigose-hoary; stem simple (6'–18' high); leaves linear; tube of the corolla much longer than the calyx (8"–1 $\frac{1}{2}$ ' long). (*Batschia longiflora*, Pursh. *L. incisum*, Lehm. *Pentalophus longiflorus*, A. DC.) — Prairies and plains, from W. Illinois and Wisconsin westward. May.

6. **MERTÉNSIA**, Roth. SMOOTH LUNGWORT.

Corolla trumpet-shaped or bell-funnel-shaped, longer than the deeply 5-cleft or 5-parted calyx, naked, or with 5 small glandular folds or appendages in the open throat. Anthers oblong or arrow-shaped. Style long and thread-form. Nutlets ovoid, fleshy when fresh, smooth or wrinkled, obliquely attached next the base by a prominent internal angle; the scar small. — Smooth! or soft-hairy perennial herbs, with pale and entire leaves, and handsome purplish-blue (rarely white) flowers, in loose and short paniced or corymbed raceme-like clusters, only the lower one leafy-bracted: pedicels slender. (Named for *Prof. Francis Charles Mertens*, a German botanist.)

§ 1. *Corolla perfectly naked in the throat; the broad trumpet-mouthed limb almost entire: filaments slender, protruding, much longer than the anthers.*

1. **M. Virginica**, DC. (VIRGINIAN COWSLIP OR LUNGWORT.) Very smooth, pale, erect (1°–2° high); leaves obovate, veiny, those of the root (4'–6' long) petioled; corolla trumpet-shaped, 1' long, many times exceeding the calyx, rich purple-blue, rarely white; lobes of the disk one on each side of the ovary. (*Pulmonaria Virginica*, L.) — Alluvial banks, W. New York to Wisconsin, Virginia, and southward. May. — Cultivated for ornament.

§ 2. *Corolla with 5 glandular folds or appendages at the throat; the limb 5-lobed.*

2. **M. maritima**, DON. (SEA LUNGWORT.) Spreading or decumbent, smooth, glaucous; leaves fleshy, ovate or obovate, the upper surface becoming papillose; corolla bell-funnel-form, twice the length of the calyx (3" long); filaments longer and narrower than the anthers; nutlets flattened. — Sea-coast, on rocks and sand, Cape Cod to Maine and northward: scarce. June–Aug. (Eu.)

3. **M. paniculata**, DON. *Roughish and more or less hairy, erect* (1°–2° high), loosely branched; leaves ovate and ovate-lanceolate, taper-pointed, ribbed, thin; corolla (6" long) somewhat funnel-form, 3–4 times the length of the lance-linear acute divisions of the calyx; filaments broader and shorter than the anthers. — Shore of L. Superior and northward and westward. July.

7. **MYOSÒTIS**, L. SCORPION-GRASS. FORGET-ME-NOT.

Corolla salver-form, the tube about the length of the 5-toothed or 5-cleft calyx; the throat with 5 small and blunt arching appendages opposite the rounded lobes; the latter convolute in the bud! Stamens included, on very short filaments. Nutlets smooth, compressed, fixed at the base; the scar minute. — Low and mostly soft-hairy herbs, with entire leaves, those of the stem sessile, and with small flowers in naked (so called) racemes, which are entirely bractless, or occasionally with one or two small leaves next the base, prolonged and straightened in fruit. Flowering through the season. (Name composed of *μύς*, mouse, and *οἶς, ὠτός*, ear, in allusion to the aspect of the short and soft leaves in some species; one popular name is MOUSE-EAR.)

* *Calyx open in fruit, its hairs appressed, none of them hooked nor glandular.*

1. **M. palustris**, Withering. (TRUE FORGET-ME-NOT.) Perennial; stems ascending from an oblique creeping base (9'–20' high), loosely branched,

smoothish; leaves rough-pubescent, oblong-lanceolate or linear-oblong; calyx moderately 5-cleft, shorter than the spreading pedicels; corolla (rather large in the genuine plant) pale blue with a yellow eye. — Nat. from Eu. near Boston, escaping from gardens. — Varies into smaller-flowered forms, among which high authorities rank *M. cæspitosa*, and (with yet more reason) the intermediate

Var. *laxa*. (*M. laxa*, *Lehm.*) Creeping base of the stem short; flowers a third or half smaller; pedicels longer. — Wet places, northward. (Eu.)

* * *Calyx closing or the lobes erect in fruit, clothed with spreading hairs, a part of them minutely hooked or glandular at the apex: corolla small: root annual or biennial.*

2. *M. arvensis*, Hoffm. Hirsute with spreading hairs, erect or ascending (6'–15' high); leaves oblong-lanceolate, acutish; racemes naked at the base and stalked; corolla blue, rarely white; pedicels spreading in fruit and longer than the 5-cleft equal calyx. ♀ (*M. intermedia*, *Link.* *M. scorpioides*, var. *arvensis*, *L.*) — Fields, &c.: not very common, perhaps not indigenous. (Eu.)

3. *M. verna*, Nutt. Bristly-hirsute, branched from the base, erect (4'–12' high); leaves obtuse, linear-oblong, or the lower spatulate-oblong; racemes leafy at the base; corolla very small, white, with a short limb; pedicels in fruit erect and appressed at the base, usually abruptly bent outwards near the apex, rather shorter than the deeply 5-cleft unequal (somewhat 2-lipped) very hispid calyx. (*M. inflexa*, *Engelm.*) — Dry hills: rather common. May–July.

4. *M. versicolor*, Pers. More slender than the last, simple at the base; racemes loose, mostly naked at the base; flowers almost sessile; corolla pale yellow changing to blue or violet; calyx deeply and equally 5-cleft. — Fields, Delaware, *W. M. Canby*. (Nat. from Eu.)

8. ECHINOSPÉRMUM, Swartz. STICKSEED.

Corolla salver-form, short, nearly as in *Myosotis*, but imbricated in the bud, the throat closed with 5 short scales. Stamens included. Nutlets erect, fixed laterally to the base of the style or central column, triangular or compressed, the back armed with 1–3 marginal rows of prickles which are barbed at the apex, otherwise naked. — Rough-hairy and grayish herbs, with small blue flowers in bracted (so called) racemes; ours annuals or biennials, flowering all summer. (Name compounded of *ἐχίμος*, a hedgehog, and *σπέρμα*, seed.)

1. *E. LAPPULA*, *Lehm.* Stem upright, branched above (1°–2° high); the short pedicels erect; leaves lanceolate, rough-hairy; nutlets each with a double row of prickles at the margins, and rough-tubercled on the back. — Waste places: common. (Nat. from Eu.)

2. *E. Redóvskii*, *Lehm.* Nutlets with a single marginal row of stout prickles, and granulate-roughened on the back: otherwise much like the last. (*E. patulum*, *Hook.*) — St. Paul's, Minnesota, and on the plains westward.

9. CYNÓGLÓSSUM, Tourn. HOUND'S-TONGUE.

Corolla funnel-form; the tube about the length of the 5-parted calyx; the throat closed with 5 obtuse scales; the lobes rounded. Stamens included. Nut-

lets depressed or convex, oblique, fixed near the apex to the base of the style, roughened all over with short barbed or hooked prickles. — Coarse herbs, with a strong scent and petioled lower leaves; the mostly panicled (so-called) racemes naked above, usually bracted at the base. Fl. all summer. (Name from *κύων*, a dog, and *γλῶσσα*, tongue; from the shape and texture of the leaves.)

1. **C. OFFICINALE**, L. (COMMON HOUND'S-TONGUE.) Biennial; clothed with short soft hairs, leafy, panicled above; upper leaves lanceolate, closely sessile by a rounded or slightly heart-shaped base; racemes nearly bractless; corolla reddish-purple (rarely white); nutlets flat on the broad upper face, somewhat margined. — Waste grounds and pastures: a familiar and troublesome weed; the large nutlets adhering to the fleece of sheep, &c. (Nat. from Eu.)

2. **C. Virginicum**, L. (WILD COMFREY.) Perennial; roughish with spreading bristly hairs; stem simple, few-leaved (2° – 3° high); stem-leaves lanceolate-oblong, clasping by a deep heart-shaped base; racemes few and corymbed, raised on a long naked peduncle, bractless; corolla pale blue; nutlets strongly convex. — Rich woods: rather common, especially westward. — Flowers intermediate in size between the other two.

3. **C. Morisoni**, DC. (BEGGAR'S LICE.) Biennial; stem hairy, very broadly branched, leafy (2° – 4° high); leaves oblong-ovate, taper-pointed, also tapering at the base, thin, minutely downy underneath and roughish above; racemes panicled, forking, diverging, hairy, leafy-bracted at the base; flowers very small; corolla white or pale blue (minute); pedicels reflexed in fruit; nutlets convex, the prickles with barbed points. (*Myosotis Virginica*, L. *Echino-spermum*, *Lehm.*) — Copses: a common and vile weed.

10. HELIOTRÏPIUM, Tourn. HELIOTROPE.

Corolla salver-shaped, short, 5-lobed; the sinuses more or less plaited in the bud; the throat open. Anthers nearly sessile. Style short: stigma conical, or capitate. Nutlets 4, when young united by their whole inner faces into a 4-celled ovary, but separating when ripe, each 1-seeded. — Herbs or low shrubby plants, the small flowers in one-sided spikes; in summer. (The ancient name, from *ἥλιος*, the sun, and *τροπή*, a turn.)

1. **H. EUROPEUM**, L. Erect annual (6'–18' high), hoary-pubescent; leaves oval, long-petioled; lateral spikes single, the terminal in pairs; calyx spreading in fruit, hairy. — Waste places, southward: scarce. (Adv. from Eu.)

2. **H. Curassavicum**, L. Apparently annual, glabrous; stems ascending; leaves lance-linear or spatulate, thickish, pale, almost veinless; spikes in pairs. — Sandy shores, Norfolk, Virginia, S. W. Illinois, and southward.

11. HELIÓPHYTUM, Cham., DC. INDIAN HELIOTROPE.

Corolla constricted at the throat. Style very short. Nutlets 2, each 2-celled, i. e. 4, in pairs, and sometimes a pair of empty false cells besides: otherwise nearly as in *Heliotropium*. (Name, *ἥλιος*, sun, and *φυτόν*, plant.)

1. **H. ÍNDICUM**, DC. Erect and hairy annual; leaves petioled, ovate or oval and somewhat heart-shaped; spikes single; fruit 2-cleft, mitre-shaped, with an empty false cell before each seed-bearing cell. (*Heliotropium Indicum*, L.) — Waste places, S. Illinois, and southward. (Adv. from India.)

ORDER 73. **HYDROPHYLLACEÆ.** (WATERLEAF FAMILY.)

Herbs, commonly hairy, with mostly alternate leaves, regular 5-merous and 5-androus flowers, in aspect between the foregoing and the next order; but the ovary entire and 1-celled with 2 parietal 4-many-ovuled placentæ, or rarely 2-celled by the union of the placentæ in the axis; style 2-cleft or 2 separate styles; fruit a 2-valved 4-many-seeded pod. — Seeds mostly reticulated or pitted. Embryo small in copious albumen. — Flowers chiefly blue or white, in one-sided cymes or racemes, which are mostly bractless and coiled from the apex when young, as in the Borage Family. (A small order of plants of no marked properties, some cultivated for ornament.)

Tribe I. HYDROPHYLLÆ. Ovary and pod 1-celled. Seeds amphitropous, pitted or reticulated, and with cartilaginous albumen. Leaves usually cut-toothed, lobed or pinnate. Style 2-cleft.

* Ovary lined with the dilated and fleshy placentæ, which enclose the ovules and seeds (in our plants these are only 4 in number) like an inner pericarp.

1. **Hydrophyllum.** Stamens exerted: anthers linear. Calyx unchanged in fruit.

2. **Nemophila.** Stamens included: anthers short. Calyx with appendages at the sinuses.

3. **Ellisia.** Stamens included. Calyx destitute of appendages, enlarged in fruit.

** Ovary with narrow parietal placentæ, in fruit projecting inwards more or less.

4. **Phacelia.** Corolla-lobes imbricated in the bud. Calyx destitute of appendages.

Tribe II. HYDROLEÆ. Ovary and pod 2-celled, the placentæ often projecting from the axis far into the cells. Seeds anatropous: albumen fleshy. Leaves undivided.

5. **Hydrolea.** Corolla between wheel-shaped and bell-shaped. Styles 2. Leaves entire.

1. **HYDROPHÝLLUM, L.** WATERLEAF.

Calyx 5-parted, sometimes with a small appendage in each sinus, early open in the bud. Corolla bell-shaped, 5-cleft; the lobes convolute in the bud; the tube furnished with 5 longitudinal linear appendages opposite the lobes, which cohere by their middle, while their edges are folded inwards, forming a nectariferous groove. Stamens and style mostly exerted: filaments more or less bearded: anthers linear. Ovary bristly-hairy (as is usual in the family); the 2 fleshy placentæ expanded so as to line the cell and nearly fill the cavity, soon free from the walls except at the top and bottom, each bearing a pair of ovules on the inner face. Pod ripening 1-4 seeds, spherical. — Perennials, with petioled ample leaves, and white or pale blue cymose-clustered flowers. (Name formed of ὕδωρ, *water*, and φύλλον, *leaf*; of no obvious application to these plants.)

* Calyx with minute if any appendages: rootstocks creeping, scaly-toothed.

1. **H. macrophýllum, Nutt.** *Rough-hairy; leaves oblong, pinnate and pinnatifid; the divisions 9-13, ovate, obtuse, coarsely cut-toothed; peduncle very long; calyx-lobes lanceolate-pointed from a broad base, very hairy.* — Rocky, shaded banks, Ohio, Indiana, Kentucky, and southward. July. — Root-leaves 1° long; flowers crowded in a globular cluster.

2. **H. Virgínicum, L.** *Smoothish (1° - 2° high); leaves pinnately divided; the divisions 5-7, ovate-lanceolate or oblong, pointed, sharply cut-toothed, the lowest mostly 2-parted, the uppermost confluent; peduncles longer than the petioles of the upper leaves, forked; calyx-lobes narrowly linear, bristly-ciliate.* — Damp woods. June - Aug.

3. **H. Canadense**, L. *Nearly smooth* (1° high); *leaves palmately 5-7-lobed, rounded, heart-shaped at the base, unequally toothed*; those from the root sometimes with 2-3 small and scattered lateral leaflets; *peduncles much shorter than the long petioles, forked, the crowded (nearly white) flowers on very short pedicels*; calyx-lobes linear-awl-shaped, nearly smooth. — Damp rich woods, W. New England to the mountains of Virginia and northward. June - Aug. — Rootstocks thickened and very strongly toothed in 2 rows by the persistent bases of the stout petioles: leaves 3' - 5' broad.

* * *Calyx with a small reflexed appendage in each sinus: stamens sometimes not exerted (probably two forms of flowers, as in some Borraginaceæ, &c.).*

4. **H. appendiculatum**, Michx. *Hairy*; stem-leaves palmately 5-lobed, rounded, the lobes toothed and pointed, the lowest pinnately divided; cymes rather loosely flowered; pedicels (at length slender) and calyx bristly-hairy. — Open woods, New York to Virginia, Wisconsin, and westward. June, July.

2. NEMÓPHILA, Nutt. NEMOPHILA.

Calyx 5-parted, and with a reflexed tooth or appendage in each sinus, more or less enlarged in fruit. Corolla bell-shaped or almost wheel-shaped; the lobes convolute in the bud; the tube mostly with 10 small folds or scales inside. Stamens included: anthers ovoid or heart-shaped. Placentæ (bearing each 2-12 ovules), pod, and seeds as in *Hydrophyllum*. — Diffuse and fragile annuals, with opposite or partly alternate pinnatifid or lobed leaves, and one-flowered peduncles; the corolla white, blue, or marked with purple. (Name composed of *νέμος, a grove*, and *φιλέω, to love*.) Some handsome species are garden annuals.

1. **N. microcalyx**, Fisch. & Meyer. *Small, roughish-pubescent*; stems diffusely spreading (2' - 8' long); leaves parted or deeply cleft into 3-5 roundish or wedge-obovate sparingly cut-lobed divisions, the upper leaves all alternate; peduncles opposite the leaves, shorter than the long petioles; flowers minute; corolla white, longer than the calyx; placentæ each 2-ovuled; pod 1-2-seeded. (*Ellisia microcalyx, Nutt. Nemophila evanescens, Darby.*) — Moist woods, Virginia (near Washington), and southward. April - June.

3. ELLÍSIA, L. ELLISIA.

Calyx 5-parted, without appendages, enlarged and foliaceous in fruit. Corolla bell-shaped or cylindraceous, not longer than the calyx, 5-lobed above; the lobes imbricated or convolute in the bud, the tube with 5 minute appendages within. Stamens included. Placentæ (each 2-ovuled), fruit, and seeds much as in *Hydrophyllum*. — Delicate and branching annuals, with lobed or divided leaves, the lower opposite, and small whitish flowers. (Named for *John Ellis*, a distinguished naturalist, an English correspondent of Linnæus.)

1. **E. Nyctèlea**, L. *Minutely or sparingly roughish-hairy, divergently branched* (6' - 12' high); leaves pinnately parted into 7-13 lanceolate or linear-oblong sparingly cut-toothed divisions; peduncles solitary in the forks or opposite the leaves, 1-flowered; calyx-lobes triangular-ovate lanceolate, pointed, about the length of the cylindraceous (whitish) corolla (in fruit becoming nearly $\frac{1}{2}$ ' long); pod pendulous. — Shady damp places, W. New Jersey and E. Pennsylvania to Virginia. May - July.

2. **E. ambigua**, Nutt. Later peduncles sometimes 2-3-flowered; corolla rather more bell-shaped, the upper part of tube 5-angular: otherwise like the preceding, of which it is probably a variety. — Illinois and westward.

4. PHACÈLIA, Juss. (Phacelia & Eutoca, R. Br.)

Calyx 5-parted; the sinuses naked. Corolla open-bell-shaped, 5-lobed; the lobes imbricated in the bud. Filaments slender, often (with the 2-cleft style) exserted: anthers ovoid or oblong. Ovary with 2 narrow linear placentaë adherent to the walls, in fruit usually projecting inwards more or less, the two often forming an imperfect partition in the ovoid 4-many-seeded pod. (Ovules 2-30 on each placenta.) — Perennial or mostly annual herbs, with either simple, lobed, or divided leaves, and commonly handsome (blue, purple, or white) flowers in one-sided raceme-like clusters. (Name from *φάκελος*, a fascicle.)

§ 1. PHACELIA. *Seeds and ovules only 4 (two on each placenta): corolla with narrow folds, appendages, or scales within, the lobes entire.*

1. **P. bipinnatifida**, Michx. Biennial; stem upright, much branched, hairy (1°-2° high); leaves long-petioled, pinnately 3-5-divided; the divisions or leaflets ovate or oblong-ovate, acute, coarsely and often sparingly cut-lobed or pinnatifid; racemes elongated, loosely many-flowered, glandular-pubescent; pedicels about the length of the calyx, spreading or recurved. — Shaded banks, in rich soil, Ohio to Illinois and southward. May, June. — Corolla bright blue, 6" broad, with 5 pairs of longitudinal folds, covering as many externally keeled deep grooves. Stamens bearded below; these, with the style, are either somewhat included (*P. brevistylis*, Buckley) or exserted in different individuals.

§ 2. COSMÁNTHUS. (*Cosmanthus*, Nolte. Sect. *Eucosmanthus*, A. DC., in part.) *Seeds and ovules only 4: corolla naked within; its lobes beautifully fringe-toothed: filaments villous-bearded below: leaves pinnatifid, the upper clasping at the base: flowers long-pedicelled: annuals or spring biennials.*

2. **P. Purshii**, Buckley. Sparsely hairy; stem erect or ascending, branched (8'-12' high); lobes of the stem-leaves 5-9, oblong or lanceolate, acute; raceme many-flowered; calyx-lobes lance-linear; corolla light blue, varying to white (about ½' in diameter). (*P. fimbriata*, Pursh., not of Michx. *Cosmanthus fimbriatus*, Nolte, &c.) — Moist wooded banks, W. Pennsylvania to Illinois and southward. April-June.

3. **P. fimbriata**, Michx. Slightly hairy, slender; stems spreading or ascending (5'-8' long), few-leaved; lowest leaves 3-5-divided into roundish leaflets; the upper 5-7-cleft or cut-toothed, the lobes obtuse; raceme 3-10-flowered; calyx-lobes linear-oblong, obtuse, becoming spatulate; corolla white (3' - 4" broad). — Woods, high mountains of Virginia, and southward. May.

§ 3. EÛTOCA. (*Eutoca*, R. Br.) *Seeds (or at least the ovules) several or many, rarely only 3 or 4 on each placenta: corolla usually with small and inconspicuous folds or appendages within, its lobes entire. (Ours are annuals or biennials.)*

4. **P. parviflora**, Pursh. Somewhat hairy, slender, diffusely spreading (3'-8' high); leaves pinnately cleft or the lower divided into 3-7 short lobes; racemes solitary, loosely 5-15-flowered; pedicels filiform, at length several times

longer than the oblong calyx-lobes; corolla bluish-white (3''-4'' broad); *pod few-seeded*.—Shaded banks, S. Penn. to Virginia and southward. April-June.

5. **P. Franklinii**, Gray. Soft-hairy; *stem erect* (6'-15' high), rather stout; leaves pinnately parted into many lanceolate or oblong-linear lobes, which are crowded and often cut-toothed or pinnatifid; *racemes short, dense, crowded into an oblong spike*; calyx-lobes linear; corolla blue; *pod many-seeded*. (*Eutoca Franklinii*, R. Br.)—Shores of Lake Superior, especially on Isle Royale; thence northward and westward.

5. HYDRÒLEA, L. HYDROLEA.

Calyx 5-parted. Corolla short-campanulate or almost wheel-shaped, 5-cleft. Filaments dilated at the base. Styles 2, distinct. Pod globular, 2-celled, and the cells often partly divided into 2 by the projection of the many-seeded placenta, thin-walled, 2-4-valved or bursting irregularly. Seeds minute, striate-ribbed.—Herbs or scarcely shrubby plants, growing in water or wet places (whence the name, from ὕδωρ, *water*), with entire leaves, often having spines in their axils, and clustered blue flowers.

1. **H. affinis**, n. sp. Glabrous throughout; stem ascending from a creeping base, armed with small axillary spines; leaves lanceolate, tapering into a very short petiole; flowers in small axillary leafy-bracted clusters; divisions of the calyx lance-ovate, equalling the corolla and the irregularly-bursting globose pod.—Banks of the Ohio in S. Illinois, *Dr. Vasey* (and of the Mississippi at Memphis, *A. Fendler*: also E. Texas, *C. Wright*: in addenda to ed. 2, referred to *H. quadrivalvis*, *Walt.*, of the Southeastern States, from which it is distinguished by the smoothness and the broader sepals.

ORDER 74. POLEMONIACEÆ. (POLEMONIUM FAMILY.)

Herbs, with alternate or opposite leaves, regular 5-merous and 5-androus flowers, the lobes of the corolla convolute (in one tribe imbricated) in the bud, a 3-celled ovary and 3-lobed style; the pod 3-celled, 3-valved, loculicidal, few-many-seeded; the valves usually breaking away from the triangular central column.—Seeds amphitropous, the coat frequently mucilaginous when moistened and emitting spiral threads. Embryo straight in the axis of copious albumen. Calyx persistent, usually imbricated. Corolla with a 5-parted border. Anthers introrse. (Inspid and innocent plants; many are ornamental in cultivation.)

Tribe I. POLEMONIÆ. Calyx 5-cleft. Corolla convolute in the bud. Filaments filiform, inserted on the tube of the corolla: cells of the anther parallel, opening lengthwise. Flowers cymose-panicled or clustered.

1. **Polemonium**. Calyx and corolla open-bell-shaped. Filaments slender, equal.

2. **Phlox**. Calyx narrow. Corolla salver-shaped, with a long tube, including the unequally inserted filaments.

Tribe II. DIAPENSIÆ. Calyx of 5 sepals. Corolla imbricated in the bud, and with the broad and flat filaments in the sinuses. Anthers opening transversely.

3. **Diapensia**. Anther-cells pointless, opening by an obliquely transverse line.

4. **Pyxidantha**. Anther-cells awn-pointed underneath, opening straight across.

1. **POLEMŌNIUM**, Tourn. GREEK VALERIAN.

Calyx bell-shaped. Stamens equally inserted at the summit of the very short tube of the open-bell-shaped or short funnel-form corolla: filaments slender, declined, hairy-appendaged at the base. Pod few-several-seeded. — Perennials, with alternate pinnate leaves, the upper leaflets sometimes confluent; the (blue or white) corymbose flowers nearly bractless. (An ancient name, from *πόλεμος*, war, of doubtful application.)

1. **P. réptans**, L. Smooth throughout; stems weak and spreading (6'–10' high, never creeping as the name denotes); leaflets 7–11, ovate-lanceolate or oblong; corymbs few-flowered; flowers nodding; lobes of the calyx rather shorter than the tube; *stamens and style not protruding* beyond the corolla, which is light blue, about $\frac{1}{2}$ wide; *pods about 3-seeded*. — Woods, W. New York to Wisconsin and southward. May, June.

2. **P. cærùleum**, L. (JACOB'S LADDER.) Stem erect (1°–3° high); leaflets 9–21, linear-lanceolate, oblong- or ovate-lanceolate, mostly crowded; flowers numerous, in a thyrsus or contracted panicle; lobes of the calyx longer than the tube; *stamens and style mostly exerted* beyond the bright blue corolla, which is nearly 1' broad; pod several-seeded. — Swamps, about the sources of the Susquehanna, New York: East of Charlottesville, Schoharie Co., *Dr. E. C. Howe*. Elk Creek, near Delhi, Delaware Co., *B. D. Gilbert*. Head of Little Lakes, Warren, Herkimer Co., *G. W. Clinton*. Warren Co., New Jersey, *A. P. Garber*. Wild far northwestward. July. (Eu.)

2. **PHLOX**, L. PHLOX.

Calyx narrow, somewhat prismatic, or plaited and angled. Corolla salverform, with a long tube. Stamens very unequally inserted in the tube of the corolla, included. Pod ovoid, with (sometimes 2 ovules but ripening only) a single seed in each cell. — Perennials (except a few Southern species, such as *P. Drummondii* of the gardens), with opposite and sessile perfectly entire leaves, the floral often alternate. Flowers cymose, mostly bracted; the open clusters terminal or crowded in the upper axils. (*Φλόξ*, *flame*, an ancient name of *Lychnis*, transferred to this North American genus.) Most of the species are cultivated in gardens.

* *Stem strictly upright: panicle pyramidal or oblong, many-flowered: peduncles and pedicels very short: lobes of the corolla entire.* (Very common in gardens.)

1. **P. paniculàta**, L. Stem stout (2°–4° high), smooth; leaves oblong-lanceolate and ovate-lanceolate, pointed, large, tapering at the base, the upper often heart-shaped at the base; *panicle ample, pyramidal-corymbed; calyx-teeth awn-pointed*. (*P. undulata*, *Ait.*, &c.) — Var. **ACUMINÀTA** (*P. acuminata*, *Pursh*) has the broader and taper-pointed leaves beneath downy, like the stem, which is sometimes rough-hairy and spotted below. — Rich woods, from Penn. to Illinois, and southward. June, July. — Flowers pink-purple varying to white.

2. **P. maculàta**, L. (WILD SWEET-WILLIAM.) Smooth, or barely roughish; *stem spotted with purple*, rather slender (1°–2° high); lower leaves lanceolate, the upper nearly ovate-lanceolate, tapering to the apex from the broad

and rounded or somewhat heart-shaped base; *panicle narrow, oblong, leafy below; calyx-teeth triangular-lanceolate, short, scarcely pointed; corolla purple* (sometimes white, when it is *P. suaveolens*, *Ait.*). Lower branches of the panicle rarely elongated, so as to become pyramidal, when it is *P. pyramidalis*, *Smith.* — Rich woods and river-banks, N. Penn. to Wisconsin, and southward. June.

* * *Stems ascending or upright, often from a decumbent base: flowers in terminal corymbed cymes: the whole plant smooth and glabrous: lobes of the corolla round and entire: calyx-teeth short, triangular-lanceolate.*

3. **P. Carolina**, L. Stems ascending ($\frac{1}{2}^{\circ}$ – 2° high), often from a prostrate base; *leaves oblong-lanceolate, or the upper ovate-lanceolate, and sometimes heart-shaped at the base, acute or pointed; flowers crowded, short-peduncled; calyx-teeth acute.* — Var. *ovata*, *Benth.*, has broader leaves (*P. ovata*, L.). — Var. *nitida*, *Benth.*, has narrower leaves (*P. nitida*, *Pursh*), and verges to the next. — Woods, W. Pennsylvania to Michigan, Virginia, and southward. June, July. — Corolla 1' long; the limb 1' broad, pink-purple.

4. **P. glaberrima**, L. Stems slender, erect (1° – 3° high); *leaves linear-lanceolate or rarely oblong-lanceolate, very smooth (except the rough and sometimes revolute margins), tapering gradually to a point (3'–4' long); cymes few-flowered and loosely corymbed; flowers peduncled (pink or whitish); calyx-teeth sharp-pointed.* (*P. carnea*, *Sims.* *P. revoluta*, *Aikin.*) — Prairies and open woods, Ohio and Wisconsin to Virginia and southward. July.

* * * *Stems ascending (or in No. 5 erect) from a spreading or prostrate base, more or less clammy-pubescent, as well as the calyx and the oblong, lanceolate, or linear leaves: flowers in terminal corymbed cymes, mostly pedicelled: calyx deeply cleft, the teeth linear-awl-shaped or setaceous.*

5. **P. pilosa**, L. Stems slender, nearly erect (1° – $1\frac{1}{2}^{\circ}$ high), usually hairy, as are the *lanceolate or linear leaves, which commonly taper to a sharp point; cymes at length open; calyx-teeth slender awl-shaped and awn-like, longer than the tube, loose or spreading; lobes of the pink-purple or rose-red (rarely white) corolla obovate, entire.* (*P. aristata*, *Michx.* *P. aristata & pilosa* in part, *Benth.* in *DC.*) — Borders of thickets and prairies, New Jersey to Wisconsin and southward. May, June. — Leaves 1'– $2\frac{1}{2}'$ long, 1''–3'' wide.

6. **P. procumbens**, *Lehm.* Stems ascending ($\frac{1}{2}^{\circ}$ – $1\frac{1}{2}^{\circ}$ high), mostly simple; *leaves broadly linear, lanceolate or ovate-oblong, abruptly acute or blunt ($\frac{1}{2}'$ – $1\frac{1}{2}'$ long, on sterile shoots often ovate); cyme mostly compact and sessile, leafy-bracted; calyx-teeth awl-shaped or linear, sharp-pointed, but seldom awned, rather longer than the tube, straight; lobes of the corolla obovate and entire (or rarely notched), purple, pink, or sometimes white.* (*P. pilosa*, *Walt.*, *Michx.*, *Ell.*, *Benth.* in part, not L. *P. pilosa*, var. ? *Walteri*, Ed. 2. *P. Walteri*, *Chapman.* *P. procumbens*, *Lehm.* is a small form.) — Dry hills and barrens, Kentucky, Virginia, and southward. May, June.

7. **P. réptans**, *Michx.* *Runners creeping, bearing roundish-obovate smoothish and thickish leaves; flowering stems (4'–8' high) and their oblong or ovate obtuse leaves ($\frac{1}{2}'$ long) pubescent, often clammy; cyme close, few-flowered; calyx-teeth linear-awl-shaped, about the length of the tube; lobes of the reddish-purple corolla round-obovate, mostly entire.* — Damp woods, Pennsylvania, Kentucky, and

southward. May, June. — Flowers showy: tube of the corolla an inch long; the limb nearly as broad.

8. **P. divaricata**, L. Stems spreading or ascending from a decumbent base (9' - 18' high); leaves oblong-ovate or the lower oblong-lanceolate (1½' long), acutish; cyme corymbose-panicled, spreading, loosely-flowered; peduncles slender; calyx-teeth slender awl-shaped, much longer than the tube; lobes of the pale lilac or bluish corolla obcordate or wedge-obovate and notched at the end, or often entire (var. *Laphamii*, Wood), ½' - ¾' long, equalling or longer than the tube, with rather wide sinuses between them. — Rocky damp woods, mountains of Virginia to N. New York, Wisconsin, and northward. May.

9. **P. bifida**, Beck. Stems ascending, branched (5' - 8' high); leaves linear, becoming nearly glabrous (½' - 1½' long, 1½'' wide); flowers few, on slender peduncles; calyx-teeth awl-shaped, about the length of the tube; lobes of the pale purple corolla 2-cleft to or below the middle (4'' long), equalling the tube, the divisions linear-oblong. — Prairies of Illinois, Mead (and Missouri). May.

* * * * *Stems creeping and tufted in broad mats, the short flowering shoots ascending, glandular-pubescent; the rigid narrow leaves crowded and fuscicled.*

10. **P. subulata**, L. (GROUND OR MOSS PINK.) Depressed, in broad mats; leaves awl-shaped, lanceolate, or narrowly linear (3'' - 6'' long); cymes few-flowered; calyx-teeth awl-shaped, rigid; corolla pink-purple or rose-color with a darker centre (sometimes white); the lobes wedge-shaped, notched, rarely entire. (*P. setacea*, L.) — Dry rocky hills and sandy banks, S. New York to Michigan and southward. April, May. — Common in cultivation.

3. DIAPÉNSIA, L. DIAPENSIA.

Calyx of 5 concave imbricated sepals. Corolla bell-shaped, 5-lobed; the lobes rounded. Filaments broad and flat, adherent to the corolla up to the sinuses, short: anthers adnate, of 2 ovoid pointless cells, diverging below, each opening therefore by a transverse-descending line. Pod enclosed in the calyx, cartilaginous; the cells few-seeded. — An alpine dwarf evergreen, growing in very dense convex tufts, with the stems imbricated below with cartilaginous narrowly spatulate mostly opposite leaves, terminated by a scape-like 1-flowered peduncle, 3-bracted under the calyx. Corolla white (½' wide). (Ancient Greek name of the Sanicle, of obscure meaning, strangely applied by Linnæus to this plant.)

1. **D. Lappónica**, L. — Alpine summits of the White Mountains, New Hampshire, and Adirondack Mountains, N. New York. July. (Eu.)

4. PYXIDANTHÈRA, Michx. PYXIDANTHERA.

Anther-cells awn-pointed at the base, opening by a strictly transverse line. Otherwise much as in *Diapensia*. — A small prostrate and creeping evergreen, with narrowly oblanceolate and awl-pointed crowded leaves, which are mostly alternate on the sterile branches, and somewhat hairy near the base. Flowers solitary and sessile, very numerous, white or rose-color. (Name from *πυξίς*, a small box, and *ἀνθήρα*, anther, the anther opening as if by a lid.)

1. **P. barbulata**, Michx. — Sandy pine barrens of New Jersey and southward. April, May.

ORDER 75. **CONVOLVULACEÆ.** (CONVOLVULUS FAMILY.)

Chiefly twining or trailing herbs, often with some milky juice, with alternate leaves (or scales) and regular 5-anded flowers; a calyx of 5 imbricated sepals; a 5-plaited or 5-lobed corolla convolute or twisted in the bud; a 2-celled (rarely 3-celled) ovary, or in one tribe 2 separate pistils, with a pair of erect ovules in each cell, the cells sometimes doubled by a false partition between the seeds, so becoming 4-celled; the embryo large, curved or coiled in mucilaginous albumen. — Fruit a globular 2-6-seeded pod. Flowers mostly showy, on axillary peduncles: pedicels articulated, often 2-bracted. (Many are cultivated for ornament, and one, the Sweet Potato, for its edible farinaceous roots: those of several species are cathartic; e. g. Jalap.) — There are three suborders, or rather strongly marked tribes.

Tribe I. CONVULVULÆ. Leafy plants, mostly twiners. Ovary 2-4-celled. Pod usually septifragal. Embryo with broad and leaf-like cotyledons, crumpled in the seed.

* Style single and undivided.

— Calyx naked, i. e. not enclosed or surrounded by leafy bracts.

1. **Quamoclit.** Stamens and style exserted. Corolla salver-shaped or nearly so. Stigma capitate-2-lobed. Pod 4-celled; the cells 1-seeded.
2. **Ipomœa.** Stamens included. Corolla funnel-form or bell-shaped. Stigma capitate, often 2-3-lobed. Pod 2-3-celled; the cells 2-seeded.
3. **Convolvulus.** Stigmas 2, elongated, linear. Otherwise much as in *Ipomœa*.
— — Calyx surrounded and enclosed by a pair of broad leafy bracts.
4. **Calystegia.** Stigmas 2, linear or oblong. Pod imperfectly 2-celled, 4-seeded.
* * Style single and 2-cleft, or styles 2, rarely more. Prostrate or spreading herbs.
5. **Bonamia.** Styles 2 and undivided, or a single one 2-cleft: stigmas capitate.
6. **Evolvulus.** Styles 2, and each 2-cleft: stigmas obtuse.

Tribe II. DICHONDREÆ. Creeping plants. Ovaries as well as styles 2 or more. Embryo, &c. as in the preceding tribe.

7. **Dichondra.** Corolla bell-shaped, 5-cleft. Pistils 2, one-seeded.

Tribe III. CUSCUTINEÆ. Leafless parasitic twiners. Embryo spiral, slender, destitute of cotyledons. Ovary 2-celled. Parts of the flower rarely in fours!

8. **Cuscuta.** The only genus of the group.

1. QUÁMOCLIT, Tourn. CYPRESS-VINE.

Sepals mostly mucronate or awned. Corolla cylindrical-tubular, with a small spreading border, not twisted in the bud. Stamens and style protruded. Stigma capitate-2-lobed. Pod 4-celled; the cells 1-seeded. — Annual twiners, with red or crimson flowers; in summer. (An aboriginal, probably Mexican, name.)

1. **Q. COCCINEA**, Mœnch. Leaves heart-shaped, acuminate, entire, or angled; sepals awn-pointed; corolla light scarlet (1' long). (*Ipomœa coccinea*, L.) — River-banks, &c., Ohio to Illinois, Virginia, and southward. (Nat. from Trop. Amer. or Ind.)

2. **Q. VULGARIS**, Choisy. (CYPRESS-VINE.) Leaves pinnately parted into linear-thread-shaped delicate parallel lobes; peduncles 1-flowered; corolla narrow, scarlet-red, and a white variety. — Sparingly escaped from gardens southward. (Adv. from India.)

2. IPOMŒA, L. MORNING-GLORY.

Calyx naked at the base. Corolla bell-shaped, funnel-form, &c., twisted in the bud. Stamens included. Stigma capitate, often 2-3-lobed. Pod 2-celled, or in one group 3-celled; the cells 2-seeded. (Name, according to Linnæus, from ἵψ, ἰρός, a *Bindweed* [which it is not], and ὅμοιος, *like*.)

§ 1. PHÁRBITIS, Choisy. Pod 3- (rarely 4-) celled; the cells 2-seeded.

1. **I. PURPŪREA**, Lam. (COMMON MORNING-GLORY.) Annual; stems retrorsely hairy; leaves heart-shaped, acuminate, entire; peduncles long umbellately 3-5-flowered; calyx bristly-hairy below; corolla funnel-form (2' long), purple, varying to white. (*Convolvulus purpureus*, L. *Pharbitis hispida*, Choisy.)—Around dwellings, escaping from cultivation. (Adv. from Trop. Amer.)

2. **I. NIL**, Roth. (SMALLER M.) Stems retrorsely hairy; leaves heart-shaped, 3-lobed, the lobes acute or acuminate; peduncles short, or rather long, 1-3-flowered; calyx densely hairy below; corolla white and purple or pale blue (1'-1½' long). (*Conv. Nil.* & *C. hederæceus*, L.)—Banks and near dwellings, from Maryland southward. (Adv. from Trop. Amer.?)

§ 2. IPOMŒA, Choisy. Pod 2-celled; the cells 2-seeded.

3. **I. lacunosa**, L. Annual; rather smooth; stem twining and creeping, slender; leaves heart-shaped, pointed, entire or angled-lobed; peduncles short, 1-3-flowered; sepals lance-oblong, pointed, bristly-ciliate or hairy, half the length of the sharply 5-lobed (white, ½'-¾' long) corolla. (*C. micranthus*, Riddell.)—Woods and fields, Penn. to Illinois and southward. Aug.

4. **I. pandurata**, Meyer. (WILD POTATO-VINE. MAN-OF-THE-EARTH.) Perennial, smooth or nearly so when old, trailing or sometimes twining; leaves regularly heart-shaped, pointed, occasionally some of them contracted at the sides so as to be fiddle-shaped; peduncles longer than the petioles; 1-5-flowered; sepals smooth, ovate-oblong, very obtuse; corolla open-funnel-form (3' long), white with purple in the tube.—Sandy fields and banks, from Connecticut to Illinois and southward. June-Aug.—Stems long and stout, from a huge root, which often weighs 10-20 pounds. Flowers opening in bright sunshine.

3. CONVŪLVULUS, L. BINDWEED.

Calyx naked at the base. Corolla open funnel-form or bell-shaped. Stamens included. Style 1: stigmas 2, linear, often revolute. Pod 2-celled; the cells 2-seeded.—Stems twining, procumbent, or often erect-spreading. Flowers mostly opening at dawn. (Name from *convolvere*, to entwine.)

1. **C. ARVENSIS**, L. (BINDWEED.) Perennial; stem procumbent or twining, and low; leaves ovate-oblong, arrow-shaped, with the lobes at the base acute; peduncles mostly 1-flowered; bracts minute, remote; corolla (9" long) white or tinged with reddish.—Fields, near the coast: likely to become a troublesome weed. June. (Nat. from Eu.)

4. CALYSTĒGIA, R. Br. BRACTED BINDWEED.

Calyx enclosed in 2 large and mostly heart-shaped leafy bracts: sepals equal. Corolla bell-funnel-form, the border obscurely 5-lobed or entire. Stamens in-

cluded. Style 1: stigmas 2, linear or oblong. Pod imperfectly 2-celled or 1-celled, 4-seeded. — Perennials, with heart-shaped or arrow-shaped leaves, and axillary 1-flowered peduncles; fl. in summer. (Name from κάλυξ, *calyx*, and στέγω, *to cover*, alluding to the bracts enclosing the calyx.)

1. **C. sepium**, R. Br. (HEDGE BINDWEED.) Stem *twining or sometimes trailing extensively*; leaves triangular-halberd-shaped or arrow-shaped, acute or pointed, the lobes at the base obliquely truncate and often somewhat toothed or sinuate-lobed; peduncles 4-angled; corolla white, or in the American plant more commonly light rose-color ($1\frac{1}{2}' - 2'$ long): the typical form *glabrous* throughout. (Convolvulus sepium, and C. repens, L.) — Varies greatly, often slightly pubescent: Var. PUBESCENS is a downy form, in the young state approaching the next. (C. Catesbyana, Pursh.) — Common, especially along the moist banks of streams. (Eu.)

2. **C. spithamæa**, Pursh. *Downy; stem low and mostly simple, upright or ascending* ($6' - 12'$ long); leaves oblong, with or without a heart-shaped or auricled base; corolla white ($2'$ long). — Dry, mostly sandy ground: not rare.

5. BONÁMIA, Thouars. (Breweria, R. Br. & Stylisma, Raf.)

Styles 2, or rarely 3, simple and distinct, or else united into one below: stigmas depressed-capitate. Otherwise as Convolvulus and Evolvulus. — Perennial prostrate or diffusely spreading herbs (or in warmer regions sometimes shrubby); flowers small; in summer: corolla more or less hairy or silky outside. (Named for Francis Bonamy, author of a Flora of Nantes.)

1. **B. humistrata**, Gray. (Proceed. Amer. Acad. 5, p. 337.) *Sparsely hairy or nearly smooth*; leaves varying from oblong with a somewhat heart-shaped base to linear, mucronate; peduncles 1-7-flowered; bracts shorter than the pedicels; *sepals pointed, glabrous or nearly so; corolla white; filaments hairy; styles united at the base*. (Convolvulus humistratus, Walt., who well distinguishes this from the next. Stylisma evolvuloides, Choix., in part. S. humistrata, Chapm.) — Dry pine barrens, Virginia (probably not in Ohio), and southward.

2. **B. aquatica**, Gray. *Minutely soft-downy and somewhat hoary; sepals silky; corolla pink or purple; filaments smooth; styles separate almost to the base*: otherwise nearly as in the last. (Conv. aquaticus, Walt. Stylisma aquatica, Chapm.) — Margin of ponds, S. Virginia? and southward.

3. **B. Pickeringii**, Gray. *Soft-pubescent or smoothish; leaves very narrowly linear or the lowest linear-spatulate, tapering to the base, nearly sessile; peduncles 1-3-flowered; bracts resembling the leaves, mostly exceeding the flowers; sepals hairy; filaments* (scarcely hairy) *and styles* (which are *united far above the middle*) *exserted from the open white corolla*. (Stylisma Pickeringii, Ed. 2.) — Rather dry sandy pine-barrens, New Jersey and southward. — Stems $1^{\circ} - 3^{\circ}$ long: leaves $1' - 1\frac{1}{2}'$ long. Corolla $4'' - 5''$ broad.

6. EVÓLVULUS, L. EVOLVULUS.

Calyx of 5 sepals, naked at the base. Corolla open funnel-form or almost wheel-shaped. Styles 2, each 2-cleft; stigmas obtuse. Pod 2-celled; the cells

2-seeded. — Low and small herbs or suffrutescent plants, mostly diffuse, never twining (hence the name, from *evolve*, to unroll, in contrast with *Convolvulus*).

1. **E. argenteus**, Pursh. Many-stemmed from a somewhat woody base, dwarf, silky-villous all over; leaves crowded, broadly lanceolate, sessile, or the lower oblong-spatulate and short-petioled, about $\frac{1}{2}$ ' long; flowers almost sessile in the axils; corolla purple, 3" broad. — Potosi lead-mines, Missouri, probably also on the Illinois side of the Mississippi: common westward.

7. DICHÓNDRÁ, Forst. DICHONDRA.

Calyx 5-parted. Corolla broadly bell-shaped, 5-cleft. Stamens included. Styles, ovaries, and the utricular 1-2-seeded pods 2, distinct. Stigmas thick. — Small and creeping perennial herbs, soft-pubescent, with kidney-shaped entire leaves, and axillary 1-flowered bractless peduncles. Corolla small, yellowish or white. (Name composed of *δῖς*, *double*, and *χόνδρος*, *roundish mass*; from the fruit.)

1. **D. rèpens**, Forst.: var. **Carolinénsis**, Choisy. Leaves round-kidney-shaped, pubescent, green both sides; corolla not exceeding the calyx (1" - 1 $\frac{1}{2}$ " long). (*D. Carolinensis*, *Michx.*) — Moist ground, Virginia, near Norfolk, and southward. (Widely diffused in the Southern hemisphere.)

8. CÚSCUTA, Tourn. DODDER.

Calyx 5- (rarely 4-) cleft, or of 5 sepals. Corolla globular-urn-shaped, bell-shaped, or somewhat tubular, the spreading border 5- (rarely 4-) cleft. Stamens furnished with a scale-like often fringed appendage at their base. Ovary 2-celled, 4-ovuled: styles distinct, or rarely united. Pod mostly 4-seeded. Embryo thread-shaped, spirally coiled in the rather fleshy albumen, destitute of cotyledons! sometimes with a few alternate scales (belonging to the plumule): germination occurring in the soil. — Leafless herbs, chiefly annuals, yellowish or reddish in color, with thread-like stems, bearing a few minute scales in place of leaves; on rising from the ground becoming entirely parasitic on the bark of herbs and shrubs on which they twine, and to which they adhere by means of papillæ developed on the surface in contact. Flowers small, cymose-clustered, mostly white; usually produced late in summer and in autumn. (Name of uncertain, supposed to be of Arabic, derivation.)

The following account of our species is contributed by DR. ENGELMANN, whose monograph of the whole genus is published in Transactions of the St. Louis Academy of Science.

§ 1. *Stigmas elongated: pod opening regularly around the base by circumcissile dehiscence, leaving the partition behind.* (*Natives of the Old World.*)

1. **C. EPFLINUM**, Weihe. (FLAX DODDER.) Stems very slender, low, flowers globular, sessile in dense scattered heads; corolla 5-parted, short-cylindrical, scarcely exceeding the broadly ovate acute divisions of the calyx, left surrounding the pod in fruit; stamens shorter than the limb; scales short, broad, crenulate, shorter than the globose ovary. — Flax-fields; in Europe very injurious: sparingly introduced with flax-seed into the Northern States. June. (Adv. from Eu.)

§ 2. *Stigmas capitate: pods indehiscent, rarely bursting irregularly.*

* *Flowers more or less pedicelled: the scaly bracts few and distant: calyx 4-5-cleft.*

+ *Corolla cylindrical, in fruit covering the top of the pod.*

2. **C. tenuiflora**, Engelm. Much branched, twining high, pale-colored; flowers at length peduncled and in rather loose cymes; tube of the corolla (ventricose after flowering) twice the length of its obtuse spreading lobes and of the ovate obtuse calyx-lobes; scales ovate, cut-fringed; stamens shorter than the lobes of the corolla; pod depressed, membranaceous, thin, yellowish. (*C. Cephalánthi*, Engelm.) — Swamps, New Jersey to Illinois and westward; on *Cephalanthus* and other shrubs, and on various tall herbs. — Flower the narrowest of all our Northern species.

3. **C. inflexa**, Engelm. Flowers peduncled, in umbel-like cymes, 1" long; tube of the mostly 4-cleft fleshy corolla as long as the ovate acutish and minutely crenate erect inflexed lobes and the acute keeled calyx-lobes; scales minute and few-toothed, appressed; pod depressed, somewhat umbonate, of a thicker texture, brown, its top covered with the remains of the corolla. (*C. Córyli*, Engelm. *C. umbrösa*, Beyrich, and Ed. 2.) — Prairies and barrens, in rather dry soil, on *Hazels*, *Ceanothus*, and other shrubs or herbs; from Western Virginia and Illinois southward and westward.

4. **C. decora**, Chois., altered by Engelm. Flowers larger than in No. 3, from 1½" to nearly 2" long, loosely paniculate, broadly campanulate; corolla 5-cleft; the lobes ovate-lanceolate, acute; the scales large, broadly oval; pod enveloped by the remains of the corolla. (*C. indecöra*, Chois. *C. neuropetala*, Engelm. *C. pulcherrima*, Scheele.) — Wet prairies, S. W. Illinois and southward; on various shrubs and herbs. — The name changed by Dr. Engelmann, because this is not a homely but the handsomest of our species.

+ + *Corolla bell-shaped, persistent at the base of the ripe pod.*

5. **C. arvensis**, Beyrich. Low; flowers small, 5-parted, peduncled in loose umbel-like cymes; tube of the corolla included in or little exceeding the broad-lobed calyx, shorter than its lanceolate acuminate spreading or reflexed lobes; stamens much shorter than the lobes of the corolla; scales ovate, fimbriate, converging and often exceeding the tube; pod globose, thin, yellowish. (*C. pentágona*, Engelm.) — In fields, prairies, and barrens, from S. New York (*C. F. Austin*) to Virginia, Illinois, and southwestward; on smaller herbs, and flowering (in June and July) earlier than any other of our species. — Stems low, scarcely over a foot high; flowers smaller than in any of our species, and quite variable: when with a large 5-angled calyx it is *C. pentágona* (Virginia): with a small one, it is var. *microcalyx* (Illinois): with a large and hemispherical one, var. *calycina* (Texas): with a fleshy verrucose calyx, it is *C. verrucösa*, Engelm. (Texas.)

6. **C. chlorocarpa**, Engelm. Low, orange-colored; flowers mostly 4-cleft, about 1" long, short-pedicelled, in scattered clusters; corolla open bell-shaped, the tube nearly the length of the acute lobes and calyx-teeth; stamens as long as the lobes; scales small, appressed, incised, sometimes almost wanting; the thick styles as long as the large depressed ovary; pod depressed, thin, greenish-yellow. (*C. polygonörum*, Engelm.) — Low grounds, on *Polygonum*

and other herbs, Delaware to Wisconsin, and southwestward. — The large ovary fills the shallow tube of the corolla.

7. **C. Gronovii**, Willd. Stems coarse, climbing high; flowers mostly 5-cleft, peduncled, in close or mostly open paniculate cymes; corolla bell-shaped, the tube longer than (or sometimes only as long as) the ovate obtuse entire spreading lobes; scales large, converging, copiously fringed, confluent at the base; pod globose, umbonate, brown. (*C. Americana*, Pursh, &c. *C. vulgivaga*, Engelm. *C. umbrösa*, Torr.) — Low, damp grounds, especially in shady places; everywhere common both east and west, and the principal species northward and eastward: chiefly on coarser herbs and low shrubs. — The close-flowered forms occur in the Northeastern States; the loosely-flowered ones westward and southward; a form with 4-parted flowers was collected in Connecticut. *C. Saurüri*, Engelm., is a form with more open flowers, of a finer texture, in the Mississippi valley.

8. **C. rostrata**, Shuttleworth. Stems coarse, climbing high; flowers (2''–3'' long) 5-parted, peduncled, in umbel-like cymes; corolla deep bell-shaped, the tube twice as long as the ovate obtuse teeth of the calyx and its ovate obtuse entire spreading lobes; the large scales fimbriate, confluent at the base; styles slender, as long as the acute ovary; the large pod pointed. — Shady valleys of the Alleghanies, from Maryland and Virginia southward; on tall herbs, rarely on shrubs. Flowers and fruit larger than in any other of our species.

* * *Flowers sessile in compact and mostly continuous clusters: calyx of 5 separate sepals surrounded by numerous similar bracts: remains of the corolla borne on the top of the globose somewhat pointed pod.* (*Lepidánche*, Engelm.)

9. **C. compacta**, Juss. Stems coarse; bracts (3–5) and sepals orbicular, concave, slightly crenate, appressed, nearly equalling or much shorter than the cylindrical tube of the corolla; stamens shorter than the oblong obtuse spreading lobes of the latter; scales pinnatifid-fringed, convergent, confluent at the base. *C. coronata*, Beyrich (*C. compacta*, Choisy) is the Eastern and Southern form, with a smaller, slenderer, more exerted corolla. *C. (Lepidanche) adpressa*, Engelm., is the Western form, with a larger, shorter, nearly included corolla. Both grow almost entirely on shrubs; the first from N. New York, and New Jersey southward; the latter from Western Virginia to the Mississippi and Missouri, in fertile shady bottoms. The clusters in fruit are sometimes fully 2' in diameter.

10. **C. glomerata**, Choisy. Flowers very densely clustered, forming knotty masses closely encircling the stem of the foster plant, much imbricated with scarious oblong bracts, their tips recurved-spreading; sepals nearly similar, shorter than the oblong-cylindrical tube of the corolla; stamens nearly as long as the oblong-lanceolate obtuse spreading or reflexed lobes of the corolla; scales large, fringed-pinnatifid; styles slender, longer than the pointed ovary; the pointed pod mostly 1–2-seeded. (*Lepidanche Compositarum*, Engelm.) — Moist prairies, Ohio to Wisconsin and southward: growing commonly on tall Compositæ. — The orange-colored stems soon disappear, leaving only the close matted coils of flowers, appearing like whitish ropes twisted around the stems.

ORDER 76. SOLANACEÆ. (NIGHTSHADE FAMILY.)

Herbs (or rarely shrubs), with a colorless juice and alternate leaves, regular 5-merous and 5-androus flowers, on bractless pedicels; the corolla imbricate, convolute, or valvate in the bud, and mostly plaited; the fruit a 2-celled (rarely 3-5-celled) many-seeded pod or berry. — Seeds campylotropous or amphitropous. Embryo mostly slender and curved in fleshy albumen. Calyx usually persistent. Stamens mostly equal, inserted on the corolla. Style and stigma single. Placentæ in the axis, often projecting far into the cells. (Foliage rank-scented, and with the fruits mostly narcotic, often very poisonous, while some are edible.) — A large family in the tropics, but very few indigenous in our district. It shades off into Scrophulariaceæ, from which the plaited regular corolla and 5 equal stamens generally distinguish it.

* Corolla wheel-shaped, 5-parted or 5-lobed; the lobes valvate and their margins usually turned inwards in the bud. Anthers connivent. Fruit a berry.

1. **Solanum.** Anthers opening by pores or chinks at the tip.

* * Corolla various, not wheel-shaped, nor valvate in the bud. Anthers separate.

← Fruit a berry, enclosed in the bladderly-inflated calyx. Corolla widely expanding.

2. **Physalis.** Calyx 5-cleft. Corolla 5-lobed or nearly entire. Berry juicy, 2-celled.

3. **Nicandra.** Calyx 5-parted. Corolla nearly entire. Berry dry, 3-5-celled.

← ← Fruit a berry with the unaltered calyx persistent at its base.

4. **Lycium.** Corolla funnel-form or tubular, not plaited. Berry small, 2-celled.

← ← ← Fruit a pod.

5. **Hyoscyamus.** Calyx urn-shaped, enclosing the smooth 2-celled pod, which opens by the top falling off as a lid. Corolla and stamens somewhat irregular.

6. **Datura.** Calyx prismatic, 5-toothed. Pod prickly, naked, more or less 4-celled, 4-valved. Corolla funnel-form.

7. **Nicotiana.** Calyx tubular-bell-shaped, 5-cleft. Pod enclosed in the calyx, 2-celled.

1. SOLANUM, Tourn. NIGHTSHADE.

Calyx and the wheel-shaped corolla 5-parted or 5-cleft (rarely 4-10-parted), the latter plaited in the bud, and valvate or induplicate. Stamens exerted: filaments very short: anthers converging around the style: opening at the tip by two pores or chinks. Berry usually 2-celled. — Herbs, or shrubs in warm climates, the larger leaves often accompanied by a smaller lateral (raméal) one; the peduncles also mostly lateral and extra-axillary. — A vast genus, chiefly in the warmer regions, including the POTATO (*S. TUBEROSUM*) and the EGG-PLANT (*S. MELONGÈNA*); while the TOMATO (*LYCOPÉRSICUM ESCULÉNTUM*) is hardly of a distinct genus. (Name of unknown derivation.)

* *Anthers blunt. (Plants not prickly, smooth or nearly so.)*

1. **S. DULCAMARA, L.** (BITTERSWEET.) *Stem shrubby, scarcely climbing; leaves ovate-heart-shaped, the upper halberd-shaped, or with two ear-like lobes at the base; flowers (purple or blue) in small cymes; berries oval, red. — Moist banks and around dwellings. June-Sept. (Nat. from Eu.)*

2. **S. NIGRUM, L.** (COMMON NIGHTSHADE.) *Annual, low, much branched and often spreading, rough on the angles; leaves ovate, wavy-toothed; flowers*

(very small, white) in *small and umbel-like lateral clusters*, drooping; berries globular, black.— Shaded grounds and fields: common. July–Sept.— A homely weed, said to be poisonous. (Nat. from Eu.)

* * *Anthers elongated, lanceolate, pointed.* (Plants mostly prickly.)

3. **S. Carolinense**, L. (HORSE-NETTLE.) Perennial, low (1° high); stem erect, prickly; leaves ovate-oblong, acute, sinuate-toothed or angled, roughish with stellate pubescence, prickly along the midrib, as also the calyx; flowers (pale blue or white, large) in simple loose racemes; berries globular, orange-yellow.— Sandy soil, Connecticut to Illinois and southward. June–Aug. (S. *Virginianum*, L., is not here identified as distinct.)

2. PHÝSALIS, L. GROUND CHERRY.

Calyx 5-cleft, reticulated and enlarging after flowering, at length much inflated and enclosing the 2-celled globular (edible) berry. Corolla between wheel-shaped and funnel-form, the very short tube marked with 5 concave spots at the base; the plaited border somewhat 5-lobed or barely 5–10-toothed. Stamens 5, erect: anthers separate, opening lengthwise.— Herbs (in this country), with the leaves often unequally in pairs, and the 1-flowered nodding peduncles extra-axillary; flowering through the summer. (Name, *φυσάλις*, a bladder, from the inflated calyx.)

* *Root annual: anthers tinged with blue or violet: stems 1°–3° high.*

+ *Corolla white, large.*

1. **P. grandiflora**, Hook. Clammy-pubescent, erect; leaves lance-ovate, pointed, entire or nearly so; corolla 1'–2' wide when expanded, almost entire, and with a woolly ring in the throat; fruiting calyx globular, apparently nearly filled by the berry.— Upper Michigan, shore of Lake Superior (*Dr. Robbins, &c.*) and northward, springing up in new clearings.

+ + *Corolla pale or greenish-yellow, small or smallish.*

2. **P. Philadelphica**, Lam. Almost glabrous, erect; leaves ovate or oblong-ovate, oblique at base, entire, repand, or very sparingly angulate-toothed; corolla brownish- or violet-spotted in the centre, 7''–10'' broad; calyx at maturity globose and completely filled by the large reddish or purple berry and open at the mouth.— Rich grounds, not rare, especially southward: also cult.

3. **P. angulata**, L. Glabrous or nearly so, erect, much branched; leaves ovate or ovate-oblong, sharply and irregularly *laciniate-toothed*; peduncles filiform; corolla unspotted, very small (3''–6'' broad when expanded); fruiting calyx conical-ovate with a truncate or sunken base, 10-angled, loosely inflated, but at length well filled by the greenish-yellow berry.— Cult. and waste grounds.

4. **P. pubescens**, L. Pubescent or clammy-hairy (rarely smoothish) *diffusely much branched or at length decumbent*; leaves ovate or heart-shaped, angulate- or repand-toothed; corolla spotted with brown-purple in the centre, 5''–6'' broad when expanded, obscurely 5–10-toothed; fruiting calyx ovate from a truncate or impressed base, pointed, sharply 5-angled, loosely enclosing the yellow or greenish berry. (*P. hirsuta*, Dunal. *P. obscura*, Michx. in part.)— Low grounds: common, especially southward and westward.

* * *Root perennial: stems mostly from slender creeping rootstocks, usually low (6' - 20' high): anthers yellow: fruiting calyx loosely inflated, 5-angled, much larger than the berry.*

+ *Wild species: corolla greenish-yellow and commonly brown or purplish in the centre, the border 5-angled or barely 5 - 10-toothed, 6'' - 12'' broad.*

5. **P. viscosa**, L. *Clammy-pubescent, diffusely much branched and widely spreading, or at first erect; leaves ovate or slightly heart-shaped, sometimes oblong, repand or obtusely toothed, rarely entire; corolla dark brown in the centre; fruiting calyx ovate, barely concave or truncate at base, sharply 5-angled; berry orange or reddish, glutinous. (P. heterophylla, Nees. P. nyc-taginea, & P. viscido-pubescent, Dunal?)* — Light or sandy soil: common.

6. **P. Pennsylvànica**, L. *Minutely hirsute-pubescent (not clammy), or nearly glabrous; leaves ovate, oblong, or oblong-lanceolate and tapering at the base, entire or sparingly repand-toothed; corolla merely darker or purplish-veiny in the centre; fruiting calyx conical or globular-ovate, pointed, and with an impressed base; berry red.* — Var. **LANCEOLATA**; the narrower-leaved and pubescent form (5' - 15' high), especially the state with a hairy calyx. (*P. lanceolata, Michx. P. maritima, M. A. Curtis.*) — Dry, often sandy soil, from Pennsylvania southward and westward, even northwestward to the Winipeg valley. — Fruiting calyx 1' - 1½' long. Shape of calyx-lobes very variable.

+ + *Introduced: corolla greenish-white, unspotted, 5-lobed.*

7. **P. ALKEKÉNGI**, L. (**STRAWBERRY TOMATO.**) More or less pubescent; sparingly branched; leaves deltoid-ovate, pointed; calyx-teeth awl-shaped; fruiting calyx broadly ovate, turning red; the berry bright red, pleasant. — Cult. and waste grounds, eastward. (Cult. & Adv. from Eu.)

3. NICÁNDRA, Adans. APPLE OF PERU.

Calyx 5-parted, 5-angled, the divisions rather arrow-shaped, enlarged and bladder-like in fruit, enclosing the 3 - 5-celled globular dry berry. Corolla with border nearly entire. Otherwise much like *Physalis*. — An annual smooth herb (2° - 3° high), with ovate sinuate-toothed or angled leaves, and solitary pale blue flowers on axillary and terminal peduncles. (Named after the poet *Nican-der* of Colophon.)

1. **N. PHYSALOÏDES**, Gærtn. — Waste grounds. (Adv. from Peru.)

4. LÝCIUM, L. MATRIMONY-VINE.

Calyx 3 - 5-toothed or cleft, not enlarging, persistent at the base of the berry. Corolla funnel-form or salver-shaped, 5-lobed, the lobes imbricated and not plaited in the bud. Stamens 5: anthers opening lengthwise. Style slender: stigma capitate. Berry small, 2-celled. Shrubby, often spiny plants, with alternate and entire small leaves, and mostly axillary small flowers. (Named from the country, *Lycia*.)

1. **L. VULGÀRE**, Dunal. (**COMMON M.**) Shrub with long sarmentose recurved-drooping branches, smooth, sparingly if at all spiny; leaves oblong- or spatulate-lanceolate, often fascicled, narrowed into a short petiole; flowers on slender peduncles fascicled in the axils; corolla short funnel-form, greenish-

purple; style and slender filaments equalling its lobes; berry oval, orange-red. (*L. Bárbarum*, *L.*, in part.) — About dwellings; and escaped into waste grounds in Pennsylvania, &c. June–Aug. (Adv. from Eu.)

5. HYOSCÿAMUS, Tourn. HENBANE.

Calyx bell-shaped or urn-shaped, 5-lobed. Corolla funnel-form, oblique, with a 5-lobed more or less unequal plaited border. Stamens declined. Pod enclosed in the persistent calyx, 2-celled, opening transversely all round near the apex, which falls off like a lid. — Clammy-pubescent, fetid, narcotic herbs, with lurid flowers in the axils of angled or toothed leaves. (Name composed of *ÿs*, *ÿós*, a hog, and *κÿάμος*, a bean; said by *Ælian* to be poisonous to swine.)

1. *H. NÿGER*, *L.* (BLACK HENBANE.) Biennial or annual; leaves clasping, sinuate-toothed and angled; flowers sessile, in one-sided leafy spikes; corolla dull yellowish, strongly reticulated with purple veins. — Escaped from gardens to roadsides. (Adv. from Eu.)

6. DATÿRA, L. JAMESTOWN-WEED. THORN-APPLE.

Calyx prismatic, 5-toothed, separating transversely above the base in fruit, the upper part falling away. Corolla funnel-form, with a large and spreading 5–10-toothed plaited border. Stigma 2-lipped. Pod globular, prickly, 4-valved, 2-celled, with 2 thick placentæ projected from the axis into the middle of the cells, and connected with the walls by an imperfect false partition, so that the pod is 4-celled except near the top, the placentæ as if on the middle of these false partitions. Seeds rather large, flat. — Rank weeds, narcotic-poisonous, with ovate leaves, and large and showy flowers on short peduncles in the forks of the branching stem; produced all summer and autumn. (Altered from the Arabic name, *Tatorah*.)

1. *D. STRAMÿNIUM*, *L.* (COMMON STRAMONIUM OR THORN APPLE.) Annual, glabrous; leaves ovate, sinuate-toothed or angled; stem green; corolla white (3' long), the border with 5 teeth. — Waste grounds: a well-known weed. (Adv. from Asia.)

2. *D. TÁTULA*, *L.* (PURPLE T.) Mostly taller; stem purple; corolla pale violet-purple. Thought to be specifically distinct from the last, on account of the behavior of the cross-breeds. (Adv. from trop. Amer.?)

7. NICOTIÀNA, L. TOBACCO.

Calyx tubular-bell-shaped, 5-cleft. Corolla funnel-form or salver-form, usually with a long tube; the plaited border 5-lobed. Stigma capitate. Pod 2-celled, 2–4-valved from the apex. Seeds minute. — Rank acrid-narcotic herbs, mostly clammy-pubescent, with ample entire leaves, and racemed or paniced flowers. (Named after *John Nicot*, who was thought to have introduced the Tobacco (*N. TABACUM*, *L.*) into Europe.)

1. *N. RÿSTICA*, *L.* (WILD TOBACCO.) Annual; leaves ovate, petioled; tube of the dull greenish-yellow corolla cylindrical, two thirds longer than the calyx, the lobes rounded. — Old fields, from New York westward and southward: a relic of cultivation by the Indians. (Adv. from Trop. Amer.)

ORDER 77. GENTIANACEÆ. (GENTIAN FAMILY.)

Smooth herbs, with a colorless bitter juice, opposite and sessile entire and simple leaves (except in Tribe II.) without stipules, regular flowers with the stamens as many as the lobes of the corolla, which are convolute (rarely imbricated and sometimes valvate) in the bud, a 1-celled ovary with 2 parietal placentæ, or nearly the whole inner face of the ovary ovuliferous; the fruit usually a 2-valved and septical many-seeded pod. — Flowers solitary or cymose. Calyx persistent. Corolla mostly withering-persistent; the stamens inserted on its tube. Seeds anatropous, with a minute embryo in fleshy albumen. (Bitter-tonic plants.)

Tribe I. GENTIANEÆ. Lobes of the corolla convolute in the bud (with the sinuses mostly plaited), or in *Obolaria* imbricated. Leaves almost always opposite or whorled, entire, those of the stem sessile. Seeds very small and numerous, with a cellular coat; in *Obolaria*, *Bartonia*, and several *Gentians*, the ovules and seeds covering the whole face of the pericarp!

* Style distinct and slender, deciduous.

1. **Sabbatia.** Corolla wheel-shaped, 5-12-parted: anthers at length recurved.

2. **Erythraea.** Corolla funnel-form or salver-shaped, 4-5-cleft: anthers soon spiral.

** Style (if any) and stigmas persistent: anthers straight.

3. **Frasera.** Corolla 4-parted, wheel-shaped; a fringed glandular spot on each lobe.

4. **Halenia.** Corolla 4-5-cleft, bell-shaped, and 4-5-spurred at the base.

5. **Gentiana.** Corolla funnel- or bell-shaped, mostly plaited in the sinuses, not spurred.

6. **Bartonia.** Corolla deeply 4-cleft, bell-shaped: no plaits. Calyx 4-parted.

7. **Obolaria.** Corolla tubular-bell-shaped, 4-lobed, with no plaits, the lobes imbricated in the bud! Calyx 2-leaved.

Tribe II. MENYANTHEÆ. Lobes of the corolla valvate in the bud, with the edges turned inwards. Stem-leaves alternate, petioled. Seed-coat hard or bony.

8. **Menyanthes.** Corolla bearded inside. Leaves 3-foliolate.

9. **Limnanthemum.** Corolla naked or bearded on the margins only. Leaves simple, rounded.

1. SABBÀTIA, Adans. AMERICAN CENTAURY.

Calyx 5-12-parted, the divisions slender. Corolla 5-12-parted, wheel-shaped. Stamens 5-12: anthers soon recurved. Style 2-parted, slender. — Biennials or annuals, with slender stems, and cymose-panicled handsome (white or rose-purple) flowers; in summer. (Dedicated to *L. Sabbati*, an early Italian botanist.)

* Corolla 5-parted, or rarely 6-7-parted.

+ Corolla white, often turning yellowish in drying: cymes corymbed, many-flowered.

1. **S. paniculata**, Pursh. Stem brachiately much-branched (1°-2° high), rather terete, but angled with 4 sharp lines; leaves linear or the lower oblong, obtuse, 1-nerved, nearly equalling the internodes; calyx-lobes linear-thread-form, much shorter than the corolla. — Low grounds, Virginia and southward.

2. **S. lanceolata**, Torr. & Gr. Stem simple (1°-3° high) bearing a flat-topped cyme; leaves ovate-lanceolate or ovate, 3-nerved, the upper acute, much shorter than the internodes; calyx-lobes longer and flowers larger than in

No. 1. (*Chironia lanceolata*, Walt. *S. corymbosa*, Baldw.)—Wet pine barrens, from New Jersey southward.

+ + Corolla rose-pink, rarely white, with a yellowish or greenish eye: stem erect, 1°–3° high, pyramidally many-flowered: branches opposite: peduncles short.

3. *S. brachiata*, Ell. Stem slightly angled, simple below (1°–2° high); leaves linear and linear-oblong, obtuse, or the upper acute; branches rather few-flowered, forming an oblong panicle; calyx-lobes nearly half shorter than the corolla. (*S. concinna*, Wood, ex char.)—Dryish grassy places, Virginia (Indiana, Wood), and southward.—Corolla rather smaller, and its lobes narrower than in the next.

4. *S. angularis*, Pursh. Stem somewhat 4-winged-angled, much branched above (1°–2½° high), many-flowered; leaves ovate, acutish, 5-nerved, with a somewhat heart-shaped clasping base; calyx-lobes one third or half the length of the corolla.—Dry ground, New York to Illinois and southward. Corolla 1½' wide; the lobes obovate.

+ + + Corolla rose-purple or white: stems (5'–20' high) slender, loosely and often alternately branched, or merely forked, terete or scarcely 4-angled: peduncles elongated and 1-flowered.

5. *S. calycosa*, Pursh. Diffusely forking, pale; leaves oblong or lance-oblong, narrowed at the base; calyx-lobes foliaceous, spatulate-lanceolate (¾'–1' long), exceeding the almost white corolla.—Marshes, E. Virginia, and southward.

6. *S. stellaris*, Pursh. Loosely branched and forking; leaves oblong- or ovate-lanceolate, or the upper linear; calyx-lobes awl-shaped-linear, varying from half to nearly the length of the bright rose-purple corolla.—Salt marshes, Massachusetts to Virginia, and southward. Too near the next.

7. *S. gracilis*, Salisb. Stem very slender, at length diffusely branched; the branches and long peduncles filiform; leaves linear, or the lower lance-linear, the uppermost similar to the setaceous calyx-lobes, which equal the rose-purple corolla. (*Chironia campanulata*, L.)—Brackish marshes, Nantucket (*Oakes*), banks of lower Delaware River (*Mr. Cooley, Dr. Diffenbaugh*), and southward.

* * Corolla 9–12-parted, large (about 2' broad). (*Lapitheia, Grisebach.*)

8. *S. chloroides*, Pursh. Stem (1°–2° high), loosely paniced above; the peduncles slender, 1-flowered; leaves oblong-lanceolate; calyx-lobes linear, half the length of the deep rose-colored (rarely white) corolla.—Borders of brackish ponds, Plymouth, Massachusetts, to Virginia, and southward.—One of our handsomest plants.

2. ERYTHRÆA, Pers. CENTAURY.

Calyx 4–5-parted, the divisions slender. Corolla funnel-form or salver-form, with a slender tube and a 4–5-parted limb. Anthers exserted, erect, twisting spirally. Style slender, single: stigma capitate or 2-lipped.—Low and small branching annuals, chiefly with rose-purple or reddish flowers (whence the name, from *έρυθρός*, red); in summer. All our Northern species were probably introduced, and occur in few localities.

1. *E. CENTAURIUM*, Pers. (CENTAURY.) Stem upright, corymbosely branched above; leaves oblong or elliptical, acutish; the uppermost linear; cymes clus-

tered, flat-topped, the flowers all nearly sessile; tube of the (purple-rose-colored) corolla not twice the length of the oval lobes. — Oswego, New York. — Plant 6'–12' high: corolla 3"–4" long. (Adv. from Eu.)

2. **E. RAMOSÍSSIMA**, Pers., var. **PULCHÉLLA**, Griseb. Low (2'–6' high); stem many times forked above and forming a diffuse cyme; leaves ovate-oblong or oval; flowers all on short pedicels; tube of the (pink-purple) corolla thrice the length of the elliptical-oblong lobes. (*E. Muhlenbergii*, Griseb., as to Penn. plant. *Éxacum pulchellum*, Pursh.) — Wet or shady places, Long Island to E. Virginia: scarce. — Flowers smaller than in No. 1. (Nat. from Eu.)

3. **E. SPICATA**, Pers. Stem strictly upright (6'–10' high); the flowers sessile and spiked along one side of the simple or rarely forked branches; leaves oval and oblong, rounded at the base, acutish; tube of the (rose-colored or whitish) corolla scarcely longer than the calyx, the lobes oblong. (*E. Pickeringii*, Oakes.) — Sandy sea-shore, Nantucket, Massachusetts, Oakes, and Norfolk, Virginia, Rugel. — Remarkable for the spike-like arrangement of the flowers. (Nat. from Eu.?)

3. FRÀSERA, Walt. AMERICAN COLUMBO.

Calyx deeply 4-parted. Corolla deeply 4-parted, wheel-shaped, each division with a glandular and fringed pit on the face. Filaments awl-shaped, usually somewhat monadelphous at the base: anthers oblong, versatile. Style persistent: stigma 2-lobed. Pod oval, flattened, 4–14-seeded. Seeds large and flat, wing-margined. — Tall and showy herbs, with a thick root, upright and mostly simple stems, bearing whorled leaves, and numerous peduncled flowers in open cymes, which are disposed in an ample elongated panicle. (Dedicated to *John Fraser*, an indefatigable collector in this country towards the close of the last century.)

1. **F. Carolinensis**, Walt. Smooth biennial or triennial (3°–8° high); leaves mostly in fours, lance-oblong, the lowest spatulate, veiny; panicle pyramidal, loosely flowered; divisions of the corolla oblong, mucronate, longer than the narrowly lanceolate calyx-lobes, each with a large and round gland on their middle; pod much flattened parallel with the flat valves. — Rich dry soil, S. W. New York to Wisconsin and southward. July. — Root very thick and bitter. Corolla 1' broad, light greenish-yellow, marked with small brown-purple dots.

4. HALÈNIA, Borkh. SPURRED GENTIAN.

Calyx 4–5-parted. Corolla short bell-shaped, 4–5-cleft, without folds or fringe, prolonged at the base underneath the erect lobes into spurs, which are glandular in the bottom. Stigmas 2, sessile, persistent on the oblong flattish pod. Seeds rather numerous, oblong. — Small and upright herbs, with yellowish or purplish paniced-cymose flowers. (Name of unknown meaning.)

1. **H. defléxa**, Grisebach. Leafy annual or biennial (9'–18' high), simple or branched above; leaves 3–5-nerved, the lowest oblong-spatulate and petioled; the others oblong-lanceolate, acute; spurs cylindrical, obtuse, curved, descending, half the length of the acutely 4-lobed corolla. (*Swértia corniculata*,

L., partly.)—Damp woods, from the northern parts of Maine, to Wisconsin, and northward. July, August.

5. GENTIÀNA, L. GENTIAN.

Calyx 4-5-cleft. Corolla 4-5-lobed, regular, usually with intermediate plaited folds, which bear appendages or teeth at the sinuses. Style short or none: stigmas 2, persistent. Pod oblong, 2-valved; the innumerable seeds either borne on placentæ at or near the sutures, or in most of our species covering nearly the whole inner face of the pod. (First shown by *Prof. H. J. Clark!*)—Flowers solitary or cymose, showy, in late summer and autumn. (Name from *Gentius*, king of Illyria, who used some species medicinally.)

§ 1. AMARELLOÏDES, Torr. & Gr. *Corolla tubular-funnel-form, without crown or plaited folds, and with the lobes naked: anthers separate, fixed by the middle, introrse in the bud, but reflexed after the flower opens: seeds wingless: annuals.*

1. **G. quinqueflora**, Lam. (FIVE-FLOWERED G.) Stem rather slender, branching (1° high); leaves ovate-lanceolate from a partly clasping and heart-shaped base, 3-7-nerved, tipped with a minute point; branches racemed or paniced, about 5-flowered at the summit; lobes of the small 5-cleft calyx awl-shaped-linear; lobes of the pale-blue corolla triangular-ovate, bristle-pointed, one fourth the length of the slender obconical tube.—Var. OCCIDENTALIS has linear-lanceolate calyx-lobes, more leaf-like, about half the length of the corolla.—Dry hilly woods, Maine to Wisconsin and southward, especially along the Alleghanies: the var. is the common form in the Western States.—Corolla nearly 1' long; in the variety proportionally shorter.

§ 2. CROSSOPÉTALUM, Frœl. *Corolla funnel-form, gland-bearing between the bases of the filaments, without crown or plaited folds; the lobes fringed or toothed on the margins: anthers as in § 1: pod somewhat stalked: seeds wingless, clothed with little scales: annuals or biennials.*

2. **G. crinita**, Frœl. (FRINGED G.) Flowers solitary on long peduncles terminating the stem or simple branches; leaves lanceolate or ovate-lanceolate from a partly heart-shaped or rounded base; lobes of the 4-cleft calyx unequal, ovate and lanceolate, as long as the bell-shaped tube of the sky-blue corolla, the lobes of which are wedge-obovate, and strongly fringed around the summit; ovary lanceolate.—Low grounds, New England to Kentucky and Wisconsin: rather common.—Plant 1°-2° high: the showy corolla 2' long.

3. **G. detónsa**, Fries. (SMALLER FRINGED G.) Stem simple or with slender branches, terminated by solitary flowers on very long peduncles; leaves linear or lanceolate-linear; lobes of the 4- (rarely 5-) cleft calyx unequal, ovate or triangular and lanceolate, pointed; lobes of the sky-blue corolla spatulate-oblong, with ciliate-fringed margins, the fringe shorter or almost obsolete at the summit; ovary elliptical or obovate.—Moist grounds, Niagara Falls to Illinois and north-westward. Passes into the last. (Eu.)

§ 3. PNEUMONÁNTHE, Necker. *Corolla bell-shaped or obconical, 5-lobed, with plaited folds which project into appendages in the sinuses: anthers erect,*

fixed by the deep sagittate base, extrorse, often cohering with each other in a ring or tube: pod stalked: perennials, mostly autumn-flowering.

* *Flowers nearly sessile, clustered or rarely solitary, 2-bracteolate.*

+ *Seeds wingless: anthers unconnected.*

4. **G. ochroleuca**, Frœl. (YELLOWISH-WHITE G.) Stems ascending, mostly smooth; the flowers in a dense terminal cluster and often also in axillary clusters; *leaves obovate-oblong*, the lowest broadly obovate and obtuse, the uppermost somewhat lanceolate, *all narrowed at the base*; calyx-lobes linear, unequal, much longer than its tube, *rather shorter than the greenish-white open corolla*, which is painted inside with green veins and lilac-purple stripes; its lobes ovate, very much exceeding the small and sparingly toothed oblique appendages; pod included in the persistent corolla. — Dry grounds, S. Pennsylvania (rare) to Virginia, and common southward.

+ + *Seeds winged: anthers connivent and usually more or cohering.*

5. **G. álba**, Muhl. Cat. ! (WHITISH G.) Stems upright, stout, and very smooth; flowers closely sessile and much crowded in a dense terminal cluster, sometimes also clustered in the upper axils; *leaves ovate-lanceolate from a heart-shaped closely clasping base*, gradually tapering to a point; calyx-lobes ovate, shorter than the top-shaped tube, and many times shorter than the tube of the corolla, reflexed-spreading; *corolla white more or less tinged with greenish or yellowish, inflated-club-shaped, at length open, its short and broad ovate lobes nearly twice the length of the toothed appendages*; pod nearly included; seeds broadly winged. (*G. flavida*, Gray, in *Sill. Jour.* *G. ochroleuca*, Sims., *Darlingt.*, *Grisebach*, in part.) — Glades and low grounds, S. W. New York to Virginia along the Alleghanies, and west to Illinois and Lake Superior. Begins to flower in July, far earlier than the two next.

6. **G. Andréwsii**, Griseb. (CLOSED G.) Stems upright, smooth; flowers closely sessile in terminal and upper axillary clusters; *leaves ovate-lanceolate and lanceolate from a narrower base*, gradually pointed, rough-margined; calyx-lobes ovate or oblong, recurved, shorter than the top-shaped tube, and much shorter than the *inflated club-shaped and truncate mostly blue corolla, which is closed at the mouth, its proper lobes obliterated*, the apparent lobes consisting of the broad fringe-toothed and notched appendages; pod finally projecting out of the persistent corolla; seeds broadly winged. (*G. Saponaria*, Frœl., not of L.) — Moist and rich soil: common, especially northward. — Corolla an inch or more in length, striped inside, the folds whitish; occasionally *pure white* throughout.

7. **G. Saponaria**, L. (SOAPWORT G.) Stem erect or ascending, smooth; the flowers clustered at the summit and more or less so in the axils; *leaves ovate-lanceolate, oblong, or lanceolate-obovate*, with rough margins, narrowed at the base; calyx-lobes linear or spatulate, acute, equalling or exceeding the tube, half the length of the corolla; *lobes of the club-bell-shaped light-blue corolla obtuse, erect or converging, short and broad, but distinct, and more or less longer than the conspicuous 2-cleft and minutely toothed appendages*; seeds acute, narrowly winged. (*G. Catesbæi*, Walt.) — Moist woods, New Jersey and S. Penn. to Virginia, Illinois, and southward: flowering late.

Var. **linearis**. Slender, nearly simple (1°–2° high); leaves linear or lance-linear (2'–3' long), acutish; appendages of the corolla shorter and less cleft, or almost entire. (G. Pneumonánthe, *Amer. auth.*, & *Ed.* 1. G. linearis, *Frœl.*) — Mountain wet glades of Maryland and Penn., to Lake Superior, Northern New York, New Hampshire (near Concord), and Maine (near Portland): beginning to blossom at midsummer. — Seems to pass on one side into G. Saponaria, on the other into G. Pneumonánthe of Europe.

8. **G. pubérula**, Michx. Stems erect or ascending (8'–16' high), mostly rough and minutely pubescent above; leaves rigid varying from linear-lanceolate to oblong-lanceolate, rough-margined (1'–2' long); flowers clustered, rarely solitary; calyx-lobes lanceolate, not longer than the tube, much shorter than the bell-funnel-form open bright-blue corolla, the spreading ovate lobes of which are acutish and twice or thrice the length of the cut-toothed appendages. (G. Catesbæi, *Ell.* G. Saponaria, var. puberula, *Ed.* 1.) — Dry prairies and barrens, Ohio to Wisconsin, and southward. Flowering near the end of summer. Corolla large for the size of the plant, 1½'–2' long. Seeds (also in G. Pneumonánthe) not covering the walls, as they do in the rest of this division.

* * Flowers 1–3, peduncled: seeds wingless: anthers separate.

9. **G. angustifolia**, Michx. Stems slender and ascending (6'–15' high); leaves linear or the lower oblanceolate, rigid; corolla open-funnel-form, azure-blue, also a greenish and white variety (2' long), about twice the length of the thread-like calyx-lobes, its ovate spreading lobes twice the length of the cut-toothed appendages. — Moist pine barrens, New Jersey, and southward.

6. BARTONIA, Muhl. (CENTAURELLA, Michx.)

Calyx 4-parted. Corolla deeply 4-cleft, destitute of glands, fringes, or folds. Stamens short. Pod oblong, flattened, pointed with a large persistent at length 2-lobed stigma. Seeds minute, innumerable, covering the whole inner surface of the pod. — Small annuals or biennials (3'–10' high), with thread-like stems, and little awl-shaped scales in place of leaves. Flowers small, white, peduncled. (Dedicated, in the year 1801, to *Prof. Benjamin Smith Barton*, of Philadelphia.)

1. **B. tenella**, Muhl. Stems branched above; the branches or peduncles mostly opposite, 1–3-flowered; lobes of the corolla oblong, acutish, rather longer than the calyx, or sometimes twice as long; anthers roundish; ovary 4-angled, the cell somewhat cruciform. — Open woods, New England to Wisconsin and southward. Aug. — *Centaurella Moseri*, *Griseb.*, is a variety with the scales and peduncles mostly alternate, and the petals acute.

2. **B. vérna**, Muhl. Stem 1–few-flowered; lobes of the corolla spatulate, obtuse, spreading, thrice the length of the calyx; anthers oblong; ovary flat. — Bogs near the coast, Virginia and southward. March. — Flowers 3''–4'' long, larger than in No. 1.

7. OBOLARIA, L. OBOLARIA.

Calyx of 2 spatulate spreading sepals, resembling the leaves. Corolla tubular-bell-shaped, withering-persistent, 4-cleft; the lobes oval-oblong, or with age

spatulate, imbricated in the bud! Stamens inserted at the sinuses of the corolla, short. Style short, persistent: stigma 2-lipped. Pod ovoid, 1-celled, the cell cruciform: the seeds covering the whole face of the walls. — A low and very smooth purplish-green perennial (3' - 8' high), with a simple or sparingly branched stem, opposite wedge-obovate leaves; the dull white or purplish flowers solitary or in clusters of three, terminal and axillary, nearly sessile; in spring. (Name from ὀβολός, a small Greek coin; to which, however, the leaves of this plant bear no manifest resemblance.)

1. **O. Virginica**, L. (*Gray, Chlor. Bor.-Am., t. 3.*) — Rich soil, in woods, from New Jersey to Illinois, and southward: rather rare.

8. MENYÁNTHES, Tourn. BUCKBEAN.

Calyx 5-parted. Corolla short funnel-form, 5-parted, deciduous, the whole upper surface white-bearded, valvate in the bud with the margins turned inward. Style slender, persistent: stigma 2-lobed. Pod bursting somewhat irregularly, many-seeded. Seed-coat hard, smooth, and shining. — A perennial alternate-leaved herb, with a thickish creeping rootstock, sheathed by the membranous bases of the long petioles, which bear 3 oval or oblong leaflets at the summit; the flowers racemed on the naked scape (1° high), white or slightly reddish. (The ancient Theophrastian name, probably from μήν, month, and άνθος, a flower, some say from its flowering for about that time.)

1. **M. trifoliata**, L. — Bogs, New England to Pennsylvania, Wisconsin, and northward. May, June. (Eu.)

9. LIMNÁNTHÉMUM, Gmelin. FLOATING HEART.

Calyx 5-parted. Corolla almost wheel-shaped, 5-parted, the divisions fringed or bearded at the base or margins only, folded inwards in the bud, bearing a glandular appendage near the base. Style short or none: stigma 2-lobed, persistent. Pod few - many-seeded, at length bursting irregularly. Seed-coat hard. — Perennial aquatics, with rounded floating leaves on very long petioles, which, in most species, bear near their summit the umbel of (polygamous) flowers, along with a cluster of short and spur-like roots, sometimes shooting forth new leaves from the same place, and so spreading by a sort of proliferous stolons: flowering all summer. (Name compounded of λίμνη, a marsh or pool, and άνθεμον, a blossom, from the situations where they grow.)

1. **L. lacunòsum**, Grisebach (partly). *Leaves entire*, round-heart-shaped (1' - 2' broad), thickish; petioles filiform; lobes of the (white) corolla broadly oval, naked, except the crest-like yellowish gland at their base, twice the length of the lanceolate calyx-lobes; style none; *seeds smooth and even*. (*Villársia lacunosa*, Vent. *V. cordata*, Ell.) — Shallow water, from Maine and N. New York to Virginia and southward.

2. **L. trachyspérnum**, Gray. *Leaves larger* (2' - 6' broad), and rounder, thicker, often wavy-margined or crenate, roughish and dark-punctate or pitted beneath; petioles stouter; *seeds glandular-roughened*. (*Menyanthes trachysperma*, Michx.) — Ponds, Maryland, (*W. M. Canby*) and southward.

ORDER 78. LOGANIACEÆ. (LOGANIA FAMILY.)

Herbs, shrubs, or trees, with opposite and entire leaves, and stipules or a stipular membrane or line between them, and with regular 4-5-merous 4-5- androus perfect flowers, the ovary free from the calyx: a connecting group between Gentianaceæ, Apocynaceæ, Scrophulariaceæ (from all which they are known by their stipules) and Rubiaceæ, from which they differ in their free ovary: our representatives of the family are all most related to the Rubiaceæ, to which, indeed, they have been appended.

* Woody twiners: leaves evergreen.

1. **Gelsemium.** Corolla large, the 5 lobes imbricated in the bud. Style slender: stigmas 4.

** Herbs.

2. **Polypremum.** Corolla 4-lobed, not longer than the calyx, imbricated in the bud.

3. **Spigelia.** Corolla 5-lobed, valvate in the bud. Style single, jointed in the middle.

4. **Mitreola.** Corolla 5-lobed, valvate in the bud. Styles 2, short, converging, united at the summit, and with a common stigma.

1. GELSÉMIUM, Juss. YELLOW (FALSE) JESSAMINE.

Calyx 5-parted. Corolla open-funnel-form, 5-lobed; the lobes imbricated in the bud. Stamens 5, with oblong sagittate anthers. Style long and slender. Stigmas 2, each 2-parted; the divisions linear. Pod elliptical, flattened contrary to the narrow partition, 2-celled, septicidally 2-valved. Seeds many or several, winged. Embryo straight in fleshy albumen; the ovate flat cotyledons much shorter than the slender radicle. — Smooth and twining shrubby plants with opposite and entire ovate or lanceolate leaves, minute stipules, and showy yellow flowers, of two sorts as to relative length of stamens and style. (*Gelsemino*, the Italian name of the Jessamine.)

1. **G. sempervirens**, Ait. (YELLOW JESSAMINE of the South.) Stem climbing high; leaves short-petioled, shining, nearly persistent; flowers in short axillary clusters; pedicels scaly-bracted; flowers very fragrant (the bright yellow corolla 1' - 1½' long); pod flat, pointed. — Low grounds, Eastern Virginia and southward. March, April.

2. POLYPRÈMUM, L. POLYPREMUM.

Calyx 4-parted; the divisions awl-shaped from a broad scarious-margined base. Corolla not longer than the calyx, almost wheel-shaped, bearded in the throat; the 4 lobes imbricated in the bud. Stamens 4, very short: anthers globular. Style 1, very short: stigma ovoid, entire. Pod ovoid, a little flattened, notched at the apex, 2-celled, loculicidally 2-valved, many-seeded. — A smooth, diffuse, much-branched, small annual, with narrowly linear or awl-shaped leaves, connected at their base across the stem by a slight stipular line; the small flowers solitary and sessile in the forks and at the ends of the branches; corolla inconspicuous, white. (Name altered from *πολύπρεμος*, *many-stemmed*.)

1. **P. procumbens**, L. — Dry fields, mostly in sandy soil, Maryland and southward; also adventive at Philadelphia. June - Oct.

3. SPIGÈLIA, L. PINK-ROOT. WORM-GRASS.

Calyx 5-parted; the lobes slender. Corolla tubular-funnel-form, 5-lobed at the summit, valvate in the bud. Stamens 5: anthers linear. Style 1, slender, hairy above, jointed near the middle. Pod short, 2-celled, twin, laterally flattened, separating at maturity from a persistent base into 2 carpels, which open loculicidally, few-seeded. — Chiefly herbs, with the pair of leaves united by means of the stipules, and the flowers spiked in one-sided cymes. (Named for *Adrian Spiegel*, latinized *Spigelius*, who wrote on botany at the beginning of the seventeenth century, and was perhaps the first to give directions for preparing an herbarium.)

1. **S. Marilandica**, L. (MARYLAND PINK-ROOT.) Stems simple and erect from a perennial root (6'–18' high); leaves sessile, ovate-lanceolate, acute; spike simple or forked, short; tube of the corolla 4 times the length of the calyx, the lobes lanceolate; anthers and style exserted. — Rich woods, Pennsylvania to Wisconsin and southward: not common northward. June, July. — Corolla $1\frac{1}{2}$ ' long, red outside, yellow within. — A well-known official anthelmintic, and a showy plant.

4. MITRÈOLA, L. MITREWORT.

Calyx 5-parted. Corolla little longer than the calyx, somewhat funnel-form, 5-lobed, valvate in the bud. Stamens 5, included. Ovary at the base slightly adnate to the bottom of the calyx, 2-celled: styles 2, short, converging and united above; the stigmas also united into one. Pod projecting beyond the calyx, strongly 2-horned or mitre-shaped, opening down the inner side of each horn, many-seeded. — Annual smooth herbs, 6'–2° high, with small stipules between the leaves, and small white flowers spiked along one side of the branches of a terminal petioled cyme. (Name, a *little mitre*, from the shape of the pod.)

1. **M. petiolàta**, Torr. & Gray. Leaves thin, oblong-lanceolate, petioled. — Damp soil, from Eastern Virginia southward.

2. **M. sessilifòlia**, Torr. & Gray, with thickish sessile and roundish leaves, probably occurs as far north as Virginia.

ORDER 79. APOCYNACEÆ. (DOGBANE FAMILY.)

Plants almost all with milky acrid juice, entire (chiefly opposite) leaves without stipules, regular 5-merous and 5-androus flowers; the 5 lobes of the corolla convolute and twisted in the bud; the filaments distinct, inserted on the corolla, and the pollen granular; the calyx entirely free from the two ovaries, which (in our genera) are distinct (and forming pods), though their styles or stigmas are united into one. — Seeds amphitropous or anatropous, with a large straight embryo in sparing albumen, often bearing a tuft of down (comose). — Chiefly a tropical family (of acrid-poisonous plants), represented in gardens by the Oleander and Periwinkle, and among wild plants by three genera: —

1. **Amsonia.** Seeds naked. Corolla with the tube bearded inside. Anthers longer than the filaments. Leaves alternate.
2. **Forsteronia.** Seeds comose. Corolla funnel-form, not appendaged. Filaments slender. Calyx glandular inside. Leaves opposite.
3. **Apocynum.** Seeds comose. Corolla bell-shaped, appendaged within. Filaments short, broad and flat. Calyx not glandular. Leaves opposite.

1. AMSÓNIA, Walt. AMSONIA.

Calyx 5-parted, small. Corolla with a narrow funnel-form tube bearded inside, especially at the throat; the limb divided into 5 long linear lobes. Stamens 5, inserted on the tube, included: anthers obtuse at both ends, longer than the filaments. Ovaries 2: style 1: stigma rounded, surrounded with a cup-like membrane. Pod (follicles) 2, long and slender, many-seeded. Seeds cylindrical, abrupt at both ends, packed in one row, naked. — Perennial herbs, with *alternate leaves*, and pale blue flowers in terminal paniced cymes. (Said to be named for a *Mr. Charles Amson.*)

1. **A. Tabernæmontana**, Walt. Loosely somewhat pubescent or hairy when young, or soon glabrous; leaves varying from ovate-lanceolate to linear-lanceolate, taper-pointed; calyx-lobes short, awl-shaped; tube of the bluish corolla little longer than the lobes, the upper part either hairy when young or glabrous. — Low grounds, Illinois, Virginia? and southward. May, June.

2. FORSTERÓNIA, Meyer. FORSTERONIA.

Calyx 5-parted, with 3–5 glands at its base inside. Corolla funnel-form, not appendaged; the limb 5-lobed. Stamens 5, included: filaments slender: anthers arrow-shaped, with an inflexed tip, adhering to the stigma. Pods (follicles) 2, slender, many-seeded. Seeds oblong, with a tuft of down. — Twining plants, more or less woody, with opposite leaves and small flowers in cymes. (Named for *Mr. T. F. Forster*, an English botanist.)

1. **F. difformis**, A. DC. Nearly herbaceous and glabrous; leaves oval-lanceolate, pointed, thin; calyx-lobes taper-pointed; corolla pale yellow. — Damp grounds, Virginia, S. Illinois, and southward. April.

3. APÓCYNUM, Tourn. DOGBANE. INDIAN HEMP.

Calyx 5-parted, the lobes acute. Corolla bell-shaped, 5-cleft, bearing 5 triangular appendages in the throat opposite the lobes. Stamens 5, inserted on the very base of the corolla: filaments flat, shorter than the arrow-shaped anthers, which converge around the ovoid obscurely 2-lobed stigma, and are slightly adherent to it by their inner face. Style none: stigma large, ovoid, slightly 2-lobed. Fruit of 2 long and slender follicles. Seeds comose with a long tuft of silky down at the apex. — Perennial herbs, with upright branching stems, opposite mucronate-pointed leaves, a tough fibrous bark, and small and pale cymose flowers on short pedicels. (An ancient name of the Dogbane, composed of *ἄπό*, *from*, and *κύων*, *a dog*, to which the plant was thought to be poisonous.)

1. **A. androsæmifolium**, L. (SPREADING DOGBANE.) Smooth, branched above; *branches divergently forking; leaves ovate, distinctly petioled;*

cymes loose, spreading, mostly longer than the leaves; corolla (pale rose-color, 4" broad) *open-bell-shaped, with revolute lobes, the tube much longer than the ovate pointed divisions of the calyx.* — *Varies, with the leaves downy underneath.* — *Borders of thickets: common northward. June, July.* — *Pods 3' - 4' long, pendent.*

2. **A. cannábinum**, L. (INDIAN HEMP.) *Stem and branches upright or ascending, terminated by erect and close many-flowered cymes, which are usually shorter than the leaves; corolla* (greenish-white) *with nearly erect lobes, the tube not longer than the lanceolate divisions of the calyx.* — *Var. GLABÉRRIMUM, DC.* Entirely smooth; leaves oblong or oblong-lanceolate, on short but manifest petioles, obtuse or rounded, or the uppermost mostly acute at both ends. — *Var. PUBÉSCENS, DC.* Leaves oblong, oval, or ovate, soft-downy underneath or sometimes on both sides, as well as the cymes. (*A. pubescens, R. Br.*) — *Var. HYPERICIFÓLIUM.* Leaves more or less heart-shaped at the base and on very short petioles, commonly smooth throughout. (*A. hypericifolium, Ait.*) — *River-banks, &c.: common. July, Aug.* — *Plant 2° - 3° high, much more upright than the last; the flowers scarcely half the size.*

ORDER 80. ASCLEPIADACEÆ. (MILKWEED FAMILY.)

Plants with milky juice, and opposite or whorled (rarely scattered) entire leaves; the follicular pods, seeds, anthers (connected with the stigma), sensible properties, &c., just as in the last family; from which they differ in the commonly valvate corolla, and in the singular connection of the anthers with the stigma, the cohesion of the pollen into wax-like or granular masses, &c., as explained under the first and typical genus.

Tribe I. ASCLEPIADEÆ. Filaments monadelphous. Pollen-masses 10, waxy, fixed to the stigma in pairs by a gland, hanging vertically.

1. **Asclepias.** Calyx and corolla reflexed, deeply 5-parted. Crown of 5 hooded fleshy bodies (nectaries, *L.*), with an incurved horn rising from the cavity of each.
2. **Acerates.** Calyx and corolla reflexed or merely spreading. Crown as in No. 1, but without a horn inside.
3. **Enslenia.** Calyx and corolla erect. Crown of 5 membranaceous flat bodies, terminated by a 2-cleft tail or awn.
4. **Vincetoxicum.** Calyx and wheel-shaped corolla spreading. Crown a fleshy 5-10-lobed ring or disk.

Tribe II. GONOLOBEÆ. Filaments monadelphous. Pollen-masses 10, affixed to the stigma in pairs, horizontal.

5. **Gonolobus.** Corolla wheel-shaped. Crown a wavy-lobed fleshy ring.

Tribe III. PERIPLOCEÆ. Filaments distinct or nearly so. Pollen-masses granular, separately applied to the stigma.

6. **Periploca.** Corolla wheel-shaped, with 5 awned scales in the throat.

1. ASCLÉPIAS, L. MILKWEED. SILKWEED.

Calyx 5-parted, persistent; the divisions small, spreading. Corolla deeply 5-parted; the divisions valvate in the bud, reflexed, deciduous. Crown of 5 hooded bodies (nectaries, L.) seated on the tube of stamens, each containing

an incurved horn. Stamens 5, inserted on the base of the corolla. filaments united in a tube which encloses the pistil: anthers adherent to the stigma, each with 2 vertical cells, tipped with a membranaceous appendage, each cell containing a flattened pear-shaped and waxy pollen-mass; the two contiguous pollen-masses of adjacent anthers forming pairs which hang by a slender prolongation of their summits from 5 cloven glands that grow on the angles of the stigma extricated from the cells by the agency of insects, and directing copious pollen-tubes into the point where the stigma joins the apex of the styles. Ovaries 2, tapering into very short styles: the large depressed 5-angled fleshy mass which takes the place of stigma common to the two. Follicles 2, one of them often abortive, soft, ovate or lanceolate. Seeds anatropous, flat, margined, downwardly imbricated all over the large placenta, which separates from the suture at maturity, furnished with a long tuft of silky hairs (*coma*) at the hilum. Embryo large, with broad foliaceous cotyledons in thin albumen. — Perennial upright herbs, with thick and deep roots: peduncles terminal or lateral and between the petioles, bearing simple many-flowered umbels: flowering in summer. (The Greek name of *Æsculapius*, to whom the genus is dedicated.)

* *Leaves opposite (or some of them in No. 5–7 in threes or fours.)*

← *Stem simple or nearly so, leafy to the top, and bearing lateral umbels as well as a terminal one: leaves ovate or oblong: flowers whitish, pinkish, or dull purple.*

↔ *Pods beset with soft spinous projections: flowers 6'–9' long when open, greenish-purple, numerous in dense umbels.*

1. **A. Cornūti**, Decaisne. (COMMON MILKWEED OR SILKWEED.) Stem tall and stout; *leaves oval-oblong (4'–8' long), contracted at base into a short petiole, pale, minutely downy beneath, as well as the peduncles, &c.; hoods of the crown ovate, obtuse, with a lobe or tooth on each side of the short and stout claw-like horn; pods ovate, covered with weak spines and woolly.* (A. *Syriaca*, L., but the plant belongs to this country only.) — Rich ground, everywhere.

2. **A. Sullivantii**, Engelm. *Very smooth throughout, tall; leaves ovate-oblong with a somewhat heart-shaped base, nearly sessile; hoods obovate, entire, obtusely 2-tined at the base outside; flowers larger (9' long) and more purple than in the preceding; pods obscurely soft-spiny, chiefly on the beak, ovate-lanceolate.* — Low grounds, Columbus, Ohio (*Sullivant*) to Illinois.

↔ *Pods even, not warty-roughened, mostly glabrous.*

3. **A. phytolaccoides**, Pursh. (POKE-MILKWEED.) Stem (3°–5° high) smooth; *leaves broadly ovate, or the upper oval-lanceolate and pointed at both ends, short-petioled, smooth or slightly downy underneath (5'–8' long); lateral umbels several; pedicels loose and nodding, numerous, long and slender (1'–3' long), equalling the peduncle, many times longer than the ovate-oblong divisions of the (greenish) corolla; hoods of the crown (white) truncate, the margins 2-toothed at the summit, the horn with a long projecting awl-shaped point; pods minutely downy.* — Moist copses; flowering early in summer. — Flower 6' long.

4. **A. purpurascens**, L. (PURPLE M.) Stem rather slender (1°–3° high); *leaves elliptical or ovate-oblong, the upper taper-pointed, minutely velvety-downy underneath, smooth above, contracted at the base into a short petiole; pedicels*

shorter than the peduncle, 3-4 times the length of the dark purple lanceolate-ovate divisions of the corolla; hoods of the crown oblong, abruptly narrowed above; the horn broadly scythe-shaped, with a narrow and abruptly inflexed horizontal point. (*A. amœna*, L., *Michx.*) — Borders of woods, &c., New England to Illinois and southward. — Flowers 6'' long.

5. **A. variegata**, L. (VARIEGATED M.) Nearly glabrous (1°-2° high); leaves ovate, oval, or obovate, somewhat wavy, contracted into short petioles; pedicels (numerous and crowded) and peduncle short, downy; divisions of the corolla ovate (white); hoods of the crown orbicular, entire, purplish or reddish, the horn semilunar with a horizontal point; pods slightly downy. (*A. nivea*, L., in part. *A. hybrida*, *Michx.*) — Dry woods, S. New York to Wisconsin and southward. July. — Remarkable for its compact umbels of nearly white flowers. Leaves 4-5 pairs, the middle ones sometimes whorled.

6. **A. ovalifolia**, Decaisne in DC. Low (6'-18' high), soft-downy, especially the lower surface of the ovate or lanceolate-oblong acute short-petioled leaves; umbels loosely 10-18-flowered, either sessile or peduncled; pedicels slender; hoods of the crown oblong, obtuse, yellowish, with a small horn, about the length of the oval greenish-white divisions of the corolla (which are tinged with purple outside); pods downy. (*A. lanuginosa*, Ed. 1, probably not of Nutt. *A. Vaseyi*, *Carey*). — Prairies and oak-openings, N. Illinois, *Vasey*, Wisconsin, *Lapham*, and northwestward. June. — Leaves 1½'-3' long, smoothish above, the upper sometimes scattered; the middle rarely in threes. Flower 4''-5'' long.

7. **A. quadrifolia**, Jacq. (FOUR-LEAVED M.) Nearly smooth; stem slender (1°-2° high), mostly leafless below, bearing usually one or two whorls of four in the middle and one or two pairs of ovate or ovate-lanceolate taper-pointed petioled leaves (2'-4' long); pedicels slender; divisions of the (pale pink) corolla oblong; hoods of the white crown elliptical-ovate, the incurved horn short and thick; pods linear-lanceolate, smooth. — Dry woods and hills: not very common. June. — Flowers 4'' long.

+ + *Stem branching, leafy to the top, bearing lateral as well as terminal umbels: leaves petioled: flowers small (3'' long): pods smooth and glabrous.*

8. **A. perennis**, Walt. Nearly glabrous; stems (1°-2° high) persistent or somewhat woody at the base; leaves lanceolate or lanceolate-ovate, tapering to both ends, thin, rather slender-petioled; flowers white, small; the small hoods of the crown shorter than the needle-shaped horn; seeds sometimes destitute of a coma! (*A. parviflora*, *Pursh*, and Ed. 2.) — Low grounds, S. Indiana, Illinois, and southward.

9. **A. incarnata**, L. (SWAMP MILKWEED.) Smooth, or nearly so, in the typical form, the stem with two downy lines above and on the branches of the peduncles (2°-3° high), very leafy; leaves oblong-lanceolate, acute or pointed, obtuse or obscurely heart-shaped at base; flowers rose-purple; hoods of the crown scarcely equalling the slender needle-pointed horn. — Var. *PULCHRA* has broader and shorter-petioled leaves, more or less hairy-pubescent, as well as the stem. (*A. pulchra*, *Willd.*) — Wet grounds; the smooth form very common northward; the hairy variety more so southward. — Milky juice scanty.

+ + + *Stem perfectly simple, producing only a single conspicuously-peduncled terminal umbel of dull-colored largish (6" long) flowers: hoods and lobes of the corolla broad: pods smooth: whole plant glabrous or nearly so, and pale or glaucous: leaves closely sessile, transversely veiny.*

10. **A. obtusifolia**, Michx. Stem tall (2° – 3° high); leaves wavy, oblong with a heart-shaped clasping base, very obtuse or retuse ($2\frac{1}{2}'$ – $5'$ long); peduncle $3'$ – $12'$ long; corolla pale greenish purple; hoods of the crown truncate and somewhat toothed at the summit, shorter than the slender awl-pointed horn. — Sandy woods and fields: not rare, especially southward.

11. **A. Meádii**, Torr. (in Ed. 2, addend.) Stem slender (1° – 2° high); leaves ovate or oblong-ovate, not wavy, obtuse or acutish ($1\frac{1}{2}'$ – $2\frac{3}{4}'$ long); peduncle only twice the length of the upper leaves; pedicels rather short; corolla greenish-white; hoods of the crown rounded-truncate at summit, and with a sharp tooth at each margin, somewhat exceeding the stouter horn; pod unknown. — Augusta, Illinois, Dr. S. B. Mead. Clinton, Iowa, Dr. Vasey. June.

+ + + + *Stem simple or mostly so (2° – 4° high), bearing 2–5 panicled umbels on a naked terminal peduncle, and sometimes single axillary ones: flowers pink-red, rather large (over 6" long): crown conspicuously elevated above the base of the corolla: pods smooth: whole plant glabrous or nearly so.*

12. **A. rubra**, L. Leaves ovate or lanceolate and tapering from a rounded or heart-shaped base to a very acute point, sessile or nearly so ($2'$ – $6'$ long, $\frac{1}{2}'$ – $2\frac{1}{2}'$ wide), bright green; umbels many-flowered; divisions of the corolla and hoods of the crown oblong-lanceolate, purple-red; the horn long and slender. (*A. laurifolia*, Michx. *A. acuminata*, Pursh.) — Wet pine-barrens, &c., New Jersey and Penn. to Virginia and southward.

13. **A. paupércula**, Michx. Stem slender (2° – 4° high); leaves elongated lanceolate or linear ($5'$ – $10'$ long), tapering to both ends, slightly petioled; umbels 5–12-flowered; divisions of the red corolla narrowly oblong; the bright orange hoods broadly oblong, obtuse, much exceeding the incurved horn. — Wet pine-barrens on the coast, New Jersey, Virginia, and southward.

* * *Leaves scattered, or some opposite: milky juice little or none: flowers orange-red.*

14. **A. tuberosa**, L. (BUTTERFLY-WEED. PLEURISY-ROOT.) Roughish-hairy; stems erect or ascending, very leafy, branching at the summit, and bearing the umbels in a terminal corymb; leaves varying from linear to oblong-lanceolate, sessile or slightly petioled; divisions of the corolla oblong (greenish-orange); hoods of the crown narrowly oblong, bright orange, scarcely longer than the nearly erect and slender awl-shaped horns; pods hoary. (*A. decumbens*, L.) — Dry hills and fields: common, especially southward. — Plant 1° – 2° high, leafy to the summit, usually with numerous and corymbed short-peduncled umbels of showy flowers.

* * * *Leaves nearly all whorled, rarely alternate, crowded: flowers white, small.*

15. **A. verticillata**, L. (WHORLED M.) Smoothish; stems slender, simple or sparingly branched, very leafy to the summit; leaves very narrowly linear, with revolute margins ($2'$ – $3'$ long, $1''$ wide), 3–6 in a whorl; umbels small, lateral and terminal; divisions of the corolla ovate (greenish-white);

hoods of the crown roundish-oval, about half the length of the incurved claw-shaped horns; pods smooth. — Dry hills: common, especially southward.

2. ACERÀTES, Ell. GREEN MILKWEED.

Nearly as in *Asclepias*; but the hoods of the crown destitute of a horn (whence the name, from *a* privative and *κέρας*, *-aros*, *a horn*). — Flowers greenish. Leaves varying from opposite to irregularly alternate, short-petioled or sessile. Pollen-masses slender-stalked.

§ 1. *Divisions of the corolla reflexed, oblong: hoods of the crown erect and concave: umbels compactly many-flowered: pods not mucronate, slender.*

* *Crown not elevated; its hoods oblong, nearly equalling the anthers.*

1. **A. viridiflora**, Ell. Minutely soft-downy, becoming smoothish; stems ascending (1° – 2° high); leaves varying from oval to linear, thick ($1\frac{1}{2}'$ – $4'$ long), umbels nearly sessile, lateral, dense and globose; flower (when the corolla is reflexed) nearly $\frac{1}{2}'$ long, short-pedicelled. — Dry soil: common, especially southward. July–Sept.

2. **A. lanuginosa**, Decaisne. *Hairy*, low ($5'$ – $12'$ high); leaves lanceolate or ovate-lanceolate; umbel solitary and terminal, peduncled; flowers not larger than in the next; pedicels slender. (Probably *Asclepias lanuginosa*, Nutt.: certainly *A. Nuttalliana*, Torr. *Acerates monocephala*, Lapham, in Ed. 2, addend.) — Prairies, Wisconsin (Lapham) and westward. July.

* * *Crown short-stalked, i. e. elevated above the base of the corolla; its hoods oval, strongly concave, and decidedly shorter than the tips of the anthers.*

3. **A. longifolia**, Ell. Minutely roughish-hairy or smoothish; stem erect (1° – 3° high), very leafy; leaves mostly alternate-scattered, linear ($3'$ – $7'$ long); umbels lateral, on peduncles of about the length of the slender pedicels; flowers $3''$ long when expanded. — Moist prairies, Ohio to Wisconsin and southward. July–Oct.

§ 2. **ANANTHERIX**, Nutt. *Divisions of the corolla ascending or barely spreading: hoods of the crown widely spreading and somewhat incurved, slipper-shaped and laterally compressed, the cavity divided at the apex by a crest-like partition: umbels solitary and terminal or corymbed, loosely-flowered: pods oblong or ovate, often somewhat mucronate with soft spinous projections.*

4. **A. paniculata**, Decaisne. Almost glabrous; stems short (1° high); leaves alternate, short-petioled, elongated-oblong, $1'$ – $2'$ wide; umbels several in a cluster, short-peduncled; flowers large ($1'$ in diameter), green, with a purplish crown. — Prairies, Illinois (Vasey, Bebb), and southward. June.

3. ENSLÉNIA, Nutt. ENSLENIA.

Calyx 5-parted. Corolla 5-parted; the divisions erect, ovate-lanceolate. Crown of 5 free membranaceous leaflets, which are truncate or obscurely lobed at the apex, where they bear a pair of flexuous awns united at their base. Anthers nearly as in *Asclepias*: pollen-masses oblong, obtuse at both ends, fixed below the summit of the stigma to the descending glands. Pods oblong-lanceolate, smooth. Seeds with a tuft, as in *Asclepias*. — A perennial twining herb,

smooth, with opposite heart-ovate and pointed long-petioled leaves, and small whitish flowers in raceme-like clusters, on slender axillary peduncles. (Dedicated to *A. Enslin*, an Austrian botanist who collected in the Southern United States early in the present century.)

1. *E. álvida*, Nutt. — River-banks, Ohio to Illinois, and southward: common. July — Sept. — Climbing 8° — 12° high: leaves 3' — 5' wide.

4. VINCETÓXICUM, Mœnch. VINCETOXICUM.

Calyx 5-parted. Corolla 5-parted, wheel-shaped. Crown flat and fleshy, disk-like, 5 — 10-lobed, simple. Anthers smooth, pods and seeds much as in *Asclepias*. — Herbs, often twining. (Name composed of *Vinca*, the Periwinkle, and *toxicum*, poison.)

1. *V. NIGRUM*, Mœnch. (BLACK V.) More or less twining, nearly smooth; leaves ovate or lance-ovate; flowers small, dark purple, in an axillary cluster, on a peduncle shorter than the leaves. — Cambridge, Mass., &c.: a weed escaping from gardens. (Adv. from Eu.)

5. GONÓLOBUS, Michx. GONOLOBUS.

Calyx 5-parted. Corolla 5-parted, wheel-shaped, sometimes reflexed-spreading; the lobes convolute in the bud. Crown a small and fleshy wavy-lobed ring in the throat of the corolla. Anthers horizontal, partly hidden under the flattened stigma, opening transversely. Pollen-masses 5 pairs, horizontal. Pods turgid, mostly muricate with soft warty projections, sometimes ribbed. Seeds with a coma. — Twining herbs or shrubs (ours herbaceous), with opposite heart-shaped leaves, and corymbose-umbelled greenish or dark purple flowers, on peduncles rising from between the petioles. (Name composed of *γωνος*, an angle, and *λοβός*, a pod, from the angled or ribbed follicles of some species.)

1. *G. lævis*, Michx. Nearly glabrous, or the stems and petioles sparingly hirsute and finely puberulent; calyx and corolla glabrous, the latter tapering-conical in the bud, the expanded divisions lanceolate, yellowish-green; pods ribbed, smooth. (*Vincetoxicum gonocarpos*, Walt. *Periploca late-scandens*, Clayt. *G. macrophyllus*, Michx., also *Decaisne*, excl. syn. Jacq. & Bot. Mag. *G. tiliaefolius*, *Decaisne*.) — River-banks, Virginia, to Illinois and southward. July.

2. *G. obliquus*, R. Br. Stems, petioles and often the ribs of the leaves beneath hirsute with spreading viscid hairs; calyx and corolla pubescent or puberulent outside, the latter narrow-conical-oblong in the bud, its divisions ligulate-linear or lanceolate, obtuse, dark dull crimson-purple within; pods copiously muricate, ribless. (*Cynanchum obliquum*, Jacq., 1786. *C. discolor*, Sims, Bot. Mag. *Gonolobus hirsutus*, Ed. 2, &c. *G. discolor*, R. & S. *G. macrophyllus*, *Decaisne* in part.) — River-banks, Penn. to Virginia. Aug., Sept. — Lobes of the corolla nearly 6" long. Pod 5' long.

3. *G. hirsutus*, Michx. (*Apocynum hirsutum*, Pluk.; perhaps *Periploca Carolinensis*, Dill., and *P. late-scandens* fl. ferrugineo, Clayt.; *Vincetoxicum acanthocarpos*, Walt.; and clearly *Cynanchum Carolinense*, Jacq.) Known from the last by its short-ovate flower-buds, and the oval or oblong divisions of the purple corolla (only about 3" long); perhaps occurs in S. E. Virginia.

6. PERÍPLOCÁ, L. PERIPLOCA.

Calyx 5-parted. Corolla 5-parted, wheel-shaped, with 5 awned scales in the throat. Filaments distinct: anthers coherent with the apex of the stigma, bearded on the back: pollen-masses 5, each of 4 united, singly affixed directly to the glands of the stigma. Stigma hemispherical. Pods smooth, widely divergent. Seeds with a silky tuft. — Twining shrubby plants, with smooth opposite leaves, and paniced-cymose flowers. (Name from περιπλοκή, a coiling round, in allusion to the twining stems.)

1. **P. GRÆCA**, L. Leaves ovate or ovate-lanceolate, shorter than the loosely-flowered cymes; divisions of the brownish-purple corolla linear-oblong, very hairy above. — Near Rochester, &c., New York. Probably hardly established. Aug. (Adv. from Eu.)

ORDER 81. OLEACEÆ. (OLIVE FAMILY.)

Trees or shrubs, with opposite and pinnate or simple leaves, a 4-cleft (or sometimes obsolete) calyx, a regular 4-cleft or nearly or quite 4-petalous corolla, sometimes apetalous; the stamens only 2 (rarely or accidentally 3 or 4); the ovary 2-celled, with 2 (rarely more) ovules in each cell. — Seeds anatropous, with a large straight embryo in hard fleshy albumen, or without albumen. — The Olive is the type of the true Oleaceæ, to which belongs the LILAC (SYRINGA), &c.; while the JESSAMINE (JASMINUM) represents another division of the order.

Tribe I. OLEINEÆ. Fruit a drupe or berry. Flowers perfect or polygamous, with both calyx and corolla; the latter valvate in the bud. Ovules suspended. Leaves simple, mostly entire.

1. **Ligustrum.** Corolla funnel-form, its tube longer than the calyx, 4-cleft.

2. **Olea.** Corolla short, bell-shaped or salver-shaped; the limb 4-parted.

3. **Chionanthus.** Corolla 4-parted or 4-petalous, the divisions or petals long and linear.

Tribe II. FRAXINEÆ. Fruit dry and winged (a samara). Flowers diœcious or polygamous, mostly apetalous, and sometimes also without a calyx. Ovules suspended. Leaves odd-pinnate.

4. **Fraxinus.** The only genus of the Tribe.

Tribe III. FORESTIEREÆ. Fruit a drupe or berry. Flowers diœcious or perfect, apetalous. Ovules suspended. Leaves simple.

5. **Forestiera.** Flowers diœcious, from a scaly catkin-like bud. Stamens 2-4.

1. LIGÚSTRUM, Tourn. PRIVET.

Calyx short-tubular, 4-toothed, deciduous. Corolla funnel-form, 4-lobed; the lobes ovate, obtuse. Stamens 2, on the tube of the corolla, included. Stigma 2-cleft. Berry spherical, 2-celled, 2-1-seeded. — Shrubs, with entire leaves on short petioles, and small white flowers in terminal thyrsoïd panicles. (The classical name.)

1. **L. VULGARE**, L. (COMMON PRIVET or PRIM.) Leaves elliptical-lanceolate, very smooth, thickish, deciduous; berries black. — Used for low hedges: naturalized in copses by the agency of birds in E. New England and Pennsylvania. June. (Nat. from Eu.)

2. ÒLEA, Tourn. OLIVE.

Calyx short, 4-toothed, rarely entire. Corolla with a short bell-shaped tube and a 4-parted spreading limb. Stamens 2. Drupe with a bony stone, 2-1-seeded. — Shrubs or trees, with opposite and coriaceous mostly entire leaves, and perfect, or (in our species) polygamous or diœcious small white flowers, in panicles or corymbs. (The classical name of the Olive, O. EUROPEÆA.)

1. *O. Americana*, L. (DEVIL-WOOD.) Leaves oblong-lanceolate, smooth and shining (3' - 6' long); fruit spherical. — Moist woods, coast of S. Virginia, and southward. May. — Tree 15° - 20° high.

3. CHIONÁNTHUS, L. FRINGE-TREE.

Calyx 4-parted, very small, persistent. Corolla of 4 long and linear petals, which are barely united at the base. Stamens 2 (rarely 3 or 4), on the very base of the corolla, very short. Stigma notched. Drupe fleshy, globular, becoming 1-celled, 1-3-seeded. — Low trees or shrubs, with deciduous and entire petioled leaves, and delicate flowers in loose and drooping graceful panicles, from lateral buds. (Name from χιών, *snow*, and άνθος, *blossom*, alluding to the light and snow-white clusters of flowers.)

1. *C. Virginica*, L. Leaves oval, oblong, or obovate-lanceolate; flowers on slender pedicels; drupe purple, with a bloom, ovoid (6" - 8" long). — Riverbanks, S. Pennsylvania and southward: very ornamental in cultivation. June. — Petals 1' long, narrowly linear, acute, varying to 5-6 in number.

4. FRÁXINUS, Tourn. ASH.

Flowers polygamous or (in our species) diœcious. Calyx small and 4-cleft, toothed, or entire, or obsolete. Petals 4, slightly cohering in pairs at the base, or only 2, oblong or linear, or altogether wanting in our species. Stamens 2, sometimes 3 or 4: anthers linear or oblong, large. Style single: stigma 2-cleft. Fruit a 1-2-celled samara or *key-fruit*, flattened, winged at the apex, 1-2-seeded. Cotyledons elliptical: radicle slender. — Light timber-trees, with petioled pinnate leaves of 3-15 either toothed or entire leaflets; the small flowers in crowded panicles or racemes from the axils of last year's leaves. (The classical Latin name, thought to be derived from φράξις, a *separation*, from the facility with which the wood splits.)

* *Fruit winged from the apex only, barely margined or quite terete towards the base: calyx minute, persistent: corolla none: leaflets stalked.*

1. *F. Americana*, L. (WHITE ASH.) Branchlets and petioles glabrous; leaflets 7-9, ovate- or lance-oblong, pointed, pale and either smooth or pubescent underneath, somewhat toothed or entire; fruit terete and marginless below, above extended into a lanceolate, oblanceolate, or wedge-linear wing. (*F. acuminata*, and *F. juglandifolia*, Lam. *F. epiptera*, Michx.) — Rich or moist woods: common. April, May. — A large forest tree, with gray furrowed bark, smooth gray branchlets, and rusty-colored buds. (The figure of the fruit in Michaux's *Sylva* is misplaced, apparently interchanged with that of the Green Ash.)

2. **F. pubescens**, Lam. (RED ASH.) *Branchlets and petioles velvety-pubescent*; leaflets 7-9, ovate or oblong-lanceolate, taper-pointed, almost entire, pale or more or less pubescent beneath; *fruit acute at the base, flattish and 2-edged, the edges gradually dilated into the long (1½' - 2') oblanceolate or linear-lanceolate wing.* (F. *tomentosa*, Michx.) — With No. 1: rare west of the Alleghanies: a smaller tree, less valuable for timber: passes by gradations into the next.

3. **F. viridis**, Michx. f. (GREEN ASH.) *Glabrous throughout*; leaflets 5-9, ovate or oblong-lanceolate, often wedge-shaped at the base and serrate above, *bright green both sides*; *fruit acute at the base, striate, 2-edged or margined, gradually dilated into an oblanceolate or linear-spatulate wing, much as in No. 2.* (F. *cóncolor*, Muhl. F. *juglandifolia*, Willd., DC., and Ed. 1, but not of Lam.) — Near streams, New England to Wisconsin and southward; most common westward. — A small or middle-sized tree. (The figure of the fruit given in Michaux's *Sylva* evidently belongs to *F. Americana*.)

* * *Fruit winged all round the seed-bearing portion.*

+ *Calyx wanting, at least in the fertile flowers, which are entirely naked!*

4. **F. sambucifolia**, Lam. (BLACK OR WATER ASH.) *Branchlets and petioles glabrous*; leaflets 7-11, *sessile, oblong-lanceolate, tapering to a point, serrate, obtuse or rounded at the base, green and smooth both sides, when young with some rusty hairs along the midrib*; *fruit linear-oblong or narrowly elliptical, blunt at both ends.* — Swamps, Penn. to Kentucky, and everywhere northward. April, May. — Small tree; its tough wood separable into thin layers, used for coarse basket-work, &c. Bruised leaves with the odor of Elder.

+ + *Calyx present, persistent at the base of the fruit.*

5. **F. quadrangulata**, Michx. (BLUE ASH.) *Branchlets square, at least on vigorous shoots, glabrous*; leaflets 7-9, short-stalked, oblong-ovate or lanceolate, pointed, sharply serrate, green both sides; *fruit narrowly oblong, blunt, and of the same width at both ends, or slightly narrowed at the base, often notched at the apex (1½' long, ¼' - ½' wide).* — Dry or moist rich woods, Ohio to Wisconsin, Illinois, and Kentucky. — Tree large, with timber like No. 1.

6. **F. platycarpa**, Michx. (CAROLINA WATER-ASH.) *Branchlets terete, glabrous or pubescent*; leaflets 5-7, ovate or oblong, acute at both ends, short-stalked; *fruit broadly winged (not rarely 3-winged), oblong (9" wide), with a tapering base.* — Wet woods, Virginia and southward. March.

5. FORESTIÈRA, Poir. (ADÈLIA, Michx.)

Flowers dicecious, crowded in catkin-like scaly buds from the axils of last year's leaves, imbricated with scales. Corolla none. Calyx early deciduous, of 4 minute sepals. Stamens 2-4: anthers oblong. Ovary ovate, 2-celled, with 2 pendulous ovules in each cell: style slender: stigma somewhat 2-lobed. Drupe small, ovoid, 1-celled, 1-seeded. — Shrubs, with opposite and often fascicled deciduous leaves and small flowers. Fertile peduncles short, 1-3-flowered. (Named for *M. Forestier*, a French physician.)

1. **F. acuminata**, Poir. *Glabrous*; leaves thin, oblong-ovate or ovate-lanceolate, acuminate at both ends, often serrulate; drupe oblong, usually pointed. — Wet river-banks, W. Illinois and southward. April.

DIVISION III. APÉTALOUS EXÓGENOUS PLANTS.

Corolla none; the floral envelopes in a single series (calyx), or sometimes wanting altogether.

ORDER 82. **ARISTOLOCHIAEÆ.** (BIRTHWORT FAMILY.)

Twining shrubs, or low herbs, with perfect flowers, the conspicuous lurid calyx valvate in the bud and coherent (at least at the base) with the 6-celled ovary, which forms a many-seeded 6-celled pod or berry in fruit. Stamens 6-12, more or less united with the style: anthers adnate, extrorse.—Leaves petioled, mostly heart-shaped and entire. Seeds anatropous, with a large fleshy rhapshe, and a minute embryo in fleshy albumen. A small family of bitter-tonic or stimulant, sometimes aromatic plants.

1. **ÁSARUM**, Tourn. ASARABACCA. WILD GINGER.

Calyx regular; the limb 3-cleft or parted. Stamens 12, with more or less distinct filaments, their tips usually continued beyond the anther into a point. Pod rather fleshy, globular, bursting irregularly. Seeds large, thick.—Stemless herbs, with aromatic-pungent creeping rootstocks, bearing 2 or 3 scales, then one or two kidney-shaped or heart-shaped leaves on long petioles, and terminated by a short-peduncled flower, close to the ground; in spring. (An ancient name, of obscure derivation.)

§ 1. *Calyx-tube wholly coherent with the ovary, the tips inflexed in bud: filaments slender, united only with the base of the style, much longer than the short anthers: styles united into one, which is barely 6-lobed at the summit, and with 6 radiating thick stigmas: leaves unspotted, a single pair, with the peduncle between them.*

1. **A. Canadense**, L. Soft-pubescent; leaves membranaceous, kidney-shaped, more or less pointed (4' - 5' wide when full grown); calyx bell-shaped, with the upper part of the short-pointed lobes widely and abruptly spreading, brown-purple inside; at each sinus is usually a small awl-shaped appendage (abortive petal).—Hillsides in rich woods: common, especially northward.

§ 2. *Calyx-tube inflated bell-shaped, somewhat contracted at the throat, only its base coherent with the lower half of the ovary; the limb 3-cleft, short: filaments very short or none: anthers oblong-linear: styles 6, fleshy, diverging, 2-cleft, each bearing a thick extrorse stigma below the cleft: leaves thickish, persistent, usually only one each year, the upper surface often whitish-mottled: peduncle very short: rootstocks clustered, ascending.*

2. **A. Virginicum**, L. Nearly glabrous; leaves round-heart-shaped (about 2' wide); calyx short, reticulated within; anthers pointless.—Virginia, and southward, in and near the mountains.

3. **A. arifolium**, Michx. Leaves halberd-heart-shaped (2' - 4' long); calyx oblong-tubular, with very short and blunt lobes; anthers obtusely short-pointed.—Virginia and southward.

2. ARISTOLÒCHIA, Tourn. BIRTHWORT.

Calyx tubular; the tube variously inflated above the ovary, mostly contracted at the throat. Stamens 6; the sessile anthers wholly adnate to the back of the short and fleshy 3-6-lobed or angled stigma. Pod naked, 6-valved. Seeds very flat. — Twining, climbing, or sometimes upright perennial herbs or shrubs, with alternate leaves and lateral or axillary greenish or lurid-purple flowers. (Named from reputed medicinal properties.)

§ 1. *Calyx-tube bent like the letter S, enlarged at the two ends, the small limb obtusely 3-lobed: anthers contiguous in pairs (making 4 cells in a row under each of the three truncate lobes of the stigma): low herbs.*

1. **A. Serpentaria**, L. (VIRGINIA SNAKEROOT.) Stems (8' - 15' high) branched at the base, pubescent; leaves ovate or oblong from a heart-shaped base, or halberd-form, mostly acute or pointed; flowers all next the root, short-peduncled. — A narrow-leaved variety is *A. sagittata*, *Muhl.*, *A. hastata*, *Nutt.*, &c. — Rich woods, Connecticut to Indiana and southward: not common except near the Alleghany Mountains. July. — The fibrous, aromatic-stimulant root is well known in medicine.

§ 2. *Calyx-tube strongly curved like a Dutch pipe, contracted at the mouth, the short limb obscurely 3-lobed: anthers contiguous in pairs under each of the 3 short and thick lobes of the stigma: twining shrubs: flowers from one or two of the superposed accessory axillary buds.*

2. **A. Siphon**, L'Her. (PIPE-VINE. DUTCHMAN'S PIPE.) Nearly glabrous; leaves round-kidney-shaped; peduncles with a clasping bract; calyx (1½' long) with a brown-purple abrupt flat border. — Rich woods, Penn. to Kentucky, and southward, along the mountains. May. — Stems sometimes 2' in diameter, climbing trees: full-grown leaves 8' - 12' broad.

3. **A. tomentosa**, Sims. Downy or soft-hairy; leaves round-heart-shaped, very veiny (3' - 5' long); calyx yellowish, with an oblique dark purple closed orifice and a rugose reflexed limb. — Rich woods, from S. Illinois southward. June.

ORDER 83. NYCTAGINACEÆ. (FOUR-O'CLOCK FAMILY.)

Herbs (or in the tropics often shrubs or trees), with mostly opposite and entire leaves, stems tumid at the joints, a delicate tubular or funnel-form calyx which is colored like a corolla, its persistent base constricted above the 1-celled 1-seeded ovary, and indurated into a sort of nut-like pericarp; the stamens few, slender, and hypogynous; the embryo coiled around the outside of mealy albumen, with broad foliaceous cotyledons. — Represented in our gardens by the FOUR-O'CLOCK, or MARVEL OF PERU (*MIRABILIS JALÁPA*), in which the calyx is commonly mistaken for a corolla, the cup-like involucre of each flower exactly imitating a calyx; — and by a single

1. OXÝBAPHUS, Vahl. OXYBAPHUS.

Flowers 1 - 5 in the same 5-lobed membranaceous broad and open involucre, which enlarges and is thin and reticulated in fruit. Calyx with a very short

tube and a bell-shaped (rose or purple) deciduous limb, plaited in the bud. Stamens mostly 3. Style filiform: stigma capitate. Fruit achenium-like, several-ribbed or angled. — Herbs, abounding on the western plains, with very large and thick perennial roots, opposite leaves, and mostly clustered small flowers. (Name *ὄξυβάφον*, a *vinegar-saucer*, or small shallow vessel; from the shape of the involucre.)

1. *O. nyctagineus*, Sweet. Nearly smooth; stem repeatedly forked (1° – 3° high); leaves varying from ovate, or somewhat heart-shaped to lanceolate; involucre 3–5-flowered. — Rocky places, from Wisconsin and Illinois southward and westward. June–Aug.

ORDER 84. PHYTOLACCÆÆ. (POKEWEED FAMILY.)

Plants with alternate entire leaves and perfect flowers, having the general characters of Chenopodiaceæ, but usually a several-celled ovary composed of as many carpels united in a ring, and forming a berry in fruit;—represented only by the typical genus

1. PHYTOLACCA, Tourn. POKEWEED.

Calyx of 5 rounded and petal-like sepals. Stamens 5–30. Ovary of 5–12 carpels, united in a ring, with as many short separate styles, in fruit forming a depressed-globose 5–12-celled berry, with a single vertical seed in each cell. Embryo curved in a ring around the albumen. — Tall and stout perennials, with large petioled leaves, and terminal racemes which become lateral and opposite the leaves. (Name compounded of *φυτόν*, *plant*, and the French *lac*, lake, in allusion to the crimson coloring matter resembling that pigment which the berries yield.)

1. *P. decandra*, L. (COMMON POKE OR SCOKE. GARGET. PIGEON-BERRY.) Stamens 10: styles 10. — Low grounds. July–Sept. — A smooth plant, with a rather unpleasant odor, and a very large poisonous root, often 4'–6' in diameter, sending up stout stalks (which are in early spring sometimes eaten as a substitute for Asparagus), at length 6° – 9° high. Calyx white: ovary green; the long racemes of dark-purple berries filled with crimson juice, ripe in autumn.

ORDER 85. CHENOPODIÆÆ. (GOOSEFOOT FAMILY.)

Chiefly herbs, of homely aspect, more or less succulent, with mostly alternate leaves, and no stipules nor scarious bracts, minute greenish flowers, with the free calyx imbricated in the bud; the stamens as many as its lobes, or occasionally fewer, and inserted opposite them or on their base; the 1-celled ovary becoming a 1-seeded thin utricle or rarely an achenium. Embryo coiled into a ring around the mealy albumen, when there is any, or else conduplicate, or spiral. — Calyx persistent, mostly enclosing the fruit. Styles or stigmas 2, rarely 3–5. (Mostly inert or innocent, weedy plants: several are pot-herbs, such as Spinach and Beet.)

- * Embryo coiled into a ring around copious central albumen. Leaves flat, not spiny nor fleshy.
- Flowers all alike and perfect, or merely polygamous by the want of stamens in some of them, clustered or paniced. Calyx obvious. Seed-coat crustaceous.
- 1. **Cycloloma.** Calyx 5-cleft, in fruit surrounded by a horizontal continuous membranaceous wing. Seed horizontal.
- 2. **Chenopodium.** Calyx 3-5-cleft or parted, the lobes naked or merely keeled in fruit. Seed horizontal, rarely vertical.
- 3. **Blitum.** Calyx of 3-5 sepals, mostly juicy or fleshy in fruit. Seed vertical.
- — Flowers monœcious or diœcious, and of 2 distinct sorts; the staminate with a regular calyx, clustered, the clusters mostly spiked.
- 4. **Atriplex.** Fertile flowers without calyx, enclosed between a pair of appressed bracts.
 - — — Flowers all perfect and alike, single in the axil of bracts, naked or 1-sepalled.
- 5. **Corispermum.** Fruit oval, flattened: pericarp adherent to the seed. Leaves linear.
- ** Embryo narrowly horseshoe-shaped or conduplicate: no albumen. Stem fleshy, jointed: leaves reduced to opposite fleshy scales or teeth. Flowers densely spiked, perfect.
- 6. **Salicornia.** Flowers sunk in hollows of the axis of the fleshy spike. Calyx utricle-like.
 - * * * Embryo coiled into a spiral: albumen mostly none. (Leaves alternate.)
- 7. **Suaeda.** Embryo flat-spiral. Calyx wingless. Leaves succulent.
- 8. **Salsola.** Embryo conical-spiral. Calyx in fruit horizontally winged. Leaves spinescent.

1. CYCLOLÒMA, Moquin. WINGED PIGWEED.

Flowers perfect, bractless. Calyx 5-cleft, with the concave lobes strongly keeled, enclosing the depressed fruit, at length appendaged with a broad and continuous horizontal scarious wing. Stamens 5. Styles 3. Seed horizontal, flat. Embryo encircling the mealy albumen. — An annual and much-branched coarse herb, with alternate sinuate-toothed petioled leaves, and small paniced clusters of sessile flowers. (Name composed of *κύκλω*, *round about*, and *λώμα*, *a border*, from the encircling wing of the calyx in fruit.)

1. **C. platyphýllum**, Moquin. (*Salsola platyphylla*, *Michx.*) — Illinois, on sandy banks of the Mississippi, and northwestward.

2. CHENOPÓDIUM, L. GOOSEFOOT. PIGWEED.

Flowers perfect, all bractless. Calyx 5-cleft, rarely 2-4-cleft or parted, with the lobes sometimes keeled, but not appendaged nor becoming succulent, more or less enveloping the depressed fruit. Stamens mostly 5: filaments filiform. Styles 2, rarely 3. Seed horizontal (sometimes vertical in Nos. 3, 7-9), lenticular; the coat crustaceous: embryo coiled partly or fully round the mealy albumen. — Weeds, usually with a white mealiness, or glandular. Flowers sessile in small clusters collected in spiked panicles. (Named from *χῆν*, *a goose*, and *ποῦς*, *foot*, in allusion to the shape of the leaves.) — Our species are all annuals (except the last two), flowering through late summer and autumn, growing around dwellings, in manured soil, cultivated grounds, and waste places.

§ 1. *Smooth or mealy, never glandular nor sweet-scented: embryo a complete ring.*

1. **C. POLYSPÉRMUM**, L. Low, often spreading, green and *wholly destitute of mealiness throughout; leaves all entire*, oblong or ovate and on slender petioles; flowers very small, the thin lobes of the calyx very incompletely enclosing the fruit; seed obtuse-edged. — In and around Boston: scarce. (Adv. from Eu.)

2. **C. ALBUM**, L. (LAMB'S-QUARTERS. PIGWEED.) Erect (1° – 3° high), *mealy and pale, sometimes green* and the mealiness obscure; *leaves varying from rhombic-ovate to lanceolate* or the uppermost even linear, acute, *all or only the lower more or less angulate-toothed*; clusters spikèd-panicled, mostly dense; seed with acute or bluntish margins. — Common, especially in cult. ground: extremely variable. — The genuine *C. album* is considerably whitish-mealy, at least the inflorescence, which is dense; the calyx with strongly keeled lobes, and completely enclosing the fruit. A green form with somewhat entire leaves and less dense inflorescence is *C. viride*, L. (Nat. from Eu.)

Var. **BOSCIANUM**. Loosely branched, more slender, the mealiness obscure or slight and only on the inflorescence, which is laxer, the flowers smaller; calyx incompletely covering the fruit, its lobes moderately or slightly if at all keeled; leaves inclined to be entire. (*C. Boscianum*, Moquin. *C. Berlandièri*, Moquin, an intermediate form. *C. polyspermum*, var. *spicatum*, Ed. 2.) — More shady places, Pennsylvania and southward. In some forms appears as if a distinct species; seemingly indigenous southwestward.

3. **C. GLAUCUM**, L. (OAK-LEAVED GOOSEFOOT.) Low (5'–12' high), *spreading, glaucous-mealy, leaves sinuately pinnatifid-toothed*, oblong, obtuse, pale green above; clusters spikèd, small; calyx-lobes not at all keeled; *seed sharp-edged, often vertical*. — Streets of towns: rather scarce. Brackish borders of Onondaga Lake. (Nat. from Eu.)

4. **C. ÚRBICUM**, L. Rather pale or dull green, nearly destitute of mealiness, with erect branches (1° – 3° high); *leaves triangular, acute, coarsely and sharply many-toothed*; *spikes erect, crowded in a long and narrow racemose panicle*; calyx-lobes not keeled; *seed with rounded margins*. — Var. **RHOMBIFOLIUM**, Moquin (*C. rhombifolium*, Muhl.), is a form with the leaves more or less wedge-shaped at the base, and with longer and sharper teeth. — Not rare eastward. (Nat. from Eu.)

5. **C. MURALE**, L. Resembles No. 4, but less erect, loosely branched (1° – $1\frac{1}{2}^{\circ}$ high); *leaves rhomboid-ovate, acute, coarsely and sharply unequally toothed*, thin, bright green; *spikes or racemes diverging, somewhat corymbed*; calyx-lobes scarcely keeled; *seed sharp-edged*. — Boston to Illinois: rare. (Adv. from Eu.)

6. **C. HÝBRIDUM**, L. (MAPLE-LEAVED GOOSEFOOT.) Bright green throughout; stem widely much branched (2° – 4° high); *leaves thin* (2'–8' long), somewhat triangular and heart-shaped, taper-pointed, *sinuate-angled*, the angles extended into a *few large and pointed teeth*; racemes diffusely and loosely panicled, leafless; calyx not fully covering the fruit, its lobes keeled; *seed sharp-edged*, the thin pericarp adhering closely to it. — Common. Heavy-scented, like *Stramonium*. (Nat. from Eu.)

§ 2. **BOTRYOIS**, Moquin. (*AMBRINA*, Moquin, in part.) *Not mealy, but more or less viscid-glandular and pleasant-aromatic: seed frequently vertical, obtuse-edged: embryo forming only two thirds or three quarters of a ring.*

7. **C. BÓTRYIS**, L. (JERUSALEM OAK. FEATHER GERANIUM.) Glandular-pubescent and viscid; *leaves slender-petioled*, oblong, obtuse, *sinuate-pinnatifid*; *racemes cymose-diverging, loose, leafless*; fruit not perfectly enclosed. — Escaped from gardens. (Adv. from Eu.)

8. **C. AMBROSIOIDES**, L. (MEXICAN TEA.) Smoothish; leaves slightly petioled, oblong or lanceolate, repand-toothed or nearly entire, the upper tapering to both ends; spikes densely flowered, leafy, or intermixed with leaves; fruit perfectly enclosed in the calyx. — Waste places: common, especially southward. (Nat. from Trop. Amer.) — Passes into

Var. **ANTHELMINTICUM**. (WORMSEED.) Root perennial (?); leaves more strongly toothed, the lower sometimes almost laciniate-pinnatifid; spikes mostly leafless. (*C. anthelminticum*, L.) — Common in waste places southward. (Nat. from Trop. Amer.)

9. **C. MULTIFIDUM**, L. Glandular-puberulent, diffusely branched; leaves once or twice pinnatifid, pale; flowers small in axillary clusters; calyx only 5-cleft, compressed and completely closed over the glandular-dotted utricle; seed always vertical. (*Roubièva multifida*, Moquin, & Ed. 2.) — Waste places, City of New York (the station now seemingly extinct), and Philadelphia. Introduced in ballast from South America, not permanently established.

3. BLITUM, Tourn. BLITE.

Flowers perfect, bractless. Calyx 3–5-parted, becoming fleshy or berry-like in fruit; the genus also made to include some with calyx unchanged in fruit. Stamens 1–5: filaments filiform. Styles or stigmas 2. Seed vertical, compressed-globular; the embryo coiled into a ring quite around the albumen. — Herbs, with petioled triangular or halberd-shaped and mostly sinuate-toothed leaves. (The ancient Greek and Latin name of some insipid pot-herb.)

§ 1. **MOROCÁRPUS**, Mœnch. *Glabrous annuals or biennials, not mealy: flowers in axillary heads, the upper ones often spiked: calyx in fruit commonly becoming fleshy or berry-like, nearly enclosing the utricle.*

1. **B. marítimum**, Nutt. (COAST BLITE.) Stem angled, much branched; leaves thickish, triangular-lanceolate, tapering below into a wedge-shaped base and above into a slender point, sparingly and coarsely toothed, the upper linear-lanceolate; clusters scattered in axillary leafy spikes; calyx-lobes 2–4, rather fleshy; stamen 1; seed shining, the margin acute. — Salt marshes, New Jersey to Massachusetts: salt springs, at Syracuse, New York (*G. W. Clinton*), and north-westward. Probably a variety of *B. rubrum* of Eu.

2. **B. capitatum**, L. (STRAWBERRY BLITE.) Stem ascending, branching; leaves triangular and somewhat halberd-shaped, sinuate-toothed; clusters simple (large), interruptedly spiked, the upper leafless; stamens 1–5; calyx berry-like in fruit; seed ovoid, flattish, smooth, with a very narrow margin. — Dry rich ground, common from W. New York to Lake Superior, and northward. June. — The calyx becomes pulpy and bright red in fruit, when the large clusters look like Strawberries. (Eu.)

§ 2. **AGATHÓPHYTON**, Moquin. *Somewhat mealy: root perennial: flowers in clusters crowded in a terminal spike: calyx not fleshy, shorter than the half-naked fruit. Intermediate between Blitum and Chenopodium.*

3. **B. Bonus-Henricus**, Reichenbach. (GOOD-KING-HENRY.) Leaves triangular-halberd-form; stamens 5. (*Chenopodium Bonus-Henricus*, L.) — Around dwellings: scarce. (Adv. from Eu.)

4. **ÁTRIPLEX**, Tourn. ORACHE.

Flowers monœcious or diœcious; the staminate like the flowers of *Chenopodium*, only sterile by the abortion of the pistil; the fertile flowers consisting simply of a naked pistil enclosed between a pair of appressed foliaceous (ovate or halberd-shaped) bracts, which are enlarged in fruit, and sometimes united. Seed vertical. Embryo coiled into a ring around the albumen. In one section, to which the Garden Orache belongs, there are also some fertile flowers with a calyx, like those of *Chenopodium*, but without stamens, and with horizontal seeds. — Herbs usually mealy or scurfy with bran-like scales, with triangular or halberd-shaped angled leaves, and spiked-clustered flowers; in summer and autumn. (The ancient Latin name, of obscure meaning.)

1. **A. pátula**, L. Erect or diffusely spreading, annual, *scurfy, green or rather hoary*, branching; leaves alternate or partly opposite, petioled, varying from triangular and halberd-form to lance-linear; fruiting bracts ovate-triangular or rhombic, entire or 1-2-toothed below, united to near the middle, their flat faces either even or sparingly warty-muricate; radicle inferior or somewhat ascending. — The two extreme forms are, Var. **HASTÁTA** (*A. hastata*, L.), with the leaves nearly all triangular-halberd-shaped, entire or sparingly toothed. — Var. **LITTORÁLIS** (*A. littoralis*, L.), with lanceolate or linear mostly entire leaves. — Salt marshes, brackish river-banks, &c., Virginia to Maine, and sparingly on the Great Lakes, and northward. The plant on the shore is more scurfy and hoary; more inland, sometimes far from saline soil, it is greener and thinner-leaved. (Eu.)

2. **A. arenária**, Nutt. *Silvery-mealy* annual, diffusely spreading; leaves oblong, narrowed at the base, nearly sessile; fruiting bracts broadly wedge-shaped, united, 2-3-toothed at the summit, and with a few prickly points on the sides; radicle superior. (*Obione arenaria*, *Moquin*, & Ed. 2.) — Sandy beaches, Massachusetts to Virginia and southward.

5. **CORISPÉRMUM**, Ant. Juss. BUG-SEED.

Flowers perfect, single and sessile in the axil of the upper leaves reduced to bracts, usually forming a spike. Calyx of a single delicate sepal on the inner side. Stamens 1 or 2, rarely 5. Styles 2. Fruit oval, flat, with the outer face rather convex and the inner concave, sharp-margined, a *caryopsis*, i. e. the thin pericarp adherent to the vertical seed. Embryo slender, coiled around a central albumen. — Low branching annuals, with narrow linear alternate 1-nerved leaves. (Name formed of *κόρις*, a bug, and *σπέρμα*, seed.)

1. **C. hyssopifólium**, L. Somewhat hairy when young, pale; floral leaves or bracts awl-shaped from a dilated base or the upper ovate and pointed, scarious-margined; fruit wing-margined. — Sandy beaches of the Great Lakes from Buffalo, a recent immigrant (*G. W. Clinton*), Chicago (*Dr. Scammon*, &c.), to Lake Superior and northwestward. Aug. - Oct. (Eu.)

6. **SALICÓRNIA**, Tourn. GLASSWORT. SAMPHIRE.

Flowers perfect, 3 together immersed in each hollow of the thickened upper joints, forming a spike; the two lateral sometimes sterile. Calyx small and

bladder-like, with a toothed or torn margin, at length spongy and narrowly wing-bordered, enclosing the flattened thin utricle. Stamens 1 or 2. Styles 2, united at base. Seed vertical. Embryo thick, conduplicate: no albumen. — Low saline plants, with succulent leafless jointed stems, and opposite branches; the flower-bearing branchlets forming the spikes. (Name composed of *sal*, salt, and *cornu*, a horn; saline plants with horn-like branches.)

§ 1. *Annuals: spikes very thick and fleshy: flowers and seeds deeply immersed.*

1. **S. herbacea**, L. Erect or at length spreading (6'–12' high), green; scales obscure and very blunt, making a truncate barely emarginate termination of the joints of stem or elongated spike; middle flower much higher than the lateral ones; seed oval or oblong. — Salt marshes of the coast and interior salt springs. Aug.–Oct. (Eu.)

2. **S. Virgínica**, L. (pl. Clayt.) Erect, less branched, naked below (2'–9' high), turning red in age; spike shorter and thicker; scales mucronate-pointed and conspicuous, especially when dry; middle flower little higher than the lateral ones; seed round-oval. (*S. mucronata*, *Lagasca?* 1818, *Bigelow*, and Ed. 2.) — Salt marshes, coast of Virginia to Maine. Sept., Oct. (Eu.?)

§ 2. *Perennial: spikes less thick, and flowers less immersed; middle one hardly higher.*

3. **S. fruticosa**, L., var. **ambigua**. (*S. ambigua*, *Michx.*) Numerous tufted stems (3'–12' long) decumbent or ascending from a hard and rather woody creeping base or rootstock, greenish, turning lead-colored; the cylindrical joints rather strongly notched at the end; seed round-oval. — Sandy wet beaches, &c., Massachusetts to Virginia and southward. Aug.–Oct. (Eu.)

7. SUÆDA, Forskal. SEA BLITE.

Flowers perfect, solitary or clustered in the axils of the leaves. Calyx 5-parted, not appendaged, fleshy, becoming somewhat inflated and closed over the fruit (utricle). Stamens 5. Stigmas 2 or 3. Seed vertical or horizontal, with a flat-spiral embryo, dividing the scanty albumen (when there is any) into 2 portions. — Fleshy maritime plants, with alternate nearly terete linear leaves. (An Arabic name.) CHENOPODINA, Moquin was founded for those species, like ours, which have horizontal seeds, — a wholly insufficient and inconstant difference.

1. **S. marítima**, Dumortier. Annual, smooth, diffusely much branched; leaves slender (1' long), acute; calyx-lobes keeled; stigmas 2; seed horizontal. (*Chenopodina marítima*, Moquin.) — Salt marshes of the sea-shore, and on the northwest plains. Aug. (Eu.)

9. SALSÒLA, L. SALTWORT.

Flowers perfect, with 2 bractlets. Calyx 5-parted, persistent and enclosing the depressed fruit in its base; its divisions at length horizontally winged on the back, the wings forming a broad and circular scarious border. Stamens mostly 5. Styles 2. Seed horizontal, without albumen, filled by the embryo, which is coiled in a conical spiral (cochleate). — Herbs, or slightly shrubby branching plants of the sea-shore, with fleshy and rather terete or awl-shaped leaves, and

sessile axillary flowers. (Name from *sal*, salt; in allusion to the alkaline salts these plants copiously contain.)

1. **S. Kàli**, L. (COMMON SALTWORT.) Annual, diffusely branching, bushy, rough or smoothish; leaves all alternate, awl-shaped, prickly-pointed; flowers single; calyx with the converging lobes forming a sort of beak over the fruit, the large rose or flesh-colored wings nearly orbicular and spreading.—Sandy sea-shore: common. Aug. (Eu.)

ORDER 86. AMARANTACEÆ. (AMARANTH FAMILY.)

Weedy herbs, with nearly the characters of the last family, but the flowers mostly imbricated with dry and scarious persistent bracts; these often colored, commonly 3 in number; the one-celled ovary sometimes many-ovuled. (The greater part of the order tropical, but several have found their way northward as weeds.)

* Anthers 2-celled: filaments separate. Ovule and seed solitary.

1. **Amarantus**. Flowers monœcious or polygamous, all with a calyx of 3 or 5 distinct erect sepals, not falling off with the fruit.
2. **Montella**. Flowers diœcious. Calyx none in fertile flowers. Utricle thin, circumcissile.
3. **Acnida**. Flowers diœcious. Fruit fleshy, indehiscent, 3-5-angled.

* * Anthers 1-celled. Ovule and seed solitary.

4. **Iresine**. Calyx of 5 sepals. Filaments united below into a cup.
5. **Fœrlichia**. Calyx 5-cleft at the apex. Filaments united throughout into a tube.

1. AMARÁNTUS, Tourn. AMARANTH.

Flowers monœcious or polygamous, 3-bracted. Calyx of 5, or sometimes 3, equal erect sepals, glabrous. Stamens 5, rarely 2 or 3, separate: anthers 2-celled. Stigmas 2 or 3. Fruit an ovoid 1-seeded utricle, 2-3-beaked at the apex, mostly longer than the calyx, opening transversely or sometimes bursting irregularly. Embryo coiled into a ring around the albumen.—Annual weeds, of coarse aspect, with alternate and entire petioled leaves, and small green or purplish flowers in axillary or terminal spiked clusters; in late summer and autumn. (*Αμάραντος*, *unfading*, because the dry calyx and bracts do not wither. The Romans, like the Greeks, wrote *Amarantus*, which the early botanists incorrectly altered to *Amaranthus*.)

§ 1. *Utricle thin, circumcissile, the top falling away as a lid: flowers polygamous.*

* *Flowers in terminal and axillary simple or mostly paniced spikes: stem erect (1°-6° high): leaves long-petioled: stamens and sepals 5.*

+ RED AMARANTHS. *Flowers and often leaves tinged with crimson or purple.*

1. **A. HYPOCHONDRIACUS**, L. Smooth or smoothish; leaves oblong-lanceolate, acute or pointed; *spikes very obtuse*, thick, crowded, the terminal one elongated and interrupted; *bracts long-awned*; *fruit 2-3-cleft at the apex*, longer than the calyx.—Rarely spontaneous around gardens. (Virginia, ex L.; but doubtless adv. from Trop. Amer.)

2. **A. PANICULATUS**, L. Stem mostly pubescent; leaves oblong-ovate or ovate-lanceolate; *spikes numerous and slender, paniced*, erect or spreading; *bracts awn-pointed*; *fruit 2-3-toothed at the apex*, longer than the calyx.—Flowers

small, green, tinged with red, or sometimes crimson as in *A. caudatus*, *L.*, the PRINCE'S FEATHER of the gardens. (*A. sanguineus*, *L.*)—In gardens, &c. (Adv. from Trop. Amer.)

+ + GREEN AMARANTHS, PIGWEED. *Flowers green, rarely a little reddish.*

3. **A. RETROFLÉXUS**, *L.* Roughish and pubescent, or smoothish; leaves dull green, long-petioled, ovate or rhombic-ovate, undulate; spikes crowded in a stiff or glomerate panicle; bracts awn-pointed, rigid, exceeding the calyx. — Var. **CHLORÓSTACHYS** (*A. chlorostachys*, *Willd.*) is smoother, with brighter green leaves and less thick and crowded spikes, apparently passing into var. **HÝBRIDUS** (*A. hybridus*, *L.*), which is smooth and more loosely paniced, — perhaps not in our district. — Cultivated and manured soil, gardens, &c. Probably indigenous southwestward. (Adv. from Trop. Amer.?)

* * *Flowers crowded in close and small axillary clusters: stems low, spreading or ascending: stamens and sepals 3, or the former only 2.*

4. **A. ÁLBUS**, *L.* Smooth, pale green; *stems whitish*, mostly spreading next the ground; leaves long-petioled, obovate and spatulate-oblong, very obtuse or retuse; flowers greenish; sepals mucronate, half the length of the rugose fruit, much shorter than the rigid pungently pointed bracts. — Waste grounds, near towns, and roadsides: common. (Nat. from Trop. Amer.?)

§ 2. *Utricle thinnish, bursting or imperfectly circumcissile: flowers monœcious.*

5. **A. SPINÓSUS**, *L.* (THORNY AMARANTH.) Smooth, bushy-branched; stem reddish; leaves rhombic-ovate or ovate-lanceolate, dull green, a pair of spines in their axils; upper clusters sterile, forming long and slender spikes; the fertile globular and mostly in the axils; flowers yellowish-green, small. — Waste grounds, Pennsylvania, Ohio, and southward. (Nat. from Trop. Amer.)

§ 3. **EÛXOLUS**, *Raf.* *Utricle rather fleshy, remaining closed or bursting irregularly: no spines: bracts inconspicuous.*

6. **A. LÍVIDUS**, *L.* Smooth, somewhat succulent, much branched (1°–3° high); *leaves livid-purplish*, long-petioled, ovate or oval; flower-clusters greenish, sessile in the axils and crowded in a terminal interrupted spike; *stamens 3; sepals mostly 3*, rather shorter than the ovate smoothish fruit. (*Euxolus lividus*, *Raf.*) — Coast of Virginia, *Clayton*. Probably an introduced species, and to include *A. oleraceus*, *L.*, and the next.

7. **A. PÙMILUS**, *Raf.* Low or prostrate; leaves more fleshy and obovate, emarginate, the ribs stouter and transverse; flower-clusters small and axillary; *stamens and sepals 5*, the latter half the length of the obscurely 5-ribbed fruit: probably a maritime form of the preceding. (*Euxolus pumilus*, *Raf.*) — Sandy beaches, Rhode Island to Virginia and southward.

8. **A. VÍRIDIS**, *L.* Smooth or minutely pubescent, spreading or ascending (6'–18' high); *leaves pale green*, ovate or ovate-oblong, long-petioled; *flowers much smaller* than in the preceding, in axillary clusters and usually also in a terminal spike; *sepals and stamens 3*, the latter thin, shorter than the small globose-ovate roughish fruit. (*Euxolus deflexus*, *Ed. 2*; but that has a larger and more elongated smooth 3-nerved utricle.) — Streets of Albany, New York: depauperate form with the terminal spike undeveloped. (Adv. from Eu.)

2. **MONTELIA**, Moquin (under ACNIDA).

Flowers diœcious, 2-3-bracted. Staminate flowers of 5 thin oblong and mucronate-tipped sepals, longer than the bracts, and as many stamens with oblong anthers; the cells of the latter united only at the middle. Pistillate flowers without any calyx, their lanceolate awl-pointed bracts longer than the 1-ovuled ovary: stigmas 2-4, very long, bristle-awl-shaped, plumose-hispid. Fruit a thin and membranaceous globular utricle, smooth and even, opening transversely around the middle; the upper part falling off like a lid. Radicle of the annular embryo inferior. — An annual glabrous herb, mostly tall, with lanceolate or oblong-ovate alternate leaves, on long petioles, and small clusters of greenish flowers, usually crowded into elongated and paniced interrupted spikes. (Probably a personal name.)

1. **M. tamariscina**. (*Amarantus tamariscinus*, Nutt. *A. altissimus* & *Miamensis*, Riddell. *Acnida altissima*, Michx. herb. *A. rusocarpa*, Moquin, &c.) — Low grounds and moist sandy shores, Vermont to Wisconsin, Illinois, and southward, especially westward. Aug., Sept. — Var. **CONCATENATA** is a form with the lower clusters in the fertile plant forming thickish distant heads (5"–6" in diameter) in the axils of the leaves; the stems often low and spreading or decumbent. — A very variable plant, as to inflorescence, height (1°–6° high), the size and shape of the leaves (1'–5' long, the petioles often of the same length), the bracts more or less awl-shaped, equalling or exceeding the fruit (which is that of *Amarantus*): but all are forms of one species. The sterile plant is *Acnida rusocarpa*, Michx., or was mixed with it in Michaux's collection, but the fruit is neither obtuse-angled, rugose, nor indehiscent. That name is unmeaning, perhaps a misprint of *ruscocarpa*.

3. **ACNIDA**, L. WATER-HEMP.

Fruit a fleshy and indehiscent utricle, 3-5-angled, the angles often rugose or tubercled-crested. Stigmas 3-5, shorter than the ovary, linear-awl-shaped. Flowers in rather loose paniced spikes. Otherwise as in the preceding genus. (Name formed of a privative and *κνιδη*, a nettle.)

1. **A. cannábina**, L. Annual, tall (2°–6° high); leaves elongated-lanceolate or ovate-lanceolate, long-petioled; fruit globular (1½"–2" long), much exceeding the pointless bracts. — Salt marshes on the coast, Massachusetts to Virginia and southward. Aug.–Oct. — Probably the only species; for *A. rusocarpa*, Michx., is certainly to be divided between this and *Montelia tamariscina*; and *A. tuberculata*, Moquin, must be one or the other.

4. **IRESIÑE**, P. Browne. IRESINE.

Flowers mostly polygamous or diœcious, 3-bracted. Calyx of 5 sepals. Stamens mostly 5: filaments slender, united into a short cup at the base: anthers 1-celled, ovate. Fruit a globular utricle, not opening. — Herbs, with opposite petioled leaves, and minute scarious-white flowers, crowded into clusters or spiked and branching panicles; the calyx, &c. often bearing long wool (whence the name, from *ειρσειώνη*, a branch entwined with fillets of wool borne in processions at festivals.)

1. **I. celosioides**, L. Nearly glabrous annual, erect, slender (2° – 4° high); leaves ovate-lanceolate; panicles narrow, naked; bracts and calyx silvery-white, the latter woolly at the base. — Dry banks, Ohio to Illinois and southward. Sept.

5. **FRÆLÍCHIA**, Mœnch. (OPLOTHÈCA, Nutt.)

Flowers perfect, 3-bracted. Calyx tubular, 5-cleft at the summit, below 2–5-crested lengthwise, or tubercled and indurated in fruit, enclosing the indehiscent thin utricle. Filaments united into a tube, bearing 5 oblong 1-celled anthers, and as many sterile strap-shaped appendages. — Hairy or woolly herbs, with opposite sessile leaves, and spiked scarious-bracted flowers. (Named for *J. A. Frölich*, a German botanist of the last century.)

1. **F. Floridana**, Moquin. Root annual; stem leafless above (1° – 2° high); leaves lanceolate, silky-downy beneath; spikelets crowded into an interrupted spike; calyx very woolly. — Illinois, and southward. Aug.

ORDER 87. **POLYGONACEÆ**. (BUCKWHEAT FAMILY.)

Herbs, with alternate leaves, and stipules in the form of sheaths (ochreæ, these sometimes obsolete) above the swollen joints of the stem; the flowers mostly perfect, with a more or less persistent calyx, a 1-celled ovary bearing 2 or 3 styles or stigmas, and a single erect orthotropous seed. Embryo curved or straightish, on the outside of the albumen, or rarely in its centre; the radicle pointing from the hilum and to the apex of the dry seed-like fruit. Stamens 4–12, inserted on the base of the 3–6-cleft calyx. Leaves usually entire. (The watery juice often acrid, sometimes agreeably acid, as in Sorrel; the roots, as in Rhubarb, sometimes cathartic.) — West of the Mississippi are a great number of ERIOGONEÆ, having their flowers surrounded by an involucre. Our few genera are all of the true Polygonaceæ, except the anomalous Brunnichia.

* Stipular sheaths (*ochreæ*) manifest. Ovule erect from the base of the cell.

→ Sepals 5, sometimes 4, somewhat equal and erect in fruit.

1. **Polygonum**. Embryo curved around one side of the albumen: cotyledons narrow.

2. **Fagopyrum**. Embryo in the albumen, its very broad cotyledons twisted-plaited.

← Sepals 4–6, the outer row reflexed, the inner erect and enlarging in fruit.

3. **Oxyria**. Sepals 4. Stigmas 2. Fruit 2-winged, samara-like.

4. **Rumex**. Sepals 6. Styles 3. Fruit 3-angled, enclosed by the inner sepals.

** Stipules obsolete. Ovule hanging from the apex of a slender stalk.

5. **Brunnichia**. Calyx 5-parted, in fruit with a wing decurrent on the pedicel. Tendril-climber.

1. **POLYGONUM**, L. KNOTWEED.

Calyx mostly 5-parted; the divisions often petal-like, all erect in fruit, withering or persistent and surrounding the lenticular or 3-angular achenium. Stamens 4–9. Styles or stigmas 2 or 3. Embryo placed in a groove on the outside of the albumen and curved half-way around it; the radicle and usually the cotyl-

edons slender. — Pedicels jointed. Ours all herbaceous, flowering through late summer and early autumn. (Name composed of *πολύ*, many, and *γόνυ*, knee, from the numerous joints.)

§ 1. **BISTORTA**, Tourn. *Calyx petal-like, deeply 5-cleft: stamens 8 or 9: styles 3, slender: achenium 3-sided: stems low and simple from a thick and woody creeping rootstock: flowers in a spike-like raceme.*

1. **P. viviparum**, L. (ALPINE BISTORT.) Smooth, dwarf (4'–8' high), bearing a linear spike of flesh-colored flowers (or often little red bulblets in their place); leaves lanceolate. — Alpine summits of the White Mountains, New Hampshire, shore of Lake Superior, and northward. (Eu.)

§ 2. **PERSICARIA**, Tourn. *Calyx petal-like, 5-parted: stigmas capitate: achenium lenticular, or (when there are 3 stigmas) 3-sided cotyledons: accumbent and albumen hard and horny except in No. 2: roots fibrous: flowers crowded in spikes or spike-like racemes.*

* Sheaths some of them with an abrupt spreading and more or less foliaceous (sometimes deciduous) border: tall branching annuals, with paniced and nodding dense cylindrical spikes: flowers rose or flesh-color: achenium flat.

2. **P. orientale**, L. (PRINCE'S FEATHER.) *Soft-hairy; leaves ovate or oblong, pointed, distinctly petioled; flowers large, bright rose-color; stamens 7; style 2-cleft; cotyledons incumbent; albumen floury.* — Sparingly escaped from gardens into waste grounds. (Adv. from India.)

3. **P. Carey**, Olney. Stem (3°–5° high) and peduncles glandular-bristly; leaves lanceolate, roughish; flowers purplish; stamens mostly 5; style 2-parted. — Shady swamps, Rhode Island to the mountains of Pennsylvania, and northward.

* * Sheaths all cylindrical and truncate, without a border.

+ Annuals: spikes oblong or linear, densely flowered: flowers rose or flesh-color, or occasionally varying to white, slightly or not at all glandular-dotted: stamens 6–8: styles 2 or 2-cleft and achenium flattened, except sometimes in No. 7, which alone has the sheaths at all or more than slightly ciliate.

4. **P. Pennsylvanicum**, L. Stem (1°–3° high), smooth below, the branches above, and especially the peduncles, beset with bristly-stalked glands; leaves lanceolate, roughish on the midrib and margins (1½'–5' long); spikes oblong, obtuse (1'–2' long), erect, thick; stamens mostly 8, somewhat exerted; style 2-cleft; achenium with flat sides. — Moist soil, in open waste places: common.

5. **P. incarnatum**, Ell. Nearly glabrous (3°–6° high); the peduncles, &c. often minutely rough with scattered sessile glands; sheaths wholly naked and glabrous; leaves rough on the midrib and margins, elongated-lanceolate (4'–12' long, 1'–3' wide below), tapering gradually from towards the base to a narrow point; spikes linear, nodding, becoming slender (1½'–3' long); stamens 6 and styles 2, both included; achenium with concave sides. (P. nodosum, var. incarnatum, Ed. 2. P. lapathifolium, Amer. authors, &c.) — Wet borders of ponds and streams; rather common everywhere, especially southward and westward. — Flowers smaller than in the last, lighter rose-color, or flesh-color, varying to white.

6. **P. lapathifolium**, Ait., is lower, with shorter and much less pointed leaves; sheaths often somewhat hairy or ciliolate; spikes oblong and blunt; flowers

pale or whitish; *style 2-cleft, or not parted to the base.* We have, sparingly, var. *INCANUM*, Koch (*P. incanum*, Willd.), and only a depauperate form of it, 3'-6' high; leaves lanceolate, obtuse, white-downy beneath; spikes barely 6'' long, erect. — Borders of Cayuga Lake, New York (*Chickering* and *Brewer*), of Lake Superior (*O. B. Wheeler*) and farther northward (*Bourgeau*). — The true *P. nodosum*, *Pers.*, which connects this species with the next, has not been detected. (Eu.)

7. **P. PERSICÀRIA**, L. (LADY'S THUMB.) Nearly smooth and glabrous (12'-18' high); *sheaths more or less bristly-ciliate*; leaves lanceolate, pointed, roughish, usually marked with a dark triangular or lunar spot near the middle; *spikes ovoid or oblong, dense, erect, on smooth (or at least not glandular) peduncles*; stamens mostly 6; *styles half 2-3-cleft*; achenium gibbous-flattened or sometimes triangular, smooth and shining. — Waste and damp places: very common. — Flowers greenish-purple. (Nat. from Eu.)

+ + *Annuals or perennials: spikes slender, filiform, loosely-flowered or interrupted: flowers greenish or whitish, sometimes purplish-tinged: sepals dotted with conspicuous glands: leaves pellucid-dotted: sheaths fringed with bristles: whole herbage pungently acrid!*

8. **P. Hydrópiper**, L. (COMMON SMARTWEED OR WATER-PEPPER.) *Annual*, 1°-2° high, smooth; *spikes nodding, usually short or interrupted*; flowers mostly greenish; stamens 6; *style 2-3-parted*; *achenium dull, minutely striate, either flat or obtusely triangular.* — Moist or wet grounds; apparently introduced eastward, but indigenous northward. (Eu.)

9. **P. àcre**, H. B. K. (WATER SMARTWEED.) *Perennial*, nearly smooth; stems rooting at the decumbent base, 2°-5° high; leaves larger and longer than in the last, taper-pointed; *spikes erect; flowers whitish, sometimes flesh-color*; stamens 8; *style mostly 3-parted, and the achenium sharply triangular, smooth and shining.* (*P. punctatum*, Ell. *P. hydropiperoides*, Pursh.) — Wet places: common, especially southward.

+ + + *Perennials, not acrid, aquatic or amphibious: sepals not glandular-dotted.*

10. **P. hydropiperoides**, Michx. (MILD WATER-PEPPER.) Stem smooth (1°-3° high), branching; the narrow *sheaths hairy, fringed with bristles*; leaves narrowly lanceolate; sometimes oblong; *spikes erect, slender, sometimes filiform, often interrupted at the base (1'-2½' long)*; flowers small, flesh-color or nearly white; *stamens 8; style 3-cleft; achenium sharply triangular, smooth and shining.* (*P. mite*, *Pers.*, not of *Schrank.*) — Wet places, and in shallow water: common, especially southward.

11. **P. amphibium**, L. (WATER PERSICARIA.) Stems somewhat simple, and usually terminated by a single ovate or oblong and *very dense spike (6'' thick)* of rather large *bright rose-red flowers*; sheaths not fringed with bristles; *stamens 5; style 2-cleft*; achenium biconvex; leaves varying greatly, from elliptical or cordate-oblong to lance-ovate or lanceolate. — The typical plant is var. *AQUÁTICUM*, Willd. (Var. *natans*, Mœnch. *P. coccineum*, Muhl., &c.), in water, nearly glabrous; leaves long-petioled, floating, elliptical or oblong, sometimes heart-shaped at base; spikes short. — Var. *TERRÉSTRE*, Willd., grows in shallow water, or in wet soil, or even "in sandy prairies" in Illinois

(*Dr. Mead*), either almost glabrous or strigose-hirsute; leaves short-petioled, varying from elongated-lanceolate to lance-ovate and taper-pointed; spikes cylindrical (1'-3' long) often in pairs. — Common. (Eu.)

§ 3. **TOVÀRIA**, Adans. *Calyx rather herbaceous (greenish), unequally 4-parted: stamens 5: styles 2, distinct, rigid and persistent on the smooth lenticular achenium: cotyledons oblong, accumbent: perennial: flowers loosely disposed in naked long and slender spikes.*

12. **P. Virginianum**, L. Almost smooth; stem terete, upright (2° – 4° high); leaves ovate, or the upper ovate-lanceolate, taper-pointed, rounded at the base, short-petioled, rough-ciliate (3'–6' long); sheaths cylindrical, truncate, hairy and fringed; flowers 1–3 from each bract, somewhat curved, the styles in fruit obliquely bent down, minutely hooked at the tip. — Thickets in rich soil: common.

§ 4. **PSEUDO-POLYGONÉLLA**. *Characters of the next section, except that the sheaths are truncate, naked and rigid; the flowers racémed, solitary from each truncate bract; pedicels jointed below the middle; calyx expanding and petaloid.*

13. **P. articulatum**, L. (JOINTWEED.) Annual; stem upright, paniculately branched (4'–12' high), slender; leaves linear-thread-form, deciduous; flowers rose-color, crowded in slender and spike-like paniced racemes, on recurved pedicels twice the length of the crowded joint-like bracts; stamens 8; achenium triangular, smooth and shining. (*Polygonella articulata*, *Meisner*, but has not the enlarged connivent inner sepals, nor the axile embryo of that genus.) — Dry, sandy soil; along the coast and the Great Lakes, and in intermediate places in New York. — Singular for its many-jointed spikes or racemes, which are 1'–3' long; the lower bracts tooth-pointed on one side. Three inner filaments dilated at the base.

§ 4. **AVICULÀRIA**, Meisn. *Calyx more or less petal-like, 5-parted: stamens 3–8; the filaments awl-shaped, 3 of them broader at the base: stigmas 3, globose, nearly sessile: achenium 3-sided: cotyledons incumbent: albumen horny: flowers inconspicuous, greenish-white, 2 or 3 together or sometimes solitary in the axils of the small leaves, appearing nearly sessile, sometimes more or less spiked along the leafless summits of the branches: ours all annuals or nearly so: sheaths scariosus, usually 2–3-cleft or cut-fringed and torn.*

14. **P. aviculàre**, L. (KNOTGRASS. GOOSE-GRASS. DOOR-WEED.) *Prostrate or spreading; leaves sessile, lanceolate or oblong, pale; stamens chiefly 5 in the American, 8 in the European plant; achenium dull, minutely granular under a lens, enclosed in the calyx.* — The commonest weed, in yards, waste places, &c. (Eu.)

Var. **eréctum**, Roth. Stems upright or ascending; leaves broader (oblong or oval) and larger. (*P. eréctum*, L.) — In richer soil or more shaded places: common.

15. **P. marítimum**, L. (COAST KNOTGRASS.) *Prostrate, glaucous, with a hard and sometimes woody and perennial root; stems very short-jointed; scariosus sheaths large; leaves thickened, elliptical-lanceolate or narrow oblong; flowers larger than in the last; achenium very smooth and shining.* (*P. glaucum*,

Nutt. *P. aviculare*, var. *littorale*, (*Ed.* 2.) — Sandy sea-beach, Mass. to Virginia and southward: at the north apparently only annual. (Eu.)

16. *P. ramosissimum*, Michx. *Stems erect or ascending, much branched (2°–4° high), rigid, many-striate; leaves lanceolate or linear, tapering into a petiole; sheaths mostly short; calyx (drying yellowish) 6-parted and with 6 or 3 stamens, or 5-parted and with 4 or 5 stamens; achenium very smooth and shining.* — Sandy shores and banks of streams, Michigan to Illinois and southward. Salt marshes, Rhode Island, *Olney*. — Larger leaves 2' long.

17. *P. tenue*, Michx. *Stem slender, upright, sparingly branched (6'–12' high), sharp-angled; leaves sessile, narrowly linear, very acute; sheaths capillary fringed; flowers axillary and loosely spiked on the filiform branches; achenium smooth and shining.* — Dry soil, and rocky hills.

§ 6. *TINIÀRIA*, Meisn. *Calyx 5-parted (rarely 4-parted), greenish tinged with rose-color or white: stamens mostly 8: styles or capitate stigmas 3 and achenium 3-sided (except in No. 18): leaves heart-shaped or arrow-shaped, petioled: sheaths semicylindrical.*

* *Annuals, erect or somewhat climbing by the reflexed prickles which beset the angles of the stem and petioles: divisions of the (pale rose-colored or white) calyx not keeled: bracts chaff-like.*

18. *P. arifolium*, L. (HALBERD-LEAVED TEAR-THUMB.) *Stem grooved-angled; leaves halberd-shaped, taper-pointed, long-petioled; flowers somewhat racemed (few); peduncles glandular-bristly; calyx often 4-parted; stamens 6; styles 2, very short; achenium lenticular (large).* — Low grounds.

19. *P. sagittatum*, L. (ARROW-LEAVED TEAR-THUMB.) *Stem 4-angled; leaves arrow-shaped, short-petioled; flowers capitate; peduncles smooth; stamens mostly 8; styles 3, slender; achenium sharply 3-angled.* — Low grounds: common. — Slender, smooth except the angles of the stem and midrib beneath: these are armed with a line of fine and very sharp saw-toothed prickles, which cut the hand drawn against them.

* * *Stems twining, not prickly: calyx with the 3 outer divisions keeled, at least in fruit: flowers in loose paniced racemes: bracts like the stipules.*

20. *P. CONVÓLVULUS*, L. (BLACK BINDWEED.) *Annual; stems twining or procumbent, low, roughish, the joints naked; leaves halberd-heart-shaped, pointed; flowers in small interrupted corymbose racemes; outer calyx-lobes keeled; achenium smoothish.* — Cult. and waste grounds: common. (Nat. from Eu.)

21. *P. cilinòde*, Michx. *Perennial, minutely downy; the sheaths fringed at the base with reflexed bristles; leaves heart-shaped and slightly halberd-shaped, taper-pointed; racemes paniced; calyx-lobes obscurely keeled; achenium very smooth and shining.* — Copses and rocky hills. Climbing 3°–9° high.

22. *P. dumetòrum*, L. (CLIMBING FALSE BUCKWHEAT.) *Perennial, smooth; sheaths naked; leaves heart-shaped or slightly halberd-shaped, pointed; racemes interrupted, leafy; the 3 outer calyx-lobes strongly keeled and in fruit winged; achenium smooth and shining.* (Eu.)

Var. *scàndens* (*P. scandens*, L.) has more paniced flowers, and usually much broader wings to the fruiting calyx than the European. — Moist thickets. Twining 8°–12° high over bushes.

2. FAGOPYRUM, Tourn. BUCKWHEAT.

Calyx petal-like, equally 5-parted, withering and nearly unchanged in fruit. Stamens 8. Styles 3: stigmas capitate. Achenium 3-sided, longer than the calyx. Embryo large, in the centre of the albumen, which it divides into 2 parts, with very broad and foliaceous plaited and twisted cotyledons. — Annuals, with triangular-heart-shaped or halberd-shaped leaves, semicylindrical sheaths, and corymbose racemes or panicles of white flowers, often tinged with green or rose-color. (Name, *φηγός*, *the beech*, and *πυρός*, *wheat*, from the shape of the grain being that of the beech-nut; whence also the English name *Buckwheat*, from the German *Buchſt*, *beech*.)

1. *F. ESCULENTUM*, Moench. (BUCKWHEAT.) Smoothish; flower with 8 honey-bearing yellow-glands interposed between the stamens; the fruit acute and entire. (*Polygonum Fagopyrum*, *L.*) — Old fields, remaining as a weed after cultivation, and escaping into copses. June — Sept. (Adv. from Eu.)

3. OXYRIA, Hill. MOUNTAIN SORREL.

Calyx herbaceous, of 4 sepals; the two outer smaller and spreading, the two inner broader and erect (but unchanged) in fruit. Stamens 6. Stigmas 2, sessile, tufted. Achenium lenticular, thin, flat, much larger than the calyx, surrounded by a broad and veiny wing. Seed flattened in the opposite direction from the wing. Embryo straight, occupying the centre of the albumen, slender. — Low alpine perennials, with round-kidney-form and long-petioled leaves chiefly from the root, obliquely truncate sheaths, and small greenish flowers clustered in paniced racemes on a slender scape. (Name from *ὄξύς*, *sour*, in allusion to the acid leaves, like those of Sorrel.)

1. *O. digyna*, Campdera. Leaves all round-kidney-form, usually notched at the end; fruit orbicular. — Alpine region of the White Mountains, New Hampshire, *Oukes*, &c., and high northward. (Eu.)

4. RUMEX, L. DOCK. SORREL.

Calyx of 6 sepals; the 3 outer herbaceous, sometimes united at the base, spreading in fruit; the 3 inner larger, somewhat colored, increasing after flowering and convergent over the 3-angled achenium, veiny, often bearing a grain-like tubercle on the back (in fruit called *valves*). Stamens 6. Styles 3: stigmas tufted. Embryo slightly curved, lying along one side of the albumen, slender. — Coarse herbs, with small and homely (mostly green) flowers, which are crowded and commonly whorled in paniced racemes; the petioles somewhat sheathing at the base. (The ancient Latin name of these plants; of unknown etymology.)

§ 1. DOCK. *Flowers perfect or monœciously polygamous: herbage not sour: none of the leaves halberd-shaped. (Flowering through the summer.)*

* *Perennials, 2°–7° high: valves not bearing bristles.*

+ *Valves (large, 3" broad, thin) all naked or one with a small grain.*

1. *R. PATIÉNTIA*, L. (PATIENCE DOCK.) A very tall species, with ovate-oblong and lanceolate leaves, those from the root 2°–3° long, and one of the

heart-shaped nearly entire valves bearing a small grain, or its midrib thickened at the base, was found spontaneous at Amherst, Mass., by *Prof. Tuckerman*, in the form with undulate leaves, *R. orientalis*, *Bernh.* (Adv. from Eu.)

2. **R. LONGIFOLIUS**, DC. (*R. domesticus*, *Hartm.*, and too near *R. aquaticus*, *L.*), known by the rounded somewhat heart-shaped valves all without a grain, — is indigenous at the northwest, perhaps in Northern Wisconsin. (Eu.)

+ *Valves (smaller) one or more of them conspicuously grain-bearing.*

+ *Indigenous: leaves not wavy, none heart-shaped, except the lowest of No. 6.*

3. **R. orbiculatus**, Gray. (GREAT WATER-DOCK.) Tall and stout (5°–6° high); racemes upright in a large compound panicle, nearly leafless; whorls crowded; *pedicels capillary, nodding, about twice the length of the fruiting calyx; the valves orbicular or round-ovate, very obtuse, obscurely heart-shaped at base, membranaceous, finely reticulated, entire or repand-denticulate (2''–3'' broad), all grain-bearing; leaves oblong-lanceolate, rather acute at both ends, transversely veined, and with obscurely erose-crenulate margins (the lowest, including the petiole, 1°–2° long, the middle rarely truncate or obscurely cordate at base).* (*R. Hydrolápathum*, var. ? *Americanus*, *Ed. 2.* *R. Britannica*, *Pursh?* *Bigelow*, *Torr.* Fl. N. Y. *R. aquaticus*, *Pursh?*) — Wet places: rather common northward. — Root yellow. Leaves occasionally abruptly contracted a little below the enlarging apex. Valves very much rounder, thinner, larger in proportion to the grain, and more reticulated than in the European *R. Hydrolapathum*, resembling those of *R. longifolius* except in bearing a conspicuous grain.

4. **R. Británnica**, L. (PALE DOCK.) Rather tall (2°–6° high); racemes spike-like and paniced, nearly leafless; whorls crowded; *pedicels nodding, shorter than the fruiting calyx; the valves broadly ovate or obscurely heart-shaped, obtuse or acutish, entire, membranaceous, loosely reticulated (about 2'' broad), one with a conspicuous grain, the others with a small grain or thickened midrib, or naked; leaves ovate-lanceolate or oblong-lanceolate, acute, pale, thickish, obscurely veiny (those of the stem 3'–6' long, contracted at base into a short petiole), none heart shaped.* (*R. Claytonii*, *Campdera*, which name is to be adopted if we reject that inconsiderately assigned by Linnæus, who transferred the obscure *Herba Britannica* of the old writers to a Virginian species. *R. xanthorhizus*, *Hoffmansegg*, ex *Meisner*. *R. altissimus*, *Wood.*) — Moist grounds, New York to Illinois and southward.

5. **R. salicifolius**, Weinmann. (WHITE DOCK.) Rather low (1°–3° high); root white; *leaves narrowly or linear-lanceolate, or the lowest oblong; whorls much crowded; pedicels much shorter than the fruiting calyx; valves deltoid-ovate, obtusish or acutish (about 1½'' long), one, two or sometimes all three with a conspicuous often very large grain: otherwise nearly as in the preceding.* (*R. pallidus*, *Bigelow.*) — Salt marshes, coast of New England; also far westward and northward.

6. **R. verticillatus**, L. (SWAMP DOCK.) Rather tall (3°–5° high); racemes nearly leafless, elongated, loose; the whorls crowded or the lower ones distant; fruit-bearing *pedicels slender, club-shaped, abruptly reflexed, 3–4 times longer than the fruiting calyx; the valves dilated-rhomboid, obtusely somewhat pointed, strongly rugose-reticulated, each bearing a very large grain; leaves lanceolate or*

oblong-lanceolate, rather obtuse, thickish, pale-green, the lowest often heart-shaped at the base. — Wet swamps: common.

→ → *Naturalized weeds: lower leaves mostly heart-shaped at the base.*

7. **R. CRISPUS**, L. (CURLLED DOCK.) Smooth; leaves with strongly wavy-curved margins, lanceolate, acute, the lower truncate or scarcely heart-shaped at the base; whorls crowded in prolonged wand-like racemes, leafless above; valves round-heart-shaped, obscurely denticulate or entire, mostly all of them grain-bearing. — A very common weed in cultivated and waste grounds. Stem 3°–4° high, from a deep spindle-shaped yellow root. There is a hybrid of this with the next. (Nat. from Eu.)

8. **R. OBTUSIFOLIUS**, L. (BITTER DOCK.) Stem roughish; lowest leaves ovate-heart-shaped, obtuse, rather downy on the veins underneath, somewhat wavy-margined, the upper oblong-lanceolate, acute; whorls loose and distant; valves ovate-halberd-shaped, and with some sharp awl-shaped teeth at the base, strongly reticulated, one of them principally grain-bearing. — Fields, &c. (Nat. from Eu.)

9. **R. CONGLOMERATUS**, Murray. (SMALLER GREEN DOCK.) Leaves oblong, pointed, slightly wavy-margined, the lower heart-shaped at the base; whorls distant, leafy; pedicels very short; valves linear-oblong, rather broader next the base; obtuse, entire, each bearing a single reddish grain. (*R. acutus*, Smith.) — Moist places; sparingly introduced. (Nat. from Eu.)

10. **R. SANGUINEUS**, L. (BLOODY-VEINED DOCK.) Leaves lanceolate, wavy-margined, the lowest heart-shaped at the base; whorls distant, in long and slender leafless interrupted spikes; pedicels very short; valves narrowly oblong, obtuse, entire, one at least grain-bearing; veins of the leaf red, or, in var. *viridis*, green. — Waste and cultivated grounds. (Nat. from Eu.)

* * *Annuals, low: valves bearing long awns or bristles.*

11. **R. MARITIMUS**, L. (GOLDEN DOCK.) Minutely pubescent, diffusely branched; leaves lance-linear, wavy-margined, the lower auricled or heart-shaped at base; whorls excessively crowded in leafy and compact or interrupted spikes; valves rhombic-oblong, lance-pointed, each bearing 2–3 long awn-like bristles on each side, and a large grain on the back. (Also *R. persicarioides*, L.) — Seashore, Virginia to Massachusetts: also Illinois and westward. — Plant 6'–12' high; remarkable for the crowded and almost orange-colored fruiting calyx, the bristles usually longer than the width of the valves. (Eu.)

§ 2. **SORREL**. *Flowers dioecious, small, in a terminal naked panicle: herbage sour: some leaves halberd-shaped: smooth perennials, flowering in spring.*

12. **R. ENGELMANNI**, Ledeb. Stem simple, 1°–2° high; leaves nearly as in the next; pedicels jointed at or below the middle; valves of the fruiting calyx round-heart-shaped, thin, finely reticulated, naked, many times larger than the achenium. (*R. hastulatus*, Baldwin.) — S. W. Illinois, thence southward and westward.

13. **R. ACETOSÉLLA**, L. (FIELD OR SHEEP SORREL.) Low (6'–12' high); leaves lance-halberd-form, at least those of the root, the narrow lobes entire; pedicels jointed with the flower; valves scarcely enlarging in fruit, ovate, naked. — An abundant weed in waste places and all sterile and worn fields. — The fertile panicles usually turn reddish in summer. (Nat. from Eu.)

5. BRUNNÍCHIA, Banks. BRUNNICHIA.

Calyx 5-parted; the divisions somewhat petal-like, oblong, connivent and coriaceous in fruit. Stamens 8: filaments capillary: styles 3, slender: stigmas depressed-capitate. Ovule hanging on the summit of a slender erect funiculus: the seed erect, 6-grooved. Embryo in one of the angles of the mealy albumen, somewhat curved. Achenium obtusely triangular, partly 3-celled, enclosed in the indurated calyx, its base and almost the whole length of the pedicel winged on one side. (Named for *F. Brunnich*, a Danish naturalist.)

1. *B. cirrhosa*, Banks. — A somewhat shrubby smooth plant, with grooved stems, climbing by tendrils extended from the ends of the branches. Leaves ovate or heart-shaped, pointed, entire: petioles dilated at base and partly clasping; but no distinct sheath or stipules. Flowers greenish, 2-5 in a fascicle from the axil of an awl-shaped bract, these crowded in axillary and terminal racemes: pedicel jointed near the base. Fruiting calyx with the wing 1' long. — Pulaski Co., S. W. Illinois, *Dr. F. Brendel*: also southward.

ORDER 88. LAURÀCÆ. (LAUREL FAMILY.)

Aromatic trees or shrubs, with alternate simple leaves mostly marked with minute pellucid dots, and flowers with a regular calyx of 4 or 6 colored sepals, imbricated in 2 rows in the bud, free from the 1-celled and 1-ovuled ovary, and mostly fewer than the stamens; anthers opening by 2 or 4 uplifted valves. — Flowers clustered. Style single. Fruit a 1-seeded berry or drupe. Seed anatropous, suspended, with no albumen, filled by the large almond-like embryo. — A well-marked family, very numerous in the tropics, represented in our district by only five species, in four genera.

* Flowers perfect, panicled: stamens 12, three of them sterile.

1. *Persea*. Calyx persistent. Anthers 4-celled, those of three stamens turned outward.

** Flowers diœcious, or nearly so: stamens in the male flowers 9.

2. *Sassafras*. Flowers in corymb- or umbel-like racemes. Anthers 4-celled, 4-valved.

3. *Lindera*. Flowers in umbel-like clusters. Anthers 2-celled, 2-valved.

4. *Tetranthera*. Flowers few in involucrate umbels. Anthers 4-celled, 4-valved.

1. PÉRSEA, Gærtn. ALLIGATOR PEAR.

Flowers perfect, with a 6-parted calyx, which persists at the base of the berry-like fruit. Stamens 12, in four rows, the 3 of the innermost row sterile and reduced to a sort of glands: the rest bearing 4-celled anthers (i. e. each of the two proper cells is divided transversely into two), opening by as many uplifted valves; the anthers of 3 stamens turned outward, the others introrse. — Trees, with persistent entire leaves, and small panicled flowers. (An ancient name of some Oriental tree.)

1. *P. Carolinensis*, Nees. (RED BAY.) Hoary at least when young with a fine down; leaves oblong, pale, soon becoming smooth above; peduncle bearing few flowers in a close cluster; sepals downy, the outer shorter; berries dark blue, on a red stalk. (*Laurus Carolinensis*, *Catesb.* *L. Borbonia*, *L.*) — Swamps, Delaware, Virginia, and southward. May. — A small tree.

2. SASSAFRAS, Nees. SASSAFRAS.

Flowers diœcious, with a 6-parted spreading calyx; the sterile kind with 9 stamens inserted on the base of the calyx in 3 rows, the 3 inner with a pair of stalked glands at the base of each; anthers 4-celled, 4-valved: fertile flowers with 6 short rudiments of stamens and an ovoid ovary. Drupe ovoid (blue), supported on a club-shaped and rather fleshy reddish pedicel. — Trees, with spicy-aromatic bark, very mucilaginous twigs and foliage; the latter deciduous, often lobed. Flowers greenish-yellow, naked, in clustered and peduncled corymbed racemes, appearing with the leaves, involucre with scaly bracts. Leaf-buds scaly. (The popular name, of Spanish origin.)

1. **S. officinale**, Nees. Leaves ovate, entire, or some of them 3-lobed, soon glabrous. (*Laurus Sassafras*, *L.*) — Rich woods: common. April. — Tree 15° – 50° high, with yellowish-green twigs.

3. LÍNDERA, Thunberg. WILD ALLSPICE. FEVER-BUSH.

Flowers polygamous-diœcious, with a 6-parted open calyx; the sterile kind with 9 stamens in 3 rows, the inner filaments 1 – 2-lobed and gland-bearing at the base; anthers 2-celled and 2-valved: fertile flowers with 15 – 18 rudiments of stamens in 2 forms, and a globular ovary. Drupe obovoid, red, the stalk not thickened. — Shrubs, with deciduous leaves, and honey-yellow flowers in almost sessile lateral umbel-like clusters, appearing before the leaves (in our species); the clusters composed of smaller clusters or umbels, each of 4 – 6 flowers and surrounded by an involucre of 4 deciduous scales. Leaf-buds scaly. (Named for *John Linder*, a Swedish botanist of the early part of the 18th century, Benzoin, *Nees*, adopted in Ed. 2, is a much later name.)

1. **L. Benzoin**, Meisner. (SPICE-BUSH. BENJAMIN-BUSH.) *Nearly smooth; leaves oblong-obovate*, pale underneath. (*Laurus Benzoin*, *L. Benzoin odoriferum*, *Nees*, & Ed. 2.) — Damp woods: rather common. March, April. — Shrub 6° – 15° high.

2. **L. melissæfolia**, Blume. Young branches and buds *pubescent; leaves oblong, obtuse or heart-shaped* at the base, downy beneath; umbels few. (*Laurus melissæfolia*, *Walt.* *L. diospyroides*, *Michx.* *Benzoin melissæfolium*, *Nees.*) — Low grounds, Virginia, Illinois? and southward. April.

4. TETRANTHÈRA, Jacq. TETRANTHERA.

Flowers diœcious, with a 6-parted deciduous calyx; the sterile with 9 stamens in 3 rows; their anthers all introrse, 4-celled, 4-valved: fertile flowers with 12 or more rudiments of stamens and a globular ovary. Drupe globular. — Shrubs or trees, with entire leaves, and small flowers in axillary clustered umbels. (Name composed of *τέτρα*, *four*, and *ἀνθηρά*, *anther*.)

1. **T. geniculata**, Nees. (POND SPICE.) Flowers (yellow) appearing before the deciduous oblong leaves, which are hairy on the midrib beneath; branches forked and divaricate, the branchlets zigzag; involucre 2 – 4-leaved, 2 – 4-flowered; fruit red. (*Laurus geniculata*, *Michx.*) — Swamps, Virginia and southward. April.

ORDER 89. **THYMELEACEÆ.** (MEZEREUM FAMILY.)

Shrubs, with acrid and very tough (not aromatic) bark, entire leaves, and perfect flowers with a regular and simple colored calyx, bearing usually twice as many stamens as its lobes, free from the 1-celled and 1-ovuled ovary, which forms a berry-like drupe in fruit, with a single suspended anatropous seed. Embryo large: albumen little or none. — A small family, represented in cultivation by DAPHNE MEZEREUM, and one or two other species; in North America only by a single species.

1. **DÍRCA**, L. LEATHERWOOD. MOOSE-WOOD.

Calyx petal-like, tubular-funnel-shaped, truncate, the border wavy or obscurely about 4-toothed. Stamens 8, long and slender, inserted on the calyx above the middle, protruded, the alternate ones longer. Style thread-form: stigma capitate. Drupe oval (reddish). — A much-branched bush, with jointed branchlets, oval-obovate alternate leaves, at length smooth, deciduous, on very short petioles, the bases of which conceal the buds of the next season. Flowers light yellow, preceding the leaves, 3 or 4 in a cluster from a bud of as many dark-hairy scales, forming an involucre, from which soon after proceeds a leafy branch. (*Δίρκη*, the name of a fountain near Thebes, applied by Linnæus to this North American genus, for no imaginable reason, unless because the bush frequently grows near mountain rivulets.)

1. **D. palústris**, L. — Damp rich woods, seldom in swamps, New England to Penn., Kentucky, and (especially) northward. April. — Shrub 2°–5° high; the wood white, soft, and very brittle; but the fibrous bark remarkably tough, used by the Indians for thongs, whence the popular names. In Northern New England also called **WICOPY**.

ORDER 90. **ELÆAGNACEÆ.** (OLEASTER FAMILY.)

Shrubs or small trees, with silvery-scurfy leaves and mostly diœcious flowers; further distinguished from the Mezereum Family by the erect or ascending albuminous seed, and the calyx-tube becoming pulpy and berry-like in fruit, and strictly enclosing the achenium; and from the following or by the calyx-tube not cohering with the ovary, &c. A small family, represented by only three North American species, only one strictly within our limits.

1. **SHEPHÉRDIA**, Nutt. SHEPHERDIA.

Flowers diœcious; the sterile with a 4-parted calyx (valvate in the bud) and 8 stamens, alternating with as many processes of the thick disk; the fertile with an urn-shaped 4-cleft calyx, enclosing the ovary (the orifice closed by the teeth of the disk), and becoming berry-like in fruit. Style slender: stigma 1-sided. — Leaves opposite, entire, deciduous; the small flowers nearly sessile in their axils on the branchlets, clustered, or the fertile solitary. (Named for *John Shepherd*, formerly curator of the Liverpool Botanic Garden.)

1. **S. Canadensis**, Nutt. (CANADIAN SHEPHERDIA.) Leaves elliptical or ovate, nearly naked and green above, silvery-downy and scurfy with rusty scales underneath; fruit yellowish-red, insipid. — Rocky or gravelly banks, Vermont to Wisconsin and northward. May. — Shrub, 3°–6° high; the branchlets, young leaves, yellowish flowers, &c., covered with the rusty scales.

2. **S. argentea**, Nutt., the BUFFALO-BERRY of Upper Missouri, has narrower leaves, tapering at base, silvery on both sides, and edible, acid, scarlet fruit: probably in Minnesota: sometimes cultivated.

ELÆÁGNUS ARGÉNTEA, Pursh, the SILVER-BERRY, with oval silvery leaves, and silvery and mealy edible fruit, differing from Shepherdia in its perfect flowers with only 4 stamens, — abounds not far beyond our northwestern limits.

ORDER 91. SANTALÁCEÆ. (SANDALWOOD FAMILY.)

Herbs, shrubs, or trees, with entire leaves; the 4–5-cleft calyx valvate in the bud, its tube coherent with the 1-celled ovary, which contains 2–4 ovules suspended from the apex of a stalk-like free central placenta which rises from the base of the cell, but the (indehiscent) fruit always 1-seeded. — Seed destitute of any proper seed-coat. Embryo small, at the apex of copious albumen: radicle directed upward: cotyledons cylindrical. Stamens equal in number to the lobes of the calyx, and inserted opposite them into the edge of the fleshy disk at their base. Style 1. A small order, the greater part belonging to warm regions, here represented only by the two following genera.

1. COMÁNDRA, Nutt. BASTARD TOAD-FLAX.

Flowers perfect. Calyx bell-shaped or soon urn-shaped, lined above the ovary with an adherent disk which has a 5-lobed free border. Stamens inserted on the edge of the disk between its lobes, opposite the lobes of the calyx, to the middle of which the anthers are connected by a tuft of thread-like hairs. Fruit drupe-like or nut-like, crowned by the persistent calyx-lobes, the cavity filled by the globular seed. — Low and smooth perennials, with herbaceous stems from a rather woody base or root, alternate and almost sessile leaves, and greenish-white flowers in terminal or axillary small umbel-like clusters. (Name from κόμη, *hair*, and ἄνδρες, for *stamens*, in allusion to the hairs on the calyx-lobes which are attached to the anthers.)

1. **C. umbellata**, Nutt. Leaves oblong, pale (1' long); peduncles several and corymbose-clustered at the summit of the stem, several-flowered; calyx-tube conspicuously continued beyond the ovary, forming a neck to the globular-urn-shaped fruit; the lobes oblong; style slender; fruit dry. — Dry ground: common. May, June. — Stems 8'–10' high, very leafy. Root forming parasitic attachments to the roots of trees (as shown by Mr. Stauffer).

2. **C. livida**, Richardson. Peduncles axillary, 3–5-flowered, shorter than the oval leaves; calyx-tube not continued beyond the ovary, the lobes ovate; style short; fruit pulpy when ripe, red. — Sandy shores of Lake Superior, and northward. — Leaves larger than in the last.

2. **PYRULÀRIA**, Michx. OIL-NUT. BUFFALO-NUT.

Flowers diœcious or polygamous. Calyx 4-5-cleft, the lobes recurved; a tuft of hairs at their base in the male flowers. Stamens 4 or 5, on very short filaments, alternate with as many rounded glands. Fertile flowers with a pear-shaped ovary invested by the adherent tube of the calyx, naked at the flat summit: disk with 5 glands: style short and thick: stigma capitate-flattened. Fruit fleshy and drupe-like, pear-shaped; the globose endocarp thin. Embryo small: albumen very oily. — Shrubs or trees, with alternate short-petioled and deciduous leaves; the small greenish flowers in short and simple spikes or racemes. (Name a diminutive of *Pyrus*, from the fruit, which in the original species looks like a small pear.)

1. **P. oleifera**, Gray. Shrub straggling (3°-12° high), minutely downy when young, at length nearly glabrous; leaves obovate-oblong, acute or pointed at both ends, soft, very veiny, minutely pellucid-punctate; spike small and few-flowered, terminal; calyx 5-cleft. (*P. pùbera*, Michx.; a little older than the other specific name, but much less appropriate. *Hamiltônia oleifera*, Muhl.) — Rich woods, mountains of Pennsylvania, and southward through the Alleghanies. May. — Whole plant imbued with an acrid oil, especially the fruit, which is an inch long.

ORDER 92. **LORANTHACEÆ**. (MISTLETOE FAMILY.)

Shrubby plants with coriaceous greenish foliage, parasitic on trees, represented in the northern temperate zone chiefly by the Mistletoe and its near allies; distinguished from the preceding family more by the parasitic growth and habit, and by the more reduced flowers, than by essential characters: represented by an American genus nearly allied to *Viscum*, or true Mistletoe, viz.

1. **PHORADÉNDRON**, Nutt. FALSE MISTLETOE.

Flowers diœcious, in short and catkin-like jointed spikes, usually several under each short and fleshy bract or scale, and sunk in the joint. Calyx globular, 3- (rarely 2-4-) lobed: in the staminate flowers a sessile anther is borne on the base of each lobe, and is transversely 2-celled, each cell opening by a pore or slit: in the fertile flowers the calyx-tube adheres to the ovary: stigma sessile, obtuse. Berry 1-seeded, pulpy. Embryo small, half imbedded in the summit of mucilaginous albumen. — Yellowish-green woody parasites on the branches of trees, with jointed much-branched stems, thick and firm persistent leaves (or only scales in their place), and axillary small spikes of flowers. (Name composed of *φωρ*, a thief, and *δένδρον*, tree; because these plants steal their food from the trees they grow upon.)

1. **P. flavescens**, Nutt. (AMERICAN MISTLETOE.) Leaves obovate or oval, somewhat petioled, longer than the spikes in their axils, yellowish; berries white. (*Viscum flavescens*, Pursh.) — New Jersey to Illinois and southward, on various deciduous-leaved trees.

ORDER 93. SAURURACEÆ. (LIZARD'S-TAIL FAMILY.)

Herbs, with jointed stems, alternate entire leaves, and perfect flowers in spikes, entirely destitute of any floral envelopes, and with 3-5 more or less separate or united ovaries. — Ovules few, orthotropous. Embryo heart-shaped, minute, contained in a little sac at the apex of the albumen. — An offshoot of the (tropical) Pepper Family, and represented only by

1. SAURURUS, L. LIZARD'S-TAIL.

Stamens mostly 6 or 7, hypogynous, with distinct filaments. Fruit somewhat fleshy, wrinkled, of 3-4 pistils united at the base. Stigmas recurved. Seeds usually solitary, ascending. — Perennial marsh herbs, with heart-shaped converging-ribbed petioled leaves, without distinct stipules; flowers (each with a small bract) crowded in a slender wand-like and naked peduncled terminal spike or raceme (its appearance giving rise to the name, from *σαῦρος*, a lizard and *οὐρά*, tail). Bract adnate to or borne on the pedicel.

1. *S. cernuus*, L. Flowers white, in a dense spike nodding at the end; bract lanceolate; filaments long and capillary. — Swamps: common. June-Aug.

ORDER 94. CERATOPHYLLACEÆ. (HORNWORT FAMILY.)

Aquatic herbs, with whorled finely dissected leaves, and minute axillary and sessile monœcious flowers without floral envelopes, but with an 8-12-cleft involucre in place of a calyx, the fertile a simple 1-celled ovary, with a suspended orthotropous ovule: seed filled by a highly developed embryo with 4 cotyledons! and a conspicuous plumule. — Consists only of the genus

1. CERATOPHYLLUM, L. HORNWORT.

Sterile flowers of 12-24 stamens, with large sessile anthers. Fruit an acheneum, beaked with the slender persistent style. — Herbs growing under water, in ponds or slow-flowing streams: the sessile leaves cut into thrice-forked thread-like rigid divisions (whence the name from *κέρας*, a horn, and *φύλλον*, leaf).

1. *C. demersum*, L. — Var. *COMMUNE* has a smooth marginless fruit beaked with a long persistent style, and with a short spine or tubercle at the base on each side. — Var. *ECHINATUM* (*C. echinatum*, Gray) has the fruit mostly larger (3" long), rough-pimpled on the sides, the narrowly winged margin spiny-toothed. — Slow streams and ponds: common, but rarely seen in fruit. Probably there is only one species. (Eu.)

ORDER 95. CALLITRICHACEÆ. (WATER-STARWORTS.)

Small annuals or perennials, mostly aquatics, with opposite entire leaves and axillary monœcious flowers without any proper floral envelopes, and with a 4-lobed and 4-celled 4-seeded fruit; — consisting only of the genus Callitriche (regarded by many botanists, perhaps with good reason, as repre-

senting the most reduced form of the Halorageæ, p. 174. The so-called perfect flower is considered to be a staminate and a pistillate, or two staminate and one pistillate naked flowers in the same axil, each of a single stamen or pistil.)

The elaboration of our species is contributed by DR. G. ENGELMANN.

1. CALLÍTRICHE, L. WATER-STARWORT.

Flowers monœcious, solitary or 2 or 3 together in the axil of the same leaf, wholly naked or between a pair of membranaceous bracts. Sterile flower a single stamen: filament bearing a heart-shaped 4-celled anther, which by confluence becomes one-celled, and opens by a single slit. Fertile flower a single 4-celled ovary, either sessile or pedicelled, bearing 2 distinct and filiform sessile, usually persistent stigmas. Ovule solitary in each cell. Fruit nut-like, compressed, 4-lobed, 4-celled, separating at maturity into as many closed 1-seeded portions. Seed anatropous, suspended, filling the cell: embryo slender, straight or slightly curved, in the axis and nearly the length of the oily albumen. — Smooth, or beset with minute stellate scales (visible only under the microscope), with spatulate or linear leaves, both forms often occurring on the same stem. (Name from *καλός*, *beautiful*, and *θρίξ*, *hair*, from the almost capillary and usually tufted stems of the commoner species.)

§ 1. TERRESTRIAL SPECIES. *Small annuals, forming tufts on merely moist soil; destitute of stellate scales and of bracts: leaves uniform, very small, obovate or wedge-shaped, 3-nerved, crowded, provided with stomata: filament not lengthening: carpels connate.*

1. **C. Austini**, Engelm. Fruit small, broader than high, deeply notched above and below, on a pedicel often nearly of its own length; lobes of the fruit narrowly winged and with a deep groove between them, wings denticulate; persistent stigmas shorter than the fruit, spreading or reflexed; leaves obovate. — On damp soil in open woods, fields and roads, New York and New Jersey (*C. F. Austin*) to Illinois, Missouri, Texas, Mexico, and South America. April–June. — Half an inch or an inch high: leaves 1"–2" long: fruit $\frac{1}{4}$ " in diameter.

C. PEPLOIDES, Nutt. and **C. NUTTALLII**, Torr. (*C. pedunculosa*, Nutt.), — the former with subsessile curiously gibbous fruit, the latter with long-peduncled fruit with eversed keels, — are southwestern species of this section.

§ 2. AMPHIBIOUS SPECIES. *Perennials? with elongated stems (occasionally quite terrestrial as in the former, or wholly submersed as in the next section): leaves with stellate scales and stomata, the floating ones obovate and 3-nerved, the submersed linear: flowers usually between a pair of bracts, rarely naked: pollen shed only in the air; the filament elongating afterwards: carpels in fruit connate.*

2. **C. verna**, L. Fruit ($\frac{1}{2}$ " long) higher than broad, obovate, slightly obcordate, usually thicker at the base than upwards, sessile, its lobes sharply keeled or very narrowly winged upwards, and with a wide groove between them; stigmas shorter than the fruit, almost erect, usually deciduous; floating leaves crowded in a tuft, obovate, narrowed into a petiole. — Common in stagnant waters, from Pennsylvania and New Jersey north and northwestward. April–

Aug. — From a few inches to a foot and more in length : upper leaves 3"–5" long ; lower ones twice as long. A terrestrial form with smaller, narrower, and more uniform leaves (*C. brevifolia*, *Pursh*), occurs where the waters recede in summer and fall. (Eu.)

3. *C. heterophylla*, *Pursh*. Fruit smaller, as broad or broader than high, deeply emarginate, thick, almost ventricose, sessile or nearly so, its lobes obtusely angled, with a small groove between them ; stigmas as long as the fruit, erect, persistent ; floating leaves crowded in a tuft, broadly spatulate, often retuse, abruptly narrowed into a long petiole. (*C. Asagrayi*, *Hegelmaier*.) — Stagnant water, New York to Illinois and southward : common. April–Aug. — Similar to the last, of same size, but with smaller leaves (2"–4" long), and fruit scarcely larger than in No. 1, but much thicker. A terrestrial form (which comes also under *C. brevifolia*, *Pursh*) and a submerged one, with linear leaves often an inch long (*C. linearis*, *Pursh*), are not rare.

§ 3. SUBMERSED SPECIES. *Perennials, entirely under water, with crowded and uniform linear 1-nerved leaves, without scales or stomata : flowers bractless, fertilized under water : filament not elongating : carpels separate nearly to the axis.*

4. *C. autumnalis*, *L.* Fruit large, flattened, circular, deeply and narrowly notched, sessile or nearly so, its lobes broadly winged, and with a very deep and narrow groove between them ; stigmas very long, reflexed, deciduous ; leaves all linear from a broader base, retuse or notched at tip. — In the St. Lawrence River, near Ogdensburgh (*G. W. Clinton*), Lake Superior (*C. G. Loring, Jr.*), and northwestward. Aug.–Sept. — Stems 3"–6" high : leaves 2"–5" long : fruit variable in size, usually 1" or more in diameter. (Eu.)

ORDER 96. **PODOSTEMACEÆ.** (RIVER-WEED FAMILY.)

Aquatics, growing on stones in running water, some with the aspect of Sea-weeds or others of Mosses or Liverworts ; the minute naked flowers bursting from a spathe-like involucre as in Liverworts, producing a 2–3-celled many-seeded ribbed pod ; — represented in North America by

1. **PODOSTEMON**, *Michx.* RIVER-WEED.

Flower solitary, pedicelled, from a tubular sac-like involucre, destitute of floral envelopes. Stamens borne on one side of the stalk of the ovary, with their long filaments united into one for more than half their length, and 2 short sterile filaments, one on each side : anthers 2-celled. Stigmas 2, awl-shaped. Pod oval, 8-ribbed, 2-celled, 2-valved. Seeds minute, very numerous on a thick persistent central placenta, destitute of albumen. — Leaves 2-ranked. (Name from *ποῦς*, *foot*, and *στῆμων*, *stamen* ; the two stamens being apparently raised on a stalk by the side of the ovary.)

1. *P. ceratophyllus*, *Michx.* Leaves rigid or horny, dilated into a sheathing base, above mostly forked into thread-like or linear lobes. — Not rare in the bottom of shallow streams. July–Sept. — A small olive-green plant, of firm texture, resembling a Sea-weed, tenaciously attached to loose stones, in the manner of a *Fucus*, by fleshy disks or processes in place of roots.

ORDER 97. **EUPHORBIAEÆ.** (SPURGE FAMILY.)

Plants usually with a milky acrid juice, and monœcious or diœcious flowers, mostly apetalous, sometimes achlamydeous (occasionally polypetalous or monopetalous); the ovary free and usually 3-celled, with a single or sometimes a pair of ovules hanging from the summit of each cell; stigmas or branches of the style as many or twice as many as the cells; fruit commonly a 3-lobed pod, the lobes or carpels separating elastically from a persistent axis and elastically 2-valved; seed anatropous; embryo straight, almost as long as and the flat cotyledons mostly as wide as the fleshy or oily albumen. Stipules often present. — A vast family in the warmer parts of the world (the acrid juice poisonous); most numerously represented in Northern countries by the genus *Euphorbia*, which has very remarkable reduced flowers enclosed in an involucre that imitates a calyx. Our last genus belongs to the Box-Family, which some botanists of late separate from the Euphorbiaceæ, on account of the rhaphe being on the outer or dorsal side of the suspended ovule, &c.

* Seeds and ovules only one in each cell.

— Staminate and pistillate flowers both destitute of calyx as well as corolla, and contained in the same cup-shaped involucre, which imitates a calyx,—the whole liable to be mistaken for a single flower.

1. **Euphorbia.** Involucre surrounding many staminate flowers (each of a single naked stamen) and one pistillate flower (a 3-lobed pistil).

— — Staminate and pistillate flowers both with a calyx, not involucre.

++ Stamens erect in the bud.

2. **Jatropha.** Flowers cymose or paniced. Calyx corolla-like, 5-cleft; the lobes imbricated in the bud. Stamens 10 or more.

3. **Stillingia.** Flowers in a spike, pistillate at the base. Calyx 2-3-parted, the lobes imbricated in the bud. Stamens 2 or 3. Stigmas or branches of the style 3, simple.

4. **Acalypha.** Flowers spiked or glomerate, the pistillate in the axil of bracts. Calyx 3-5-parted; in staminate flowers valvate in the bud. Stamens mostly 8: anthers with 2 separate pendulous cells. Styles or stigmas 3, dissected.

5. **Tragia.** Flowers in racemes, pistillate at the base. Calyx in staminate flowers valvate in the bud. Anther-cells united. Styles united at the base, simple.

++ ++ Stamens inflexed in the bud.

6. **Croton.** Flowers spiked or glomerate. Ovary and fruit 3- (rarely 2-4-) celled.

7. **Crotonopsis.** Flowers scattered on the branchlets. Ovary and fruit 1-celled.

** Seeds and ovules 2 in each cell. Calyx imbricated in the bud.

8. **Phyllanthus.** Flowers axillary. Stamens mostly 3, and usually monadelphous.

9. **Pachysandra.** Flowers spiked. Calyx 4-parted. Stamens 4, separate.

1. **EUPHÓRBIA, L.** SPURGE.

Flowers monœcious, included in a cup-shaped 4-5-lobed involucre (*flower* of older authors) resembling a calyx or corolla, and usually bearing large thick glands (with or without petal-like margins) at its sinuses. Sterile flowers numerous and lining the base of the involucre, each from the axil of a little bract, and consisting merely of a single stamen jointed on a pedicel like the filament: anther-cells globular, separate. Fertile flower solitary in the middle of

the involucre, soon protruded on a long pedicel, consisting of a 3-lobed and 3-celled ovary with no calyx, or a mere vestige. Styles 3, each 2-cleft; the stigmas therefore 6. Pod separating into 3 one-seeded carpels, which split elastically into 2 valves. Seed often caruncled. — Plants (herbs in the United States), with a milky acrid juice. Peduncles terminal, often umbellate-clustered; in the first section mostly appearing lateral, but not really axillary. (Named after *Euphorbus*, physician to King Juba.)

Genus newly elaborated for this work by DR. GEORGE ENGELMANN.

A. APPENDICULATÆ. *Glands of the involucre with petal-like, usually white or rose-colored, and entire or toothed margins or appendages; these almost obsolete in No. 1.*

§ 1. **ANISOPHYLLUM.** *Leaves all similar, opposite, on short petioles, small, oblique at base, furnished with awl-shaped or scaly and often fringed persistent stipules: stems much branched, spreading or usually procumbent: involucre solitary in the forks of the branches or in terminal or pseudo-lateral clusters, small, always with 4 glands: seeds without a caruncle: all our species are annuals, flowering throughout summer and autumn.*

* *Seeds smooth and even, ash-colored: leaves entire, glabrous, as is the whole plant, and pale green or glaucous.*

1. **E. polygonifolia**, L. Prostrate-spreading; leaves oblong-linear, obtuse, mucronate, slightly cordate or obtuse at base (4"–8" long); stipules setaceously divided; peduncles in the forks of the branches, as long as the petioles; lobes of the involucre longer than the minute not appendaged glands; pods obtusely angled; seeds ovate (over 1" long, the largest of this section). — Sandy shores of the Atlantic and of the Great Lakes.

2. **E. Géyeri**, Engelm. Procumbent; leaves oblong-ovate, obtuse, slightly mucronate, mostly acutish at base, lowermost cordate (3"–6" long); stipules setaceously divided; peduncles as long as petioles, at length in loose foliaceous lateral clusters; glands of the involucre with narrow white or red appendages; pods acutely angled; seeds ovate, acute at one end ($\frac{1}{2}$ " long). — Sandy soil, Illinois (*Geyer, Vasey*) to Wisconsin and Minnesota (*T. J. Hale*).

The nearly allied *E. PETALOÏDEA*, Engelm., of Kansas and Nebraska, extends into Western Missouri and Iowa, and may cross the upper Mississippi; it is distinguished by its half-erect spreading growth; longer, narrower, and retuse or emarginate leaves; peduncles longer than petioles; larger involucre, the broadly campanulate appendages much larger and conspicuous; capsule obtusely angled; seeds nearly a line long.

3. **E. sérpens**, H. B. K. Stems filiform, prostrate, and often rooting; leaves round-ovate, obtuse or cordate at base (only $\frac{1}{2}$ "–1 $\frac{1}{2}$ " long); stipules membranaceous, triangular; peduncles much longer than petioles, at length in loose foliaceous lateral clusters; glands of the very small involucre with minute crenulate appendages; pods acutely angled; seeds obtusely angled ($\frac{1}{3}$ " long or less). (*E. herniarioides*, Nutt. and Ed. 2.) — In rich soil, Illinois, especially in the alluvions of the larger rivers, and southwestward: also adventive on ballast sand-banks of the Delaware near Philadelphia. (*I. Martindale, C. F. Parker.*)

* * *Seeds minutely roughened, or transversely wrinkled, or pitted, ash-colored, or (in the last species) blackish: leaves more or less serrulate, smooth or often hairy.*

4. **E. serpyllifolia**, Pers. Glabrous, prostrate-spreading; *leaves obovate-oblong*, narrowed at the very oblique base, sharply serrulate towards the obtuse apex (3'' - 6'' long, often with a red spot); stipules lanceolate, fimbriate; peduncles as long or longer than petioles, at length in loose foliaceous lateral clusters; glands of the small involucre with narrow somewhat toothed appendages; pods sharply angled; *seeds acutely quadrangular, slightly cross-wrinkled* and often pitted (nearly $\frac{3}{4}$ '' long). — Wisconsin and Minnesota, and southwestward.

5. **E. glyptosperma**, Engelm. Glabrous (or very rarely puberulent), erect-spreading; *leaves linear-oblong*, mostly falcate, very unequal at base, slightly serrulate towards the obtuse apex (2'' - 5'' long); stipules lanceolate, setaceously divided; peduncles as long as petioles, in dense foliaceous lateral clusters; glands of the very small involucre with narrow crenulate appendages; pods sharply angled; *seeds sharply 4-angled and with 5 or 6 sharp transverse wrinkles* ($\frac{1}{2}$ '' long). — Wisconsin (*Hale, Lapham*) to Illinois, opposite St. Louis, where is also the pubescent form (*Riehl*), and southwestward.

6. **E. maculata**, L. Prostrate; stems puberulent or hairy; *leaves oblong-linear*, very oblique at base, serrulate upwards, more or less pubescent or sometimes smoothish (4'' - 6'' long), usually with a brown-red spot in the centre; stipules lanceolate, fimbriate; peduncles as long as petioles, in dense foliaceous lateral clusters; glands of the small involucre minute, with narrow slightly crenate (usually red) appendages; pods acutely angled, puberulent; seeds ovate ($\frac{2}{3}$ '' long), sharply 4-angled and with about 4 shallow grooves across the concave sides. (*E. thymifolia, Pursh*, not *L. E. depréssa, Torr.*) — Open places, roadsides, &c.: everywhere.

7. **E. humistrata**, Engelm. Procumbent, puberulent or hairy; *leaves elliptical or obovate*, very oblique at base, serrulate towards the apex, sparsely hairy underneath (4'' - 9'' long, sometimes with a brown spot above); stipules lanceolate, fimbriate; peduncles rather shorter than petioles, in dense scarcely foliaceous lateral clusters; *involucre cleft on the back*, its (red or white) appendages truncate or crenate; pods sharply angled, puberulent; *seeds ovate, obtusely angled, minutely roughened* ($\frac{1}{2}$ '' long). — Rich soil, in the alluvions of the Mississippi and lower Ohio and their tributaries; and also southward.

8. **E. hypericifolia**, L. Smooth or with scattered hairs, ascending or erect (1° - 2° high); leaves oblique at the obtuse or slightly cordate base, ovate-oblong or oblong-linear, sometimes falcate, serrate ($\frac{1}{2}$ '' - 1 $\frac{1}{2}$ '' long), often with a red spot or red margins; stipules triangular; peduncles longer than the petioles, collected in loose leafy cymes at the end of the branches; *appendages of the involucre entire*, larger and white, or smaller and sometimes red; *pod glabrous, obtusely angled; seeds ovate, obtusely angled, wrinkled and tubercled* ($\frac{1}{2}$ '' long), blackish. (This character is from the U. S. plant, *E. Preslii, Gussone* and *Boissier*, the *E. hypericifolia*, var. *communis, Engelm.* The West Indian original *E. hypericifolia, L.* (found also in Florida) has more compact and usually lateral inflorescence, smaller flowers and small reddish seeds.) — Open places, cultivated soil: very common.

§ 2. **PETALOMA.** *Uppermost or floral leaves with conspicuous white petal-like margins, whorled or opposite, the others scattered: erect annuals, with the leaves equal at the base and entire, and with lanceolate deciduous stipules: involucre mostly 5-lobed, collected in an umbel-like inflorescence: no caruncle to the seed.*

9. **E. MARGINATA**, Pursh. Stem stout (2° – 3° high), erect, hairy; leaves sessile, ovate or oblong, acute; umbel with 3 dichotomous rays; glands of the involucre with broad white appendages. — Frequently cultivated in gardens for its showy broadly white-margined floral leaves: native of the plains of Kansas and Nebraska.

§ 3. **TITHYMALOPSIS.** *Only the uppermost or floral leaves whorled or opposite: erect perennials, with entire leaves equal at the base: stipules none: involucre mostly 5-lobed, in the forks of the branches and terminal, in an umbelliform inflorescence: seeds without caruncle.*

10. **E. corollata**, L. Glabrous or sometimes sparingly hairy (2° – 3° high); leaves ovate, lanceolate, or linear, entire, obtuse; umbel 5- (3–7-) forked, and the forks again 2–3- (or rarely 5-) forked; involucre long-peduncled; pod slender-pedicelled, smooth; seeds thick (1" long or more), ash-colored, the surface slightly uneven. — Rich or sandy soil, New York to Wisconsin and southward. July–Oct. — Conspicuous for the showy false lobes of the involucre, which appear like five white petals, the true lobes minute and incurved.

B. EXAPPENDICULATÆ. *Glands of the involucre without petaloid appendages.*

§ 4. **POINSETTIA.** *Involucres in terminal clusters, 4–5-lobed, with few (or often solitary) cup-shaped glands: seed without a caruncle: ours erect annuals, with variable, entire, dentate, or sinuate leaves, all of them or only the upper ones opposite; the uppermost often colored, especially at the base: stipules small and glandular.*

11. **E. dentata**, Michx. Erect or ascending, hairy (1° high); leaves ovate, lanceolate, or linear, petioled, coarsely toothed (1'–2' long), *only the lowest ones alternate*, all others opposite, upper ones often paler at base; involucre almost sessile, with 5 oblong dentate lobes, and one or sometimes more *short-stalked glands*; seeds ovate-globular, slightly tubercled. — Rich soil, Ohio to Illinois and southward. July–Sept.

12. **E. heterophylla**, L. Erect (1° – 3° high), glabrous; *leaves alternate*, petioled, ovate-fiddle-shaped and sinuate-toothed, or lanceolate or linear and entire, often only those of the branches linear; the upper ones usually with a red base; involucre about the length of the peduncle, with 5 ovate incised lobes and a single or few and *almost sessile glands*; seeds nearly globular, tubercled. (*E. cyathophora*, Jacq.) — Slopes and rocky soil, W. Illinois and southward. July–Sept.

§ 5. **TITHYMALUS.** *Involucres in a terminal dichotomous or commonly umbelliform inflorescence, 5- or usually 4-lobed, with as many flat or convex entire or crescent-shaped glands: seeds mostly carunculate: ours ascending or erect, and glabrous (except No. 15) annuals or perennials; with entire or serrulate leaves, and no stipules.*

* *Perennials with entire leaves, all or only the upper ones opposite: involucre long-*

peduncled in a dichotomous inflorescence, mostly with 5 obtuse glands: seeds without caruncle.

3. **E. Ipecacuánhæ**, L. Stems many from a very long perpendicular root, erect or diffusely spreading (5'–10' long), forking from near the base; leaves varying from obovate or oblong to narrowly linear, almost sessile, glabrous; peduncles elongated ($\frac{1}{2}$ '–1' long); pod long-pedicelled, obtusely angled, nearly smooth; seed ovate, white, sparsely marked with impressed dots. — Sandy soil, near the coast, New York to Virginia and southward. May–July.

* * *Perennials or mostly annuals, with serrulate or rarely entire scattered leaves, only the floral leaves in the umbelliform inflorescence whorled or opposite and of different shape: glands of the involucre mostly 4, transversely oval, obtuse.*

+ *Seeds smooth and even: pod warty or rough.*

14. **E. Darlingtonii**, Gray. Tall perennial (2°–4° high); leaves entire, minutely downy beneath; those of the stem lanceolate-oblong from a narrow base; the floral ones oval, very obtuse; the upper roundish-dilated with a truncate base; umbel 5–8-rayed, then simply forked; pod minutely warty; large globular seed with a small caruncle. (*E. nemoralis*, Darl., not of Kitaibel.) — Copses, Penn. and southward along the mountains. July–Sept.

15. **E. PLATYPHYLLA**, L. Erect annual (8'–18' high); upper stem-leaves lanceolate-oblong, acute, cordate at base, minutely serrulate, mostly with scattered hairs beneath; floral ones triangular-ovate, subcordate; umbel 5-rayed; involucre with ciliate lobes and large sessile glands; styles longer than the ovary, united at the base, slightly 2-cleft; pod covered with depressed warts. — Along the Great Lakes and the St. Lawrence to L. Champlain. June–Aug. (Adv. from Eu.)

16. **E. obtusata**, Pursh. Erect annual (1°–2° high); leaves oblong-spatulate, minutely serrulate, smooth, all obtuse; upper ones cordate at base; floral ones ovate, dilated, barely mucronate; umbel once or twice divided into 3 rays, then into 2; involucre with naked lobes and small stipitate glands; styles distinct and longer than the ovary, rect, 2-cleft to the middle; pod beset with long warts. — Damp woods, Virginia to Illinois. May–July.

+ + *Seeds rugose or reticulated: leaves serrulate: annuals.*

17. **E. dictyospérma**, Fischer & Meyer. Stem erect (8'–18' high); leaves oblong- or obovate-spatulate, smooth, all obtuse and obtusely serrate; upper ones cordate at base; floral ones roundish-ovate or obscurely heart-shaped, slightly mucronate; umbels once or twice 3-forked, then 2-forked; involucre with nearly naked lobes and small almost sessile glands; styles shorter than the ovary, spreading or recurved; pod warty; seeds delicately reticulated. (*E. Arkan-sana*, Engelm. & Gray.) — Prairies and roadsides, Kentucky (*Short*), and west and southwestward. May–July.

18. **E. HELIOSCÓPIA**, L. Stems ascending (6'–12' high), stout; leaves all obovate and very rounded or retuse at the end, finely serrate, smooth or a little hairy, those of the stem wedge-shaped; umbel divided into 5 rays, then into 3, or at length simply forked; glands orbicular, stalked; pod smooth and even; seeds with coarse honeycomb-like reticulations. — Waste places, especially eastward and along the Great Lakes. July–Sept. (Nat. from Eu.)

* * * *Perennials or annuals; ours with entire and scattered leaves, only the floral ones in the umbel-like inflorescence whorled or opposite and of different shape: glands of the involucre mostly 4, crescent-shaped or 2-horned.*

— *Seeds smooth and dark-colored: perennials, with running rootstocks.*

19. **E. ÉSULA**, L. Stems clustered (1° high); leaves lanceolate or linear; the floral (yellowish) broadly heart-shaped, mucronate; umbel divided into many rays, then forking; glands short-horned (brown); pods smoothish and granular. — Essex County, Massachusetts, Oakes. June. (Adv. from Eu.)

20. **E. CYPARISSIAS**, L. Stems densely clustered (6' – 10' high); stem-leaves linear, crowded, the floral ones heart-shaped; umbel many-rayed; glands crescent-shaped; pods granular. — Escaped from gardens to roadsides, in a few places in New England. (Adv. from Eu.)

— *Seeds sculptured, ash-colored: pod smooth: annuals or biennials.*

21. **E. PÉPLUS**, L. Erect or ascending (5' – 10' high); leaves petioled, thin, round-obovate, the upper floral ones ovate; umbel 3-rayed, then forking; glands long-horned; lobes of the pod 2-wing-crested on the back; seeds 2-grooved on the inner face, pitted on the back (scarcely over half a line long). — Waste places eastward: not common. July, Aug.

22. **E. commutata**, Engelm. Stems branched from a commonly decumbent base (6' – 12' high); leaves obovate, obtuse; the upper all sessile, the upper floral ones roundish-dilated, broader than long; umbel 3-forked; glands with slender horns; pod obtusely angled, crestless; seeds ovate, pitted all over (a line long). (*E. Ohioica*, Steudel & Hochstetter.) — Along streams and shady slopes, from Virginia towards the mountains to Kentucky, Wisconsin, and westward. May, June. — Leaves often persistent over the winter on sterile shoots, turning red. Larger in all its parts than *E. Peplus*, with which it has been confounded; but the characters of the pod and seeds readily distinguish it.

* * * * *A glabrous annual or biennial, with entire opposite and decussate leaves, an umbelliform inflorescence, and short-horned glands: seeds carunculate.*

23. **E. LÁTHYRIS**, L. Stem stout (2° – 3° high); leaves thick, linear or oblong, the floral ones oblong-ovate and heart-shaped; umbel 4-rayed, then forking. — Sparingly escaped from gardens, where it is common. (Adv. from Eu.)

2. **JÁTROPHA**, L. (CNIDOSCOLUS, Pohl., Ed. 2.)

Flowers monœcious, rarely diœcious, in a terminal open forking cyme; the fertile ones usually in the lower forks. Calyx corolla-like, in the staminate flowers often salver-shaped, 5-lobed; in the pistillate, 5-parted, imbricated or convolute in the bud. Corolla of 5 distinct or apparently united petals, or in our species none. Glands of the disk opposite the calyx-lobes. Stamens 10 – 30, in 2 or more whorls: filaments monadelphous at the base. Ovary mostly 3-celled: styles 3, united below, their summits once or twice forked. Pod 3-celled, 3-seeded, separating into 3 two-valved carpels. Seed carunculate. — Perennial herbaceous or shrubby plants, chiefly tropical, with alternate mostly long-petioled palmately-veined leaves, and stipules. — Our species is of the section **CNIDOSCOLUS**; of plants mostly armed with stinging bristles. (Name said by Linnæus to be formed of *ιατρόν*, a remedy, and *φάγω*, to eat.)

1. **J. ùrens**, L., var. **stimulòsa**, J. Müller. (TREAD-SOFTLY. SPURGE-NETTLE.) Herbaceous, from a long perennial root, branching (6'–2° high); leaves roundish-heart-shaped, 3–5-lobed; the divisions toothed, cut, or even pin-natifid, often discolored; flowers slender; filaments 10, monadelphous only at the woolly base, or the outer set almost distinct. (*J. stimulosa*, *Michx.* *Cnidoscolus stimulosus*, *Ed.* 2.) — Dry sandy soil, Virginia on the coast, and southward. June–Sept.

3. STILLÍNGIA, Garden. STILLINGIA.

Flowers monœcious, aggregated in a terminal spike. Petals and glands of the disk none. Calyx 2–3-cleft or parted; the divisions imbricated in the bud. Stamens 2 or 3: anthers adnate, turned outwards. Style thick: stigmas 3, diverging, simple. Pod 3-celled, 3-lobed, 3-seeded. Seed carunculate. — Smooth upright plants, with the alternate leaves mostly 2-glandular at the base; the fertile flowers few at the base of the dense sterile spike (rarely separate); the bract for each cluster with a gland on each side. (Named for *Dr. B. Stillingfleet.*)

1. **S. sylvática**, L. Herbaceous (1°–3° high); leaves almost sessile, oblong-lanceolate, serrulate; glands of the spike saucer-shaped. — Sandy and dry soil, E. Virginia and southward. June–Sept.

4. ACALÛPHA, L. THREE-SEEDED MERCURY.

Flowers monœcious; the sterile very small, clustered in spikes, with the few or solitary fertile flowers at their base, or sometimes in separate spikes. Calyx of the sterile flowers 4-parted and valvate in the bud; of the fertile 3–5-parted. Corolla none. Stamens 8–16: filaments short, monadelphous at the base: anther-cells separate, long, often worm-shaped, hanging from the apex of the filament. Styles 3, the upper face or stigmas cut-fringed (usually red). Pod separating into 3 globular carpels which split into 2 valves, rarely of only one carpel. — Herbs (ours annuals), or in the tropics often shrubs, with the appearance of Nettles or Amaranths; the leaves alternate, petioled, with stipules. Clusters of sterile flowers with a minute bract; the fertile surrounded by a large and leaf-like cut-lobed persistent bract. (*Ἀκαλήφῃ*, an ancient name of the Nettle.)

* *Fruit smooth or merely pubescent: seeds nearly smooth.*

1. **A. Virginica**, L. Leaves ovate or oblong-ovate, obtusely and sparsely serrate, long-petioled; sterile spike rather few-flowered, mostly shorter than the deeply palmately-cleft fruiting bracts. — Fields and open places: common. July–Sept. — A homely weed, 1°–2° high, smoothish or rather hairy, often turning purplish in autumn. Fertile flowers 1–3 in each axil, along with the small and short-peduncled sterile spike: bracts large and leaf-like, unequally cut into 5–9 lanceolate lobes. — Passes by various forms (belonging to var. *INTERMÈDIA*, *J. Müll.*), into

Var. **grácilens**, with lanceolate or even linear leaves, less toothed and shorter-petioled; the slender sterile spike often 1' long, and much surpassing the less cleft or few-toothed fruiting bracts. (*A. gracilens*, *Ed.* 1 & 2. *A. Virginiana*, var. *graciliscens*, *J. Müll.* *A. monococca*, *Engelm.*, a reduced form with

two cells of the ovary abortive.)—Sandy dry soil, Rhode Island to Illinois, and common southward.

* * *Fruit echinate with soft bristly green projections: seeds rough-wrinkled.*

2. **A. Caroliniana**, Walt., Ell. Leaves thin, ovate-cordate, sharply and closely serrate-toothed, abruptly acuminate, long-petioled; sterile spikes short, axillary; the fertile ones mostly terminal and elongated, its bracts deeply cut into many linear lobes. (*A. ostryæfolia*, *Riddell.*)—New Jersey (Princeton, *Torrey*), Ohio, and southward.

5. TRÀGIA, Plumier. TRAGIA.

Flowers monœcious, in racemes, apetalous. *Ster. Fl.* Calyx 3-5- (chiefly 3-) parted, valvate in the bud. Stamens 2 or 3: filaments short: anther-cells united. *Fert. Fl.* Calyx 3-8-parted, persistent. Style 3-cleft or 3-parted; the branches 3, simple. Pod 3-celled, 3-lobed, bristly, separating into three 2-valved 1-seeded carpels. Seeds not carunculate.—Erect or climbing plants (perennial herbs in U. S.), pubescent or hispid, sometimes stinging, with mostly alternate stipulate leaves; the small-flowered racemes terminal or opposite the leaves; the sterile flowers above, the few fertile at the base, all with small bracts. (Named for the early herbalist *Bock*, latinized *Tragus*.)

1. **T. innócuæ**, Walt. *Erect*, paniculate-branched, *softly hairy-pubescent* (6'-12' high); *leaves* varying from obovate-oblong to narrowly linear, *acute at the base*, obtusely or sinuately few-toothed or lobed, sometimes entire, *short-petioled or sessile*, paler beneath; stamens 2. (*T. ùrens*, *L.*, having been discarded as a false name, the next oldest, and a good one, is adopted, rather than the recent one of *T. discolor*, *Müller.*)—Dry sandy soil, E. Virginia and southward. May—Aug.

2. **T. urticæfólia**, Michx. *Erect or reclining* or slightly twining, hirsute with stinging hairs; *leaves ovate-lanceolate or triangular-lanceolate*, or the lower ovate, *all somewhat cordate or truncate at the base*, coarsely cut-toothed, *short-petioled*.—Virginia (*Pursh*), and common southward.

3. **T. macrocárpa**, Willd. *Twining*, somewhat hirsute, not stinging; *leaves deeply cordate*, ovate, sharply serrate (3' long), all but the uppermost *long-petioled* (pod $\frac{1}{2}$ ' broad). (*T. cordata*, *Michx.*)—Kentucky (*Michaux*), and southward.—Apparently quite distinct.

6. CRÓTON, L. CROTON.

Flowers monœcious, rarely diœcious, mostly in terminal spike-like racemes or spikes. *Ster. Fl.* Calyx 5- (rarely 4-6-) parted; the divisions lightly imbricated or nearly valvate in the bud. Petals usually present, but mostly small or rudimentary, hypogynous, as many as the divisions of the calyx. Glands or lobes of the disk as many as and alternate with the petals. Receptacle usually hairy. Stamens 5 or more: filaments with the anthers inflexed in the bud. *Fert. Fl.* Calyx 5-10-cleft or parted, nearly as in the staminate flowers: but petals none or minute rudiments. Ovary mostly 3-celled, rarely 2-celled, with a single ovule in each cell: styles as many, from once to thrice 2-cleft. Pod 3- (rarely 2-4-) celled, separating into as many 2-valved 1-seeded carpels. Seeds

carunculate. — Stellate-downy, or scurfy, or hairy and glandular plants, mostly strong-scented; the sterile flowers above; the fertile usually at the base of the same spike or cluster. Leaves alternate, or sometimes imperfectly opposite, with or without obvious stipules. (*Κροτών*, the Greek name of the Castor-oil Plant, of this family.)

§ 1. **GEISELÈRIA**, Klotzsch. *Sterile flowers mostly with a 4-parted calyx, as many ovate-lanceolate petals, a 4-rayed disk, and 8 stamens: fertile flowers with a 5-parted calyx, and very minute awl-shaped rudiments of petals; the 3 styles 2-cleft.*

1. **C. glandulosus**, L. Annual, rough-hairy and glandular (1° – 2° high), somewhat umbellately branched; leaves oblong or linear-oblong, obtusely toothed, the base with a saucer-shaped gland on each side; fertile flowers capitate-clustered at the base of the sterile spike, sessile in the forks and terminal. — Open waste places, Virginia, Illinois, and southward, July–Sept.

§ 2. **PILINÓPHYTUM**, Klotzsch. *Sterile flowers with the calyx equally 5-parted, as many glands alternate with the petals, and 10–14 stamens: fertile flowers with a 7–12-parted calyx and without petals; the 3 styles twice or thrice 2-parted.*

2. **C. capitatus**, Michx. Annual, densely soft-woolly and somewhat glandular (1° – 2° high), branched; leaves long-petioled, lance-oblong or elongated-oblong, rounded at the base, entire; petals obovate-lanceolate, densely fimbriate; fertile flowers several, capitate-crowded at the base of the short terminal sterile spike. — Barrens of Illinois, Kentucky, and southward. Pine barrens of New Jersey, *Knieskern!* July–Sept.

§ 3. **GYNAMBLÒSIS**, Torr. (*Engelmannia*, Klotzsch.) *Sterile flowers with an unequally 3–5-parted calyx, and as many petals and scale-like glands; the stamens varying from 3–11: fertile flowers with an equally 5-parted calyx, and with no petals, 5 glands, and a 2–3-celled ovary, crowned with as many sessile 2-parted stigmas.*

3. **C. monanthógynus**, Michx. Annual, whitish-stellate-pubescent and rusty-glandular; stems (1° – 2° high), slender, erect, below often umbellately 3–4-forked, then repeatedly 2–3-forked or alternately branched; leaves oblong-ovate or narrowly oblong, entire, often acutish (6''–12'' long, about twice the length of their petioles); flowers in the forks, the sterile few on the summit of a short and erect peduncle, the fertile few and clustered or mostly solitary on short recurved peduncles; stamens 3–8; ovary 2-celled; fruit often by abortion 1-celled and 1-seeded; the seed broadly oval. (*C. ellipticum*, Nutt. *Engelmannia Nuttalliana*, Klotzsch. *Gynamblosis monanthogyna*, Torr.) — Barrens and dry prairies, from Illinois and Kentucky southward and westward. June–Sept.

(*C. EURTÉGYNUS*, as it may be named, is the related Texan species, — with more silvery down, rounder leaves on longer petioles, 7–12 stamens, more peduncled fertile flowers, and a 3-celled ovary generally ripening 3 oblong-oval seeds, — mentioned by Torrey as a possible variety of this, and taken by Baillon and Müller for *C. ellipticus* of Nuttall.)

7. CROTONÓPSIS, Michx. CROTONOPSIS.

Flowers monœcious, in very small terminal or lateral spikes or clusters, the lower fertile. *Ster. Fl.* Calyx equally 5-parted. Petals 5, spatulate. Stamens 5, opposite the petals: filaments distinct, inflexed in the bud, enlarged at the apex. *Fert. Fl.* Calyx unequally 3-5-parted. Petals none. Glands (petal-like scales) 5, opposite the sepals. Ovary 1-celled, simple, 1-ovuled, bearing a twice or thrice forked style. Fruit dry and indehiscent, small, 1-seeded. — A slender low annual, with alternate or opposite short-petioled linear or elliptical-lanceolate leaves, which are green and smoothish above, but silvery hoary with starry hairs and scurfy with brownish scales underneath, as well as the branches, &c. (Name compounded of *Κρότων*, and *ὄψις*, *appearance*, for a plant with the aspect and general character of Croton.)

1. *C. linearis*, Michx. — Dry sandy soil, New Jersey (*Knieskern*, *C. E. Smith*), Bristol, Pennsylvania (*E. Diffenbaugh*), Illinois, and southward. July-Sept. The form with shorter and broader leaves is *C. elliptica*, *Willd.*, and *C. argentea*, *Pursh.*

8. PHYLLÁNTHUS, L. PHYLLANTHUS.

Flowers monœcious, axillary. Calyx usually 5-6-parted, imbricated in the bud. Petals none. Stamens mostly 3, erect in the bud, often united. Ovules 2 in each cell of the ovary. Pod depressed; each carpel 2-valved, 2-seeded. Seeds not carunculate. — Leaves alternate, 2-ranked, with small stipules. (Name composed of *φύλλον*, *leaf*, and *ἄνθος*, *blossom*, because the flowers in a few species are borne upon leaf-like dilated branches.)

1. *P. Carolinensis*, Walt. Annual, low and slender, branched; leaves obovate or oval, short-petioled; flowers commonly 2 in each axil, almost sessile, one staminate, the other fertile; calyx 6-parted; stamens 3; styles 3, each 2-cleft; glands of the disk in the fertile flowers united in a cup. — Gravelly banks, E. Penn. to Illinois and southward. July-Sept.

9. PACHYSÁNDRA, Michx. PACHYSANDRA.

Flowers monœcious, in naked spikes. Calyx 4-parted. Petals none. *Ster. Fl.* Stamens 4, separate: filaments long-exserted, thick and flat: anthers oblong-linear. *Fert. Fl.* Ovary 3-celled: styles 3, thick, awl-shaped, recurved, stigmatic down their whole length inside. Ovules a pair in each cell, suspended, with the rhaphe dorsal (turned away from the placenta). Pod deeply 3-horned, 3-celled, splitting into 3 at length 2-valved 2-seeded carpels. — Nearly glabrous, low and procumbent, perennial herbs, with matted creeping rootstocks, and alternate, ovate or obovate, coarsely toothed leaves, narrowed at the base into a petiole. Flowers each 1-3-bracted, the upper staminate, a few fertile ones at the base, unpleasantly scented: sepals greenish or purplish: filaments white (the size and thickness of the latter giving the name, from *παχύς*, *thick*, and *ἄνδρα*, used for stamen).

1. *P. procumbens*, Michx. Stems (6'-9' long) bearing several approximate leaves at the summit on slender petioles, and a few many-flowered spikes along the base; the intervening portion naked, or with a few small scales.

(*P. erecta*, Raf., *Baillon*, is the same.) — Woods, mountains of Kentucky, W. Virginia, and southward. March — May.

ORDER 98. EMPÉTRACEÆ. (CROWBERRY FAMILY.)

Low shrubby evergreens, with the foliage, aspect, and compound pollen of Heaths, and the drupaceous fruit of Arctostaphylos, but the divided or laciniate stigmas, &c. of some Euphorbiaceæ: — probably only an apetalous and polygamous or diœcious degenerate form of Ericaceæ, — comprising three genera, two of which occur within the limits of this work, and the third farther south.

1. ÉMPETRUM, Tourn. CROWBERRY.

Flowers polygamous, scattered and solitary in the axils of the leaves (inconspicuous), scaly-bracted. Calyx of 3 spreading and somewhat petal-like sepals. Stamens 3. Style very short: stigma 6-9-rayed. Fruit a berry-like drupe, with 6-9 seed-like nutlets, each containing an erect anatropous seed. Embryo terete, in the axis of copious albumen, with a slender inferior radicle and very small cotyledons. (An ancient name, from *ἐν*, upon, and *πέτρος*, a rock.)

1. *E. nigrum*, L. (BLACK CROWBERRY.) Procumbent and spreading; leaves linear-oblong, scattered; fruit black. — Alpine summits of the mountains of New England and New York, Lake Superior, and northward. (Eu.)

2. CORÈMA, Don. (BROOM-CROWBERRY.)

Flowers diœcious or polygamous, collected in terminal heads, each in the axil of a scaly bract, and with 5 or 6 thin and scarious imbricated bractlets, but no proper calyx. Stamens 3, rarely 4, with long filaments. Style slender, 3- (or rarely 4-5-) cleft: stigmas narrow, often toothed. Drupe small, with 3 (rarely 4-5) nutlets. Seed, &c. as in the last. — Diffusely much-branched little shrubs, with scattered or nearly whorled narrowly linear heath-like leaves. (Name *κόρημα*, a broom, from the bushy aspect.)

1. *C. Conrădii*, Torrey. Diffusely branched, nearly smooth; drupe very small, dry and juiceless when ripe. (*Empetrum*, Torr. *Tuckermănia*, *Klotzsch*. *Oakësia*, *Tuckermann*.) — Sandy pine barrens and dry rocky places, New Jersey; Long Island; Plymouth and Cape Cod, Massachusetts; Bath, and islands of Penobscot Bay, Maine. (Also Newfoundland.) April. — Shrub 6'-9' high: the sterile plant handsome in flower, on account of the tufted purple filaments and brown-purple anthers. (*Gray*, *Chlor. Bor.-Am.* t. 1.)

ORDER 99. URTICĂCEÆ. (NETTLE FAMILY.)

Plants with stipules, and monœcious or diœcious, or rarely (in the Elm Family) perfect flowers, furnished with a regular calyx, free from the 1-celled (rarely 2-celled) ovary which forms a 1-seeded fruit; the embryo in the albumen when there is any, its radicle pointing upwards; stamens as

many as the lobes of the calyx and opposite them, or sometimes fewer. Cotyledons usually broad. Stipules often deciduous. — A large order (far the greater part tropical), comprising the following suborders, viz. : —

SUBORDER I. **ULMACEÆ.** ELM FAMILY.

Flowers perfect or monœciously polygamous. Filaments straight or moderately incurved in the bud. Styles or stigmas 2. Fruit a samara or drupe. Seed suspended. — Trees, with a watery juice (no active or noxious properties), and alternate leaves.

* Fruit dry, winged or crested (a samara): anthers extrorse.

1. **Ulmus.** Flowers sometimes perfect. Ovary 2-ovuled. Fruit 1-celled, winged.
2. **Planera.** Flowers polygamous. Ovary 1-ovuled. Fruit wingless, nut-like.

* * Fruit a drupe: anthers introrse.

3. **Celtis.** Flowers polygamous. Ovary 1-ovuled. Cotyledons curved and crumpled.

SUBORDER II. **ARTOCARPEÆ.** BREAD-FRUIT & FIG FAMILY.

Flowers monœcious or diœcious, crowded in catkin-like spikes or heads; the calyx, &c. becoming fleshy or juicy in fruit, but the 1- (rarely 2-) celled ovary ripening as a dry achenium. Styles or stigmas commonly 2. — Mostly trees or shrubs, with a milky or yellow (acid or poisonous) juice, and alternate (rough or smooth) leaves. — Stamens inflexed in the bud, and elastically spreading when the flower opens, in the Tribe **MOREÆ.**

4. **Morus.** Fertile and sterile flowers in separate spikes. Calyx berry-like in fruit.

SUBORDER III. **URTICEÆ.** NETTLE FAMILY.

Flowers monœcious or diœcious. Filaments transversely wrinkled and inflexed in the bud, straightening or spreading elastically when the flower opens. Style or stigma simple. Ovary always 1-celled, with an erect orthotropous ovule, forming an achenium in fruit. Embryo straight in the axis of albumen. — Herbs (or in the tropics often shrubs or trees), with a watery bland juice, a tough fibrous bark, and opposite or alternate leaves: many are armed with stinging hairs.

* Calyx in the fertile flowers of 2-5 separate or nearly separate sepals.

+ Plant beset with stinging bristles.

5. **Urtica.** Sepals 4 in both sterile and fertile flowers. Achenium straight and erect, enclosed by the 2 inner and larger sepals. Stigma capitate-tufted. Leaves opposite.
6. **Laportea.** Sepals 5 in the sterile flowers, 4 in the fertile, or apparently only 2, the two exterior minute and obscure. Achenium very oblique and bent down, nearly naked. Stigma long and awl-shaped. Leaves alternate.

+ + Plant wholly destitute of stinging bristles.

7. **Pilea.** Sepals 3 or 4, those of the fertile flowers unequal, all or all but one small. Achenium partly naked, straight and erect. Stigma pencil-tufted.

* * Calyx of the fertile flowers tubular or cup-shaped, enclosing the achenium.

8. **Behmeria.** Flowers monœcious, glomerate, the clusters spiked, not involucrate. St. le long and thread-shaped, stigmatic down one side.
9. **Parietaria.** Flowers polygamous, in involucrate-bracted clusters. Stigma tufted.

SUBORDER IV. CANNABINEÆ. (HEMP FAMILY.)

Flowers diœcious; the sterile racemed or paniced; the fertile in clusters or catkins. Filaments short, not inflexed in the bud. Fertile calyx of one sepal, embracing the ovary. Stigmas 2, elongated. Ovary 1-celled, with an erect orthotropous ovule, forming a glandular achenium in fruit. Seed with no albumen. Embryo coiled or bent. — Herbs, with a watery juice, mostly opposite and lobed or divided leaves, and a fibrous inner bark (yielding bitter and narcotic products).

10. **Cannabis**. Fertile flowers spiked-clustered. Anthers drooping. Leaves 5–7-divided.

11. **Humulus**. Fertile flowers in a short spike forming a membranaceous catkin in fruit. Anthers erect. Leaves 3–5-lobed.

1. ÚLMUS, L. ELM.

Calyx bell-shaped, 4–9-cleft. Stamens 4–9, with long and slender filaments. Ovary 2-celled, with a single anatropous ovule suspended from the summit of each cell, rarely 1-celled: styles 2, short, diverging, stigmatic along the inner edge. Fruit (by obliteration) a 1-celled and 1-seeded membranaceous samara, winged all around. Albumen none: embryo straight; the cotyledons large. — Flowers polygamous, purplish or yellowish, in lateral clusters, in our species preceding the leaves, which are strongly straight-veined, short-petioled, and oblique or unequally somewhat heart-shaped at the base. Stipules small, caducous. (The classical Latin name.)

* *Flowers appearing nearly sessile: fruit orbicular, not ciliate: leaves very rough above.*

1. **U. fúlva**, Mich. (SLIPPERY OR RED ELM.) Buds before expansion soft-downy with rusty hairs (large); leaves ovate-oblong, taper-pointed, doubly serrate (4'–8' long, sweet-scented in drying), soft-downy underneath or slightly rough downwards; branchlets downy; calyx-lobes and stamens 7–9; fruit (8''–9'' wide) with the cell pubescent. — Rich soil from W. New England to Lake Superior and southward. March, April. — A small or middle-sized tree, with tough reddish wood, and a very mucilaginous inner bark.

* * *Flowers on slender drooping peduncles or pedicels, which are jointed above the middle: fruit ovate or oval, fringed-ciliate: leaves smooth above, or nearly so.*

2. **U. Americana**, L. (pl. Clayt.), Willd. (AMERICAN OR WHITE ELM.) Buds and branchlets glabrous; branches not corky; leaves obovate-oblong or oval, abruptly pointed, sharply and often doubly serrate (2'–4' long), soft-pubescent beneath, or soon glabrous; flowers in close fascicles; calyx with 7–9 roundish lobes; fruit glabrous except the margins ($\frac{1}{2}$ ' long), its sharp points incurved and closing the notch. — Moist woods, especially along rivers, in rich soil. April. — A large and well-known ornamental tree, variable, usually with spreading branches and drooping branchlets.

3. **U. racemosa**, Thomas. (CORKY WHITE ELM.) Bud-scales downy-ciliate, and somewhat pubescent, as are the young branchlets; branches often with corky ridges; leaves nearly as in the last, but with veins more simple and straight; flowers racemed; fruit much as in the last, but rather larger. — River-banks, W. New England to Wisconsin and southward. April.

4. **U. alata**, Michx. (WHAHOO OR WINGED ELM.) *Bud-scales and branchlets nearly glabrous; branches corky-winged*, at least some of them; leaves downy beneath, ovate-oblong and oblong-lanceolate, acute, thickish, small (1' - 2½' long); calyx-lobes obovate; fruit downy on the face, at least when young. — Virginia to Illinois, and southward. March. — Wood fine-grained, valuable.

2. PLÁNERA, Gmelin. PLANER-TREE.

Flowers monœciously polygamous. Calyx 4 - 5-cleft. Stamens 4 - 5. Ovary ovoid, 1-celled, 1-ovuled, crowned with 2 spreading styles which are stigmatose down the inner side, in fruit becoming coriaceous and nut-like, not winged. Albumen none: embryo straight. — Trees with small leaves, like those of Elms, the flowers appearing with them, in small axillary clusters. (Named for *J. J. Planer*, a German botanist.)

1. **P. aquática**, Gmel. Nearly glabrous; leaves ovate-oblong, small; fruit stalked in the calyx, beset with irregular rough projections. — Wet banks, Kentucky (*Michaux*) and southward. April.

3. CÉLTIS, Tourn. NETTLE-TREE. HACKBERRY.

Flowers monœciously polygamous. Calyx 5 - 6-parted, persistent. Stamens 5 - 6. Ovary 1-celled, with a single suspended ovule: stigmas 2, long and pointed, recurved. Fruit a globular drupe. Embryo curved, nearly enclosing a little gelatinous albumen: cotyledons folded and crumpled. — Leaves pointed, petioled. Stipules caducous. Flowers greenish, axillary, the fertile solitary or in pairs, peduncled, appearing with the leaves; the lower usually staminate only, in little fascicles, or racemose along the base of the branches of the season. (An ancient Greek name for the Lotus; the fruit of the European Nettle-tree supposed to have been the food of the *Lotophagi*.)

1. **C. occidentális**, L. (SUGARBERRY. HACKBERRY.) *Leaves reticulated*, ovate, cordate-ovate and ovate-lanceolate, taper-pointed, usually conspicuously and sharply so, more or less oblique at the base, glabrous, *sharply serrate*, sometimes sparingly so, or soft-pubescent beneath, at least when young; fruit on a peduncle from once to twice the length of the petiole, reddish or yellowish, turning dark purple at maturity, its peduncle once or twice the length of the petiole. (Also *C. Audibertiana*, *Spach.*, &c.) — Woods and river-banks, New England to Wisconsin and southward. April, May. — A small or middle-sized tree, with the aspect of an Elm, bearing sweet and edible fruits as large as bird-cherries, at first obovate, ripe in autumn; the flesh thin. — Var. **PUMILA**. Low and straggling (4° - 10° high); leaves thin when mature, and smooth, slightly acuminate. (*C. pumila*, *Pursh.*) River-banks, on rocks, from Maryland southward. — Var. **CRASSIFOLIA**. A tall or low tree; leaves thicker, usually serrate all round, and with a long tapering point, dull above, pale beneath. (*C. crassifolia*, *Lam.*) — Common southward and westward.

2. **C. Mississipiénsis**, Bosc. *Leaves entire, very long taper-pointed*, rounded at the base, mostly oblique, thin, and smooth; fruit small. (*C. integrifolia*, *Nutt.*) — W. Kentucky, Illinois? and southwestward. — Even this probably runs into the last.

4. MÖRUS, Tourn. MULBERRY.

Flowers monœcious or diœcious; the two kinds in separate axillary and catkin-like spikes. Calyx 4-parted; lobes ovate. Stamens 4: filaments elastically expanding. Ovary 2-celled, one of the cells smaller and disappearing: styles 2, thread-form, stigmatic down the inside. Achenium ovate, compressed, covered by the succulent berry-like calyx, the whole fertile spike thus becoming a thickened oblong and juicy (edible) aggregate fruit.—Trees with milky juice and rounded leaves: sterile spikes rather slender. (*Mopéa*, the ancient name.)

1. **M. rubra**, L. (RED MULBERRY.) *Leaves* heart-ovate, serrate, rough above, downy underneath, pointed (on young shoots often lobed); flowers frequently diœcious; *fruit* dark purple, long.—Rich woods, New England to Illinois and southward. May.—Small tree, ripening its blackberry-like fruit in July.

2. **M. álba**, L. (WHITE MULBERRY.) *Leaves* obliquely heart-ovate, acute, serrate, sometimes lobed, smooth and shining; *fruit* whitish.—Spontaneous near houses: introduced for feeding silk-worms. (Adv. from Eu.)

5. URTICA, Tourn. NETTLE.

Flowers monœcious, or rarely diœcious; clustered, the clusters mostly in racemes, spikes, or loose heads. *Ster. Fl.* Sepals 4. Stamens 4, inserted around the cup-shaped rudiment of a pistil. *Fert. Fl.* Sepals 4, in pairs; the 2 outer smaller and spreading; the 2 inner flat or concave, in fruit membranaceous and enclosing the straight and erect ovate flattened achenium. Stigma sessile, capitate and pencil-tufted.—Herbs armed with stinging hairs. Leaves opposite: stipules in our species distinct. Flowers greenish; in summer. (The classical Latin name; from *uro*, to burn.)

* *Perennials: flower-clusters in branching paniced spikes, often diœcious.*

1. **U. grácilis**, Ait. *Sparingly bristly*, slender (2°–6° high); *leaves* ovate-lanceolate, pointed, serrate, 3–5-nerved from the rounded or scarcely heart-shaped base, almost glabrous, the elongated slender petioles sparingly bristly; spikes slender and loosely paniced. (*U. procera*, Willd.)—Fence-rows and moist ground: common. Stings few.

2. **U. dioica**, L. *Very bristly and stinging* (2°–3° high); *leaves* ovate, heart-shaped, pointed, very deeply serrate, downy underneath as well as the upper part of the stem; *spikes* much branched.—Waste places and roadsides, chiefly eastward. (Nat. from Eu.)

** *Annuals: flower-clusters chiefly axillary and shorter than the petiole, androgynous.*

3. **U. urens**, L. *Leaves* elliptical or ovate, very coarsely and deeply serrate with long spreading teeth, the terminal not longer than the lateral ones; *flower-clusters* 2 in each axil, small and loose.—Waste grounds, near dwellings, eastward: scarce. Plant 8'–12' high, with sparse stings. (Nat. from Eu.)

4. **U. chamædryoides**, Pursh. *Leaves* ovate and mostly heart-shaped, the upper ovate-lanceolate, coarsely serrate-toothed; *flower-clusters* globular, 1–2 in each axil, and spiked at the summit. (*U. gracilis* & *U. verna*, Raf. *U. purpurascens*, Nutt.)—Alluvial shaded soil, Kentucky and southward.—Slender, 6'–30' high, sparsely beset with stings.

6. **LAPORTEA**, Gaudichaud. WOOD-NETTLE.

Flowers monœcious or diœcious, clustered, in loose cymes; the upper widely spreading and chiefly or entirely fertile; the lower mostly sterile. *Ster. Fl.* Sepals and stamens 5, with a rudiment of an ovary. *Fert. Fl.* Calyx of 4 sepals, the two outer or one of them usually minute, and the two inner much larger. Stigma elongated awl-shaped, hairy down one side, persistent. Achenium ovate, flat, extremely oblique, reflexed on the winged or margined pedicel, nearly naked.—Perennial herbs, with stinging hairs, large alternate serrate leaves, and axillary stipules. (Named for *M. Laporte*.)

1. **L. Canadensis**, Gaudichaud. Leaves ovate, pointed, strongly feather-veined (3'–7' long), long-petioled; fertile cymes divergent; stipule single, 2-cleft. (*U. Canadensis* and *U. divaricata*, *L.*)—Moist rich woods. July–Sept.—Stem 2°–3° high.

7. **PÍLEA**, Lindl. RICHWEED. CLEARWEED.

Flowers monœcious or diœcious. *Ster. Fl.* Sepals and stamens 3–4. *Fert. Fl.* Sepals 3, oblong, more or less unequal: a rudiment of a stamen commonly before each in the form of a hooded scale. Stigma sessile, pencil-tufted. Achenium ovate, compressed, erect, partly or nearly naked.—Stingless, mostly glabrous and low herbs, with opposite leaves and united stipules; the staminate flowers often mixed with the fertile. (Named from the shape of the larger sepal of the fertile flower in the original species, like the *pileus*, or felt cap, of the Romans, which partly covers the achenium. In our solitary species the three sepals are nearly equal, small, and not hooded.)

1. **P. pumila**, Gray. (RICHWEED. CLEARWEED.) Low (3'–18' high); stems smooth and shining, pellucid; leaves ovate, coarsely toothed, pointed; 3-ribbed and veiny; flower-clusters much shorter than the petioles; sepals of the fertile flowers lanceolate, scarcely unequal. (*Urtica pumila*, *L.* *Dubrucelia pumila*, *Gaudichaud*. *Adice pumila*, *Raf.*)—Cool and moist shaded places. July–Sept.

8. **BŒHMÈRIA**, Jacq. FALSE NETTLE.

Flowers monœcious or diœcious, clustered; the sterile much as in *Urtica*; the fertile with a tubular or urn-shaped entire or 2–4-toothed calyx enclosing the ovary. Style elongated awl-shaped, stigmatic and papillose down one side. Achenium elliptical, closely invested by the dry and persistent compressed calyx.—No stings. (Named after *G. R. Böhmer*, Professor at Wittenberg in the last century.)

1. **B. cylindrica**, Willd. Perennial, smoothish; stem (1°–3° high) simple; leaves chiefly opposite, oblong-ovate or ovate-lanceolate, pointed, serrate, 3-nerved, long-petioled; stipules distinct; flowers diœcious, or the two kinds intermixed, the small clusters densely aggregated in simple and elongated axillary spikes, the sterile interrupted, the fertile often continuous.—A state with alternate leaves is *B. lateriflora*, *Muhl.*—Moist or shady ground: very common throughout.

9. **PARIETÀRIA**, Tourn. PELLITORY.

Flowers monœciously polygamous; the staminate, pistillate, and perfect intermixed in the same involucre-bracted cymose axillary clusters; the sterile much as in the last; the fertile with a tubular or bell-shaped 4-lobed and nerved calyx, enclosing the ovary and the ovoid achenium. Style slender or none: stigma pencil-tufted. — Homely, diffuse or tufted herbs, not stinging, with alternate entire 3-ribbed leaves, and no stipules. (The ancient Latin name, because growing on old walls.)

1. **P. Pennsylvànica**, Muhl. Low, annual, simple or sparingly branched, minutely downy; leaves oblong-lanceolate, thin, veiny, roughish with opaque dots; flowers shorter than the leaves of the involucre; stigma sessile. — Shaded rocky banks, Vermont to Wisconsin and southward. June — Aug.

10. **CÁNNABIS**, Tourn. HEMP.

Flowers dioecious; the sterile in axillary compound racemes or panicles, with 5 sepals and 5 drooping stamens. Fertile flowers spiked-clustered, 1-bracted: the calyx of a single sepal enlarging at the base and folded round the ovary. Embryo simply curved. — A tall roughish annual, with digitate leaves of 5-7 linear-lanceolate coarsely toothed leaflets, the upper alternate; the inner bark of very tough fibres. (The ancient name, of obscure etymology.)

1. **C. SATIVA**, L. (HEMP.) — Waste and cultivated ground. (Adv. from Eu.)

11. **HÛMULUS**, L. HOP.

Flowers dioecious; the sterile in loose axillary panicles, with 5 sepals and 5 erect stamens. Fertile flowers in short axillary and solitary spikes or catkins: bracts foliaceous, imbricated, each 2-flowered, in fruit forming a sort of membranaceous strobile. Calyx of a single sepal, embracing the ovary. Achenia invested with the enlarged scale-like calyx. Embryo coiled in a flat spiral. — Twining rough perennials, with stems almost prickly downwards, mostly opposite heart-shaped and palmately 3-7-lobed leaves, with persistent ovate stipules between the petioles. (Name thought to be a diminutive of *humus*, moist earth, from the alluvial soil where the Hop spontaneously grows.)

1. **H. Lùpulus**, L. (COMMON HOP.) Leaves mostly 3-5-lobed, and commonly longer than the petioles; bracts, &c., smoothish; the fruiting calyx, achenium, &c., sprinkled with yellow resinous grains, giving the bitterness and aroma of the hop. — Alluvial banks: common northward and westward, where it is clearly indigenous, July. (Eu.)

ORDER 100. **PLATANÀCÆ**. (PLANE-TREE FAMILY.)

Trees, with watery juice, alternate palmately-lobed leaves, sheathing stipules, and monœcious flowers in separate and naked spherical heads, destitute of calyx or corolla; the fruit merely club-shaped 1-seeded nutlets, furnished with bristly down along the base: consists only of the following genus (of uncertain relationship).

1. **PLÁTANUS**, L. PLANE-TREE. BUTTONWOOD.

Sterile flowers of numerous stamens, with club-shaped little scales intermixed: filaments very short. Fertile flowers in separate catkins, consisting of inversely pyramidal ovaries mixed with little scales. Style rather lateral, awl-shaped or thread-like, simple. Nutlets coriaceous, small, tawny-hairy below, containing a single orthotropous pendulous seed. Embryo in the axis of thin albumen. (The ancient name, from *πλατύς*, broad.)

1. **P. occidentális**, L. (AMERICAN PLANE OR SYCAMORE.) Leaves mostly truncate at base, angularly sinuate-lobed or toothed, the short lobes sharp-pointed; fertile heads solitary, hanging on a long peduncle. — Alluvial banks: very common, especially westward. May. — A very large and well-known tree, with a white bark, separating early in thin brittle plates.

ORDER 101. **JUGLANDACEÆ**. (WALNUT FAMILY.)

Trees, with alternate pinnate leaves, and no stipules; flowers monœcious, the sterile in catkins (aments) with an irregular calyx adnate to the bract; the fertile solitary or in a small cluster or spike, with a regular 3-5-lobed calyx adherent to the incompletely 2-4-celled but only 1-ovuled ovary. Fruit a kind of dry drupe, with a crustaceous or bony nut-shell, containing a large 4-lobed orthotropous seed. Albumen none. Cotyledons fleshy and oily, sinuous or corrugated, 2-lobed: radicle short, superior. Petals sometimes present in the fertile flowers. — A small family of important trees, consisting chiefly of the two following genera.

1. **JUGLANS**, L. WALNUT.

Sterile flowers in long and simple lateral catkins from the wood of the preceding year; the calyx adherent to the entire bracts or scales, unequally 3-6-cleft. Stamens 12-40: filaments free, very short. Fertile flowers solitary or several together on a peduncle at the end of the branches, with a 4-toothed calyx, bearing 4 small petals at the sinuses. Styles 2, very short: stigmas 2, somewhat club-shaped and fringed. Fruit with a fibrous-fleshy indehiscent epicarp, and a mostly rough irregularly furrowed endocarp or nut-shell. — Trees, with strong-scented or resinous-aromatic bark, few-scaled or almost naked buds (3 or 4 superposed, and the uppermost far above the axil), odd-pinnate leaves of many serrate leaflets; and the embryo sweet and edible. Pith in plates. (Name contracted from *Jovis glans*, the nut of Jupiter.)

1. **J. cinèrea**, L. (BUTTERNUT.) Leaflets oblong-lanceolate, pointed, rounded at the base, downy, especially underneath, the *petioles and branchlets downy with clammy hairs*; fruit oblong, clammy, pointed, the nut deeply sculptured and rough with ragged ridges, 2-celled at the base. — Rich woods: common. May: fruit ripe in Sept. — Tree 30°-50° high, with gray bark and widely spreading branches; wood lighter brown than in the next.

2. **J. nigra**, L. (BLACK WALNUT.) Leaves ovate-lanceolate, taper-pointed, somewhat heart-shaped or unequal at the base, smooth above, the lower surface

and the *petioles minutely downy*; *fruit spherical*, roughly dotted, the nut corrugated, 4-celled at top and bottom. — Rich woods; rare in the Eastern, very common in the Western States. May: the fruit ripe in Oct. — A large and handsome tree, with brown bark, and valuable purplish-brown wood turning blackish with age. Seed less oily than the butternut, more so than in the European Walnut (*J. RÈGIA*).

2. CÁRYA, Nutt. HICKORY.

Sterile flowers in slender lateral and clustered catkins: calyx naked, adherent to the bract, unequally 2-3-parted. Stamens 3-10: filaments short or none, free. Fertile flowers 2-5 in a cluster or short spike, on a peduncle terminating the shoot of the season: calyx 4-toothed: petals none. Stigmas sessile, 2 or 4, large, papillose, persistent. Fruit with a 4-valved, firm and at length dry exocarp, falling away from the smooth and crustaceous or bony endocarp or nut-shell, which is incompletely 2-celled, and at the base mostly 4-celled. — Fine timber-trees, with hard and very tough wood, and scaly buds, from which in spring are put forth usually both kinds of flowers, the sterile below and the fertile above the leaves. Nuts ripen and fall in October. (*Καρύα*, an ancient name of the Walnut.)

§ 1. *Sterile catkins fuscicled (no common peduncle or sometimes a very short one) from separate lateral scaly buds near the summit of shoots of the preceding year: bud-scales few: fruit elongated-oblong: the thin-shelled nut 2-celled below: seed sweet: leaflets short-stalked, numerous.*

1. *C. olivæfórmis*, Nutt. (PECAN-NUT.) Minutely downy, becoming nearly smooth; leaflets 13-15, oblong-lanceolate, tapering gradually to a slender point, falcate, serrate; nut olive-shaped. — River bottoms, from Illinois southward. — A large tree; its delicious nuts well-known.

§ 2. *Sterile catkins in threes (or rarely more) on a common peduncle from the axil of the inner scales of the common bud, therefore at the base of the shoot of the season, which, then bearing 3 or 4 leaves, is terminated by the fertile flowers: fruit globular or oval: nut 4-celled at base: leaflets sessile or nearly so.*

* *Bud-scales numerous, about 10, successively enwrapping; the inner ones accrescent, becoming thin and membranaceous and rather tardily deciduous: husk of the fruit splitting promptly into 4 thick (but variable in this respect) and when dry hard or woody valves: seed sweet and delicious. (The hickory nuts of the market.)*

2. *C. álba*, Nutt. (SHELL-BARK OR SHAG-BARK HICKORY.) Bark of trunk shaggy, exfoliating in rough strips or plates; inner bud-scales becoming large and conspicuous, persistent till the flowers are fully developed; *leaflets* 5, when young minutely downy beneath, finely serrate, the three upper obovate-lanceolate, the *lower pair much smaller* and oblong-lanceolate, all taper-pointed; fruit globular or depressed; *nut white*, flattish-globular, barely mucronate, the shell thinnish. — Large and handsome tree, furnishing most valuable wood and the principal hickory nuts of the market.

3. *C. microcárpa*, Nutt. (SMALL-FRUITED HICKORY.) Nut as in the preceding, but smaller (7"-9" long), and the husk much thinner; while the foliage resembles that of No. 6; *the leaflets* 5-7, *oblong-lanceolate* glandular under-

neath, and the bark of the trunk is said to be close: an uncertain species. — New York to Penn. and southwestward.

4. **C. sulcàta**, Nutt. (WESTERN SHELL-BARK HICKORY.) Bark, &c. as in No. 1; *leaflets* 7–9, more downy beneath; *fruit* oval or ovate, 4-ribbed above the middle, the husk very thick; *nut* large (1½'–2' long) and usually angular, dull white or yellowish, thick-walled, usually strongly pointed at both ends. — Pennsylvania to Wisconsin and southward. — Seed as sweet as in No. 1. Heart-wood light-colored.

5. **C. tomentosa**, Nutt. (MOCKER-NUT. WHITE-HEART HICKORY.) Bark close, rough, but not shaggy and exfoliating on old trunks; catkins, shoots, and lower surface of the leaves *tomentose* when young, resinous-scented; *leaflets* 7–9, lance-obovate or the lower oblong-lanceolate, pointed; *fruit* globular or ovoid, with a very thick and hard husk; *nut* globular, not compressed, 4-ridged towards the slightly pointed summit, brownish, very thick-shelled, 1' in diameter or smaller. — Dry woods, New England to Virginia, Kentucky and southward. — Wood, &c. as in the last: seed more oily. — A var. **MÁXIMA**, of Nuttall bears "fruit as large as an apple," the husk exceedingly thick.

* * *Bud-scales* numerous or few; *husk* of the fruit thin and rather friable at maturity, 4-valved only to the middle or tardily to near the base: seed more or less bitter: bark of old trunk not exfoliating.

6. **C. porcina**, Nutt. (PIG-NUT or BROOM H.) Bud-scales nearly as in No. 5, but smaller, caducous; shoots, catkins, and leaves *glabrous* or nearly so; *leaflets* 5–7, oblong- or obovate-lanceolate and taper-pointed, serrate; fruit pear-shaped, oblong, or oval; *nut* oblong or oval (1½' to nearly 2' long), with a thick bony-shell; the oily seed at first sweet in taste, then bitterish. (*C. glabra*, Torr. & Ed. 2. *Juglans glabra*, Wang., Muhl., &c., is much the oldest name, but not quite clear in application. The ordinary forms of the present species are *J. ovata* and *J. obcordata*, Wang.) — Woodlands: common. — Wood very tough: heart-wood reddish or dark-colored: bark of trunk rough.

7. **C. amara**, Nutt. (BITTER-NUT or SWAMP H.) Scales of the small yellowish buds about 6, valvate in pairs, caducous in leafing; catkins and young herbage more or less pubescent, soon becoming almost glabrous; *leaflets* 7–11, lanceolate or oblong-lanceolate; fruit globular, narrowly 6-ridged; *nut* globular, short-pointed, white (barely 1' long), thin-walled; seed at first sweet-tasted, soon extremely bitter. — Moist soil: common. — Wood less valued, husk and nut-shell thinner and less hard than in other species: bark of trunk close and smooth.

ORDER 102. CUPULIFERÆ. (OAK FAMILY.)

Trees or shrubs, with alternate and simple straight-veined leaves, very deciduous stipules, and monœcious flowers; the sterile in catkins (aments) (or capitate-clustered in the Beech); the fertile solitary, clustered or spiked, furnished with an involucre which forms a cup or covering to the 1-celled and 1-seeded nut. Ovary more or less 2–7-celled, with 1 or 2 pendulous anatropous ovules in each cell; but all the cells and ovules except one

disappearing in the fruit. Calyx adherent to the ovary, the minute teeth crowning its summit. Seed with no albumen, filled with the embryo: cotyledons thick and fleshy, in many edible: radicle short, superior. Stipules forming the bud-scales. Leaves usually conduplicate in the bud.

Tribe I. QUERCINEÆ. Sterile flowers with a distinct 4-7-lobed calyx, including 5-20 stamens: filaments slender, free, exerted: anthers 2-celled. Fertile flowers one or few enclosed in a cupule consisting of bracts variously consolidated. Ovary imperfectly 3-7-celled, crowned with 3-8 (mostly 6) calyx-teeth, forming a nut (*gland*), in fruit surrounded or enclosed by the indurated scaly or prickly cupule.

* Sterile flowers in slender catkins.

1. **Quercus.** Cupule 1-flowered, scaly, and entire: nut hard and terete.

2. **Castanea.** Cupule 2-4-flowered, forming a prickly hard bur, 2-4-valved when ripe.

** Sterile flowers in a small head.

3. **Fagus.** Cupule 2-flowered, 4-valved, containing 2 sharply triangular nuts.

Tribe II. CARPINEÆ. Sterile flowers destitute of true calyx, consisting of several stamens included under and more or less adnate to a bract: filaments short: anthers 1-celled. Fertile flowers in a short spike, catkin, or head, two together under each fertile bract, each with one or more bractlets, which form a foliaceous or membranaceous involucre to the nut. Ovary imperfectly 2-celled, 2-ovuled. Seed-coat single.

* Bract of staminate flower furnished with a pair of bractlets inside.

4. **Corylus.** Involucre enclosing the large bony nut, leafy-coriaceous.

** Bract of staminate flower simple: nut small, achenium-like.

5. **Ostrya.** Each ovary and nut included in a bladder and closed bag.

6. **Carpinus.** Each nut subtended by an enlarged leafy bractlet.

1. QUÉRCUS, L. OAK.

Sterile flowers in slender and naked catkins: bracts caducous: calyx 2-8-parted or lobed: stamens 5-12: anthers 2-celled. Fertile flowers scattered or somewhat clustered, consisting of a nearly 3-celled and 6-ovuled ovary, with a 3-lobed stigma, enclosed by a scaly bud-like involucre which becomes an indurated cup (*cupule*) around the base of the rounded nut or acorn. Cotyledons remaining underground in germination: radicle very short, included.—Flowers greenish or yellowish. Sterile catkins single or often several from the same lateral scaly bud, filiform and hanging in all our species. (The classical Latin name.) All flower in spring, and shed their nuts in October.

§ 1. ANNUAL-FRUITED; i. e. *acorns perfected in the autumn of the first year, borne therefore on the wood of the season, usually in the axil of the leaves, and often raised on a peduncle; the kernel commonly sweet-tasted: lobes or teeth of the leaves if any not bristle-pointed: abortive ovules persistent under the seed: sterile flowers mostly 4-6-parted and 8-androus.*

* WHITE OAKS. *Leaves lyrate or sinuate-pinnatifid, pale beneath.*

1. **Q. álba, L.** (WHITE OAK.) *Mature leaves smooth, pale or glaucous underneath, bright green above, obovate-oblong, obliquely cut into 3-9 oblong or linear and obtuse mostly entire lobes; cup hemispherical-saucer-shaped, rough or tubercled at maturity, naked, much shorter than the ovoid or oblong acorn.—Rich woods: common.—A valuable large tree. Lobes of the leaves short and broad 3-5, or 5-9 deep and narrow. Acorn about 1' long; kernel edible.*

2. **Q. obtusiloba**, Michx. (POST-OAK. ROUGH OR BOX WHITE-OAK.) *Leaves grayish or yellowish-downy underneath, pale and rough above, thickish, sinuately cut into 5-7 rounded divergent lobes, the upper ones much larger and often 1-3-notched; cup deep saucer-shaped, naked, one third or half the length of the ovoid acorn.* (*Q. stellata*, Willd.) — Sandy or sterile soil: common, especially southward. — A small tree, with very durable wood. Acorn 6''-9'' long.

3. **Q. macrocarpa**, Michx. (BUR-OAK. OVER-CUP OR MOSSY-CUP WHITE-OAK.) *Leaves obovate or oblong, lyrate-pinnatifid or deeply sinuate-lobed, or nearly parted, irregular, downy or pale beneath; the lobes sparingly and obtusely toothed, or the smaller ones entire; cup deep, thick and woody, conspicuously imbricated with hard and thick pointed scales, the upper ones awned, so as usually to make a mossy-fringed border; acorn broadly ovoid (1'-1½' long), half immersed in or entirely enclosed by the cup.* — Rich soil, W. New England to Wisconsin, Kentucky, and southwestward. — A handsome, middle-sized tree. Cup very variable, especially in size, from 9'' to 2' across.

Var. **olivæformis** (*Q. olivæformis*, Michx.) is apparently a mere state of this (figured by Michaux with unripe or imperfect fruit), with narrower and more deeply lobed leaves, and oblong acorns and cups.

* * CHESTNUT-OAKS. *Leaves coarsely sinuate-toothed, but not lobed, except slightly in No. 4, whitish and more or less downy beneath: cup hoary, hemispherical or a little depressed, about half the length of the oblong-ovoid edible acorn.*

4. **Q. bicolor**, Willd. (SWAMP WHITE OAK.) *Leaves obovate or oblong-obovate, wedge-shaped at base, coarsely sinuate-crenate and often rather pinnatifid than toothed, soft-downy and white-hoary underneath, the main primary veins 6-8 pairs, lax and little prominent; fruiting peduncle much longer than the petiole; upper scales of the cup awn-pointed, sometimes forming a mossy-fringed margin; acorn scarcely 1' long.* (*Q. Prinus*, var. *tomentosa*, Michx. *Q. Prinus*, *discolor*, Michx. f. & Ed. 2.) — Low ground: common. A tall tree.

5. **Q. Prinus**, L. (CHESTNUT-OAK.) *Leaves varying, obovate or oblong, with an obtuse or acute base, undulately crenate-toothed, minutely downy beneath, the main primary ribs 10-16 pairs, straight, prominent beneath; fruiting peduncles shorter than the petioles, often very short; cup thick (6''-12'' wide), mostly tuberculate with hard and stout scales; acorn 1' or less in length.* — Dry or moist ground: common southward, scarce northward: a middle-sized or small tree.

Var. **monticola**, Michx. (ROCK CHESTNUT-OAK.) (*Q. montana*, Willd.) Connects with the next variety, but has large acorns; the cup is figured and described as top-shaped: but I have not seen the like when the acorn is well grown. — From Vermont southward along the upper country. A small tree.

Var. **acuminata**, Michx. (YELLOW CHESTNUT-OAK.) (*Q. Castanea*, Muhl. & Ed. 2.) *Leaves slender-petioled, often oblong or even lanceolate, usually acute or pointed, mostly obtuse or roundish at the base, almost equably and rather sharply toothed; cup hemispherical, thin, of small appressed scales, 5''-7'' broad; acorn 7''-9'' long.* — Rich soil, W. New England to Wisconsin: common in the Middle States. — Leaves more like those of Chestnut than any other; the primary veins very straight, impressed above, prominent beneath. A middle-sized tree.

Var. **hùmilis**, Marsh. (DWARF CHESTNUT- or CHINQUAPIN-OAK.) (*Q. pumila*, Michx. *Q. Prinus*, var. *Chincapin*, Michx. *f.*, A. DC. *Q. prinoides*, Willd. & Ed. 2.) Foliage as in one other of the preceding forms; acorns and cups similar, but mostly smaller (abundant, sessile or nearly so); but is a shrub, only 2°–4° high; seemingly therefore a distinct species, but no good character is found. — Poor soil, sandy barrens, &c., S. New England and New York to Wisconsin and southward.

* * * LIVE OAKS. *Leaves coriaceous, evergreen, entire or rarely spiny-toothed.*

6. **Q. vírens**, Ait. (LIVE OAK.) *Leaves small, oblong or elliptical, hoary beneath as well as the branchlets; peduncle usually conspicuous, 1–3-fruited; cup top-shaped; acorn oblong; cotyledons completely united into one mass.* — Coast of Virginia and southward. Farther south becoming a large tree: timber invaluable.

§ 2. BIENNIAL-FRUITED; i. e. *acorns perfected in the autumn of the second year, therefore on old wood below the leaves of the season: peduncles short and thick or none: kernel bitter; the abortive ovules at the apex of the seed: calyx of sterile flowers 2–5-parted: stamens 3–5.*

* *Leaves entire or with a few teeth, or somewhat 3–5-lobed at the summit, coriaceous, inclined to be persistent southward, but none of them really evergreen at the north, the tips or lobes commonly bristle-pointed: acorns globular, small, at most only 6'' long. (Intermediate forms, in certain cases probably hybrids, occur between all these species and some of the next section.)*

← *Leaves not dilated upwards, generally entire: acorn globose.*

7. **Q. cinèrea**, Michx. (UPLAND WILLOW-OAK.) Hoariness and shape of the leaves as in No. 6, but commonly more lance-oblong or lanceolate, and rather more *downy beneath*, and the shallow cups and globular acorns as in the next. — Dry pine-barrens, from E. Virginia southward.

8. **Q. Phéllus**, L. (WILLOW-OAK.) *Leaves linear-lanceolate, narrowed to both ends, soon glabrous, light green (3'–4' long); cup saucer-shaped.* — Sandy low woods, Long Island and New Jersey to Kentucky and southward. — Tree 30°–50° high, remarkable for the willow-like leaves.

9. **Q. imbricària**, Michx. (LAUREL or SHINGLE OAK.) *Leaves lanceolate-oblong, thickish, smooth and shining above, downy underneath, the down commonly persistent; cup between saucer-shaped and top-shaped.* — Barrens and open woodlands, New Jersey to Wisconsin and southward. — Tree 30°–50° high; the wood used for shingles in the Western States, whence the specific name.

← ← *Leaves thick, widening or often much dilated upwards, when they are more or less sinuate or somewhat 3–5-lobed: acorns globular-ovoid.*

10. **Q. aquática**, Catesby. (WATER-OAK.) *Leaves glabrous and shining, obovate-spatulate or narrowly wedge-form, with a long tapering base, and an often obscurely 3-lobed summit, varying to oblanceolate; cup saucer-shaped or hemispherical.* — Wet grounds, around ponds, &c., Maryland to Virginia and southward. — Tree 30°–40° high; running into many varieties, especially southward; the leaves on seedlings and strong shoots often incised or sinuate-pinnatifid; then mostly bristle-pointed.

11. **Q. nigra**, L. (BLACK-JACK or BARREN OAK.) *Leaves broadly wedge-shaped*, but sometimes rounded or obscurely cordate at the base, *widely dilated* and somewhat 3-lobed (rarely 5-lobed) at the summit, occasionally with one or two lateral conspicuously bristle-tipped lobes or teeth, rusty-pubescent beneath, shining above, large (4'–9' long); *cup top-shaped*, coarse-scaly; acorn short-ovoid. (*Q. ferruginea*, Michx.) — Dry sandy barrens, New York to Illinois, and southward. — Tree 8°–25° high.

+ + + *Anomalous or occasional, probably some or all of them hybrid forms, derived wholly or in part from the foregoing species.*

Q. TRIDENTATA, Engelm., arranged by DC. as a var. of *Q. nigra*, is just intermediate between it and *Q. imbricaria*. — Near St. Louis, *Engelmann*.

Q. QUINQUELOBA, Engelm., is intermediate between, and probably derived from *Q. nigra*, and *Q. coccinea*, var. *tinctoria*. — St. Louis, *Engelmann*.

Q. LEANA, Nutt. (LEA'S OAK), by its characters and by the foliage of the second generation, communicated by *Mr. David Christy*, is pretty clearly a hybrid between *Q. imbricaria* and *Q. coccinea*, var. *tinctoria*. — S. Ohio and Illinois: two or three trees known.

Q. HETEROPHYLLA, Michx. (BARTRAM'S OAK), lately rediscovered in Delaware and New Jersey, by *T. Meehan*, *C. E. Smith*, &c. — has lacinate leaves like those of vigorous young shoots of *Q. aquatica*, to which De Candolle refers it as a variety. It is as likely to be a state of *Q. Phellos*, with dilated and toothed or cut leaves.

* * BLACK and RED OAKS. *Leaves pinnatifid or lobed, and slender-petioled, not coriaceous, the tips of the lobes or teeth conspicuously bristle-pointed.*

+ *Mature leaves soft-downy beneath: cup saucer-shaped with a somewhat top-shaped base, about half the length of the fully developed small acorn.*

12. **Q. ilicifolia**, Wang. (BEAR or BLACK SCRUB-OAK.) *Dwarf* (3°–8° high), straggling; *leaves obovate, wedge-shaped at base, angularly about 5- (3-7-) lobed, white-downy beneath*; acorn ovoid, globular, 5''–6'' long. — Sandy barrens and rocky hills, New England to Ohio and Kentucky. (*Q. Banistèri*, Michx.) — Leaves 2'–4' long, thickish, with short and triangular spreading lobes.

13. **Q. falcata**, Michx. (SPANISH OAK.) *Leaves grayish-downy or fulvous underneath*, obtuse or rounded at the base, 3–5-lobed above; *the lobes prolonged, mostly narrow and more or less scythe-shaped*, especially the terminal one, entire or sparingly cut-toothed; acorn globose, 4''–5'' long. — Dry or sandy soil, New Jersey to Illinois and southward. — A small or large tree, extremely variable in foliage; bark excellent for tanning. (*Q. triloba*, Michx., seems to be a confusion of this and *Q. nigra*.)

+ + *Mature leaves glabrous on both sides or nearly so; oval, oblong or somewhat obovate in outline, all except some of No. 14 varying from moderately sinuate-pinnatifid to deeply pinnatifid, turning various shades of red or crimson in late autumn: large trees; the wood reddish, coarse-grained. (Apparently these species all naturally intercross.)*

14. **Q. coccinea**, Wang. (SCARLET OAK.) *Cup top-shaped, or hemispherical with a conical scaly base* (7''–9'' broad), coarsely scaly, covering half or more of the broadly or globular-ovoid acorn. — Leaves in the ordinary forms, at least on

full-grown trees, bright green, shining above, turning red in autumn, deeply pinnatifid, the lobes divergent and sparingly cut-toothed; acorns 6" - 9" long, the kernel and the scar in the cup whitish or yellowish; bark of the trunk gray, the interior reddish. — Moist or dry soil: common.

Var. **tinctòria**. (QUERCITRON, YELLOW-BARKED, OR BLACK OAK. (*Q. tinctoria*, *Bartram.*) Leaves, especially on young trees, often less deeply pinnatifid, sometimes barely sinuate, more membranaceous, commonly retaining some pubescence on the lower surface, turning brownish, orange, or dull red in autumn; bark of trunk darker-colored and rougher on the surface, thicker, and internally orange, much more valuable for the tanner and dyer; cup sometimes less top-shaped, rather hemispherical with a conical base, the scar inside orange-colored, the kernel yellowish. But the shape of the acorn-cup and the character of the bark do not always coincide: and in the figure of the younger Michaux, and in one of the two by the elder, the cup is just that of true *coccinea*. The foliage, in general, approaches that of *Q. rubra*. — Rich and poor soil.

Var. **ambigua**. (GRAY OAK.) (*Q. ambigua*, or *borealis*, *Michx. f.*) Found along our northeastern borders to Lake Champlain and northward, figured and briefly characterized as with the foliage of *Q. rubra* and the fruit of *Q. coccinea*. The acorn in rising more out of the cup, also approaches the former. The Oak of Lake Superior, with "wood better than that of *Q. rubra*" (*Dr. Robbins*) has cup and acorn still more like this last.

A hybrid *Q. COCCINEA-ILICIFOLIA* is found by *Dr. Robbins* at Northbridge, S. Massachusetts.

15. **Q. rubra**, L. (RED OAK.) *Cup saucer-shaped* or flat, with a narrow raised border (9" - 12" in diameter), of rather fine closely appressed scales, sessile or on a very short and abrupt narrow stalk or neck, *very much shorter than the oblong-ovoid or ellipsoidal acorn*, which is 1' or less in length; leaves rather thin, moderately (rarely very deeply) pinnatifid, turning dark red after frost; bark of trunk dark gray, smoothish. — Common both in rich and poor soil. — Timber coarse and poor. In Illinois and southward occurs a form with a deeper cup, more or less conical at base.

Var. **runcinata**, Engelm. Leaves with less deep and more ascending lobes; fruit nearly half smaller; acorn 7" - 9" long by 6" broad; cup with a convex or obscurely top-shaped base: approaches the next therefore in fruit, but not in foliage. St. Louis, in company with *Q. rubra*, *palustris* and *imbriaria*; probably a hybrid.

16. **Q. palustris**, Du Roi. (SWAMP SPANISH, OR PIN OAK.) *Cup flat-saucer-shaped*, sometimes contracted into a short scaly base or stalk; fine-scaled (5" - 7" broad), *very much shorter than the ovoid-globose acorn*, which is 5" - 7" long; leaves deeply pinnatifid with divergent lobes and broad rounded sinuses. — Low grounds: rather common. — A middle-sized tree: timber accounted better than of the last.

2. CASTÀNEA, Tourn. CHESTNUT.

Sterile flowers interruptedly clustered in long and naked cylindrical catkins: calyx mostly 6-parted: stamens 8-20: filaments slender: anthers 2-celled.

Fertile flowers few, usually 3 together in an ovoid scaly prickly involucre: calyx with a 6-lobed border crowning the 3-7-celled 6-14-ovuled ovary: abortive stamens 5-12: stigmas bristle-shaped, as many as the cells of the ovary. Nuts coriaceous, ovoid, enclosed 2-3 together or solitary in the hard and thick very prickly 4-valved involucre. Cotyledons very thick, somewhat plaited, cohering together, remaining underground in germination. — Leaves strongly straight-veined. Flowers appearing later than the (undivided straight-veined) leaves; the catkins axillary near the end of the branches, cream-color; the fertile flowers at the base of the upper ones. (The classical name, from that of a town in Thessaly.)

1. **C. véscá**, L. (CHESTNUT.) *Leaves oblong-lanceolate, pointed, serrate with coarse pointed teeth, when mature smooth and green both sides; nuts 2 or 3 in each involucre, therefore flattened on one or both sides: in the American tree, var. AMERICÁNA, Michx., leaves acute at the base, nuts smaller and sweeter.* — Rocky or hilly woods, Maine to Michigan and Kentucky and southwards, especially along the Alleghanies. June, July. — A large tree, with light coarse-grained wood. (Eu.)

2. **C. pùmila**, Michx. (CHINQUAPIN.) *Leaves oblong, acute, serrate with pointed teeth, whitened-downy underneath; nut solitary, not flattened.* — Sandy woods, from (Long Island?) S. Penn. and Ohio, southward, where it abounds. June. — Shrub or tree 6°-20° high. Involucres small, often spiked; the ovoid pointed nut scarcely half as large as a common chestnut, very sweet.

3. FÁGUS, Tourn. BEECH.

Sterile flowers in small heads on drooping peduncles, with deciduous scale-like bracts: calyx bell-shaped, 5-7-cleft: stamens 8-16: filaments slender: anthers 2-celled. Fertile flowers usually in pairs at the apex of a short peduncle, invested by numerous awl-shaped bractlets, the inner grown together at their bases to form the 4-lobed involucre: calyx-lobes 6, awl-shaped: ovary 3-celled with 2 ovules in each cell: styles 3, thread-like, stigmatic along the inner side. Nuts sharply 3-sided, usually 2 in each urn-shaped and soft-prickly coriaceous involucre, which divides to below the middle into 4 valves. Cotyledons thick, folded and somewhat united; but rising and expanding in germination. Trees, with a close and smooth ash-gray bark, a light horizontal spray, and undivided strongly straight-veined leaves, which are open and convex in the tapering bud, and plaited on the veins. Flowers appearing with the leaves, the yellowish staminate flowers from the lower, the pistillate from the upper axils of the leaves of the season. (The classical Latin name, from *φάγω*, to eat, in allusion to the esculent nuts.)

1. **F. ferrugínea**, Ait. (AMERICAN BEECH.) *Leaves oblong-ovate, taper-pointed, distinctly and often coarsely toothed; petioles and midrib soon nearly naked; prickles of the fruit mostly recurved or spreading.* (*F. ferrugínea* and *F. sylvéstris*, Michx. f.) — Woods: common, especially northward, and along the Alleghanies southward. May. — Leaves longer, thinner, and less shining than in the European Beech, most of the silky hairs usually early deciduous; the very straight veins all running into the salient teeth.

4. **CÓRYLUS**, Tourn. HAZEL-NUT. FILBERT.

Sterile flowers in drooping cylindrical catkins consisting of 8 (half-) stamens with 1-celled anthers, their short filaments and pair of scaly bractlets cohering more or less with the inner face of the bract or scale of the catkin. Fertile flowers several in a scaly bud or ovoid catkin, each a single ovary in the axil of a scale or bract, and accompanied by a pair of lateral bractlets; the ovary tipped with a short limb of the adherent calyx, incompletely 2-celled, with 2 pendulous ovules, one of them sterile: style short: stigmas 2, elongated and slender. Nut ovoid or oblong, bony, each enclosed in a leafy or partly coriaceous cup or involucre, consisting of the two bractlets enlarged and often grown together, lacerated at the border. Cotyledons very thick (but raised to the surface of the soil in germination), edible; the short radicle included. — Shrubs or small trees, with thinnish doubly-toothed leaves, folded lengthwise in the bud, flowering in early spring: sterile catkins single or fascicled from scaly buds of the axils of the preceding year, the fertile terminating early leafy shoots. (The classical name, probably from *kópus*, a *helmet*, from the involucre.)

1. **C. Americana**, Walt. (WILD HAZEL-NUT.) *Leaves roundish-heart-shaped*, pointed; *involucre open above* down to the globose nut, of 2 broad foliaceous cut-toothed almost distinct bracts, their base coriaceous and downy, or with glandular bristles intermixed. — Thickets: common. — Twigs and petioles often glandular-bristly. Nut smaller and thicker-shelled than the European Hazel-nut.

2. **C. rostrata**, Ait. (BEAKED HAZEL-NUT.) *Leaves ovate or ovate-oblong*, somewhat heart-shaped, pointed; *involucre of united bracts*, much prolonged above the ovoid nut into a narrow tubular beak, densely bristly. — Common northward and along the Alleghanies. — Shrub 2°–5° high, with slender and mostly smooth branches.

5. **ÓSTRYA**, Michx. HOP-HORNBEAM. IRON-WOOD.

Sterile flowers in drooping cylindrical catkins, consisting of several stamens in the axil of each bract: filaments short, often forked, or irregularly united, bearing 1-celled (half-) anthers; their tips hairy. Fertile flowers in short catkins; a pair under each deciduous bract, each of an incompletely 2-celled 2-ovuled ovary, crowned with the short bearded border of the adherent calyx, tipped with 2 long-linear stigmas, and enclosed in a tubular bractlet, which in fruit becomes a closed bladderly oblong bag, very much larger than the small and smooth nut; these inflated involucre loosely imbricated to form a sort of strobile, in appearance like that of the Hop. — Slender trees, with very hard wood, brownish furrowed bark, and foliage resembling that of Birch: leaves open and concave in the bud, more or less plaited on the straight veins. Flowers in spring, appearing with the leaves; the sterile catkins 1–3 together from scaly buds at the tip of the branches of the preceding year; the fertile single, terminating short leafy shoots of the season. (The classical name.)

1. **O. Virginica**, Willd. (AMERICAN HOP-HORNBEAM. LEVER-WOOD.) Leaves oblong-ovate, taper-pointed, very sharply doubly serrate, downy beneath, with 11–15 principal veins; buds acute; involucre sacs bristly-hairy at the base. — Rich woods: common. Hop-like fruit full grown in Aug.

6. **CARPINUS**, L. HORNBEAM. IRON-WOOD.

Sterile flowers in drooping cylindrical catkins, consisting of several stamens in the axil of a simple and entire scale-like bract: filaments very short, mostly 2-forked, the forks bearing 1-celled (half-) anthers with hairy tips. Fertile flowers several, spiked in a sort of loose terminal catkin, with small deciduous bracts, each subtending a pair of flowers, as in *Ostrya*; but the involucre-like bractlets are open, enlarged in fruiting and foliaceous, merely subtending the small ovate several-nerved nut. — Trees, or tall shrubs, with a smooth and close gray bark, in this and in the slender buds and straight-veined leaves resembling the Beech; the leaf-buds and the inflorescence as in *Ostrya*. (The ancient Latin name.)

1. **C. Americana**, Michx. (AMERICAN HORNBEAM. BLUE OF WATER BEECH.) Leaves ovate-oblong, pointed, sharply doubly serrate, soon nearly smooth; bractlets 3-lobed, halberd-shaped, sparingly cut-toothed on one side, acute. — Along streams: common. — Tree or shrub 10° – 20° high, with a ridged trunk, and very hard whitish wood; also called IRON-WOOD.

ORDER 103. **MYRICACEÆ**. (SWEET-GALE FAMILY.)

Monœcious or diœcious shrubs, with both kinds of flowers in short scaly catkins, and resinous-dotted often fragrant leaves, — differing from the Birch Family chiefly by the 1-celled ovary with a single erect orthotropous ovule, and the drupe-like nut. Involucre none. — Consists chiefly of the typical genus, from which our Sweet-Fern is not sufficiently distinct.

1. **MYRICA**, L. BAYBERRY. WAX-MYRTLE.

Flowers chiefly diœcious: the sterile in oblong or cylindrical, the fertile in ovoid catkins, from axillary scaly buds; both destitute of calyx and corolla, solitary under a scale-like bract and with a pair of bractlets. Stamens 2 – 8: filaments somewhat united below: anthers 2-celled. Ovary with 2 – 4 scales at its base, and 2 thread-like stigmas. Fruit a small globular nut, or dry drupe, coated with resinous grains or wax. (*Μυρίκη*, the ancient name of the Tamarisk or some other shrub; perhaps from *μυρίζω*, to perfume.)

1. **M. Gale**, L. (SWEET GALE.) Leaves wedge-lanceolate, serrate towards the apex; pale, later than the flowers; sterile catkins closely clustered; nuts in imbricated heads, 2-winged by the two thick ovate scales which coalesce with its base. — Wet borders of ponds, New England to Virginia in the mountains, Penn. to Wisconsin, and northward. April. — Shrub 3° – 5° high. (Eu.)

2. **M. cerifera**, L. (BAYBERRY. WAX-MYRTLE.) Leaves oblong-lanceolate, narrowed at the base, entire or wavy-toothed towards the apex, shining and resinous-dotted both sides, somewhat preceding the flowers; sterile catkins scattered, oblong; scales wedge-shaped at the base; nuts scattered and naked, bony, and incrustated with white wax. — Sandy soil on and near the sea-shore: also on Lake Erie. May. — Shrub 3° – 8° high, with fragrant leaves: the catkins sessile along the last year's branches; the fruits sometimes persistent for 2 or 3 years.

2. **COMPTONIA**, Solander. SWEET-FERN.

Flowers frequently monoëcious; the sterile in cylindrical catkins, with kidney-heart-shaped pointed scale-like bracts, and 3-6 stamens; the fertile in globular catkins, bur-like: ovary surrounded by 8 long linear-awl-shaped scales, persistent around the ovoid-oblong smooth nut: otherwise as in *Myrica*. — Leaves linear-lanceolate, pinnatifid with many rounded lobes, thin, appearing rather later than the flowers. Stipules half heart-shaped. (Named after *Henry Compton*, Bishop of London a century ago, a cultivator and patron of botany.)

1. **C. asplenifolia**, Ait. (*Myrica Comptonia*, *C. DC.*) — Sterile hills, New England to Virginia, Wisconsin, and northward. April, May. — Shrub 10-20 high, with sweet-scented fern-like leaves.

ORDER 104. **BETULACEÆ**. (BIRCH FAMILY.)

Monoëcious trees or shrubs, with alternate simple mostly straight-veined leaves, both kinds of flowers in scaly catkins, 2 or 3 under each bract, and no involucre to the naked 1-celled and 1-seeded often winged small nut, which results from a 2-celled and 2-ovuled ovary. Stipules often early deciduous. Stigmas 2, thread-like. Seed anatropous, suspended: no albumen. Cotyledons flattish, oblong, foliaceous in germination. — Comprises the two genera, Birch and Alder.

1. **BÉTULA**, Tourn. BIRCH.

Sterile flowers 3, and bractlets 2, under each shield-shaped scale or bract of the catkins, consisting each of a calyx of one scale bearing 4 short filaments with 1-celled anthers, or strictly of 2 two-parted filaments, each division bearing an anther-cell. Fertile flowers 2 or 3 under each 3-lobed bract, without bractlets or calyx, each of a naked ovary, becoming a broadly winged and scale-like nutlet or small samara crowned with the two spreading stigmas. — Outer bark usually separable in sheets, that of the branchlets dotted. Twigs and leaves often spicy-aromatic. Foliage mostly thin and light. Buds sessile, scaly. Sterile catkins long and drooping, terminal and lateral, formed in summer, remaining naked through the succeeding winter, and expanding their golden flowers in early spring, with or preceding the leaves: fertile catkins oblong or cylindrical, usually terminating very short 2-leaved early lateral branches of the season. (The ancient Latin name.)

* *Trees, with brown or yellow-gray bark on the trunk, sweet-aromatic as well as the twigs, membranaceous and straight-veined Hornbeam-like leaves with a heart-shaped or rounded base, on short petioles, and sessile very thick fruiting catkins: their about equally 3-cleft scales rather persistent: wing of fruit on each side not broader than the seed-bearing body.*

1. **B. lénta**, L. (CHERRY-B. SWEET OR BLACK BIRCH.) *Bark of trunk dark brown, close (outer layers scarcely laminate), very sweet-aromatic; leaves ovate or oblong-ovate from a more or less heart-shaped base, acuminate, sharply*

and finely doubly serrate all round, when mature shining or bright green above and glabrous except on the veins beneath; *fruiting catkins oblong-cylindrical* (1' - 1½' long), the scales with short and *divergent lobes*. (*B. carpinifolia*, Ehrh., Michx.) — Moist woods, &c. : common northward from New England to Illinois, and along the Alleghany region southward. — Rather large tree, reddish-bronze-colored on the spray, much as in the Garden Cherry : timber rose-colored, fine-grained, valuable for cabinet-work.

2. **B. lutea**, Michx. f. (YELLOW OR GRAY BIRCH.) *Bark of trunk yellowish- or silvery-gray, detaching in very thin filmy layers*, within and the twigs much less aromatic; leaves slightly or not at all heart-shaped and often narrowish towards the base, duller-green above and usually more downy on the veins beneath; *fruiting catkins oblong-ovoid* (1' or less in length, 6'' - 9'' thick), the thinner scales (5'' - 6'' long) twice as large as in No. 1, and with narrower *barely spreading lobes*. (*B. excelsa*, Amer. authors, but not of Ait., Regel, &c. The latter unaccountably fails to distinguish the present from the preceding species.) — Moist woods, New England to L. Superior and northward. — Wood whiter and less valuable : tree not higher than No. 1. Leaves 3' - 5' long.

* * *Trees, with chalky-white bark of the trunk separable in thin sheets, ovate or triangular leaves of firmer texture, on long and slender petioles : fruiting catkins cylindrical, usually hanging on rather slender peduncles ; their scales glabrous, with short diverging lobes, separating freely from the axis : wing of the fruit much broader than its body.*

3. **B. álba**, var. **populifolia**, Spach. (AMERICAN WHITE BIRCH.) *Leaves triangular* (deltoid), *very taper-pointed*, truncate or nearly so at the broad base, *smooth and shining both sides*, except the resinous glands when young. (*B. populifolia*, Ait.) — Common on poor soils, Penn. to Maine, near the coast. — A small and slender graceful tree (15° - 25° high), with bark much less separable into sheets than the next; the mostly very long-pointed leaves on petioles of fully half their length, tremulous as those of an Aspen. (Eu.)

4. **B. papyracea**, Ait. (PAPER OR CANOE BIRCH.) *Leaves ovate, taper-pointed*, heart-shaped or abrupt (or rarely wedge-shaped) at the base, *smooth above, dull underneath*. — Woods, New England to Penn. and Wisconsin, almost entirely northward, and extending far northwest. — A large tree, with fine-grained wood, and very tough durable bark splitting freely into paper-like layers. Leaves dark-green above, pale, glandular-dotted, and a little hairy on the veins underneath, sharply and unequally doubly serrate, 3 - 4 times the length of the petiole. — Var. **MINOR**, *Tuckerman*, in the alpine region of White Mountains, is a dwarf form, approaching the var. *occidentalis* of N. W. Amer. and *B. Davurica*. In this country no transitions are seen between our White and the Paper Birch. (The original *B. excelsa*, Ait., and of Regel, seems likely to belong here, or to have been mixed up with the next.)

* * * *Tree, with greenish-brown bark, somewhat laminate, and reddish twigs, ovate leaves whitish beneath, and soft-downy peduncled fruiting catkins.*

5. **B. nigra**, L. (RIVER OR RED BIRCH.) Leaves rhombic-ovate, acutish at both ends, irregularly doubly serrate, whitish and (until old) downy underneath; petioles and peduncle of nearly the same length (3'' - 7'') and with

the oblong catkin tomentose; the bracts with oblong-linear nearly equal lobes; fruit broadly winged. (*B. rubra*, *Michx. f.*) — Low river-banks, E. Massachusetts to Illinois and southward. — A rather large tree, with light-colored wood, and somewhat Alder-like leaves.

* * * * *Shrubs, with brownish bark, rounded or wedge-shaped crenate and mostly small leaves of thickish or coriaceous texture, and oblong or cylindrical glabrous and mostly erect catkins, on short peduncles.*

6. **B. pumila**, L. (LOW BIRCH.) Stems (2°–8° high) erect or ascending, not glandular; young branches and lower face of young leaves mostly soft-downy; leaves obovate, roundish, or orbicular (6"–16" long), pale beneath, veinlets on both faces finely reticulated; wing of the fruit mostly narrower than the body. — Bogs, Conn. (Canaan, *W. H. Leggett*) to Michigan and northward. — Leaves in one form resiniferous or glandular-dotted, usually not at all so. — **B. GRAYII**, Regel, recently characterized on specimens of a shrub introduced from Central Ohio into the Cambridge Botanic Garden, since lost, appears to be only a marked variety or luxuriant form of the present species, with shoots and young leaves beneath more tomentose, and wings of fruit (which are as wide as body in one Michigan specimen of *B. pumila*) here almost twice as wide!

7. **B. glandulosa**, *Michx.* (DWARF BIRCH.) Stems erect or mostly spreading (1°–4° high), or when alpine procumbent; branchlets glabrous, conspicuously dotted with resinous wart-like glands; leaves roundish wedge-obovate or sometimes orbicular (6"–9" long), green both sides, less reticulated; fruiting catkins mostly shorter and oblong or oval; wing of the fruit narrower than or sometimes equalling the body. (*B. nana*, *Ed. 2*, not of *L.* A round-leaved alpine form is *B. rotundifolia*, *Spach.*, and *B. Littelliana*, *Tuckerman.*) — High mountains, Northern New England and New York, and shore of Lake Superior and northward. — The resinous-glandular branches chiefly distinguish some of the larger forms from *B. pumila*, and the small alpine ones from *B. nana*, *L.* of Europe: probably they run together.

2. **ÁLNUS**, *Tourn.* ALDER.

Sterile catkins elongated and drooping, with 4 or 5 bractlets and 3 (rarely 6) flowers under each short-stalked shield-shaped scale; each flower usually with a 3–5-parted calyx and as many stamens: filaments short and simple: anthers 2-celled. Fertile catkins ovoid or oblong; the fleshy scales each 2–3-flowered, with a calyx of four little scales adherent to the scales or bracts of the catkin, which are thick and woody in fruit, wedge-obovate, truncate, or 3–5-lobed, and persistent. — Shrubs or small trees, with few-scaled leaf-buds, and solitary or often racemose-clustered catkins, terminating leafless branchlets or peduncles. (The ancient Latin name.)

§ 1. *Flowers developed in spring with the leaves; the sterile from catkins which have remained naked over winter; while the fertile have been enclosed in a scaly bud: fruit with a conspicuous thin wing, as in Birch.*

1. **A. viridis**, DC. (GREEN or MOUNTAIN ALDER.) Leaves round-oval, ovate, or slightly heart-shaped, glutinous and smooth or softly downy underneath, serrate with very sharp and closely set teeth, on young shoots often

cut-toothed; fertile catkins slender-stalked, clustered, ovoid. (*A. undulata*, Willd. *Betula crispa*, Michx.) — On mountains and along streams descending from them, N. New England and New York, shore of L. Superior, and northward. Also in the Alleghanies southward. Shrub 3°–8° high. (Eu.)

§ 2. *Flowers developed in earliest spring, before the leaves, from mostly clustered catkins which (of both sorts) were formed the foregoing summer and have remained naked over winter; fruit wingless or with a narrow coriaceous margin.*

2. *A. incana*, Willd. (SPECKLED OR HOARY A.) *Leaves broadly oval or ovate, rounded at the base, sharply serrate, often coarsely toothed, whitened and mostly downy underneath; stipules oblong-lanceolate; fruit orbicular.* (*A. glauca*, Michx.) — Shrub or small tree 8°–20° high, forming thickets along streams: the common Alder northward. — Var. *GLAUCA* has the leaves pale, but when old quite smooth, beneath. (Eu.)

3. *A. serrulata*, Ait. (SMOOTH A.) *Leaves obovate, acute at the base, sharply serrate with minute teeth, thickish, green both sides, smooth or often downy beneath; stipules oval; fruit ovate.* — Shrub 6°–12° high: the common Alder from S. New England to Wisconsin, Kentucky, and southward.

§ 3. *Flowers in autumn (Sept.) from catkins of the season; the fertile mostly solitary in the axils of the leaves, ripening the fruit a year later: fruit wingless.*

4. *A. maritima*, Muhl., Nutt. Sylv. t. 10. (SEA-SIDE A.) *Glabrous; leaves oblong, ovate, or obovate with a wedge-shaped base, slender-petioled, sharply serrulate, bright green, or rather rusty beneath; fruiting catkins large, ovoid or oblong (9"–12" long, 6" thick).* (*A. oblongata*, Regel, not of Willd. *A. Japonica*, Siebold & Zuccarini, according to Regel.) — Along streams, Delaware and E. Maryland, Dr. Pickering, W. M. Canby, &c. Also, what is thought to be the same species in Japan! — Tree 20° high.

ORDER 105. SALICACEÆ. (WILLOW FAMILY.)*

Diocious trees or shrubs, with both kinds of flowers in catkins, one under each bract, entirely destitute of floral envelopes (unless one or two gland-like bodies represent the calyx); the fruit a 1-celled and 2-valved pod, with 2 parietal or basal placentæ, bearing numerous seeds furnished with a long silky down. — Style short or none: stigmas 2, often 2-lobed. Seeds ascending, anatropous, without albumen. Cotyledons flattened. — Leaves alternate, undivided, with scale-like and deciduous, or else leaf-like and persistent, stipules. Wood soft and light: bark bitter.

1. SÀLIX, Tourn. WILLOW. OSIER.

Bracts (*scales*) of the catkins entire. Sterile flowers of 3–10, mostly 2, distinct or united stamens, accompanied by 1 or 2 little glands. Fertile flowers also with a small flat gland at the base of the ovary on the inner side: stig-

* This order was elaborated for the first edition by JOHN CAREY, Esq.; whose account is essentially preserved, pending the publication of Professor Andersson's monograph in the forthcoming volume of DeCandolle's Prodrromus.

mas short. — Trees or shrubs, generally growing along streams, with terete and lithe branches. Leaves mostly long and pointed, entire or glandularly toothed. Buds covered by a single scale, with an inner adherent membrane (separating in § 2). Catkins appearing before or with the leaves. (The classical name, said to be derived from the Celtic *sal*, near, and *lis*, water.)

§ 1. *Catkins lateral and sessile, appearing before the leaves in April or May: stamens 2: scales dark red or brown becoming black, more or less hairy, persistent.*

* *Ovary stalked, downy, hairy, or woolly.*

+ *Catkins ovoid or cylindrical: leaves entire or obscurely wavy-toothed, hairy or woolly, with prominent veins and more or less revolute margins. Shrubs.*

1. **S. cándida**, Willd. (HOARY WILLOW.) Leaves narrowly lanceolate, taper-pointed, or the lowest obtuse, the upper surface and young branches covered with a thin web-like wool more white and dense beneath; stipules small, lanceolate, toothed, about the length of the petioles; catkins cylindrical, closely flowered; the fertile $1\frac{1}{2}'$ – $2'$ long at maturity; ovary densely woolly; style distinct; stigmas 2-cleft; scales oblong, obtuse. (*S. incána*, Michx., not of Schrank.) — New York and New Jersey to Wisconsin and northward; in bogs. — Stems 2° – 5° high: reddish twigs smooth and shining at maturity. The whole shrub very white in exposed situations, greener in shade.

2. **S. tristis**, Ait. (DWARF GRAY W.) Leaves almost sessile, wedge-lanceolate, pointed, or the lower obtuse, grayish-woolly on both sides, the upper side becoming nearly smooth at maturity; stipules minute, hairy, very early deciduous; catkins small and very short, globular when young, loosely flowered; ovary with a long tapering beak, clothed with silvery hairs; style short; stigmas 2-lobed. — Bogs: common. — Shrub 1° – $1\frac{1}{2}^{\circ}$ high, much branched: leaves thick, $1\frac{1}{2}'$ long. Stipules seldom seen, often reduced to a mere gland. — Var. **MICROPHÝLLA** has very small and rigid contorted leaves.

3. **S. hùmilis**, Marshall. (PRAIRIE W.) Leaves petioled, lanceolate or obovate-lanceolate, acute or obtuse with an abrupt point, slightly downy above, more thickly so or sometimes grayish-woolly beneath; stipules small, semi-ovate and entire, or larger and lunate with 2–4 teeth, shorter than the petioles; catkins ovoid or oblong, small, often recurved; ovary hairy; style distinct; stigmas 2-cleft. (*S. Muhlenbergiana*, Barratt. *S. confera*, Muhl.) — Dry plains and barrens: common. — Shrub 3° – 8° high, varying much; the small forms sometimes scarcely distinguishable from No. 2; but leaves longer, thinner, and generally stipulate: the larger forms, with leaves $3'$ – $5'$ long and $\frac{3}{4}'$ – $1'$ broad, resemble those of the two next species, but retain more or less down beneath at maturity. — The species of this and the following section often bear cones on the ends of the branches, formed of closely imbricated leaves, probably occasioned by the puncture of insects.

+ + *Catkins cylindrical, large, clothed with very long glossy hairs: leaves more or less serrate, smooth and shining above, glaucous beneath and at length smooth. Shrubs or small trees.*

4. **S. discolor**, Muhl. (GLAUCOUS W.) Leaves lanceolate, ovate-lanceolate, or oblong, acute, irregularly toothed on the sides, entire at the base and apex; stipules lunate, toothed; catkins erect; scales very hairy, oblanceolate;

ovary densely silky.—Low meadows and river-banks: common.—A large shrub or small tree, 8°–15° high. Young leaves commonly obtuse and pubescent, at length becoming smooth and whitish-glaucous beneath, Stipules on vigorous shoots equalling the petiole, often inconspicuous. Young catkins 1½' long, glossy, blackish with the conspicuous scales, elongating in fruit to 2½'.—*S. eriocéphala*, Michx., admitted in former editions, is of this species.

* * *Ovary stalked, silky-hoary and shining: catkins with a few small leaf-like bracts at their base; leaves finely and evenly serrate, silky-gray or glaucous beneath, drying dark: stipules varying from linear to semilunar, toothed, very deciduous. Shrubs, 3°–10° high.*

5. **S. sericea**, Marshall. (SILKY W.) *Leaves lanceolate, pointed, downy above, grayish underneath with short silky hairs; sterile catkins small, globular; the fertile cylindrical, closely flowered; scales obtuse, round-obovate, as long as the stalk of the densely-silky ovoid ovary; stigma sessile.* (*S. grisea*, Willd.)—Sandy river-banks: common. Fertile catkins, at length 1' long; the pods not spreading or elongating in fruit, thus appearing sessile.

6. **S. petiolaris**, Smith. (PETIOLED W.) *Leaves lanceolate, pointed, smooth above, slightly silky beneath when young, at length smooth and glaucous; fertile catkins ovoid-cylindrical; scales acute, very hairy, scarcely as long as the stalk of the silky tapering ovary; style distinct; stigma 2-cleft.* (*S. rosmarinifolia*, and *S. fuscata*, Pursh.?)—Same situations as the last, which this resembles; but the mature leaves are not silky beneath, nor so blackish in drying; the scales not so dark, and clothed with longer white hair; fertile catkins shorter and broader; the pods spreading and showing the stalks.

* * * *Ovary sessile or almost so: downy: catkins bracted at the base. Small trees.*
+ *Filaments and often the reddish anthers united, so as to appear as one.*

7. **S. PURPUREA**, L. (PURPLE W.) *Leaves oblanceolate, pointed, smooth, minutely and sparingly toothed; catkins cylindrical; scales round and concave, very black; stigmas nearly sessile; ovary sessile.* (*S. Lambertiana*, Pursh.)—Low grounds.—Twigs olive-colored or reddish. (Adv. from Eu.)

+ + *Filaments separate.*

8. **S. VIMINALIS**, L. (BASKET OSIER.) *Leaves linear-lanceolate, very long, (3'–6'), taper-pointed, entire or obscurely crenate, lustrous white and satiny beneath; sterile catkins oblong, the fertile cylindrical, silky-hairy; style elongated; stigmas linear, mostly entire.*—Wet meadows.—Considered the best species for basket-work.—*S. Smithiana*, Willd. (thought to be a hybrid of this with some other species), differing principally in the somewhat broader leaves, is also occasionally met with. (Adv. from Eu.)

§ 2. *Catkins lateral, with 4–5 leafy bracts at the base, appearing with or before the leaves in May or June: inner membrane of the scales of flowering buds separating from the cartilaginous exterior, sometimes raised on the apex of the bursting catkins: ovary slender-stalked, smooth (under a lens granular, with occasionally a few short hairs at the base): stamens 2: scales dark or black, hairy, persistent.*

9. **S. cordata**, Muhl. (HEART-LEAVED W.) *Leaves lanceolate or ovate-lanceolate, heart-shaped, truncate, or even acute at base, taper-pointed, sharply*

toothed, smooth, paler and with veinlets reticulated beneath; stipules kidney-shaped or ovate, toothed, often large and conspicuous, of the length of the (when young downy) petiole, or sometimes small and almost entire; catkins appearing with the leaves, leafy at base, cylindrical, the fertile elongating in fruit; ovary lanceolate, tapering to the summit. — Var. *RIGIDA* has the leaves large and rigid, with coarser teeth, of which the lowest are somewhat elongated. (*S. rigida*, *Muhl.* *S. Torreyana*, *Barratt*, which has leaves of a deeper green beneath, appears to belong here.) — Var. *MYRICOIDES* has narrower leaves, neither heart-shaped nor truncate at the base. (*S. myricoides*, *Muhl.*) — Var. *ANGUSTATA* has lanceolate tapering finely serrate leaves, acute at the base, and a more slender style. (*S. angustata*, *Pursh*, & *Ed. 2.*) — Low or inundated places: common. — Shrub or small tree, with leaves $2\frac{1}{2}$ ' - 6' long.

§ 3. *Catkins lateral, with a few leafy bracts at the base, appearing with the leaves in May or June: ovary stalked, silky-hoary: stamens 2: scales persistent.*

10. *S. livida*, *Wahl.*, var. *occidentalis*. (*LIVID W.*) *Leaves oblong or obovate-lanceolate, acute, obscurely toothed, downy above, prominently veiny, softly hairy and glaucous beneath; stipules semilunar, toothed; catkins cylindrical, the fertile becoming loose in fruit; ovary almost linear; style scarcely any; pods slender, tapering from near the base, their very slender stalk longer than the narrow scale. (S. rostrata, Richardson, & Ed. 2. S. vagans, cinerascens, var. occidentalis, Anders.) — Moist or dry ground, New England to Penn., Illinois, and northward: common. — Shrub 3° - 15° high. A transformation of the anthers into imperfect ovaries is frequently observable in this species.*

11. *S. chlorophylla*, *Anders.* (*GREEN W.*) *Glabrous, except the catkins; leaves oblong-lanceolate or oblong, mostly entire, obtuse or acutish at both ends (1' - 2' long), shining above, pale or glaucous beneath; fertile catkins dense, short-cylindrical; ovary ovoid-conical; pod silky, very short-stalked; style slender; scale dark-colored, villous. (S. phycifolia, Ed. 2, not of L.) — Moist ravines, on the alpine summits of the White Mountains, New Hampshire (Oakes, Tuckerman), and northward. — A low spreading shrub, with leaves of a coriaceous texture when old.*

§ 4. *Catkins peduncled (long and loose), borne on the summit of lateral leafy branches of the season, appearing in May and June: scales greenish-yellow, more or less hairy, falling before the pods are ripe: filaments hairy below. Shrubs and trees, with the branches very brittle at the base.*

* *Stamens 3 - 6 or sometimes more: ovary stalked, glabrous.*

12. *S. lucida*, *Muhl.* (*SHINING W.*) *Leaves ovate-oblong or lanceolate and narrow, usually with a long tapering point, smooth and shining on both sides, serrate; stipules oblong and toothed; stamens commonly 5, rarely 7 - 10; pods tapering. — Along streams: rather common. — A beautiful species, sometimes flowering at the height of 3°, sometimes becoming a small bushy tree of 12° - 15°. Probably the same as *S. pentandra*, *L.*, of *Eu.**

13. *S. nigra*, *Marsh.* (*BLACK W.*) *Leaves narrowly lanceolate, pointed and tapering at each end, serrate, smooth (except on the petioles and midrib) and green on both sides; stipules small, deciduous; scales short and rounded, woolly; glands large, of the sterile flowers cleft, of the sterile kidney-shaped;*

stamens 3-6; pods mostly short-ovate. (*S. ambigua*, *Pursh.*)—Var. *FALCATA* has the leaves elongated, scythe-shaped, and the stipules large, broadly lunate, reflexed. (*S. falcata*, *Pursh.* *S. Purshiana*, *Spreng.* *S. ligustrina*, *Michx. f.*)—Tree 15°-25° high, with a rough black bark: frequent along streams, especially southward.

* * *Stamens mostly 2: ovary very short-stalked or nearly sessile, glabrous.*

14. **S. FRÁGILIS**, L. (BRITTLE W.) *Leaves lanceolate, taper-pointed, smooth, glaucous beneath* (slightly silky when young), serrate with inflexed teeth; stipules half-heart-shaped; pods tapering-conical.—Var. *DECIPiens* has dark brown buds, and the lowest leaves on the branches broadly obovate, very obtuse. (*S. decipiens*, *Hoffm.*)—Var. *RUSSELLIANA* has the leaves long and bright, strongly and very sharply serrate; the younger ones and upper branches of the annual shoots silky-downy towards autumn; stipules large and taper-pointed. (*S. Russelliana*, *Smith.*)—Var. *VÍRIDIS*, has long and acute flexuous erect catkins, tough pendulous branchlets, and firmer bright-green leaves. (*S. viridis*, *Fries.*)—A tall and handsome tree, with smooth polished branches; cultivated for basket-work. (Adv. from Eu.)

15. **S. ÁLBA**, L. (WHITE W.) *Leaves lanceolate or elliptic-lanceolate, pointed, toothed, clothed more or less with white and silky hairs, especially beneath*; stipules lanceolate; stigmas nearly sessile, thick and recurved.—Var. *VITELLINA* has yellow branches; leaves shorter and broader. (*S. vitellina*, *Smith & Borrer.*)—Var. *CÆRÛLEA* has the leaves nearly smooth at maturity, greenish or brownish, and greatly resembles the preceding species. (*S. cærulea*, *Smith.*)—A familiar tree, of rapid growth, attaining a height of 50°-80°. (Adv. from Eu.)

16. **S. BABYLÓNICA**, Tourn. (WEEPING WILLOW), belongs to this section, and is planted for ornament. Only the fertile plant is known in the United States.—There is also a remarkable form of it with curled or annular leaves (*S. annularis*, *Forbes*), well known in gardens as the RING-LEAVED or HOOP WILLOW.

* * * *Stamens 2: ovary stalked, mostly silky or downy; the scale narrow.*

17. **S. longifolia**, Muhl. (LONG-LEAVED W.) *Leaves linear-lanceolate, very long, tapering at each end, nearly sessile, remotely denticulate with projecting teeth, clothed with gray hairs when young*; stipules small, lanceolate, toothed; gland long, in the sterile flowers sometimes deeply 2-3-cleft; in the fertile longer than the short stalk of the ovary; stigmas very large, sessile.—River-banks, rooting extensively in sand or gravel: common, especially westward. Varying greatly, 2°-20° high.

§ 5. *Catkins peduncled, borne on the lateral (or sometimes the terminal) leafy branches of the season, appearing in June: stipules deciduous or none: scales persistent.—Small shrubs, mostly with underground spreading stems, sending up short erect or prostrate branches.*

* *Upright (1°-3° high), not alpine: pod long-stalked: stamens 2.*

18. **S. myrtilloides**, L. (MYRTLE W.) *Very glabrous, except the scales of the catkin; leaves elliptical, oblong, or linear-oblong, entire (1'-2' long) reticulated, pale or glaucous beneath, somewhat coriaceous; catkins ob-*

long, the fertile loose in fruit; stalk of the ovary slender, longer than the greenish-yellow scale. (*S. pedicellaris*, *Pursh.*, more luxuriant than the European plant.) — Cold peat-bogs, New England to Wisconsin and northward. (Eu.)

* * *Prostrate or spreading and matted, alpine.*

19. **S. Cútléri**, Tuckerman. (CUTLER'S W.) *Leaves* elliptical and pointed, or obovate and obtuse, tapering at the base, slightly toothed, *strongly veiny, smooth and shining above*, pale and rather glaucous beneath; catkins mostly lateral, oblong-cylindrical; *ovary smooth, short-stalked*; style distinct; *stamen single*; *scales oblanceolate, entire, black, covered with long silky hairs.* (*S. Uva-Ursi*, *Pursh.*, in part, & *Ed. 2.*) Alpine summits of the high mountains of Maine, New Hampshire, and New York.

20. **S. argyrocárpa**, Anders. (SILVERY-FRUITED W.) *Leaves* lanceolate, acute, or the earliest obovate and obtuse, irregularly *repand-crenate*, smooth and green above, *covered beneath when young with long and shining deciduous hairs*, at maturity smooth and glaucous; catkins ovoid, short; *ovary densely silvery-silky, stalked*; style distinct; *stamens 2-3*; gland sometimes double; scales obovate, obtuse, clothed with long hairs. (*S. repens*, *Ed. 2.* but is much nearer *S. arbuscula*, *L.*) — Moist alpine ravines of the White Mountains, New Hampshire. — Whole plant, when young, of a glossy satiny lustre; the leaves at length becoming quite smooth, with a white and prominent midrib, and slightly elevated veins.

21. **S. herbácea**, L. (HERB-LIKE W.) *Leaves roundish-oval, heart-shaped, notched at the apex*, serrate, smooth and shining, with reticulated veins; *catkins issuing from the terminal buds*, small and few-flowered; *ovary sessile, smooth*; scales smooth, ciliate. — Alpine summits of the White Mountains of New Hampshire, and high northward. — A very small herb-like species, the stems seldom rising above an inch or two from the ground. (Eu.)

2. PÓPULUS, Tourn. POPLAR. ASPEN.

Bracts (*scales*) of the catkins irregularly cut-lobed at the apex. Flowers from a cup-shaped disk which is obliquely lengthened in front. Stamens 8-30, or more: filaments distinct. Stigmas elongated. — Trees, with broad and more or less heart-shaped or ovate-toothed leaves, and mostly angular branches. Buds scaly, covered with resinous varnish. Catkins long and drooping, appearing before the leaves. (The ancient name, called *Arbor Populi*, because it was used to decorate the public walks, or on account of the constant agitation of the leaves by every impulse.)

1. **P. tremuloides**, Michx. (AMERICAN ASPEN.) *Leaves roundish-heart-shaped*, with a short sharp point, and *small somewhat regular teeth*, smooth on both sides, with downy margins; *scales cut into 3-4 deep linear divisions*, fringed with long hairs. — Woods: common. — Tree 20°-50° high, with smooth greenish-white bark. Leaf-stalk long, slender, and laterally compressed, which accounts for the continual agitation of the foliage by the slightest breeze.

2. **P. grandidentata**, Michx. (LARGE-TOOTHED ASPEN.) *Leaves roundish-ovate*, with *large and irregular sinuate teeth*, when young densely covered

with white silky wool, at length smooth both sides; *scales cut into 5-6 unequal small divisions*, slightly fringed. — Woods: common, especially northward. — A rather larger tree than the last, with a smoothish gray bark.

3. **P. heterophylla**, L. (DOWNY POPLAR.) *Branches round; leaves heart-shaped or roundish-ovate, obtuse, serrate, white-woolly when young*, at length nearly smooth, except on the elevated veins beneath. — Swamps, W. New England to Illinois and southward. — Tree 40°-60° high, with large, usually blunt leaves; the sinus, when heart-shaped, closed by the overlapping lobes which conceal the insertion of the nearly round leaf-stalk.

4. **P. monilifera**, Ait. (COTTON-WOOD. NECKLACE POPLAR.) *Young branches slightly angled, becoming round; leaves broadly deltoid, with spreading prominent nerves, slightly heart-shaped or truncate at the base, taper-pointed, serrate with cartilaginous and incurved slightly hairy teeth; fertile catkins very long; scales lacerate-fringed, not hairy; stigmas nearly sessile, toothed, dilated and very large.* — Margins of lakes and streams, W. New England to Illinois and southward, especially westward. — A large tree, 80° high or upwards; the vigorous branches decidedly angled, bearing large leaves; the more stunted round, with smaller foliage. (*P. Canadensis, Michx. f. P. lævigata, Willd.*)

5. **P. angulata**, Ait. (ANGLED COTTON-WOOD.) *Branches acutely angular or winged; leaves broadly deltoid or heart-ovate, smooth, crenate-serrate, or with obtuse cartilaginous teeth.* — Low grounds, Pennsylvania to Wisconsin and southward. — Tree large as (and probably a mere variety of) the last, and like it bearing very large heart-shaped leaves (7'-8' in length and breadth) on young plants and suckers: on full-grown trees only one fourth of that size, and commonly without the sinus.

6. **P. balsamifera**, L. (BALSAM POPLAR. TACAMAHAC.) *Branches round; leaves ovate, gradually tapering and pointed, finely serrate, smooth on both sides, whitish and reticulately veined beneath; scales dilated, slightly hairy; stamens very numerous.* — N. New England to Wisconsin, and northward. — A tall tree, growing on the borders of rivers and swamps: its large buds varnished with copious fragrant resinous matter.

Var. **candicans**. (BALM OF GILEAD.) *Leaves broader and more or less heart-shaped, pointed, serrate, whitish and reticulate-veined beneath; petiole commonly hairy.* (*P. candicans, Ait.*) — N. New England to Lake Superior and Kentucky: rare in a wild state, but common in cultivation.

P. NIGRA, L., was admitted by the elder Michaux into his Flora, without any mention of its locality. It was afterwards published by his son, under the name of *P. Hudsonica*: he, however, found it "only on the banks of the Hudson River, above Albany." Lastly, it was described as *P. betulifolia* by Pursh, who further added as its station, "about Lake Ontario" It was probably introduced from Europe, and was latterly so considered by the younger Michaux himself.

P. DILATATA, Ait., the well-known LOMBARDY POPLAR (probably a form of *P. nigra*) has been extensively introduced as an ornamental tree.

P. ALBA, L., the ABELE or WHITE POPLAR of the Old World, is occasionally planted, when it spreads widely by the root, and becomes more common than is desirable.

SUBCLASS II. GYMNOSPÉRMEÆ.

Pistil represented by an open scale or leaf, or else entirely wanting; the ovules and seeds therefore naked (without a pericarp), and fertilized by the direct application of the pollen. Cotyledons often more than two.

ORDER 106. CONIFERÆ. (PINE FAMILY.)

Trees or shrubs, with resinous juice, mostly awl-shaped or needle-shaped entire leaves, and monœcious or rarely diœcious flowers in catkins, destitute of calyx or corolla. Ovules orthotropous. Embryo in the axis of the albumen, nearly its length. (Wood destitute of ducts, composed chiefly of a homogeneous large woody fibre which is marked with circular disks on two sides.) Composes the three following Suborders:—

SUBORDER I. ABIETINÆÆ. PROPER PINE FAMILY.

Fertile flowers in catkins, consisting of open imbricated carpels in the form of scales in the axil of a bract; in fruit forming a *strobile* or *cone*. Ovules 2, adherent to the base of each carpellary scale, their orifice turned downward. Buds scaly.

* Cones maturing the year after flowering.

1. **Pinus.** Leaves 2–5 in a cluster from the axil of a scale-like primary leaf, persistent.
- ** Cones maturing the same year.
2. **Abies.** Leaves all scattered on the branches and alike, persistent.
3. **Larix.** Leaves many in a cluster, the primary ones similar to the others, deciduous.

SUBORDER II. CUPRESSINÆÆ. CYPRESS FAMILY.

Fertile flowers consisting of few carpellary scales, without bracts, bearing single or several erect ovules on their base (the orifice upward), forming a closed *strobile* or a sort of *drupe* in fruit. Buds naked.

* Flowers monœcious. Strobile dry, opening at maturity.

4. **Thuja.** Fruit of few imbricated oblong scales. Ovules 2. Leaves scale-like, closely imbricated on the flattened branches.
5. **Cupressus.** Fruit of several shield-form thick scales united in a globular woody cone. Seeds 2 or more on the stalk of each scale. Leaves scale-like or awl-shaped.
6. **Taxodium.** Fruit of several thick shield-shaped scales united in a globular woody cone. Seeds 2 on the base of each scale. Leaves linear, 2-ranked, deciduous.
- ** Flowers chiefly diœcious. Fruit berry-like, not opening.
7. **Juniperus.** Fruit composed of 3–6 coalescent 1–3-ovuled scales, becoming fleshy.

SUBORDER III. TAXINÆÆ. YEW FAMILY.

Fertile flower solitary, consisting of a naked ovule, ripening into a nut-like or *drupe*-like seed. Carpel or scale entirely wanting. Buds scaly.

8. **Taxus.** Ovule erect, encircled at the base by an annular disk, which in fruit forms a berry-like cup around the nut-like seed.

1. PĪNUS, Tourn. PINE.

Sterile catkins arranged in a spike at the base of the shoot of the same spring, involucrate by a nearly definite number of scales, consisting of numerous stamens spirally inserted on the axis, with very short filaments and a scale-like connective: anther-cells 2, opening lengthwise. Pollen of 3 united grains. Fertile catkins solitary or aggregated immediately below the terminal bud, or lateral on the young shoot, consisting of imbricated carpellary scales, each in the axil of a persistent bract, bearing a pair of inverted ovules at the base. Fruit a cone formed of the imbricated and woody carpellary scales, which are thickened at the apex (except in White Pines), persistent, spreading when ripe and dry; the 2 nut-like seeds partly sunk in excavations at the base of the scale, and in separating carrying away a part of its lining in the form of a thin and fragile wing. Cotyledons 3-12, linear. — Primary leaves of the shoots thin and chaff-like, merely bud-scales; from their axils immediately proceed the secondary leaves, which make the foliage, in the form of fascicles of 2 to 5 needle-shaped evergreen leaves, from slender buds, some thin scarious bud-scales sheathing the base of the cluster. When there are only 2 leaves in the cluster they are semicylindrical and when dry channelled or concave; when more than 2 triangular; their edges in our species serrulate. Blossoms developed in spring; the cones maturing only in the autumn of the second year. (The classical Latin name.)

The species here arranged and characterized from notes contributed by DR. ENGELMANN.

§ 1. *Leaves 2 or 3 in a close sheath: cones (except in No. 7) persistent after shedding the seed; their woody scales thickened at the end, and usually spiny-tipped.*

* *Fertile catkins and cones lateral: scales much thickened at the end: leaves rigid.*

+ *Leaves in threes.*

1. **P. Tæda**, L. (LOBLOLLY OR OLD-FIELD PINE.) *Leaves long (6' - 10'), with elongated sheaths, light green; cones elongated-oblong (3' - 5' long) and tapering; scales tipped with a stout incurved spine.* — Barren light soil, Delaware (W. M. Canby), Virginia, and southward near the coast. — Tree 50° - 100° high. Sterile catkins slender, 2' long, their involucre of 10-13 (rarely fewer) scales. Seeds with 3 strong and rough ridges on the under side: cotyledons 5-8.

2. **P. rígida**, Miller. (PITCH PINE.) *Leaves (3' - 5' long) dark green, from short sheaths; cones ovoid-conical or ovate (1' - 3½' long), often in clusters; the scales tipped with a short and stout recurved prickle.* — Sandy or spare rocky soil, Maine to W. New York and southward: common. — Tree 30° - 70° high, with very rough and dark bark, and hard resinous wood. Sterile catkins slender (½' - 1½' long), with 6-8 involucreal scales. Seed with hardly any ridges: cotyledons about 5.

+ + *Leaves in twos, or some of them in threes in No. 6.*

3. **P. púngens**, Michx. (TABLE MOUNTAIN PINE.) *Leaves stout, short (1¼' - 2½' long), crowded, bluish, the sheath short (very short on old foliage); cones ovate (3½' long), the scales armed with a strong hooked spine (¼' long).* — Alleghany Mountains, especially in the Blue Ridge, N. Carolina and Virginia

(near Charlottesville, *M. A. Curtis*) to Penn., *Prof. Porter*, as far north as Port Clinton near Reading (*T. Meehan*). — Sterile catkins 6''–9'' long, with 6–8 involucre scales. Seed with 2 or 3 slight ridges: cotyledons about 7.

4. **P. inops**, Ait. (JERSEY OR SCRUB PINE.) *Leaves short* ($1\frac{1}{2}'$ –3' long); cones oblong-conical, sometimes curved (2'–3' long), the scales tipped with a straight or recurved awl-shaped prickle. — Barrens and sterile hills, New Jersey to Kentucky and southward. A straggling tree, 15°–40° high, with spreading or drooping branchlets: young shoots with a purplish glaucous bloom. Sterile catkins linear-oblong, 5''–6'' long, with about 8 involucre scales.

5. **P. Banksiana**, Lambert. (GRAY OR NORTHERN SCRUB PINE.) *Leaves short* (1' long), oblique, divergent; cones conical, oblong, usually curved ($1\frac{1}{2}'$ –2' long), smooth, the scales pointless. (*P. rupestris*, *Michx. f.*) — Rocky banks, N. Maine, N. Michigan and Wisconsin, and northward. — A straggling shrub or low tree (5°–20° high): the sterile catkins nearly as in No. 4. Seed with 2 or 3 ridges on the inner side: cotyledons 4 or 5.

6. **P. mitis**, Michx. (YELLOW PINE.) *Leaves in pairs or sometimes in threes, from long sheaths, slender* (3'–5' long); cones ovate- or oblong-conical (barely 2' long); the scales tipped with a minute and weak prickle. (*P. variabilis*, *Pursh.*) — Dry or sandy soil, W. New England? and New Jersey to Wisconsin, and common southward. — Tree 50°–60° high, straight, producing a durable, fine-grained, moderately resinous timber, valuable for flooring, &c. Leaves more soft and slender than in any of the preceding, dark green. The western form has more rigid leaves, and more tuberculate and spiny cones. Sterile catkins linear, 6''–9'' long, with 9–12 involucre scales. Seed with 2 or 3 rough ridges: cotyledons 5–7.

* * *Fertile catkins and cones terminal, the latter deciduous after shedding the seed: their scales slightly thickened, pointless: leaves in twos, slender.*

7. **P. resinosa**, Ait. (RED PINE.) *Leaves from long sheaths, elongated* (5'–6' long), dark green; cones ovate-conical, smooth (about 2' long); sterile catkins oblong-linear (6''–9'' long), subtended by about 6 involucre scales which are early deciduous by an articulation above their base; seeds slightly ridged on the lower surface; cotyledons 6 or 7. (*P. rubra*, *Michx. f.*) — Dry woods, Mass. to Penn., Wisconsin, and northward. — Tree 50°–80° high, with reddish and rather smooth bark, and compact, not very resinous wood. — Wrongly called *Norway Pine*.

§ 2. *Leaves 5 in a (loose-scaled entirely deciduous) sheath: cones terminal, deciduous after shedding the seed; their scales slightly if at all thickened at the end and wholly destitute of prickle or point: bark smooth except on old trunks.*

8. **P. Stróbus**, L. (WHITE PINE.) *Leaves very slender, glaucous; sterile catkins oval* (4''–5'' long), with 6–8 involucre scales at base; fertile catkins long-stalked, cylindrical; cones narrow, cylindrical, nodding, often curved (4'–6' long); seed smooth; cotyledons 8–10. — Cool and damp woods; common northward to the Mississippi, and southward in the Alleghanies. — The White Pine (called in England *Weymouth Pine*) is our tallest tree, often 120°–160° in a straight column in primitive forests, and is invaluable for its soft and light white or yellowish wood, in large trunks nearly free from resin.

2. **À BIES**, Tourn. **SPRUCE. FIR.**

Sterile catkins scattered or somewhat clustered in the axils of leaves of the preceding year. Fertile catkins and cones lateral or terminal on branches of the preceding year: scales of the cone thin and even, not at all thickened nor prickly-tipped. Leaves all of one kind and foliaceous, scattered (not fascicled), short, persistent for two or more years. Flowering in spring, and cones maturing in autumn. Otherwise nearly as in *Pinus*. (The classical Latin name.) Of European species *A. excelsa*, the **NORWAY SPRUCE** is most commonly, and *A. pectinata*, the **SILVER FIR**, occasionally planted.

§ 1. **SPRUCE.** (*PICEA* of *Link*, not of *L.* **ABIES**, *L.* and others. These ancient names much transposed by moderns.) *Cones hanging from or near the end of a branch; the scales persistent on the axis: anthers tipped with a rounded recurved appendage, their cells distinct and opening lengthwise: pollen nearly as in Pine: leaves needle-shaped and 4-sided, pointing every way.*

1. **A. nigra**, Poir. (**BLACK OR DOUBLE SPRUCE.**) Leaves short (6" - 8" long), either dark green or glaucous-whitish; cones ovate or ovate-oblong (1' - 1½' long), mostly recurved, persistent, the rigid scales with a thin often eroded edge. — Swamps and cold mountain woods, New England to Wisconsin and northward, and southward along the mountains.

2. **A. alba**, Michx. (**WHITE SPRUCE.**) Leaves pale or glaucous; cones nodding, cylindrical (about 2' long), pale, deciduous, the thinner scales with an entire edge; a handsomer tree than No. 1, in aspect more like a **Balsam Fir**. — Northern New England and New York to Lake Superior, and northward.

§ 2. **HEMLOCK-SPRUCE.** (*TSUGA*, Endl.) *Cones hanging on the end of declined branches of the preceding year, persistent, small; the scales persistent on the axis: sterile catkins small, of a few capitate anthers, their short confluent cells opening transversely: pollen-grains simple: leaves flat, whitened beneath, petioled, spreading in two directions so as to seem 2-ranked.*

3. **A. Canadensis**, Michx. (**HEMLOCK SPRUCE.**) Leaves short-linear, obtuse (½' long); cones oval (6" - 8" long), of few thin scales much longer than the bracts. — Hilly or rocky woods: very common northward, and rare southward in the Alleghanies. — A large tree, when young the most graceful of Spruces, with a light and spreading spray, and delicate foliage, bright green above, silvery underneath. Timber coarse-grained and poor.

§ 3. **FIR.** (*ABIES* of *Pliny.* *PICEA*, *L.*, not of *Link.*) *Cones erect on the upper side of spreading branches; their scales and mostly exerted bracts deciduous from the persistent axis at maturity: seeds and bark of tree with balsam-bearing vesicles: anthers tipped with a 1-2-pointed appendage, their cells opening by laceration; pollen nearly as in Pine: leaves flat, with midrib prominent on the whitened lower face, mostly sessile, on horizontal branches more or less spreading in two directions, so as to seem 2-ranked.*

4. **A. balsamea**, Marshall. (**BALSAM FIR.**) Leaves narrowly linear; cones cylindrical, large (2' - 4' long, 1' thick), violet-colored; the bracts obovate, serrulate, tipped with an abrupt slender point, included or slightly projecting. — Cold damp woods and swamps, New England to Penn., Wisconsin, and north-

ward. — A slender tree, of no value as timber, when young very handsome, but short-lived. Leaves 6"–10" in length, narrower and lighter green above than those of the European *Silver Fir*. Also called *Balm-of-Gilead Fir*. *Canada balsam* is drawn from blisters in the bark of this and the next species. — A subalpine state on the Green Mountains, &c., has shorter or oblong, stunted cones, and approaches the next.

5. **A. Fraseri**, Pursh. (FRASER'S OR SOUTHERN BALSAM FIR.) *Cones small* (1'–2' long), *oblong-ovate*; the bracts *oblong-wedge-shaped*, the short-pointed upper part *much projecting and reflexed*. (*A. balsamifera*, Michx. Fl.) — Mountains of Penn., Virginia, and southward on the highest Alleghanies. — Foliage, &c. nearly as in the last.

3. LÄRIX, Tourn. LARCH.

Catkins lateral, terminating short spurs on the branches of the preceding year, short or globular, developed in early spring; the sterile from leafless buds; the fertile mostly with leaves below. Anther-cells opening transversely. Pollen-grains simple, globular. Cones as in Spruce, the scales persistent. — Leaves needle-shaped, soft, deciduous, all foliaceous, very many in a fascicle developed in early spring from lateral scaly and globular buds, and scattered along the developed shoots of the season. Fertile catkins crimson or red in flower. (The ancient name.)

1. **L. Americana**, Michx. (AMERICAN OR BLACK LARCH. TAMARACK. HACKMATACK.) Leaves short; cones ovoid (6"–9" long), of few rounded scales, arranged in $\frac{2}{3}$ order. (*Pinus pëndula*, Ait. *P. microcärpa*, Lambert.) — Swamps, New England to Penn. and Wisconsin, and (chiefly) northward. — A slender tree, with heavy, close-grained wood, horizontal branches, and more slender and usually shorter leaves than the *European Larch*; — which is a handsomer tree, and has the scales of its larger cones arranged in the order $\frac{8}{11}$.

4. THÛJA, Tourn. ARBOR VITÆ.

Flowers mostly monœcious on different branches, in very small terminal ovoid catkins. Stamens with a scale-like filament or connective, bearing 4 anther-cells. Fertile catkins of few imbricated scales, fixed by the base, each bearing 2 erect ovules, dry and spreading at maturity. Cotyledons 2. — Small evergreen trees, with very flat 2-ranked spray, on which the small and appressed persistent leaves are closely imbricated: these are of two sorts, on different or successive branchlets; the one awl-shaped; the other scale-like, blunt, short, and adnate to the branch. (*Θυϊά*, *Θύα*, or *Θυεία*, the ancient name of some resin-bearing evergreen.)

1. **T. occidentälis**, L. (AMERICAN ARBOR VITÆ.) Leaves appressed-imbricated in 4 rows on the 2-edged branchlets; scales of the cones pointless; seeds broadly winged all round. — Swamps and cool rocky banks: common from Penn. northward, where it forms extensive "cedar-swamps," and is called WHITE CEDAR: rare southward along the Alleghanies. May, June. — Tree 20°–50° high, yielding a pungent aromatic oil: wood light, but very durable.

5. CUPRÉSSUS, Tourn. CYPRESS.

Flowers monœcious on different branches, in terminal small catkins. Sterile catkins composed of shield-shaped scale-like filaments bearing 2-4 anther-cells under the lower margin. Fertile catkins globular, of shield-shaped scales bearing several erect bottle-shaped ovules. Cone globular, firmly closed, but opening at maturity; the scales thick, pointed or bossed in the middle; the few or several seeds attached to their contracted base or stalk. Cotyledons 2 or 3. — Strong-scented evergreen trees, with very small and scale-like or some awl-shaped closely appressed-imbriated leaves, and exceedingly durable wood. (The classical name.)

1. **C. thyoides**, L. (WHITE CEDAR.) Leaves minute, pale, ovate or triangular-awl-shaped, often with a small gland on the back, closely imbricated in 4 rows; anther-cells 2 under each scale; cones small (3" - 5" in diameter); seeds slightly winged. — Swamps, Massachusetts to Wisconsin, Virginia, and southward. May. — Tree 30° - 70° high; the wood and fibrous shreddy bark, as well as the foliage, much as in *Arbor Vitæ*; but the spray more slender, the leaves finer and glaucous-green.

6. TAXÒDIUM, Richard. BALD CYPRESS.

Flowers monœcious, the two kinds on the same branches. Sterile catkins spiked-panicked, of few stamens: filaments scale-like, shield-shaped, bearing 2-5 anther-cells. Fertile catkins ovoid, in small clusters, scaly, with a pair of ovules at the base of each scale. Cone globular, closed, composed of very thick and angular somewhat shield-shaped scales, bearing 2 angled seeds at their base. Cotyledons 6-9. — Trees, with narrow linear 2-ranked light and deciduous leaves; a part of the slender leafy branchlets of the season also deciduous in autumn. (Name compounded of *Τάξος*, the *Yew*, and *εἶδος*, *resemblance*, the leaves being *Yew-like*.)

1. **T. distichum**, Richard. (AMERICAN BALD CYPRESS.) Leaves linear and spreading; also some awl-shaped and imbricated on flowering branchlets. — Swamps, Delaware, to S. Illinois, and southward, where it is a very large and valuable tree. March, April.

7. JUNÍPERUS, L. JUNIPER.

Flowers diœcious, or occasionally monœcious, in very small lateral catkins. Anther-cells 3-6, attached to the lower edge of the shield-shaped scale. Fertile catkins ovoid, of 3-6 fleshy coalescent scales, each one-ovuled, in fruit forming a sort of berry, which is scaly-bracted underneath, black with white bloom. Seeds 1-3, wingless, bony. Cotyledons 2. — Evergreen trees or shrubs, with awl-shaped or scale-like rigid leaves, often of two shapes in § 2. (The classical name.)

§ 1. *Leaves all in whorls and linear-awl-shaped, prickly-pointed, free, articulated with the stem, with a midrib and rib-like margins.*

1. **J. communis**, L. (COMMON JUNIPER.) Leaves in threes, with slender prickly point, spreading, bright green except the glaucous-white upper face,

6''-9'' long; berries large. — Dry sterile hills: common. May, June. — Low shrub, ascending or spreading on the ground. (Eu.)

Var. **alpina**, L. (*J. nana*, Willd.), is a prostrate state, with shorter and less tapering, mostly ascending or incurved leaves. — Shores of upper Great Lakes, Maine, and northward. (Eu.)

§ 2. *Leaves small, mostly opposite, not articulated but connate-decurrent on the stem of two somewhat different forms, i. e. awl-shaped and loose, and scale-shaped and appressed-imbricated, the latter flattened and often with a resiniferous gland on the back, and no distinct nerve or midrib.*

2. **J. Virginiana**, L. (RED CEDAR OR SAVIN.) Scale-shaped leaves acute or acutish; *fruit small, erect* on the short supporting branchlet. — Dry, mostly sterile or rocky soil: common. May. — Shrub, small tree, or westward often a large tree, 60°-90° high; with most durable, compact, reddish and odorous wood.

3. **J. Sabina**, L., var. **procumbens**, Pursh. Scale-shaped leaves obtuse or acutish, strongly appressed; *fruit larger, nodding on the recurved peduncle-like branchlet*; stems procumbent or prostrate, sometimes extensively creeping. (*J. Virginiana*, var. *humilis*, Ed. 2.) — Rocky banks, borders of swamps, &c., Maine to Wisconsin along and near the Great Lakes, and northward. May, June. (Eu.)

8. TÁXUS, Tourn. YEW.

Flowers mostly dioecious, or sometimes monœcious, axillary from scaly buds; the sterile in small globular catkins formed of a few naked stamens: anther-cells 3-8 under a shield-like somewhat lobed connective. Fertile flowers solitary, scaly-bracted at the base, consisting merely of an erect sessile ovule; with an annular disk, which becomes cup-shaped around its base, and at length pulpy and berry-like, globular and red, and nearly enclosing the nut-like seed. Cotyledons 2. — Leaves evergreen, flat, mucronate, rigid, scattered, 2-ranked. (The classical name, probably from *τόξον*, a bow; the wood anciently used for bows.)

1. **T. baccata**, L., var. **Canadensis**. (AMERICAN YEW. GROUND HEMLOCK.) Stems diffusely spreading; leaves linear, green both sides. (*T. Canadensis*, Willd.) — Moist banks and hills, near streams, especially in the shade of evergreens: common northward, extending southward only along the Alleghanies. April. — Our Yew is a low and straggling or prostrate bush, never forming an erect trunk like that of Europe and of Northwest America. (Eu.)

CLASS II. MONOCOTYLÉDONOUS OR ENDÓGENOUS PLANTS.

Stems with no manifest distinction into bark, wood, and pith; but the woody fibre and vessels in bundles or threads which are irregularly imbedded in the cellular tissue: perennial trunks destitute of annual layers. Leaves mostly parallel-veined (nerved) and sheathing at the base, seldom separating by an articulation, almost always alternate or scattered and not toothed. Parts of the flower commonly in threes. Embryo with a single cotyledon, and the leaves of the plumule alternate.

ORDER 107. ARACEÆ. (ARUM FAMILY.)

Plants with acrid or pungent juice, simple or compound often veiny leaves, and flowers crowded on a spadix, which is usually surrounded by a spathe. — Floral envelopes none, or of 4–6 sepals. Fruit usually a berry. Seeds with fleshy albumen, or none but filled with the large fleshy embryo in Nos. 2, 4, and 5. A large family, chiefly tropical. Herbage abounding in slender raphides. — The genuine Araceæ have no floral envelopes, and are almost all monœcious or diœcious: but the genera of the second section with more highly developed flowers are not to be separated.

* Spathe surrounding or subtending the spathe: flowers naked; i. e. without perianth.

1. **Arisæma.** Flowers monœcious or diœcious, covering only the base of the spadix.
2. **Peltandra.** Flowers monœcious, covering the spadix; anthers above, ovaries below.
3. **Calla.** Flowers perfect (at least the lower ones), covering the whole of the short spadix.

Spathe open and spreading.

** Spathe surrounding the spadix in No. 4, none or imperfect in the rest: flowers with a calyx or perianth and perfect, covering the whole spadix.

4. **Symplocarpus.** Spadix globular, in a fleshy shell-shaped spathe. Stemless.
5. **Orontium.** Spadix narrow, naked, terminating the terete scape.
6. **Acorus.** Spadix cylindrical, borne on the side of a leaf-like scape.

1. ARISÆMA, Martius. INDIAN TURNIP. DRAGON-ARUM.

Spathe convolute below and mostly arched above. Flowers monœcious or by abortion diœcious, covering only the base of the spadix, which is elongated and naked above. Floral envelopes none. Sterile flowers above the fertile, each of a cluster of almost sessile 2–4-celled anthers, opening by pores or chinks at the top. Fertile flowers consisting each of a 1-celled ovary, tipped with a depressed stigma, and containing 5 or 6 orthotropous ovules erect from the base

of the cell; in fruit a 1-few-seeded scarlet berry. Embryo in the axis of albumen. — Low perennial herbs, with a tuberous rootstock or corm, sending up a simple scape sheathed with the petioles of the simple or compound veiny leaves, as if coalescent. (A play upon *Arum*, the ancient name; probably formed of ἄρον, *Arum*, and σήμα, *a sign or mark*.)

1. **A. triphýllum**, Torr. (INDIAN TURNIP.) *Leaves mostly 2, divided into 3 elliptical-ovate pointed leaflets; spadix mostly dioecious, club-shaped, obtuse, much shorter than the spathe, which is flattened and incurved-hooded at the summit.* (*Arum triphyllum*, L.) — Rich woods: common. May. — Corm turnip-shaped, wrinkled, farinaceous, with an intensely acrid juice. Spathe with the petioles and sheaths green, or often variegated with dark purple and whitish stripes or spots (*Arum atrorubens*, *Ait.*); the limb ovate-lanceolate, pointed.

2. **A. Dracontium**, Schott. (GREEN DRAGON. DRAGON-ROOT.) *Leaf usually solitary, pedately divided into 7-11 oblong-lanceolate pointed leaflets; spadix often androgynous, tapering to a long and slender point beyond the oblong and convolute pointed spathe.* (*Arum Dracontium*, L.) — Low grounds along streams. June. — Corms clustered. Petiole 1°-2° long, much longer than the peduncle. Spathe greenish, rolled into a tube, with a short erect point.

2. PELTÁNDRA, Raf. ARROW ARUM.

Spathe elongated, convolute throughout, wavy on the margin, curved at the apex. Flowers monœcious, thickly covering the long and tapering spadix throughout. Floral envelopes none. Anthers sessile, naked, covering all the upper part of the spadix, each of 5 or 6 cells imbedded in the margin of a thick and shield-shaped connective, opening by a terminal pore. Ovaries 1-celled at the base of the spadix, bearing several amphitropous or nearly orthotropous ovules at the base: stigma almost sessile. Fruit a leathery or fleshy utricle, 1-3-seeded. Seed obovate, surrounded by a tenacious jelly, the base empty, the upper part filled with a large and fleshy spherical embryo; no albumen. — A stemless herb, with arrow-shaped leaves and simple scapes from the root of thick tufted fibres. Upper part of the spathe and the sterile portion of the spadix rotting away after flowering, leaving the fleshy base firmly enclosing the globular cluster of fruit. (Name composed of πέλτη, *a target*, and ἀνήρ, *for stamen*, from the shape of the latter.)

1. **P. Virgínica**, Raf. (*P. undulata*, Raf. *Arum Virginicum*, L. *Leconia*, Torr. *Reusselæria*, Beck.) — In shallow water: common. June. — Leaves large, pointed; nerves reticulated next the margin. (It seems to have escaped attention that this plant has an *exalbuminous corm-like embryo*, nearly as in *Symplocarpus*.)

3. CÁLLA, L. WATER ARUM.

Spathe open and spreading, ovate (abruptly pointed, the upper surface white), persistent. Spadix oblong, entirely covered with flowers; the lower perfect and 6-androus; the upper often of stamens only. Floral envelopes none. Filaments slender: anthers 2-celled, opening lengthwise. Ovary 1-celled, with 5-9 erect anatropous ovules: stigma almost sessile. Berries (red) distinct, few-seeded.

Seeds with a conspicuous rhaphe and an embryo nearly the length of the hard albumen. — A low perennial herb, growing in cold bogs, with a long creeping rootstock, bearing heart-shaped long-petioled leaves, and solitary scapes. (An ancient name, of unknown meaning.)

1. **C. palústris**, L. — Cold bogs, New England to Penn., Wisconsin, and common northward. June. — Seeds surrounded with jelly. (Eu.)

4. SYMPLOCÁRPUŠ, Salisb. SKUNK CABBAGE.

Spathe hooded-shell-form, pointed, very thick and fleshy, decaying in fruit. Spadix globular, short-stalked, entirely covered with perfect flowers, which are thickly crowded and their (1-celled or abortively 2-celled) ovaries immersed in the fleshy receptacle. Sepals 4, hooded. Stamens 4, opposite the sepals, with at length rather slender filaments: anthers extrorse, 2-celled, opening lengthwise. Style 4-angled and awl-shaped: stigma small. Ovule solitary, suspended, anatropous. Fruit a globular or oval mass, composed of the enlarged and spongy spadix, enclosing the spherical seeds just beneath the surface, which is roughened with the persistent and fleshy sepals and pyramidal styles. Seeds filled by the large globular and fleshy corm-like embryo, which bears one or several plumules at the end next the base of the ovary: albumen none. — Perennial herb, with a strong odor like that of the skunk, and also somewhat alliaceous; a thick descending rootstock bearing a multitude of long and coarse fibrous roots, and a cluster of very large and broad entire veiny leaves, preceded in earliest spring by the nearly sessile spathes, which barely rise out of the ground. (Name from *συμπλοκή*, connection, and *καρπός*, fruit, in allusion to the coalescence of the ovaries into a compound fruit.)

1. **S. fœtidus**, Salisb. (*Ictodes*, Bigel.) — Bogs and moist grounds: common. — Leaves ovate and heart-shaped, 1°–2° long when grown, short-petioled. — Spathe spotted and striped with purple and yellowish-green, ovate, incurved. Fruit ripe in September, forming a roughened globular mass 2'–3' in diameter, in decay shedding the bulblet-like seeds, which are 4''–6'' in diameter, and filled with the singular solid fleshy embryo.

5. ORÓNTIUM, L. GOLDEN-CLUB.

Spathe incomplete and distant, merely a leaf-sheath investing the lower part of the slender scape, and bearing a small and imperfect bract-like blade. Flowers crowded all over the narrow spadix, perfect: the lower with 6 concave sepals and 6 stamens; the upper ones with 4. Filaments flattened: anthers 2-celled, opening obliquely lengthwise. Ovary 1-celled, with an anatropous ovule: stigma sessile, entire. Fruit a green utricle. Seed without albumen. Embryo thick and fleshy, "with a large concealed cavity at the summit, the plumule curved in a groove on the outside." (*Torr.*) — An aquatic perennial, with a deep rootstock, long-petioled and entire oblong and nerved floating leaves, and the spadix terminating the elongated scape; its rather club-shaped emersed apex as thick as the spadix. (Origin of the name obscure.)

1. **O. aquáticum**, L. — Ponds, Massachusetts to Virginia, near the coast, and southward. May.

6. **ÁCORUS**, L. SWEET FLAG. CALAMUS.

Spadix cylindrical, lateral, sessile, emerging from the side of a simple 2-edged scape which resembles the leaves, densely covered with perfect flowers. Sepals 6, concave. Stamens 6: filaments linear · anthers kidney-shaped, 1-celled, opening across. Ovary 2-3-celled, with several pendulous orthotropous ovules in each cell: stigma minute. Fruit at length dry, gelatinous inside, 1-few-seeded. Embryo in the axis of albumen. — Pungent aromatic plants, especially the thick creeping rootstocks (*calamus* of the shops), which send up 2-edged sword-like leaves, and scapes similar to them, bearing the spadix on one edge; the upper and more foliaceous prolongation sometimes considered as a kind of open spathe. (The ancient name, from *a* privative, and *κόρη*, the pupil of the eye, having been used as a remedy for sore eyes.)

1. **A. Cálamus**, L. Scape leaf-like and prolonged far beyond the (yellowish-green) spadix. — Margin of rivulets, swamps, &c. Probably truly indigenous northward. June. (Eu.)

ORDER 108. **LEMNACEÆ**. (DUCKWEED FAMILY.)

Minute stemless plants, floating free on the water, destitute of distinct stem and foliage, being merely a frond, producing one or few monœcious flowers from the edge or upper surface, and commonly hanging roots from underneath: ovules rising from the base of the cell. Fruit a 1-7-seeded utricle. Seed large; the apex or radicular extremity of the seed-coat separable as an operculum or lid (as in Cabomba, &c.) Embryo straight, surrounded by fleshy or sometimes very scanty albumen — The simplest, and some of them the smallest of flowering plants, propagating by the proliferous growth of a new individual from a cleft in the edge or base of the parent frond, remaining connected for some time or separating, also by autumnal fronds in the form of minute bulblets, which sink to the bottom of the water, but rise and vegetate in spring; the flowers (in summer) and fruit scarce, in some species hardly ever seen. Frond more or less cavernous; the upper surface furnished with stomata. — These plants may be regarded as a sort of very simplified Araceæ.

Arranged from notes contributed by C. F. AUSTIN, Esq., who is preparing a monograph of the American Lemnaceæ.

1. **LÉMNA**, L. DUCKWEED. DUCK'S-MEAT.

Flowers produced from a cleft in the margin of the frond, usually three together surrounded by a spathe; two of them staminate, consisting of a stamen only; the other pistillate, of a simple pistil; the whole therefore imitating a single diandrous flower. *Ster. Fl.* Filament slender: anther 2-celled, didymous; the cells dehiscent transversely: pollen-grains large, spherical, muricate. *Fert. Fl.* Ovary 1-celled: style and truncate or funnel-shaped stigma simple. Ovules and seeds 1-7. — Fronds producing rootlets underneath, proliferous

from a cleft in the margin towards the base, and at length stipitate; the tissue abounding with bundles of acicular raphides, as in Araceæ. (An old Greek name, of uncertain meaning.)

§ 1. LEMNA, Schleiden. *Root single: ovule one, half-anatropous or orthotropous.*

1. **L. trisúlca**, L. *Fronds oblong-lanceolate from a stalked base, thin, denticulate at the tip ($\frac{1}{2}$ ' - $\frac{3}{4}$ ' long), proliferous from one or both sides; seed half-anatropous. — Ponds and springy places; immersed and living through the winter, usually several generations remaining connected. Flowers found, for the first time in this country, by C. M. Booth, Rochester, New York. (Eu.)*

2. **L. Tórrayi**, Austin, n. sp. *Fronds oblong or obovate-oblong, usually somewhat falcate (1'' - 2 $\frac{1}{2}$ '' long), thin, faintly 1-nerved, cavernous to the apex, pale green both sides, commonly in groups of 4 - 8; utricle elongated-ovate, pointed by the long style, flattish, usually half the length of the frond; seed oblong and rather abruptly expanded below the middle, unequally cordate at the base, orthotropous, very obtuse, faintly striate when dry; testa fleshy, loosely adhering to the thickish and solid inner coat; operculum distinctly apiculate; albumen very scanty (a single layer of cells). — Pools, New Jersey (fertile) to Missouri and southward.*

3. **L. perpusilla**, Torr. *Fronds obovate or roundish-obovate, thinnish (1'' - 2'' long), obscurely 3-nerved, grouped in circular patches (of 4 - 8); utricle ovate and at length oblique, tipped with a rather long eccentric style; seed roundish or obovate-oval, obtuse, oblique in the utricle, orthotropous, slightly striate when dry; testa coriaceous, solid, firmly attached to the very thin inner coat; operculum scarcely apiculate; albumen abundant. — S. New York to Wisconsin and southward: not rare. — Var. TRINÉRVIS, Austin. *Fronds larger, distinctly 3-nerved, somewhat hyaline; utricle ovate, pointed with rather long style; seed ovate, acutish, mostly straight, unequally cordate. — New York and Penn. (fertile) to Wisconsin.**

4. **L. minor**, L. *Fronds obovate or elliptical-obovate, thickish (1'' - 2'' long), often grouped and short-stalked, obscurely 3-nerved; utricle short-urn-shaped and tipped with a short style; seed oval or ovate, half-anatropous, horizontal, parallel with the margin of the frond. — Stagnant waters: common: rare in flower. (Eu.)*

Var. **obscura**, Austin. *Fronds obovate, convex on both surfaces, often purple underneath. — New York to Virginia: sterile.*

Var. **orbiculata**, Austin. *Fronds nearly orbicular, flat above, convex and dark purple underneath, proliferous on very short included stalks. — New York to Wisconsin and southward: sterile. — This and the preceding variety have been taken for L. gibba by American authors.*

§ 2. SPIRÒDELA, Schleiden. *Roots several in a cluster: ovules 2, anatropous.*

5. **L. polyrrhiza**, L. *Fronds round-obovate (2'' - 4'' long), thick, purple and rather convex beneath, dark green above, palmately (mostly 7-) nerved. — Very common in ponds and pools; not found here in blossom. (Eu.)*

§ 3. TELMATÓPHACE, Schleiden. *Roots single: ovules and seeds 2 - 7, anatropous: albumen hardly any in the mature seed.*

6. **L. gibba**, L. *Fronds obovate or almost orbicular ($1\frac{1}{2}$ '' - 3'' long), nearly flat above, much thickened and cavernous-spongy underneath (almost hemispherical), proliferous on short and very fragile stalks, therefore seldom*

more than 2 or 3 in connection. — Occurs in Arizona and southward: but the true plant seems not to have been detected within our limits. (Eu.)

2. WÓLFFIA, Horkel, Schleiden.

Flowers central, bursting through the upper surface of the roundish entirely and loosely cellular frond, only 2; one pistillate, of a single stamen with a one-celled 2-valved anther; the other pistillate, a globular ovary, tipped with a very short style and a depressed stigma. Ovule orthotropous, rather oblique in the cell. Pollen-grains smooth. Utricle spherical. Albumen thin. — *Fronde rootless*, proliferous from a cleft at the base, the offspring soon detached: no raphides. — The simplest and smallest of flowering plants, from $\frac{1}{4}$ " — $\frac{1}{3}$ " long (a new African and Cuban species much larger), floating as little grains on the water. (Named for *John Fred. Wolff*, who wrote on *Lemna* in 1801.)

1. **W. Columbiána**, Karsten. Frond oval-orbicular, lenticular ($\frac{1}{4}$ " — $\frac{1}{3}$ " long), very cavernous, when dry concave and darker green above, sometimes purple beneath, the chink at the base appendaged by an expanded hyaline border; flowers and fruit not seen. — Floating on stagnant waters with *Lemna*; near Dutcher's Bridge, Salisbury, Connecticut, *Dr. J. W. Robbins*, who discovered it many years ago. Orange Co., New York, *C. F. Austin*. Lake Ontario, *J. A. Paine*. Also in the Southern States (Louisiana, *Riddell*, &c.): probably not rare, but overlooked. The fruiting plant recently discovered by Karsten in Venezuela, according to *Hegelmaier*, in litt.

ORDER 109. TYPHACEÆ. (CAT-TAIL FAMILY.)

Marsh or aquatic herbs, with nerved and linear sessile leaves, and monœcious flowers on a spadix or in heads, destitute of proper floral envelopes. Ovary tapering into a style and (usually elongated) 1-sided stigma. Fruit nut-like when ripe, 1-seeded, rarely 2-seeded. Seed suspended, anatropous: embryo straight in copious albumen. Root perennial. — Comprises only the two following genera.

1. TÝPHA, Tourn. CAT-TAIL FLAG.

Flowers in a long and very dense cylindrical spike terminating the stem; the upper part consisting of stamens only, intermixed with long hairs, and inserted directly on the axis; the lower or fertile part consisting of ovaries, surrounded by club-shaped bristles, which form the copious down of the fruit. Nutlets minute, very long-stalked. — Spathes merely deciduous bracts, or none. Rootstocks creeping. Leaves long, sheathing the base of the simple jointless stems, erect, thickish. Flowering in summer. (Name from *τίφος*, a fen.)

1. **T. latifolia**, L. (COMMON CAT-TAIL OR REED-MACE.) *Leaves flat*: staminate and pistillate parts of the spike approximate. Common. (Eu.)

2. **T. angustifolia**, L. (NARROW-LEAVED OR SMALL C.) *Leaves channelled towards the base, narrowly linear*: staminate and pistillate parts of the spike usually separated by an interval. A rarer and smaller plant. (Eu.)

2. SPARGANIUM, Tourn. BUR-REED.

Flowers collected in separate dense and spherical leafy-bracted heads, which are scattered along the summit of the stem; the upper ones sterile, consisting merely of stamens, with minute scales irregularly interposed; the lower or fertile larger, consisting of numerous sessile pistils, each surrounded by 3-6 scales much like a calyx. Fruit wedge-shaped or club-shaped. — Rootstocks creeping and stoloniferous: roots fibrous. Stems simple or branching, sheathed below by the base of the linear leaves. Flowering through the summer. (Name from *σπάργανον*, a fillet, from the ribbon-like leaves.)

* *Erect, with branched inflorescence of numerous heads: pistil as long as the surrounding truncate scales, attenuated into a short style bearing one or often two elongated stigmas: nuts sessile, wedge-shaped, angular: leaves for the greater part flat and merely keeled, the base triangular with concave sides.*

1. **S. eurycarpum**, Engelm. Fruit many-angled ($3\frac{1}{2}''$ – $4''$ long) when fully ripe, with a broad and depressed or retuse summit ($2\frac{1}{2}''$ – $3\frac{1}{2}''$ wide) abruptly tipped in the centre; fruit-heads 1' wide. (S. ramosum, in part, of most American botanists.) — Borders of ponds, lakes, and rivers, from New England and Pennsylvania northward and westward. — Stems stout, 2° – 4° high; heads 2 to 6 or more: the largest species known.

(S. RAMOSUM, Hudson, of Europe, has not yet been found on this continent: it is distinguished by smaller heads, and smaller, few-angled, usually 1-seeded fruit, with a conical and long-pointed summit.)

* * *Erect or rarely floating, with simple (or rarely branched) inflorescence of numerous heads; the conspicuous style longer than the spatulate denticulate scales: stigmas always single, linear or oblong: nuts attenuated at both ends, and with a stalked base, nearly terete: stems rather slender: leaves (unless floating) triangular with flat sides in the lower half.*

2. **S. simplex**, Hudson, GENUINUM. Erect (9'–15' high), slender; inflorescence simple, the lower heads supra-axillary, sessile or commonly peduncled ($7''$ – $8''$ wide); stigma linear, equal to the style; fruit more or less contracted in the middle. — New England and northward. (Eu.)

Var. **Nuttallii**. Like the last or type, but heads axillary; stigma linear-oblong, shorter than the style; fruit less contracted. (S. Americanum, Nutt.) — From Pennsylvania and New England northward and northwestward. — Inflorescence rarely branched; heads $8''$ – $9''$ wide.

Var. **androccladum**. Stoutier ($1\frac{1}{2}^{\circ}$ – 3° high); inflorescence branched below; branches bearing numerous sterile (rarely also 1 or even 2 fertile) heads; stigma linear, as long as the style; fruit larger, not contracted, long-tapering upwards and downwards. (S. ramosum, in part, of American authors.) — From New England southward and especially westward. — Heads $10''$ – $12''$ wide.

Var. **fluitans**. Leaves floating; inflorescence branched; branches bearing fertile heads below; stigma oval, shorter than the style; fruit somewhat contracted and with a short stipe. (S. fluitans, Fries.) — Ponds at the base of the White Mountains, Oakes. — Heads $6''$ – $7''$ wide. (Eu.)

Var. **angustifolium**. Leaves floating, longer and narrower than in the

last; inflorescence simple; lower fertile heads usually supra-axillary, sessile or peduncled; stigma linear, about the length of the style; fruit smaller, short-stiped, contracted in the middle. (*S. angustifolium*, *Michx.* *S. affine*, *Schnitzlein*; said to be the true *S. natans* of Linnæus.)—Mountain lakes and slow streams, New York, New England, and northward.—Heads 5"—7" wide. Dwarf states, growing nearly out of water, have shorter erect leaves. (Eu.)

* * * *Usually floating, with very slender stems and delicate always flat and narrow leaves: inflorescence simple, of few small heads: stigma oval, about as long as the short style, scarcely surpassing the oval or obovate denticulate scales: nuts oval, with a very short stipe and short point.*

3. **S. minimum**, Bauhin, Fries. Fertile heads solitary or usually 2, axillary, sessile or the lower one peduncled, fruit heads 4"—5" in diameter; nuts somewhat triangular, the lower third usually contracted; stems when out of water only 5'—6' high; when submersed longer. (*S. natans*, of older authors, but not of Linnæus, according to Fries. *S. angustifolium*, *Ed. 2.*)—Northern New England to Wisconsin and northward. (Eu.)

ORDER 110. NAIADACEÆ. (PONDWEED FAMILY.)

Immersed aquatic plants, with jointed stems and sheathing stipules within the petioles, or with sheathing bases to the leaves, inconspicuous flowers, which are naked or with a free merely scale-like calyx; the ovaries solitary or 2—4 and distinct, 1-celled, 1-ovuled. Seed without albumen, filled by the large embryo, often curved or hooked. Flowers usually bursting from a spathe, sometimes on a spadix.

* Flowers monœcious or diœcious, axillary, naked, monandrous.

1. **Naias**. Pistils solitary and naked. Stigmas 2 or 4
2. **Zannichellia**. Pistils about 4 from a cup-shaped involucre or sheath.
3. **Zostera**. Pistils and anthers alternately sessile in two rows on one side of a linear spadix enclosed in a leaf. Stigmas 2.

* * Flowers perfect, spiked or clustered.

4. **Ruppia**. Flowers naked (no perianth), each of 4 large anther-cells, and 4 ovaries.
5. **Potamogeton**. Flowers with sepals, stamens, and sessile ovaries each 4.

1. NAIAS, L. NAIAD.

Flowers diœcious or monœcious, axillary, solitary and sessile; the sterile consisting of a single stamen enclosed in a little membranous spathe: anther at first nearly sessile, the filament at length elongated. Fertile flowers consisting of a single ovary tapering into a short style: stigmas 2—4, awl-shaped: ovule erect, anatropous. Fruit a little seed-like nutlet, enclosed in a loose and separable membranous epicarp. Embryo straight, the radicular end downwards.—Slender branching herbs, growing under water, with opposite and linear leaves, somewhat crowded into whorls, spinulose-toothed, sessile and dilated at the base. Flowers very small, solitary, but often clustered with the branch-leaves in the axils; in summer. (*Naiás*, *water-nymph*; an ill-chosen name for these insignificant water-weeds.)

1. **N. mājor**, All. Leaves linear, rather broad, strongly repand-toothed, the back as well as the stem more or less beset with little spines, the sheathing base entire or nearly so; flowers diœcious; anther 4-celled, 4-valved. — New York, Onondaga Lake, *G. W. Clinton*; Lake Ontario, near Rochester, *C. M. Booth*: recent discoveries. (Eu.)

2. **N. flexilis**, Rostk. Leaves very narrowly linear and minutely serrate, as is their abrupt rounded sheathing base; flowers monœcious? (*N. Canadensis*, *Michx.* *Caulinia flexilis*, *Willd.*) — Ponds and slow streams: common. (Eu.)

2. ZANNICHÉLLIA, Micheli. HORNED PONDWEED.

Flowers monœcious, sessile, naked, usually both kinds from the same axil; the sterile consisting of a single stamen, with a slender filament bearing a 2-4-celled anther; the fertile of 2-5 (usually 4) sessile pistils in the same cup-shaped involucre, forming obliquely oblong nutlets in fruit, beaked with a short style, which is tipped by an obliquely disk-shaped or somewhat 2-lobed stigma. Seed orthotropous, suspended, straight. Cotyledon taper, bent and coiled up. — Slender branching herbs, growing under water, with opposite or alternate long and linear thread-form entire leaves, and sheathing membranous stipules. (Named in honor of *Zannichelli*, a Venetian botanist.)

1. **Z. palústris**, L. Style at least half as long as the fruit, which is flatish, somewhat incurved, even, or occasionally more or less toothed on the back (not wing-margined in our plant), nearly sessile; or, in var. *PEDUNCULATA*, both the cluster and the separate fruits evidently peduncled. — Ponds and slow streams: rather rare. July. (Eu.)

3. ZOSTÈRA, L. GRASS-WRACK. EEL-GRASS.

Flowers monœcious; the two kinds naked and sessile and alternately arranged in two rows on the midrib of one side of a linear leaf-like spadix, which is hidden in a long and sheath-like base of a leaf (spathe); the sterile flowers consisting of single ovate or oval 1-celled sessile anthers, as large as the ovaries, and containing a tuft of threads in place of ordinary pollen; the fertile of single ovate-oblong ovaries attached near their apex, tapering upward into an awl-shaped style, and containing a pendulous orthotropous ovule: stigmas 2, long and bristle-form, deciduous. Utricle bursting irregularly, enclosing an oblong longitudinally ribbed seed (or nutlet). Embryo short and thick (proper cotyledon almost obsolete), with an open chink or cleft its whole length, from which protrudes a doubly curved slender plumule. — Grass-like marine herbs, growing wholly under water, from a jointed creeping stem or rootstock, sheathed by the bases of the very long and linear, obtuse, entire, grass-like, ribbon-shaped leaves (whence the name, from *ζωστήρ*, a band).

1. **Z. marina**, L. Leaves obscurely 3-5-nerved. — Common in bays along the coast, in water of 5°-15° deep. Aug. (Eu.)

4. RÚPPIA, L. DITCH-GRASS.

Flowers perfect, 2 or more approximated on a slender spadix, which is at first enclosed in the sheathing spathe-like base of a leaf, entirely destitute of floral

envelopes, consisting of 2 sessile stamens, each with 2 large and separate anther-cells and 4 small sessile ovaries, with a single campylotropous suspended ovule: stigma sessile, depressed. Fruit of little obliquely-ovate pointed drupes, each raised on a slender stalk which appears after flowering; the spadix itself also then raised on an elongated thread-form peduncle. Embryo ovoid, with a short and pointed plumule from the upper end, by the side of the short cotyledon. — Marine herbs, growing under water, with long and thread-like forking stems, and slender almost capillary alternate leaves, sheathing at the base. Flowers rising to the surface at the time of expansion. (Dedicated to *H. B. Ruppis*, a German botanical author of the early part of the 18th century.)

1. **R. marítima**, L. Leaves linear-capillary; nut ovate, obliquely erect; fruiting peduncles capillary ($\frac{1}{2}$ '–1' long). — Shallow bays, along the whole coast: also Onondaga Lake (near salt springs), New York, *J. A. Paine*. Chiefly a narrowly leaved variety with strongly pointed fruit, approaching *R. rostellata*, *Koch*. June–Sept. (Eu.)

5. POTAMOGETON, Tourn. PONDWEED.

Flowers perfect. Sepals 4, rounded, valvate in the bud. Stamens 4, opposite the sepals: anthers nearly sessile, 2-celled. Ovaries 4 (rarely only one), with an ascending campylotropous ovule: stigma sessile or on a short style. Fruit drupe-like when fresh, more or less compressed: endocarp (*nutlet*) crustaceous. Embryo hooked, annular, or cochleate, the radicular end pointing downwards. — Herbs of fresh, or one in brackish, ponds and streams, with jointed mostly rooting stems, and 2-ranked leaves, which are usually alternate or imperfectly opposite; the submersed ones pellucid, the floating ones often dilated, and of a firmer texture. Stipules membranous, more or less united and sheathing. Spikes sheathed by the stipules in the bud, mostly raised on a peduncle to the surface of the water. (An ancient name, composed of *ποταμός*, a river, and *γείτων*, a neighbor, from their place of growth.)

The following account of the genus is contributed by DR. J. W. ROBBINS, of Uxbridge, Mass. — By *fruit*, the full-grown fresh or macerated fruit is intended; by *nutlet*, that with the fleshy outer portion or epicarp removed. All except No. 15 flower in summer: the month mentioned indicates the season of the maturation of the fruit, which, especially, should be collected.

§ 1. **DIVERSIFOLII**. *Leaves of two sorts; the floating ones coriaceous, and different in form from the more delicate submersed ones.*

* *Submersed leaves mostly with no distinction of blade and petiole, being in fact bladeless more or less flattened petioles, or phyllodia, sessile, grass-like, narrowly linear, or so attenuated as to become filiform or capillary.*

← *Stems rather stout: stipules free from the leaves: spikes all emersed, cylindrical and densely fruited: fruits fleshy and turgid, obliquely obovate: embryo coiled into less or scarcely more than one turn.*

1. **P. natans**, L. *Stem simple or sparingly branched; floating leaves all long-petioled, elliptical or ovate, somewhat cordate at base, obtuse but with a blunt point, 21–29-nerved; upper submersed leaves lanceolate, early perishing, the lower (later in the season) very slender (3'–7' long, barely a line wide);*

upper *stipules* very long, acute; peduncle about the thickness of the stem; spikes 1' - 2' long; sides of the turgid nutlet with a small deep impression in the middle; embryo-coiled into an incomplete elliptical ring. — Var. *PROLIXUS*, Koch. More slender; upper leaves lanceolate, frequently submersed, acute; stem (6° - 12°), submersed leaves (sometimes 21') and stipules very long. — Common in ponds and ditches: the var. in deep flowing water. Aug. (Eu.)

2. *P. Oakesianus*, Robbins, n. sp. Stem more slender, much branched; floating leaves smaller (1' - $1\frac{1}{2}'$ long), ovate- or oblong-elliptical, obtuse, fewer (17 - 23-) nerved; lowest submersed ones almost capillary (only $\frac{1}{4}''$ - $\frac{1}{2}''$ wide), continuing through the flowering season; spikes shorter ($\frac{3}{4}'$ - 1' long), on peduncles much thicker than the stem; fruit smaller and more acute; sides of the turgid nutlet not at all impressed; curvature of the embryo nearly circular, its apex directed to a point above its base. — Ponds, and especially pools and stagnant ditches, not rare in E. Massachusetts. Aug.

3. *P. Claytonii*, Tuckerman. Stems compressed, often simple from the creeping rootstocks; floating leaves chiefly opposite (1' - $1\frac{1}{2}'$ long), 11 - 17-nerved, oblong, tapering into a short petiole, the lower gradually narrowing and passing into the submersed ones, which are very numerous and approximate, 2-ranked, linear (2' - 5' long, and 1'' - $2\frac{1}{2}''$ wide), 5-nerved, the lateral nerves slender and nearly marginal, the space within the inner nerves coarsely cellular-reticulated; stipules very obtuse; spikes numerous, about the length of the thickened peduncle; fruit round-obovate, flattish, 3-keeled when dry; nutlet distinctly impressed on the sides; curvature of the embryo transversely oval. (*P. Nuttallii*, Chamisso?) — Still or flowing water, and even in small streams: common. — Perfectly distinct, and peculiar in the reticulation of the centre of the delicate grass-like submersed leaves. July, Aug.

+ + Characters of the preceding section, but all the parts small, slender and delicate as in the next, or as in No. 20 (and like it propagated by autumn buds), but with coriaceous floating leaves.

4. *P. Vaseyi*, Robbins, n. sp. Very delicate; stem almost capillary; floating leaves obovate (3'' - 5'' long) and about the length of their filiform petioles, with 5 nerves deeply impressed beneath, cross-veins distinct; submersed leaves filiform-linear, very attenuate (1' - 2' long, $\frac{1}{8}''$ - $\frac{1}{4}''$ wide) and acute; stipules not adnate, scarious; spikes all emersed and similar, few, interrupted-oblong, 3 - 5-flowered, on a thickish peduncle; fruit oblique round-obovate ($\frac{3}{8}''$ long) compressed, slightly sharp-margined, tipped with a distinct recurved style, the sides impressed and face acute; upper portion of the embryo circularly incurved, its apex transverse to the fruit. — Illinois, near Ringwood, McHenry Co., *Dr. G. Vasey*. Apparently also in Quinsigamond Lake, Mass.

+ + + Stems slender or filiform and much branched: floating leaves sometimes wanting: stipules adnate to the base of the leaf: spikes of two kinds: one emersed, cylindrical and many-flowered, on a club-shaped peduncle; the other submersed, globular and few-flowered: fruit flat, cochleate, with thin or scarcely any flesh, and a thin nutlet: embryo spiral.

5. *P. Spirillus*, Tuckerman. Floating leaves varying from oval to lance-oblong and lanceolate (the largest 10'' long and 4'' wide), usually obtuse,

about equalling the *rather dilated petioles*, with 5 - many nerves beneath deeply impressed; upper submersed leaves either with or without a lance-oblong or broad-linear proper blade; the numerous lower ones narrow-linear, tapering towards the obtuse apex ($3\frac{1}{2}' - 1\frac{1}{2}'$ long, $\frac{1}{4}'' - \frac{3}{8}''$ wide); stipules early lacerate; *submersed flowers usually solitary on very short erect peduncles*; *shell-shaped fruit with the back either winged and with 4-5 distinct teeth or wingless and entire*; *embryo coiled 1 $\frac{3}{4}$ turns*. — Rivers, and even far up small streams, in company with the preceding, rarely with the following, Maine (*J. Blake*) to Lake Superior and Virginia. June - Aug. — Stem less slender than the next.

6. **P. hybridus**, Michx. Floating leaves varying from oval to lance-oblong; (the largest 10'' long and 6'' wide), often acute, longer than the *filiform petioles*, with about 5 - 7 nerves beneath deeply impressed; submersed leaves very numerous, almost setaceous, (1' - 3' long, very rarely $\frac{1}{2}''$ wide); stipules obtuse; *submersed spikes 1-4-flowered, their peduncles (of their own length) frequently recurved*; *fruit minute, about 8-toothed on the margin*; *embryo coiled 1 $\frac{1}{2}$ turns*. (*P. diversifolius*, *Barton*. *P. setaceus*, *Pursh.*) — Shallow stagnant waters. June - Aug. — Emerged spikes 4'' to (in var. *spicatus*, *Engelm.*) 7'' long.

* * *Submersed leaves lanceolate, rarely oval or linear, membranaceous.*

7. **P. rufescens**, Schrader. Stem simple, floating leaves (often wanting, *P. obrutus*, *Wood*) rather thin, *wedge-oblancoate, narrowed into a short petiole*, 11 - 17-nerved; *submersed leaves almost sessile, lanceolate and lance-oblong, smooth on the margin, fewer-nerved*; stipules broad, hyaline, obtuse, upper ones acuminate; fruit obovate, lenticular, pitted when immature, with an acute margin and pointed with the rather long style; embryo incompletely annular. — Rivers and streams, New England to Pennsylvania, Illinois, and especially northward. — Aug. - Sept. — Spike often somewhat compound! (*Eu.*)

8. **P. lonchites**, Tuckerman. Stem often branching below; *floating leaves thinnish, lance-oblong or long-elliptical*, often acute, *long-petioled*, 17 - 23-nerved; *submersed leaves very long (3' - 12', by 2'' - 12'' wide), lanceolate and lance-linear, 7-15-nerved, coarsely reticulated*; peduncles somewhat thickened upwards; fruit obliquely obovate, obscurely 3-keeled when fresh and distinctly so when dry, the middle one winged above and sometimes with 3 - 5 shallow indentations; the rounded slightly curved face surmounted by the short style; nutlet with the sides scarcely impressed; upper part of the embryo circularly incurved. (*P. Americanus*, *Chumisso?*) — Rivers and ponds, New England to Illinois. Aug., Sept. — More slender than the very similar *P. fluitans* of Europe, its leaves longer petioled, the floating ones more abrupt at the base; stipules not bicarinate.

9. **P. pulcher**, Tuckerman. Stem simple, black-spotted; *leaves of three kinds*; floating ones becoming very large ($4\frac{1}{2}'$ by $3\frac{1}{2}'$) *roundish-ovate and cordate or ovate-oblong*, 25 - 37-nerved; *all alternate*; upper submersed ones (3 - 5) usually lanceolate, acute at the base and very long-acuminate, 10 - 15-nerved, very thin, cellular each side of the midrib, undulate, short-petioled; lowest (2 - 4 near the base of the stem) *thicker, plane, oval or oblong with a rounded base, or spatulate-oblong, on longer petioles*; stipules rather short and obtuse; peduncles thicker than the stem; fruit with a rounded back and angular face, pointed with the

style, distinctly 3-keeled when fresh, sharply so when dry; nutlet marked on the back by two deep furrows and in front by a sinus below the angle; sides flat; upper portion of the embryo circularly much incurved. — Ponds; found as yet only in Eastern Massachusetts and “in ponds on hills north of St. Louis,” Missouri, *Engelmann*: also Georgia, *Le Conte*. — July, Aug.

10. **P. amplifolius**, Tuckerman. Stems simple, of very variable length; floating leaves (sometimes wanting) large, oblong or lance-ovate, sometimes slightly cordate at base, abruptly acutish, 30–50-nerved, on rather long petioles; submersed leaves often very large (reaching 7' by 2'), lanceolate or oval, acute at each end, usually much recurved, undulate, mostly on short petioles; stipules very long and tapering to a point, soon becoming loose; peduncles thickened upward, in deep water much elongated; fruit rather obliquely obovate, the back rounded and bluntly keeled; nutlet slightly impressed on the sides; upper portion of the embryo incurved into a ring. — Ponds and large rivers: not rare. — Aug., Sept. — In very shallow water sometimes without membranaceous leaves, and in deep water it may have those only. — (Leaves on a radical shoot in one specimen with adnate stipules!)

11. **P. gramineus**, L. Stem slender, very branching below; floating leaves mostly thin, variable, but with a short blunt point, 9–15-nerved; submersed ones usually lanceolate or linear-lanceolate, acuminate or cuspidate, narrowed towards the base, about 7-nerved on the stem and 3-nerved on the branches; upper ones petioled, lower sessile; stipules obtuse, loose; peduncles somewhat thickened upwards; fruit small, roundish, compressed, scarcely keeled; upper portion of the embryo annular. (*P. heterophyllus*, *Schreber*.) — Still or flowing water: common. — Varies exceedingly in its submersed leaves, peduncles, and otherwise. The ordinary forms are: — Var. **GRAMINIFOLIUS**, Fries. Submersed leaves lance-linear, attenuate at each end, flaccid, sometimes more than 4' long by 3" wide; stem elongated. — Var. **HETEROPHYLLUS**, Fries. Lower leaves shorter, lanceolate, more rigid: the commonest form. The following are doubtfully referred to this species. (Eu.)

Var. ? **SPATHULÆFORMIS**. (*P. spathæformis*, *Tuckerman*, in herb.) Branches scattered; floating leaves obovate or oblong, with a larger point; submersed ones spatulate-oblong, obtuse, mucronate, sometimes recurved; spikes large and densely flowered. — Mystic Pond, near Boston, *Tuckerman*. The fruit is lacking to prove its rank.

Var. ? **MYRIOPHYLLUS**. Sending up from running rootstocks many short repeatedly dichotomous and densely leafy stems; fertile stems very slender; floating leaves small, delicate, lance-oblong, on long filiform petioles; submersed stem-leaves larger, early perishing; those of the branches (deep green) linear-oblong, very small ($\frac{3}{4}$ '–1' long), acute, sometimes minutely serrulate; spike slender, loosely-flowered, much shorter than the thickened peduncle. — Apponaug Pond, Rhode Island, without fruit.

§ 2. **CONFORMIFOLII**. Leaves all submersed and similar, mostly sessile, membranaceous and dilated, lanceolate, oblong, or oval. (Stipules obtuse, becoming loose.)

12. **P. lucens**, L. Stem thick, branching, sometimes very large; leaves more or less petioled, oval or lanceolate, mucronate, often rough-serrulate, frequently

shining; fruit roundish and compressed, with obtuse margins, slightly keeled; embryo circularly incurved above. — Ponds: not common. Aug., Sept. (Eu.)

Var. **minor**, Nolte. Smaller; upper leaves distinctly petioled and sometimes emersed, the others sessile, all usually numerous, undulate and shining.

Var. ? **Connecticutensis**. Stem flexuous; leaves all submersed, nearly sessile, lanceolate, acuminate, crisped, not shining; fruit larger, distinctly keeled; nutlet thick and hard. — Saltonstall's pond, East Haven, Connecticut, 1850.

13. **P. prælóngus**, Wulfen. Stem very long, branching, flexuous; leaves lance-oblong or lanceolate (sometimes 7' long), half-clasping, obtuse with a boat-shaped cavity at the extremity, thence splitting on pressure; stipules scarious, very obtuse; spikes rather loose-flowered; peduncles very long (sometimes reaching 20'); fruit obliquely obovate, compressed, sharply keeled when dry; style terminating the nearly straight face; curve of the embryo oval and longitudinal. — Ponds and large rivers, E. New England, and along the Great Lakes to Lake Superior. Sept., Oct. — Stem white: foliage bright green. (Eu.)

14. **P. perfoliátus**, L. Stem branching; leaves orbicular, ovate or lanceolate from a cordate-clasping base, usually obtuse and often minutely serrulate; peduncles short, cylindrical; fruit irregularly obovate, obtusely margined; embryo incurved in an oval. — Ponds and slow streams: common. Sept., Oct. (Eu.)

Var. **lanceolátus**. Larger; leaves long-lanceolate from a cordate-clasping base and acuminate, wavy, 3' to sometimes 4½' long; peduncles thickened upwards. — Along the Great Lakes. — This form seems peculiarly American.

15. **P. crispus**, L. Stem compressed; leaves linear-oblong, half-clasping, obtuse, serrulate, crisped-wavy, 3-nerved; fruit long-beaked; upper portion of the embryo incurved in a large circle. — Flowing and stagnant waters, Delaware, Penn., and New Jersey, Tatnall, Porter, Meehan. June, July. (Eu.)

§ 3. **ANGUSTIFOLIÆ**. Leaves all submersed and similar, mostly membranaceous and sessile, linear or setaceous. (No. 16, 17, and 20 are often gemmiparous, propagating by narrow terminal buds detached in autumn.)

* *Stipules free from the sheathing base of the leaf.*

16. **P. compréssus**, L. (ex Fries.) Stem branching, wing-flattened; leaves linear and grass-like (commonly 4' by 1½'), abruptly pointed, with many fine and 3 larger nerves; stipules (seen young) oblong, very obtuse; spikes cylindrical, 12-15-flowered, not half as long as the peduncle; fruit obliquely obovate, somewhat keeled and with slight teeth on the back, the sides not impressed, the face arching and terminated by the short style; summit of the large embryo lying transverse to the fruit. (*P. zosteræfolius*, Schumacher.) — Still and slow-flowing waters, New England to Penn. and Wisconsin: not common. Aug., Sept. (Eu.)

17. **P. obtusifólius**, Mertens & Koch. Stem flattened, very branching, leaves linear, tapering towards the base, obtuse and mucronate or very acute, 3- (rarely 5-) nerved; stipules elongated, very obtuse; spike ovate, continuous, 5-8-flowered, about the length of the peduncle; fruit oval, apiculate with the style, not keeled when fresh, upper portion of embryo coiled inward and lying transverse to the fruit. — Slow streams and ponds: very rare: Dillerville swamp, near Lancaster, Penn., Prof. Porter. Swamp of Beaver pond, near Central mine, and floating in Gratiot Lake, N. Michigan. Sept., Oct. (Eu.)

18. **P. Niagarënsis**, Tuckerman. *Stem flattened, very branching; leaves linear, very acute, mucronate and much attenuate at the nearly sessile base, 3-5-nerved, scarcely veiny (1½'-3' long and at most 1" wide); midrib below dilated; stipules acutish (sometimes 8" long), the numerous nerves early becoming bristles; peduncles short, club-shaped, compressed; spikes few, capitate, 8-12-flowered; fruit roundish, compressed, with a winged and toothed keel and angled face; "seed convolute-uncinate."* — Rapids above Niagara Falls, Tuckerman. Aug.

19. **P. pauciflorus**, Pursh. *Stem filiform, flattish and very branching; leaves narrow linear (1'-2' long and seldom ½" wide), acute, obscurely 3-nerved; stipules obtuse; spikes capitate, 1-4- usually 2-flowered, on short club-shaped peduncles; fruit roundish-lenticular; the back more or less crested; upper portion of the embryo incurved in a circle.* — Still or stagnant waters: common. Aug., Sept. — Its largest forms are approached by the preceding.

20. **P. pusillus**, L. *Stem slender, flattish or nearly cylindrical, branching; leaves narrow- or setaceous-linear, obtuse or acute, furnished with translucent glands on each side at the base; stipules at first obtuse; spikes interrupted or capitate, 2-8-flowered, on rather long peduncles; fruit obliquely elliptical, scarcely keeled; apex of embryo incurved and directed obliquely downwards.* — Pools and ditches: rather common, especially southward. — The principal forms are

Var. **màjor**, Fries. *Stem less branching; leaves broader (almost 1" wide), often 5-nerved; spikes interrupted.* (*P. mucronatus*, Schrader.) — This hardly passes into the following forms: rather rare. July. (Eu.)

Var. **vulgàris**, Fries. *Slender, very branching; leaves 3-nerved, often obtuse; spikes cylindrical and interrupted, or capitate and then but 1-3-flowered.* — A rare form (E. Mass.) has sometimes lanceolate floating leaves of the length of the petioles, with 5 nerves impressed beneath, as in *P. hybridus*. A Swedish specimen in Fries. Herb. Norm. exhibits the same in the following variety, though in a slighter degree. July, Aug. (Eu.)

Var. **tenuissimus**, Mertens & Koch. *Stem very slender and much branched; leaves almost setaceous, acute or cuspidate, obsolete 3-nerved; spikes interrupted or capitate.* — New England and New York: rather rare. July, Aug. — All three are rather sparingly furnished with reproductive buds: also the last two fruit freely, — the reverse of the fact in the following.

Var. ? **gemmaiparus**. *Stem filiform and very branching, leaves thicker, perfectly setaceous and usually exceedingly attenuate to the finest point, scarcely with a proper midrib; stipules long (½'-1'), obtuse; spikes very few, always interrupted, 3-6-flowered, long-peduncled; propagating buds very numerous; fruit wanting.* (*P. gemmaiparus*, Robbins in herb.) — Pools and slow-flowing waters: outlet of Mystic Pond, near Boston, Tuckerman; valley of the Blackstone from Worcester to Providence. — This plant is annual, propagated exclusively by its gemmæ, the fruit not maturing.

21. **P. Tuckermàni**, Robbins. *Very slender and delicate from a creeping rootstock, of a fine light green; stem filiform with several short and repeatedly dichotomous leaf-bearing branches; leaves thin and flat, but setaceous and tapering to near the fineness of a hair (1'-4' long and ¼" extreme width), obscurely 1-3-nerved, with a few coarse reticulations; stipules rather persistent below, ½' long, obtuse; peduncle solitary, very long, rather thickening upward; spike 4-8-*

flowered, in fruit continuous, oblong; fruit thick-lenticular, obscurely 3-keeled; nutlet slightly impressed on the sides; shell thick and hard; embryo nearly annular. — Cold ponds, White Mountains, New Hampshire, to Penn. June, July.

* * *Stipules united with the sheathing base of the leaf.*

22. **P. pectinatus**, L. *Stem filiform*, repeatedly dichotomous; leaves *se-taceous*, attenuate to the apex, 1-nerved with a few transverse veins; *spikes interrupted*, on long filiform peduncles; fruit obliquely broad-obovate, compressed, bluntly keeled; shell of nutlet very thick; embryo nearly annular. — Lake Champlain to Lake Superior, and along the coast, both in fresh and salt water. Aug. — Oct. (Eu.)

23. **P. Robbinsii**, Oakes. *Stem ascending* from a creeping base, *rigid*, very branching, *invested by the bases of the leaves and stipules*; leaves *crowded in two ranks, recurved-spreading*, narrow-lanceolate or linear (3' — 5' long and 2" — 3" wide), acuminate, *ciliate-serrulate* with translucent teeth, many-nerved; stipules obtuse when young, their nerves soon becoming bristles; *spikes numerous, loosely few-flowered*, on short peduncles. A single, rather immature fruit in Professor Tuckerman's herbarium, from Prof. Eaton, is obliquely obovate, acutish at the base, 3-keeled on the back, the middle keel winged, much arched on the thinner face, compressed and impressed on the sides, and apiculate with the rather long style; superior portion of the large embryo circularly incurved and pointing obliquely downwards. — Oozy bottoms of ponds and slow streams: common in New England: also in New York, Pennsylvania, and Ohio. Flowering in June and July. Mature fruit not yet seen.

(P. DÉNSUS, L. The plant upon which Schweinitz introduced this European species into the U. S. flora proves to be *Anacharis Canadensis*.)

ORDER III. ALISMACEÆ. (WATER-PLANTAIN FAMILY.)

Marsh herbs, with scape-like flowering stems, and perfect or monœcious flowers, not on a spadix, furnished with both calyx and corolla: sepals and petals each 3, distinct. Ovaries 3 — many, distinct or partly so, or if united separating at maturity, forming as many 1 — 2-seeded pods or achenia. Seed ascending or erect. Embryo without albumen. Stamens hypogynous, from 6 to many: anthers extrorse, 2-celled. Leaves sheathing at the base. — Comprises two very distinct suborders, viz. :—

SUBORDER I. JUNCAGINEÆ. ARROW-GRASS FAMILY.

Calyx and corolla colored alike (greenish). Carpels more or less united. Seed anatropous, with a straight embryo. Leaves petiole-like, without a blade. (Flowers perfect.)

1. **Triglochin**. Ovaries 3 — 6, united to the apex into one, but separating in fruit.

2. **Scheuchzeria**. Ovaries 3, almost distinct, forming diverging pods in fruit.

SUBORDER II. ALISMEÆ. WATER-PLANTAIN FAMILY.

Calyx green and persistent. Corolla white, deciduous. Ovaries numerous, distinct. Seed campylotropous: embryo bent double or hook-

shaped. Leaves or some of them commonly furnished with a blade. (Flowers perfect, monœcious, or sometimes diœcious.)

3. **Alisma**. Flowers perfect, with definite, mostly 6 stamens. Carpels flattened, whorled.

4. **Echinodorus**. Flowers perfect, with 6 - many stamens. Carpels capitate, ribbed.

5. **Sagittaria**. Flowers monœcious, rarely diœcious, with indefinite, rarely few stamens. Carpels capitate, flattened, winged.

1. TRIGLÒCHIN, L. ARROW-GRASS.

Sepals and petals nearly alike (greenish), ovate, concave, deciduous. Stamens 6: anthers oval, on very short filaments. Pistils united into a 3-6-celled compound ovary: stigmas sessile: ovules solitary. Pod splitting when ripe into 3-6 carpels, which separate from a persistent central axis. — Perennials, with rush-like, fleshy leaves, below sheathing the base of the wand-like naked and jointless scape. Flowers small, in a spiked raceme, bractless. (Name composed of *τρεις*, *three*, and *γλωχίϋ*, *point*, from the three points of the ripe fruit in No. 1 when dehiscent.)

1. **T. palústre**, L. *Scape* (6' - 18' high) *and leaves slender; fruit linear-club-shaped; the 3 carpels* when ripe separating from below upwards leaving a triangular axis, *awl-pointed at the base*. — Marshes, both fresh and brackish, New York to Illinois, and northward. Aug. (Eu.)

2. **T. marítimum**, L. *Scape* (12' - 20' high) *and leaves thickish, fleshy, fruit ovate or oblong, acutish, of 6 or rarely 5 carpels* which are rounded at the base and slightly grooved on the back; the edges acute. — Salt marshes along the coast, also salt springs in the interior, shore of the Great Lakes, and northward. — Var. **ELÁTUM** (*T. elatum*, *Nutt.*) grows in cold and fresh bogs, from W. New York to Wisconsin, often 2½' high, and has the angles of the carpels sharper, or almost winged. (Eu.)

2. SCHEUCHZÈRIA, L. SCHEUCHZERIA.

Sepals and petals oblong, spreading, nearly alike (greenish-yellow), but the latter narrower, persistent. Stamens 6: anthers linear. Ovaries 3, globular, slightly united at the base, 2-3-ovuled, bearing flat sessile stigmas, in fruit forming 3 diverging and inflated 1-2-seeded pods, opening along the inside. — A low bog-herb, with a creeping jointed rootstock, tapering into the ascending simple stem, which is zigzag, partly sheathed by the bases of the grass-like conduplicate leaves, and terminated by a loose raceme of a few flowers, with sheathing bracts. (Named for *John* and *John Jacob Scheuchzer*, distinguished Swiss botanists early in the 18th century.)

1. **S. palústris**, L. — Peat-bogs, New England to Pennsylvania, Illinois, and northward. July. (Eu.)

3. ALÍSMÀ, L. WATER-PLANTAIN.

Flowers perfect. Petals involute in the bud. Stamens definite, mostly 6. Ovaries many in a simple circle on a flattened receptacle, forming flattened coriaceous achenia, which are dilated and 2-3-keeled on the back. — Roots fibrous.

Leaves all from the root, several-ribbed, with connected veinlets. Scape with whorled paniced branches. Flowers small, white or pale rose-color. (The Greek name; of uncertain derivation.)

1. **A. Plantago**, L., var. **Americanum**. Root perennial; leaves long-petioled, ovate, oblong, or lanceolate, pointed, mostly rounded or heart-shaped at the base, 3-9-nerved; panicle loose, compound, many-flowered (1° - 2° long); carpels obliquely obovate, forming an obtusely triangular whorl in fruit. (*A. trivialis* and *parviflora*, *Pursh.*) — Shallow water. July-Sept. (Eu.)

4. ECHINODORUS, Richard, Engelmann.

Flowers perfect. Petals imbricated in the bud. Stamens 6-21 or more. Ovaries several or many, imbricated in a head, forming thick and ribbed achenia in fruit, often beaked with a projecting persistent style. — Habit intermediate between the preceding genus and the following. Fl. summer and autumn. (Name from *ἐχινώδης*, prickly, or from *ἐχίνοσ*, and *δορός*, a leathern bottle, applied to the ovary, which is in most species armed with the persistent style, so as to form a sort of prickly head of fruit.)

Genus elaborated for this work by DR. ENGELMANN.

1. **E. parvulus**, Engelm. Leaves lanceolate or spatulate, acute ($\frac{1}{2}$ '- $1\frac{1}{2}$ ' long, including the petiole); shoots often creeping and proliferous; scapes ($1'$ - $3'$ high) bearing a 2-8-flowered umbel; pedicels reflexed in fruit; stamens 9; styles much shorter than the ovary; achenia beakless, many-ribbed; root annual. — Margin of ponds, Michigan to Illinois and westward. — Flower $3''$ broad.

2. **E. rostratus**, Engelm. Leaves broadly heart-shaped, obtuse, nerved ($1'$ - $3'$ long, excluding the petiole); scape erect, longer than the leaves, bearing a branched panicle of proliferous umbels; stamens 12; styles longer than the ovary; achenia beaked, many-ribbed; root annual. (*Alisma rostrata*, Nutt.) — Swamps and ditches, Illinois and southward. — Plant from $3'$ to 2° high. Flower $5''$ wide. Head of fruit ovoid, $3''$ wide.

3. **E. radicans**, Engelm. Leaves somewhat truncately broadly heart-shaped, obtuse, nerved ($3'$ - $8'$ broad, long-petioled); stems or scapes prostrate, creeping (2° - 4° long), proliferous, bearing many whorls of flowers; stamens about 21; styles shorter than the ovary; achenia short-beaked, the keeled back denticulate. (*Alisma radicans*, Nutt.) — Swamps, Illinois and southward. — Flowers $6''$ - $9''$ in diameter.

5. SAGITTARIA, L. ARROW-HEAD.

Flowers monœcious, or often diœcious in No. 2, and polygamous in No. 3. Petals imbricated in the bud. Stamens indefinite, rarely few. Ovaries many, crowded in a spherical or somewhat triangular depressed head on a globular receptacle, in fruit forming flat membranaceous winged achenia. — Marsh or aquatic, perennial, stoloniferous herbs, with milky juice and fibrous roots; the scapes sheathed at the base by the bases of the long cellular petioles, of which the primary ones, and sometimes all of them, are flattened, nerved, and destitute of any proper blade (i. e. are phyllodia): when present the blade is arrow-shaped or lanceolate, nerved and with cross veinlets as in *Alisma*. Flowers

produced all summer, mostly whorled in threes, with membranous bracts; the sterile above. (Name from *sagitta*, an arrow, from the prevalent form of the leaves.)

Genus newly elaborated for this edition by DR. ENGELMANN.

* *Filaments narrow, as long as the linear-oblong anthers: scape simple or branched.*

1. **S. lancifolia**, L. Scape 2° - 5° high, with several of the lower whorls fertile; bracts ovate, acute or acuminate; pedicels slender, the fertile scarcely shorter than the sterile ones; filaments pubescent; achenia obovate-falcate, pointed with an incurved beak; leaves lanceolate or lance-oblong, rarely linear, all with a tapering base, thick or coriaceous (6' - 18' long and on a long and stout petiole, never sagittate), the nerves mostly arising from the very thick midrib. (*S. falcata*, Pursh, and Ed. 2.) — Swamps, Virginia and southward to the West Indies.

2. **S. variabilis**, Engelm. Scape ($\frac{1}{4}^{\circ}$ - 4° high) angled, with one or more of the lower whorls fertile; bracts mostly pointed; pedicels of the fertile flowers about half the length of the sterile ones; petals with white claws; filaments glabrous, nearly twice the length of the anthers; achenia broadly obovate, with a long and curved beak $\frac{1}{4}$ - $\frac{1}{2}$ its length; leaves very various, almost always sagittate. (*S. sagittifolia* Amer. auth. etc. — The European species has the fertile pedicels only $\frac{1}{2}$ or $\frac{1}{4}$ the length of the sterile; claws of the petals purpletinged; filaments not longer than the anthers; achenia almost orbicular, very broadly winged and with a short and straight beak.) — In water or wet places: very common. — Excessively variable in size and foliage: the following are the leading forms. — Var. **OBTUSA** (*S. obtusa*, Willd.) is large and diœcious; the broadly sagittate leaves obtuse, 6' - 12' long. — Var. **LATIFOLIA** (*S. latifolia*, Willd.) is large, monœcious, with broad and acute sagittate leaves. — Var. **HASTATA** is the ordinary form, with narrow halberd-shaped or sagittate leaves (including *S. hastata*, Pursh). — Var. **DIVERSIFOLIA**, with some leaves lanceolate or ovate-lanceolate, others more or less sagittate. — Var. **ANGUSTIFOLIA** has the narrow leaves with long and linear diverging lobes. — Var. **GRACILIS** (*S. gracilis*, Pursh) is the most slender form, with nearly linear leaves and lobes. — Var. **PUBESCENS** (*S. pubescens*, Muhl.): upper part of petiole and of scape and especially the orbicular-ovate obtuse bracts and sepals pubescent or woolly; leaves obtuse or acute; beak of fruit (as also sometimes in some of the other forms) horizontal, so that the fruit-head appears compact and smoothish, while usually it has a squarrose surface, from the protruding and recurved beak, New Jersey and southward. — A state with double flowers has been found in Pennsylvania and Delaware.

3. **S. calycina**, Engelm. Scape weak (3' - 9' high), at length mostly procumbent; usually only the lowest whorl fertile, with pedicels as long as those of the sterile flowers, recurved in fruit; bracts orbicular, obtuse or rarely pointed; *calyx appressed to the fruit-head and partly covering it*; filaments slightly rough, as long as the anthers; achenia obovate with a short horizontal style; leaves broadly halberd-shaped, obtuse or acutish, with wide spreading lobes, often wider than long, or lanceolate or sometimes reduced to linear phyllodia. — Maine to Delaware, Wisconsin, and southward. — Var. **SPONGIOSA**, with a loose or spongy texture and linear bladeless leaves submersed, occurs eastward;

Var. *FLÛITANS*, with lance-linear floating leaves, has been found in Missouri and westward; and Var. *GRÁNDIS*, with leaves 9' - 12' wide and 9' long, branched scape, and fruit-heads 9" diameter, grows farther south. — This species shows 9 - 12 stamens in the fertile, and some sterile pistils in the sterile flowers; and thus connects with *Echinodorus*.

* * *Filaments very short, with enlarged mostly glandular base: scape more simple.*

4. **S. heterophýlla**, Pursh. Scape weak (3' - 2° high), at length procumbent; bracts roundish, obtuse; flowers of the lowest whorl fertile and almost sessile; the sterile on long pedicels; filaments glandular-pubescent; achenia narrowly obovate with a long erect beak; leaves lanceolate or lance-oval, entire, or with one or two narrow basal sagittate appendages. — Rather common, at least southward. — Var. *ELLÍPTICA* has broad leaves (sometimes 6' long and 5' wide), either obtuse or cordate at the base, or sagittate. — Var. *RÍGIDA* (*S. rigida*, Pursh, on the Niagara and along the Great Lakes), the tallest form, has stout petioles and rigid narrowly lanceolate blades, acute at both ends. — Var. *ANGUSTIFÓLIA* has nearly linear leaves.

5. **S. graminea**, Michx. Scape very slender, erect (3' - 2° high); the lower whorls fertile; bracts rather obtuse and usually connate; pedicels all slender, the sterile and fertile of equal length; filaments glandular-pubescent; achenia small, narrowly obovate, almost beakless; leaves varying from ovate-lanceolate to linear or reduced to broad and acute phyllodia (when it is *S. acutifolia*, Pursh); scarcely ever sagittate. (*S. simplex* of Amer. authors; not of Pursh, whose plant of this name is a dioecious form of *S. variabilis*.) — Rather common, especially southward. — Flowers and fruit-heads smaller than in any of the foregoing; except in the var. *PLATYPHÝLLA*, which is found farther south, and has leaves 6' - 9' long and 3' - 4' wide; flowers 1' wide, on pedicels 1½' - 2' long.

6. **S. pusilla**, Nutt. Scape (1' - 3' high) weak, reclining in fruit; bract single, clasping; one or two whorls only, of which but a single flower is fertile, recurved in fruit; stamens about 7, with glabrous filaments; achenia obovate, with an erect beak and three notched dorsal ridges. (*Alisma subulata*, Pursh.) — Inundated shores, from Eastern New Jersey (*C. F. Austin*) and Philadelphia southward near the coast.

S. NATANS, Michx., closely allied to the last, is only found farther south; it is a larger plant with long phyllodia, or oval floating leaves, glabrous filaments, and obovate short-beaked achenia, with 5 - 9 crenate angles, — by which structure it is nearly connected with *Echinodorus*.

ORDER 112. **HYDROCHARIDACEÆ.** (FROG'S-BIT FAMILY.)

Aquatic herbs, with dioecious or polygamous regular flowers on scape-like peduncles from a spathe, and simple or double floral envelopes, which in the fertile flowers are united into a tube and coherent with the 1 - 3-celled ovary. Stamens 3 - 12, distinct or monadelphous: anthers 2-celled. Stigmas 3 or 6. Fruit ripening under water, indehiscent, many-seeded. Seeds ascending, without albumen: embryo straight.

Tribe I. STRATIOTIDÆ. Ovary 6-9-celled: stigmas 6-9.

1. **Limnobium.** Filaments 6-12, unequally united in the sterile flowers: anthers linear.

Tribe II. VALLISNERIÆ. Ovary 1-celled with 3 parietal placentæ: stigmas 3.

2. **Anacharis.** Stem leafy. Tube of the perianth of the fertile flowers thread-form.

3. **Vallisneria.** Stemless. Tube of the perianth not prolonged beyond the ovary.

1. LIMNÏBIUM, Richard. AMERICAN FROG'S-BIT.

Flowers diœcious, (or monœcious?) from sessile or somewhat peduncled spathes; the sterile spathe 1-leaved, producing about 3 long-pedicelled flowers; the fertile 2-leaved, with a single short-pedicelled flower. Calyx 3-parted or cleft; sepals oblong-oval. Petals 3, oblong-linear. Filaments entirely united in a central solid column, bearing 6-12 linear anthers at unequal heights: there are 3-6 awl-shaped rudiments of stamens in the fertile flowers. Ovary 6-9-celled, with as many placentæ in the axis, forming an ovoid many-seeded berry in fruit: stigmas as many as the cells, but 2-parted, awl-shaped (ovules orthotropous, *Torr.*).—A stemless perennial herb, floating in stagnant water, proliferous by runners, with long-petioled and round-heart-shaped leaves, which are spongy-reticulated and purplish underneath; rootlets slender, hairy. Sterile flowers rather small; the fertile larger; peduncle nodding in fruit. Petals white? (Name from *λιμόβιος*, *living in pools.*)

1. **L. Spóngia**, Richard. (*Hydrocharis Spongia*, *Bosc.* *H. cordifolia*, *Nutt.*)—Lake Ontario (*Dr. Bradley*, *Dr. Sartwell*), Illinois, *Dr. Vasey*, and in the Southern States. Aug.—Leaves 1'-2' long, faintly 5-nerved. Peduncle of the sterile flower about 3' long, thread-like; of the fertile, only 1', stout.

2. ANÁCHARIS (and ELODÆA), Richard. WATER-WEED.

Flowers polygamo-diœcious, solitary and sessile from a sessile and tubular 2-cleft axillary spathe. Sterile flowers small or minute; with 3 sepals, barely united at the base, and usually 3 similar or narrower petals: filaments short and monadelphous at the base, or none; anthers 3-9, oval. Fertile flowers either pistillate or apparently perfect: perianth extended into an extremely long and capillary tube; the limb 6-parted; the small lobes (sepals and petals) obovate, spreading. Stamens 3-9, sometimes merely short sterile filaments, without anthers, or with imperfect ones, sometimes with oblong anthers. Ovary 1-celled, with 3 parietal placentæ, each bearing a few orthotropous ovules; the capillary style coherent with the tube of the perianth: stigmas 3, large, 2-lobed or notched, exserted. Fruit oblong, coriaceous, few-seeded.—Perennial slender herbs, growing under water, with elongated branching stems, thickly beset with pellucid and veinless, 1-nerved, sessile, whorled or opposite leaves. The staminate flowers (which are rarely seen) commonly break off, as in *Vallisneria*, and float on the surface, where they expand and shed their pollen around the stigmas of the fertile flowers, which are raised to the surface by the excessively prolonged calyx-tube, varying in length according to the depth of the water. (Name formed of *άν*, *throughout*, and *ἀχαρίς*, *without charms*, being rather homely water-weeds.)

1. **A. Canadensis**, Planchon. Leaves in threes or fours, or the lower opposite, varying from linear to oval-oblong, minutely serrulate; stamens 9 in

the sterile flowers, 3 or 6 almost sessile anthers in the perfect flowers. (*Elodèa Canadensis*, *Michx.*, and *E. latifolia*, *Cuspary*, who has recently well illustrated this and the two related genera; all perhaps to be reduced to one, *Hydrilla*. *Udora Canadensis*, *Nutt.* *Anacharis Alsinastrum* (*Babington*), *Nuttallii*, and *Canadensis*, also *Apalánthe Schweinitzii*, *Planchon.*) — Slow streams and ponds: common. July. — Nat. in England, where it is very troublesome by its rapid increase, filling navigable waters; but no inconvenience of the sort is complained of here in its native country.

3. VALLISNÈRIA, Micheli. (TAPE-GRASS. EEL-GRASS.)

Flowers strictly dioecious: the sterile numerous and crowded in a head on a conical receptacle, enclosed in an ovate at length 3-valved spathe which is borne on a very short scape: stamens mostly 3. Fertile flowers solitary and sessile in a tubular spathe which is borne on an exceedingly long scape. Perianth (calyx) 3-parted in the sterile flowers; in the fertile with a linear tube coherent with the 1-celled ovary, but not extended beyond it, 3-lobed (the lobes obovate); also 3 linear small petals. Stigmas 3, large, nearly sessile, 2-lobed. Ovules very numerous, scattered over the walls, orthotropous. Fruit elongated, cylindrical, berry-like. — Stemless plants, with long and linear grass-like leaves, growing entirely under water. The staminate clusters being confined to the bottom of the water by the shortness of the scape, the flower-buds themselves spontaneously break away from their short pedicels and float on the surface, where they expand and shed their pollen around the fertile flowers, which are raised to the surface at this time: afterwards the thread-form fertile scapes (2–4 feet long, according to the depth of the water) coil up spirally, drawing the fruit under water to ripen. (Named for *Ant. Vallisneri*, an early Italian botanist.)

1. *V. spiràlis*, L. Leaves linear, thin, long and ribbon-like (1°–2° long), obscurely serrulate, obtuse, somewhat nerved and netted-veined. — Common in slow waters. Aug. (Eu.)

ORDER 113. BURMANNIACEÆ. (BURMANNIA FAMILY.)

Small annual herbs, often with minute and scale-like leaves, or those of the root grass-like; the flowers perfect, with a 6-cleft corolla-like perianth, the tube of which adheres to the 1-celled or 3-celled ovary; stamens 3 and distinct, opposite the inner divisions of the perianth; pod many-seeded, the seeds very minute. — A small, chiefly tropical family, of which only one species is found within our borders.

1. BURMANNIA, L. (TRIPTERÉLLA, *Michx.*)

Ovary 3-celled, with the thick placentæ in the axis. Filaments 3, very short. Style slender: stigma capitate-3-lobed. Pod often 3-winged. (Named for *J. Burmann*, an early Dutch botanist.)

1. *B. biflora*, L. Stem low and slender (2'–4' high), 2-flowered at the summit, or soon several-flowered; perianth (2''–3'' long) bright blue, 3-winged. (*Tripterella cærulea*, *Michx.*) — Peaty bogs, Virginia and southward.

ORDER 114. **ORCHIDACEÆ.** (ORCHIS FAMILY.)

Herbs, clearly distinguished by their perfect irregular flowers, with 6-merous perianth adnate to the 1-celled ovary, with innumerable ovules on 3 parietal placenta, and with either one or two gynandrous stamens, the pollen cohering in masses. Fruit a 1-celled 3-valved pod, with innumerable minute seeds, appearing like fine saw-dust. Perianth of 6 divisions in 2 sets; the 3 outer (*sepals*) mostly of the same petal-like texture and appearance as the 3 inner (*petals*). One of the inner set differs more or less in figure, direction, &c. from the rest, and is called the *labellum* or *lip*; only the other two taking the name of *petals* in the following descriptions. The lip is really the upper or posterior petal, i. e. the one next to the axis, but by a twist of the ovary of half a turn it more commonly is directed forward, as if anterior, i. e. is next the bract. Before the lip, in the axis of the flower, is the *column*, composed of a single stamen, or in *Cypripedium* of two stamens and a sterile rudiment of a third, variously coherent with or borne on the style or thick fleshy stigma; the anther 2-celled; each cell containing one or more masses of pollen (*pollinia* or *pollen-masses*). Stigma a broad glutinous surface, except in *Cypripedium*. — Perennials, often tuber-bearing or tuberous-rooted; some epiphytes. Leaves parallel-nerved, all alternate. Flowers often showy, commonly singular in shape, solitary, racemed, or spiked, each subtended by a bract, — in all arranged for fertilization by the aid of insects, very few capable of unaided self-fertilization. (See articles on Fertilization of Orchids, in *Sill. Jour.* 1862, 1863, &c.) — A vast family, but few in the United States.

I. Anther only one. (The 2 cells should not be mistaken for anthers!)

Tribe I. OPHRYDEÆ. Anther (of 2 separate cells) borne on and entirely adnate to the face of the stigma, erect or reclined. Pollen cohering into a great number of coarse grains, which are all fastened by elastic and cobwebby tissue into one large mass, and to a stalk that connects it with a gland or disk which was originally a part of the stigma. Flower in our species ringent, the lip with a spur beneath: one distinct gland to each pollen-mass.

1. **Orchis.** The two glands or viscid disks enclosed in a common pouch.
2. **Habenaria.** The two glands or disks naked (without any pouch or covering), either approximate or widely separated.

Tribe II. NEOTTIEÆ. Anther dorsal and erect or inclined, attached by its base only or by a short filament to the back or summit of the column, persistent. Pollen in our genera loosely cohering (mostly by some delicate elastic threads) in 2 or 4 soft masses, and soon attached directly to a viscous gland on the beak of the stigma.

3. **Goodyera.** Lip entire, free from the column, without callosities at the base.
4. **Spiranthes.** Lip ascending and embracing the column below, 2 callosities at the base.
5. **Listera.** Lip flat, spreading or pendulous, 2-lobed at the apex.

Tribe III. ARETHUSEÆ, MALAXIDEÆ, &c. Anther terminal and inverted (except in No 12) like a lid over the stigma, deciduous.

* Pollen powdery or pulpy, in 2 or 4 delicate masses: no gland.

6. **Arethusa.** Lip bearded, its base adherent to the linear column. Pollen-masses 4.
7. **Pogonia.** Lip more or less crested, free from the club-shaped column. Pollen-masses 2.

8. **Calopogon.** Lip bearded, stalked, free: column winged at the apex. Pollen-masses 4.
 * * Pollen in 4-8 smooth waxy masses,
 + Without stalks, attached directly to a large gland.
9. **Calypso.** Lip inflated and sac-like. Column winged and petal-like. Stem 1-flowered.
 + + With stalks to the 2 or 4 pollen-masses, connecting them with a gland. •
10. **Tipularia.** Lip short, flat, long-spurred beneath. Raceme many-flowered.
 + + + With stalks to the 8 pollen-masses, but no gland.
11. **Bletia.** Lip hooded, crested, spurless. Scape several-flowered.
 + + + + Without either stalks or glands to the 4 pollen-masses.
 ++ Plants green and with ordinary leaves. Sepals spreading.
12. **Microstylis.** Column minute, round: anther erect.
13. **Liparis.** Column elongated, margined at the apex: anther lid-like.
 ++ Plants tawny or purplish, leafless, or with a root-leaf only.
14. **Corallorhiza.** Perianth gibbous at base, or with a spur adherent to the ovary.
15. **Aplectrum.** Perianth not gibbous nor spurred at base. A green autumnal leaf.

II. Anthers two, or very rarely three.

Tribe IV. CYPRIPEDIÆ. The stamen which bears the anther in the rest of the order here usually forms a petal-like sterile appendage to the column. Pollen not in masses: no stalks nor gland.

16. **Cypripedium.** Lip an inflated sac. Anthers 2, one each side of the column.

1. ÓRCHIS, L. ORCHIS.

Flower ringent; the sepals and petals nearly equal, all of them (in our species) converging upwards and arching over the column. Lip turned downwards, coalescing with the base of the column, bearing a nectariferous spur at the base underneath. Anther-cells contiguous and parallel. Pollen cohering in numerous coarse waxy grains, which are collected on a cobweb-like elastic tissue into 2 large masses (one filling each anther-cell) borne on a slender stalk, the base of which is attached to a gland or sticky disk of the stigma, the two glands contained in a common little pouch or hooded fold, placed just above the orifice of the spur or nectary. Flowers showy, in a spike.—These glands stick fast to the proboscis of a butterfly or some such insect introduced into the nectar-bearing spur: when it flies to another flower, it drags out of the anther and carries with it the pollen-masses, and applies them to the stigma of the second or of several succeeding flowers, thus effecting cross-fertilization. (*Ὀρχίς*, the ancient name.)

1. **O. spectabilis, L.** (SHOWY ORCHIS.) Root of thick fleshy fibres, producing 2 oblong-ovate shining leaves (3' - 5' long), and a few-flowered 4-angled scape (4' - 7' high); bracts leaf-like, lanceolate; sepals and petals all lightly united to form the vaulted galea or upper lip, pink-purple, the ovate undivided lip white.—Rich woods, New England to Kentucky and (especially) northward. May.

2. HABENÀRIA, Willd., R. Br. REIN-ORCHIS.

Glands or viscid disks (to which the pollen-masses are attached) naked and exposed, separate, sometimes widely separated (becoming attached, some to the proboscis, others to the face or head of insects feeding upon the nectar of the

spur, the pollen thus carried from one blossom to another) : otherwise nearly as in true Orchis : the lateral sepals, however, mostly spreading. (Name from *habena*, a thong or rein, in allusion to the shape of the lip or spur of some species.) — PLATANThERA, named primarily for the species with the glands and bases of the anther-cells widely divergent, and GYMNADENIA, where these are approximate, are found to afford no wholly fixed or clear practical distinctions. Accordingly, in this edition, our species are restored to the older genus.

§ 1. GYMNADÈNIA, R. Br. *Cells of the anther parallel and approximate, their glands therefore contiguous. (Appendages of the stigma in our species two or three and much developed, oblong or club-shaped.)*

1. **H. tridentata**, Hook. Stem slender (6'–12' high), with a single oblong or oblanceolate obtuse leaf below, and 2 or 3 small ones like bracts above; spike 6–12-flowered, oblong; *flowers greenish or whitish, very small; lip wedge-oblong, truncate and with 3 short teeth at the apex*; the slender and slightly club-shaped spur curved upwards, longer than the ovary. (Orchis tridentata, Muhl. Gymnadenia tridentata, Lindl.) — Wet woods : common northward and along the Alleghanies. June, July. — Root of few fleshy fibres. Appendages of the stigma (sterile stamens?) three, oblong-club-shaped, one outside each orbicular gland and one between them, rising as high as the anther-cell, their cellular viscid summits receiving pollen in the unopened flower, and penetrated by pollen-tubes!

2. **H. integrà**, Spreng. Stem several-leaved (15' high), the 1 or 2 lower leaves elongated, oblong-lanceolate, acute; the others becoming smaller and bract-like; spike densely many-flowered, oblong-cylindrical; *flowers orange-yellow, small; lip ovate, entire or slightly crenulate or wavy, shorter than the awl-shaped descending spur*. (Orchis integrà & flava, Nutt. H. Ellióttii, Beck. Gymnadenia flava, Lindl.) — Wet pine barrens of New Jersey, Virginia, and southward. July. — Root of very fleshy fibres, one or two of them tuber-like. Appendages of the stigma two, lateral, oblong, fleshy : rostellum or middle appendage narrow.

§ 2. PERULÀRIA, Lindl. *Cells of the anther nearly parallel, separated by a broadish connective, narrowed towards the base, the margins of which are extended so as to form the sides of a deep oblong groove or cavity (more than semi-circular in cross-section), which is lined by the dilated orbicular and incurved gland. (Flowers small, greenish, slender-spurred.)*

3. **H. viréscens**, Spreng. Leaves ovate-oblong or oblong-lanceolate; the uppermost linear-lanceolate and pointed, passing into the bracts of the elongated raceme; petals ovate; flowers dull green; lip furnished with a tooth on each side and a strong nasal protuberance in the middle of the base, oblong, truncate-obtuse, about the length of the sepals, half the length of the slender club-shaped spur. (Orchis flava, L. ! but the flowers are not yellow. O. virescens, Muhl., Willd. O. fuscéscens, Pursh., not of L. O. herbiola, Pursh. O. bidentata, Ell. O. scutellata, Nutt. H. herbiola, R. Br. H. flava, Gray. Platanthera flava, Gray.) — Wet places : common. June, July. — Stem 10'–20' high; the spike at first dense, with the bracts longer than the flowers, at length

elongated and often loose, the upper bracts shorter than the flowers; which are quite small, and with scarcely a tinge of yellow, drying brownish. — The Siberian *H.* (*Perularia*) *fuscescens* is clearly distinct.

§ 3. **PLATANTHÈRA**, Richard. *Cells of the anther sometimes parallel, more commonly divergent, so that their tapering bases and the exposed glands are more or less distant. (Root a cluster of fleshy fibres, or tuberous-thickened.)*

* *Flowers greenish or white, small, numerous in a close spike: spur not longer than the entire or merely notched narrow lip: anther-cells almost parallel, wholly adnate: stem leafy.*

← *Spur short and sac-like: the 3 sepals and 2 narrow petals erect: glands small, rather widely separated. (PERISTYLUS, Lindl.)*

4. **H. viridis**, R. Br., var. **bracteata**, Reichenbach. Lower leaves obovate, the upper oblong and gradually reduced to lanceolate acute bracts 2–4 times the length of the green flowers; spike 10–30-flowered; lip oblong-linear or slightly spatulate, truncate and 2–3-toothed or lobed at the tip, more than twice the length of the spur. (*H. bracteata*, R. Br.) — Stem 6'–12' high. According to Mr. Darwin, in the common European *H. viridis* each gland is protected by a minute pouch: this is not yet verified in ours. — Damp woods, especially northward. (Eu.)

← ← *Spur slender, incurved, about as long as the entire lip: lateral sepals spreading.*

5. **H. hyperborea**, R. Br. Stem very leafy (6'–2° high) *leaves lanceolate, erect; spike dense (2'–15' long); lower bracts lanceolate, longer than the (greenish) flowers; lip and petals lanceolate, somewhat equal, the latter spreading from the base; anther somewhat overhanging the transversely dilated stigma; glands orbicular; stalk of the pollen-masses very slender and weak. (P. hyperborea, Huronensis, &c., Lindl.)* — Peat-bogs and wet cold woods: common northward. June, July. (Eu.)

6. **H. dilatata**, Gray. (That of Hook, Exot. Fl. is the preceding.) Resembles No. 5, but usually more slender, with narrower commonly *linear leaves; flowers white; lip lanceolate from a rhomboidal-dilated base, entire, its base with the bases of other petals and sepals erect-connivent, above spreading; anther-cells almost parallel; glands approximate, large and strap-shaped, vertical, nearly as long as the pollen mass and its short flat stalk together; stigma narrow; a trowel-shaped conspicuous beak (rostellum) between the bases of the anther-cells. (Orchis dilatata, Pursh.)* — Cold bogs: common northward. June, July.

* * *Flowers greenish or white and purple, few or several (5–15) in a loose spike, rather large for the size of the plant: scape or stem naked above, 1-leaved at the base (5'–9' high): spur not longer than the lip: anther-cells wholly adnate.*

7. **H. rotundifolia**, Richardson. Leaf varying from almost orbicular to oblong (1½'–3' long); *flowers rose-purple; or the lip white and spotted with purple, 3-lobed, and the larger middle lobe dilated and 2-lobed or strongly notched at the summit (4''–6'' long), exceeding the ovate-oblong petals and sepals, and the slender depending spur; anther-cells contiguous and parallel. (O. rotundifolia, Pursh.)* — Damp woods and bogs, N. Maine, Mr. Goodrich; Warren, Herkimer Co., New York, J. A. Payne; shore of L. Michigan in Wisconsin, Lewis Foote, and northward. June.

8. **H. obtusata**, Richardson. Leaf obovate or spatulate-oblong; flowers greenish-white; upper sepal very broad and rounded, the others and the petals lance-oblong; lip entire, linear or lanceolate, deflexed (3" long), about the length of the tapering and curving spur; anther-cells arcuate and widely separated. (*O. obtusata*, Pursh.) — Cold peat-bogs, &c., northeastern coast of Maine, and on mountains of New England and N. New York to Lake Superior (chiefly sub-alpine), and northward. June. (Eu.)

* * * Flowers white or greenish, numerous in a loose spike, on a naked scape, 2-leaved at the base: spur longer than the narrow entire lip: anther-cells widely diverging, their narrowed beak-like bases projecting forwards: base of the stalk of the pollen-mass laterally affixed by a short intermediate body to the back of the orbicular gland, the viscous face of which looks obliquely inward (the space between the two broad enough to receive the head of a butterfly).

9. **H. Hoókeri**, Torr. Leaves orbicular, spreading (3'–4' broad); scape mostly naked ($\frac{1}{2}^{\circ}$ – 1° high), bearing 10–20 upright sessile yellowish-green flowers in a strict spike; sepals ovate-lanceolate; lip lanceolate, pointed, incurved, longer than the lance-awl-shaped petals; spur slender, acute, about the length of the ovary (nearly 1' long). (*H. orbiculata*, Hook.) — Woods, Rhode Island to Penn., Wisconsin and northward. June. — Var. **OBLONGIFÓLIA**, J. A. Paine. Leaves oblong (3'–5' by $1\frac{1}{2}$ '–2'). Little Falls, New York, J. A. Paine.

10. **H. orbiculata**, Torr. Leaves very large (4'–8' wide), orbicular, spreading flat on the ground; scape bracted, bearing many spreading greenish-white flowers in a loose raceme; upper sepal orbicular, the lateral ovate; lip narrowly linear and slightly spatulate, obtuse, drooping, nearly thrice the length of the oblong-lanceolate and falcate obtuse petals; spur curved, slender (about $1\frac{1}{2}$ ' long), gradually thickened towards the blunt apex, twice the length of the ovary; anther-cells strongly projecting at the free beak-like base (the glands nearly $\frac{1}{4}$ ' apart). (*H. macrophylla*, Hook. *Orchis orbiculata*, Pursh.) — Rich woods, especially of Coniferæ, Maine to Pennsylvania and Lake Superior, and southward along the Alleghanies. July. — Leaves very smooth, shining above, silvery underneath. Scape 1° – 2° high.

* * * * (FRINGED ORCHIS.) Flowers several or many in an open spike, with mostly foliaceous bracts: stem (rather tall) leafy: spur thread-shaped or scarcely club-shaped, longer than the fringed, cleft, or dissected lip: anther-cells widely separated and usually diverging, their narrow beak-like bases, supported by the arms of the stigma, strongly projecting forwards or partly upwards: base of the stalk of the pollen-mass mostly affixed more or less laterally to the gland.

+ Lip pectinately fringed but undivided: flowers golden yellow or white: anther-cells widely divergent and beak-pointed, the orbicular glands as if raised on a tentacle much projecting forwards or slightly inwards: ovary long, tapering to the summit.

11. **H. cristata**, R. Br. Lower leaves lanceolate, elongated; the upper gradually reduced to sharp-pointed bracts, nearly the length of the crowded (yellow) flowers; spike oblong or cylindrical; petals rounded, crenate; lip ovate, with a lacerate-fringed margin, scarcely shorter than the slender obtuse incurved spur, which is not half the length of the ovary. — (*O. cristata*, Michx.) — Bogs, Penn. (Pursh) to Virginia and southward. July. — Flowers only a quarter as large as in the next.

12. **H. ciliaris**, R. Br. (YELLOW FRINGED-ORCHIS.) Leaves oblong or lanceolate; the upper passing into pointed bracts, which are shorter than the ovaries; spike oblong, rather closely many-flowered; flowers bright orange-yellow; lateral sepal rounded, reflexed; petals linear, cut-fringed at the apex; lip oblong, about half the length of the spur, furnished with a very long and copious capillary fringe. (*O. ciliaris*, L.) — Wet sandy places, New England to Michigan, and especially southward: rare north of New Jersey. July–Sept. — Our handsomest species, $1\frac{1}{2}^{\circ}$ – 2° high, with a short spike of very showy flowers; the lip $\frac{1}{2}$ ' long, the conspicuous fringe fully $\frac{3}{4}$ ' long on each side.

13. **H. blephariglóttis**, Hook. (WHITE FRINGED ORCHIS.) Leaves, &c. as in the last; flowers white; petals spatulate, slightly cut or toothed at the apex; lip ovate- or lanceolate-oblong, with the irregular capillary fringe of the margins usually shorter than its disk, one third the length of the spur. — Var. HOLOPÉTALA (*Platanthera holopetala*, Lindl.) has narrower petals with the toothing obsolete, and the lip less fringed. — Peat-bogs and borders of ponds, with the preceding, commonly taking its place in the northward. July. — A foot high, the flowers beautiful, but rather smaller than in the last.

← ← (GREENISH FRINGED-ORCHIS.) Lip 3-parted above the stalk-like base, the divisions cut into capillary fringes: flowers greenish- or yellowish-white: anther-cells not very divergent, the beaked bases, supported on the upper edge of the broad arms of the stigma, projecting forwards; the large glands oval or lanceolate, transverse, nearly facing each other: ovary short-tapering at the summit: the long spurs gradually thickened downward.

14. **H. leucophæa**. Leaves oblong-lanceolate; the bracts similar, rather shorter than the (large) flowers; spike commonly elongated, loose; petals obovate, minutely cut-toothed; divisions of the lip broadly wedge-shaped or fan-shaped, many-cleft to the middle into a copious thread-like fringe; spur longer than the ovary ($1'$ – $1\frac{1}{2}'$ long); glands transversely oval. (*Orchis leucophæa*, Nutt.) — Moist meadows, Ohio to Wisconsin and southwestward. July. — Stem 2° – 4° high. Lip $7''$ – $10''$ long.

15. **H. lácera**, R. Br. (RAGGED FRINGED-ORCHIS.) Leaves oblong or lanceolate; raceme loosely many-flowered; petals oblong-linear, entire; divisions of the lip narrow, deeply parted into a few long nearly capillary lobes; spur about the length of the ovary; glands lanceolate, as long as the stalk of the pollen-mass. (*O. psychodes*, Muhl., &c., not of L. *Platanthera psychodes*, Lindl. *O. lacera*, Michx.) — Bogs and moist thickets: common. July.

← ← ← (PURPLE FRINGED-ORCHIS.) Lip fan-shaped, 3-parted above the stalk-like base, the dilated divisions erosely fringed: flowers purple: anther-cells widely separated, but little divergent, their tapering bases (supported as in the preceding) strongly projecting, the orbicular glands looking obliquely forwards and downwards: ovary contracted only at the summit: the long curving spur somewhat thickened downward.

16. **H. psychódes**, Gray. Leaves oblong or lanceolate, the uppermost passing into linear-lanceolate bracts; raceme cylindrical, densely many-flowered; lower sepals round-oval, obtuse; petals wedge-obovate or spatulate, denticulate above; divisions of the spreading lip broadly wedge-shaped, many-cleft into a short fringe.

(*O. psycodes*, *L.*! *O. fimbriata*, *Pursh*, *Bigelow*. *O. incisa* and *O. fissa*, *Muhl. in Willd.* *Platanthera fimbriata*, *Lindl.*)—Wet meadows and bogs: common. July, Aug.—Flowers short-pedicelled, crowded in a spike of 4'–10' in length, small, but very handsome, fragrant; lip short-stalked, barely $\frac{1}{2}$ ' broad and not so long; the middle lobe broadest and more closely fringed, but not so deeply cleft as the lateral ones.

17. **H. fimbriata**, R. Br. Lower leaves oval or oblong, the upper few, passing into lanceolate bracts; spike or raceme oblong, loosely-flowered; lower sepals ovate, acute; petals oblong, toothed down the sides; divisions of the pendent large lip fan-shaped, more fringed. (*O. fimbriata*, *Ait.*, *Willd.*, *Hook. Exot. Fl.* &c. *O. grandiflora*, *Bigelow.*)—Wet meadows, New England to Penn. and (chiefly) northeastward. June.—Flowers fewer, paler (or lilac-purple), and 3 or 4 times larger than those of the preceding; the more ample dilated lip $\frac{3}{4}$ ' to 1' broad, with a deeper and almost capillary crowded fringe, different-shaped petals, &c.

18. **H. peramœna**, Gray. Lower leaves oblong-ovate, the upper lanceolate; spike oblong or cylindrical, densely flowered; lower sepals round-ovate; petals rounded-obovate, raised on a claw; divisions of the large lip very broadly wedge-shaped, irregularly eroded-toothed at the broadly dilated summit, the lateral ones truncate, the middle one 2-lobed. (*Platanthera peramœna*, *Ed. 2.* *P. fissa*, *Lindl.* *O. fissa*, *Pursh*, not of *Muhl.*)—Moist meadows and banks, Penn. to Illinois, and southward along the Alleghanies. Aug.—Flowers large and showy, violet-purple; the lip paler and very ample, 8''–10'' long: its divisions minutely and variably toothed, or sparingly cut along the terminal edge, but not fringed.

3. GOODYERA, R. Br. RATTLESNAKE-PLANTAIN.

Lip sac-shaped, sessile, entire, and without callosities at the base. Otherwise as *Spiranthes*.—Root of thick fibres, from a somewhat fleshy creeping rootstock, bearing a tuft of thickish petioled leaves, usually reticulated with white veining. Scape, spike, and the greenish-white small flowers usually glandular-downy. (Dedicated to *John Goodyer*, an early English botanist.)

§ 1. Lip strongly saccate-inflated and with a short spreading or recurved tip: anther short, borne on a distinct filament attached to the back of the short column, blunt: gland-bearing tip or beak of the stigma very short.

1. **G. repens**, R. Br. Small (5'–8' high) and slender; leaves ovate, more or less white-reticulated (about 1' long); flowers several, in a loose 1-sided spike; lip with an ovate recurved tip; sepals ovate.—Woods, under evergreens: common northward and through the Alleghanies. Aug. (Eu.)

2. **G. pubescens**, R. Br. Larger; leaves strongly white-reticulated; scape 6'–12' high; the numerous crowded flowers not one-sided; tip of the globular lip very short: otherwise like the preceding, and too near it.—Rich woods, eastward and southward. July, Aug.

§ 2. Lip barely saccate below, tapering and its sides involute above: anther ovate and long-pointed, borne on the base of the very short proper column, which is continued above the stigma into a conspicuous long tapering awl-shaped gland-bearing beak.

Aspect of Goodyera: structure of the flower nearly of Spiranthes, but the lip without callosities.

3. **G. Menzièsii**, Lindl. Leaves ovate-oblong, acute (2'–3' long), less white-reticulated than the preceding, some not at all so; scape 9'–12' high; flowers rather numerous in a looser often 1-sided spike; flower-buds less pubescent, elongated-ovate and pointed; lip with the saccate-conduplicate lower portion gradually tapering into the narrow barely spreading summit. (*Spiranthes decipiens*, Hook.)—Woods, Western New York to Michigan (confounded with *G. pubescens*), Lake Superior, and far westward. July.

4. SPIRÁNTHEs, Richard. LADIES' TRESSES.

Flower somewhat ringent, oblique on the ovary; the sepals and petals all narrow, mostly erect or connivent, the three upper pieces sticking together more or less, the two lower covering the base of the lip. Lip oblong, short-stalked or sessile, the lower part involute around the column, and with a callous protuberance on each side of the base; the somewhat dilated summit spreading or recurved, crisped, wavy, or rarely toothed or lobed. Column short, oblique, bearing the ovate stigma on the front, and the sessile or short-stalked (mostly acute or pointed) 2-celled erect anther on the back. Pollen-masses 2 (one in each cell), ovate, each 2-parted from the base (or even again divided) into thin and tender plates of granular pollen united by elastic threads, their summits together soon adhering to the narrow boat-shaped viscid gland of the stigma, which is set in the slender or tapering thin process or beak terminating the column, and is carried away on the proboscis of insects visiting the flower, with the pollen, to be deposited upon the stigma of another flower. After the removal of the gland, the beak is left as a 2-toothed or 2-forked tip.—Roots clustered-tuberous: stem more or less naked above, leaf-bearing below or at the base. Flowers small (ours all white or greenish-white), bent horizontally, 1–3-ranked in a spike, which is commonly more or less spirally twisted (whence the name, from *σπίρα*, a coil or curl, and *ἄνθος*, flower.)

* *Flowers in 3 ranks, crowded in a close spike. (Leaves at the root and base of the stem present at the flowering season.)*

1. **S. latifolia**, Torr. Low; naked stem or scape 4'–9' high, smooth; leaves all next the base, oblong or lance-oblong (1'–3' long, 6''–8'' wide), 3–5-nerved, contracted into a sheathing base; spike narrow (1'–2' long), flowers small; lip quadrate-oblong, thin, wavy-crisped at the very obtuse or truncate apex, the small callosities at the base oblong, marginal and adnate for their whole length; gland and beak of the stigma short. (*S. plantaginea*, Torr. in *N. Y. Fl.*, not of Lindl. *S. æstivalis*, Oakes, *Cat.*)—Moist banks, Vermont and N. New York to Michigan, Penn. (near Lancaster, *Prof. Porter*), and Delaware, *W. M. Canby*. June.—Perianth 2''–3'' long; lip yellowish on the face, not contracted in the middle nor the margins involute.

2. **S. Romanzoviàna**, Chamisso. Stem leafy below, and leafy-bracted above (5'–15' high); leaves varying from oblong-lanceolate to grassy-linear; spike dense, oblong or cylindrical (1'–4' long); perianth curved and the summit manifestly ringent, the sepals and petals all connivent in the upper portion or

galea; the *lip* ovate-oblong, contracted below the rounded wavy-crenulate much recurved summit, otherwise entire, the callosities at base globular and smooth; gland oblong-linear and the 2-horned *beak* of the stigma short. (*S. gemmípara*, Lindl. (*Neottia*, *Smith.*) *S. cernua*, in part, *Hook. & Ed. 2.*) — High and cool bogs, New York, from Herkimer and Otsego Co., to Lake Superior, and north-westward. July, Aug. — Perianth about 4'' long, pure white, smooth or smoothish. (Eu. Bantry Bay, Ireland, only.)

3. *S. cérnua*, Richard. Stem leafy below and leafy bracted above (6'–20' high); leaves linear-lanceolate, the lowest elongated (4'–12' long, 2''–4'' wide); spike cylindrical, rather dense (2'–5' long) and with the flowers either pubescent or nearly smooth; perianth horizontal or recurving, the lower sepals not upturned or connivent with the upper; lip oblong and very obtuse when outspread, but conduplicate or the margins much incurved, wavy-cripsed above the middle, especially at the flattish and recurved-spreading apex, the callosities at the base prominent, nipple-shaped, somewhat hairy; gland of the stigma linear, in a long and very slender beak. — Common in wet places, especially eastward and southward. Sept., Oct. — Very variable in size, foliage, &c.: the commoner form, with pure white sweet-scented flowers, often nearly losing its root-leaves at flowering-time: a variety in dry ground has greenish-cream colored stronger-scented flowers, and retains its root-leaves. Perianth 4''–5'' long.

* * *Flowers in one straight or spirally twisted rank.*

+ Stem bearing towards and at the base elongated leaves, which mostly persist during the flowering season.

4. *S. grámínea*, Lindl., Var. *Wálteri*. Stem 9'–2° high; lower and root-leaves linear or lance-linear (3'–8' long, 2''–4'' wide) gradually tapering to the base, the upper reduced to sheathing bracts; spike linear, dense (2'–5' long), usually much twisted, the axis, ovaries, &c. downy-pubescent; bracts ovate and gradually, or rhombic-ovate and abruptly taper-pointed, surpassing the ovary, the margins broadly hyaline; lip ovate-oblong when outspread, with rather small callosities at base, crisped at the rounded apex; anther and beak of the stigma very acute. (*Limòdorum præcox*, *Walt.* *Neóttia tórtilis*, *Pursh*, *Barton*, *Fl.*, &c. *S. tortilis*, *Chapm.*) — Wet, grassy places, S. New England to Virginia, and southward. July, Aug., at the north. — Root of fleshy or somewhat tuberous thickened fibres. Perianth 3'' long. — The original, West Indian *S. tortilis* (*Swartz*), *Richard*, has a smoother much less twisted spike, smaller bracts, and more leafless scape, the root-leaves seldom present at flowering-time: it is very like *S. brevifolia*, *Chapm.* (*S. longilábris*, *Lindl.*?). Our plant has a more acute tip to the anther and stigma than the Mexican.

+ + *Scape very slender, merely bracted; the leaves with a blade all in a cluster at the ground, ovate or oblong, abruptly contracted into a petiole, commonly withering away at or before flowering: flowers small, and whole plant gabrous or nearly so: bracts small, sharp-pointed, not longer than the pod.*

5. *S. grácilis*, Bigelow. Roots clustered, tuberous-thickened; scape 8'–18' high, bearing a slender many-flowered one-sided or twisted spike; lip oval when outspread, narrowly oblong in natural form, thickish and green above with thin white margins, the recurved obtuse or acutish apex wavy-cripsed, the callosities

at the base nipple-shaped. (Also *S. Beckii*, Lindl., at least as to the Northern plant.)—Hilly woods and sandy plains: common. July–Oct.—Perianth barely $1\frac{1}{2}$ "–2" long.

6. *S. simplex*, n. sp. Root a solitary oblong or spindle-shaped tuber; no leaves at flowering time; scape 5'–9' high, bearing a small narrow (rarely 1-sided) spike of very small flowers (perianth 1"– $1\frac{1}{2}$ " long); lip thin, white, obovate-oblong, the apex eroded and crisped, the callosities at the base slender.—Dry, sandy soil, E. Mass. (Nantucket, *Dr. Robbins*), New Jersey, and Delaware, *Wm. M. Canby*. Aug., Sept.—Spike 1'–3' long.

5. LÍSTERA, R. BROWN. TWAYBLADE.

Sepals and petals nearly alike, spreading or reflexed. Lip mostly drooping, longer than the sepals, 2-lobed or 2-cleft. Column wingless: stigma with a rounded beak. Anther borne on the back of the column at the summit, erect, ovate: pollen powdery, in 2 masses, joined to a minute gland.—Roots fibrous. Stem bearing a pair of opposite sessile leaves in the middle, and a spike or raceme of greenish or brownish-purple small flowers. (Dedicated to *Martin Lister*, an early and celebrated British naturalist.)

* Column very short. (Sepals ovate, reflexed: plants delicate, 4'–8' high.)

1. *L. cordata*, R. Brown. Leaves round-ovate, somewhat heart-shaped ($\frac{1}{2}$ '–1' long); raceme smooth; flowers minute, crowded, on pedicels not longer than the ovary; lip linear, twice the length of the sepals, 1-toothed on each side at the base, 2-cleft.—Damp cold woods; from Penn. northward. June, July. (Eu.)

2. *L. australis*, Lindl. Leaves ovate; raceme loose and slender; flowers very small, on minutely glandular-pubescent pedicels twice the length of the ovary; lip linear, 3–4 times the length of the sepals, 2-parted, the divisions linear-setaceous.—Damp thickets, New Jersey to E. Virginia and southward. June.

** Column longer, arching or straightish.

3. *L. convallarioides*, Hook. Leaves oval or roundish, and sometimes a little heart-shaped (1'– $1\frac{1}{2}$ ' long); raceme loose, pubescent; flowers on slender pedicels; lip wedge-oblong, 2-lobed at the dilated apex, and 1-toothed on each side at the base, nearly twice the length of the narrowly lanceolate spreading sepals, purplish, $\frac{1}{2}$ ' long. (*Epipáctis convallarioides*, *Swartz.*)—Damp mossy woods, along the whole Alleghany Mountains, to Penn., also Maine to Lake Superior, and northward.—Plant 4'–9' high.

6. ARETHÛSA, GRONOV. ARETHUSA.

Flower ringent; the lanceolate sepals and petals nearly alike, united at the base, ascending and arching over the column. Lip dilated and recurved spreading towards the summit. Column adherent to the lip below, petal-like, dilated at the apex. Anther lid-like, terminal, of 2 approximate cells: pollen-masses powdery-granular, 2 in each cell.—Beautiful low herbs, consisting of a sheathed scape from a globular solid bulb, terminated usually by a single large rose-purple flower. Leaf solitary, linear, nerved, hidden in the sheaths of the scape, protruding after flowering. (Dedicated to the Nymph *Arethusa*.)

1. **A. bulbosa**, L. Flower single, erect, with an entire lip recurved at the apex and bearded-crested down the face.—Bogs, Virginia to Maine, N. Wisconsin, and northward: rather scarce or local. May.—Flower 1'–2' long, very handsome, bright rose-purple; very rarely a pair of flowers.

7. POGONIA, Juss. POGONIA.

Flower irregular, the sepals and petals separate. Lip crested or 3-lobed. Column free, elongated, club-shaped, wingless. Anther terminal and lid-like, stalked: pollen-masses 2 (one in each cell), powdery-granular. (*Πογωνίας*, bearded, from the lip of some of the original species.)

§ 1. *Sepals and petals nearly equal and alike, pale rose-color, sometimes white.*

1. **P. ophioglossoides**, Nutt. Root of thick fibres; stem (6'–9' high) bearing a single oval or lance-oblong leaf near the middle and a smaller one or bract near the terminal flower, rarely one or two others with a flower in their axil; lip spatulate below, appressed to the column, beard-crested and fringed.—In bogs. June, July.—Flower 1' long, sweet-scented.—An interesting monster of this, with two additional lips, and some other petaloid parts, was found in Herkimer Co., New York, by *J. A. Paine*.

2. **P. pendula**, Lindl. Stem (3'–8' high) from oblong tubers, bearing 3 to 7 alternate ovate-clasping very small (3''–6'') leaves, the upper 1–4 with drooping flowers in their axils on slender pedicels; lip spatulate, somewhat 3-lobed, roughish or crisped above, crestless. (*Triphora pendula*, Nutt.)—Damp woods: rather scarce. Aug.—Perianth $\frac{1}{2}$ ' long, narrow.

§ 2. *Sepals linear, dingy or brownish, longer and much narrower than the erect or connivent petals: lip 3-lobed at the apex, crested down the middle, beardless: flowers solitary (or rarely a pair), terminal: root a cluster of fibres.*

3. **P. divaricata**, R. Br. Stem (1°–2° high) bearing a lanceolate leaf in the middle, and a leafy bract next the flower, which is recurved on the ovary; but the sepals ascending or diverging, spatulate-linear, longer than the lanceolate-spatulate pointed and flesh-colored petals, these about 1'–1 $\frac{1}{2}$ ' long.—Wet pine-barrens, Quaker Bridge, New Jersey (*W. H. Leggett*), Virginia, and southward. June, July.

4. **P. verticillata**, Nutt. Stem (6'–12' high), naked, except some small scales at the base, and a whorl of mostly 5 obovate or obovate-oblong sessile leaves at the summit; flower dusky purplish, on a peduncle longer than the ovary and pod; sepals more than twice the length of the petals, narrowly linear, spreading from a mostly erect base (1 $\frac{1}{2}$ '–2' long); lip with a narrow crest down the middle.—Low woods: rather rare, especially eastward. May, June.—Glaucous when young. Stalk of pod about 1 $\frac{1}{2}$ ' long, more than half the length of the leaves.

5. **P. affinis**, C. F. Austin, n. sp. Somewhat smaller than the preceding; leaves paler and rather narrower; flowers (not rarely in pairs) yellowish or greenish; peduncle much shorter than the ovary and pod; sepals not twice the length of the petals, tapering to the base; lip crested over the whole face and on the middle of the lobes.—With the last, which it nearly resembles, but is much rarer. Southern New York and Northern New Jersey, *C. F. Austin*; and Connecticut, near New Haven, *Edward Dana*.

8. CALOPÒGON, R. Br. CALOPOGON.

Flower with the ovary or stalk not twisting, therefore presenting its lip on the upper or inner side. Sepals and petals nearly alike, lance-ovate, spreading, distinct. Lip spreading, distant from the column, raised on a narrowed base or stalk, dilated at the summit, strongly bearded along the upper side. Column free, slender, winged at the apex. Anther terminal and lid-like, sessile: pollen-masses 4 (two in each cell), of soft powdery grains, lightly connected by delicate threads.—Scape from a small solid bulb, sheathed below by the base of the grass-like leaf, naked above, bearing several flowers. Bracts minute. (Name composed of *καλός*, *beautiful*, and *πάργων*, *beard*, from the bearded lip.)

1. **C. pulchéllus**, R. Br. Leaf linear; scape about 1° high, 2-6-flowered; lip as if hinged at the insertion, beautifully bearded towards the dilated summit with white, yellow, and purple club-shaped hairs.—Bogs: common. June, July.—Flowers 1' broad, pink-purple.

9. CALÝPSO, Salisb. CALYPSO.

Sepals and petals nearly similar, ascending, spreading, lanceolate, pointed. Lip larger than the rest of the flower, sac-shaped, inflated (9'' long), 2-pointed underneath the apex. Column broadly winged and petal-like, ovate, bearing the lid-like anther just below the apex: pollen-masses waxy, 2, each 2-parted, all sessile on a square gland.—A little bog-herb; the solid bulbs producing a single ovate or slightly heart-shaped thin leaf, as in *Aplectrum*, and a short (3'-5' high) scape, sheathed below, bearing a large and showy (variegated purple, pink, and yellow) flower. (Name from the goddess *Calypso*.)

1. **C. boreális**, Salisb.—Cold bogs and wet woods, the bulbs resting in moss, Northern New England to Michigan, and northward. May.—A very local and beautiful plant. Lip somewhat resembling that of a Lady's Slipper, woolly-hairy inside. (Eu.)

10. TIPULÀRIA, Nutt. CRANE-FLY ORCHIS.

Sepals and petals spreading, oblong; the latter rather narrower. Lip prolonged underneath into a thread-like ascending spur twice or thrice the length of the flower (9''-12'' long), 3-lobed; the middle lobe linear, a little wavy, as long as the petals, the side lobes short and triangular. Column narrow and wingless. Anther lid-like, terminal: pollen-masses 2, waxy, each 2-parted, connected by a linear stalk with the transverse small gland.—Herb with large solid bulbs connected horizontally, on a distinct petiole, producing in autumn a single ovate nerved and plaited leaf on a slender petiole, tinged with purple beneath; and in summer a long and naked slender scape (10'-18' high), with 1 or 2 sheaths at the base, bearing a many-flowered raceme of small greenish flowers tinged with purple. (Name from a fancied resemblance of the flowers to insects of the genus *Tipula*.)

1. **T. discolor**, Nutt. Lip blunt at the tip (which distinguishes it from a recently discovered Himalayan species).—Sandy woods, Massachusetts to Michigan and southward: very scarce. July.

11. **BLÉTIA**, Ruiz & Pavon. BLETIA.

Sepals spreading, equal, rather exceeding the petals. Lip hooded, hinged as it were with the column, crested along the upper face, often 3-lobed. Column half-cylindrical; the fleshy anther forming a lid at its apex. Pollen-masses 8, in pairs, with a stalk to each pair, waxy, becoming powdery. — Scape many-flowered from solid tubers. (Named for *Louis Blet*, a Spanish botanist.)

1. **B. aphylla**, Nutt. Leafless; scape (1° – 2° high) beset with purplish scales, the lower ones sheathing; flowers racemed, brownish-purple; lip not saccate. Rich woods, Kentucky and southward.

12. **MICRÓSTYLIS**, Nutt. ADDER'S-MOUTH.

Sepals spreading. Petals thread-like or linear, spreading. Lip auricled or ovate at the base, not tubercled, entire or nearly so. Column very small, terete, with 2 teeth or auricles at the summit and the erect anther between them. Pollen-masses 4, in one row (2 in each cell), cohering by pairs at the apex, waxy, without any stalks, threads, or gland. — Little herbs, from solid bulbs, producing simple stems or scapes, which bear in our species a single leaf, and a raceme of minute greenish flowers. (Name composed of *μικρός*, *little*, and *στυλίς*, *a column or style*.)

1. **M. monophyllos**, Lindl. Slender ($4'$ – $6'$ high); leaf sheathing the base of the stem, ovate-elliptical; *raceme spiked, long und slender; pedicels not longer than the flowers*; lip long-pointed. — Cold wet swamps, N. New England to Pennsylvania, Wisconsin, and northward. July. (Eu.)

2. **M. ophioglossoides**, Nutt. Leaf near the middle of the stem, ovate, clasping; *raceme short and obtuse; pedicels much longer than the flowers*; lip 3-toothed at the summit. — Damp woods: more common southward. July.

13. **LÍPARIS**, Richard. TWAYBLADE.

Sepals and petals nearly equal, linear, or the latter thread-like, spreading. Lip flat, entire, often bearing 2 tubercles above the base. Column elongated, incurved, margined at the apex. Anther lid-like, terminal: pollen-masses 4, in one row (2 to each cell), slightly united in pairs, without stalk, threads, or gland. — Small, low herbs, with solid bulbs, producing 2 root-leaves and a low scape, which bears a raceme of few purplish or greenish flowers. (Name from *λίπαρός*, *fat or shining*, in allusion to the smooth or unctuous leaves.)

1. **L. liliifolia**, Richard. Leaves ovate; petals thread-like, reflexed; *lip large* ($\frac{1}{2}'$ long) *wedge-obovate, abruptly short-pointed, brown-purplish*. (*Malaxis liliifolia*, Swartz.) — Moist woodlands: commonest in the Middle States. June.

2. **L. Læselii**, Richard. Leaves elliptical-lanceolate or oblong, keeled; *lip obovate or oblong* ($2''$ long), mucronate, *yellowish-green, shorter than the linear unequal petals and sepals*. (*Malaxis Correana*, Barton.) — Bogs, New England to Penn., Wisconsin, and northward: rare. June. (Eu.)

14. **CORALLORHIZA**, Haller. CORAL-ROOT.

Perianth somewhat ringent, oblique and gibbous or obscurely spurred at the base; the oblong or lanceolate sepals and petals nearly alike, the upper arching;

the lateral sepals ascending, their bases with that of the lip forming the gibbosity or short spur which is mostly adnate to the summit of the ovary: lip slightly adherent to the base of the 2-edged straightish column, bearing a pair of projecting ridges on the face below, spreading or recurved at the apex. Anther terminal, lid-like. Pollen-masses 4, obliquely incumbent, soft-waxy, free. Brownish or yellowish herbs, destitute of green foliage, with much-branched and toothed coral-like root-stocks (probably root-parasitical), sending up a simple scape, with sheaths in place of leaves, and bearing small and dull-colored flowers in a spiked raceme. (Name composed of *κοράλλιον*, *coral*, and *ρίζα*, *root*.)

§ 1. *Small spur or sac adnate to the summit of the ovary: flowers small: lip whitish or purplish, often mottled with crimson.*

1. **C. innata**, R. Brown. Plant slender, light brownish or yellowish (3' - 9' high), 5 - 12-flowered; pedicels very short; lip somewhat hastately 3-lobed above the base, the lamellæ thick and rather short; spur a very small protuberance; pod oval or elliptical (3'' - 4'' long). (*C. verna*, Nutt.)—Swamps and damp woods. May, June. — Perianth only 2' or 2½'' long. (Eu.)

2. **C. odontorhiza**, Nutt. Plant light brown or purplish; stem rather slender, bulbous-thickened at the base (6' - 16' high), 6 - 20-flowered; pedicels rather slender; lip entire, or merely denticulate, thin, broadly ovate or obovate, abruptly contracted into a claw-like base, the lamellæ a pair of short projections; the spur represented by a small cavity wholly adnate to the summit of the ovary; pod at first very acute at the base, at length short-oval (4'' long). (*C. Wistariana*, Conrad.)—Rich woods, New York to Michigan, and especially southward: rare northward. May, July. — Perianth about 3'' long.

3. **C. multiflora**, Nutt. Plant purplish, rather stout (9' - 18' high), 10 - 30-flowered; lip deeply 3-lobed, with a short narrowed base and with prominent lamellæ; spur manifest and protuberant; pod oblong (6'' - 9'' long), short-pedicelled. — Dry woods: common. July - Sept. — Perianth 2½'' - 4'' long.

§ 2. *Spur none: the broadly gibbous somewhat saccate base of the perianth wholly free from the ovary: flowers large for the genus, purple, unspotted, more expanding.*

4. **C. Macræi**, Gray. Plant purplish, stout (6' - 16' high), bearing 15 - 25 large flowers in a crowded spike, on very short pedicels; lip oval or obovate, perfectly entire, concave, barely narrowed at the base, where it bears 1 - 3 short lamellæ; all the parts of the perianth marked with 3 darker nerves; pod oblong (9'' long). — Woods, from near Lake Erie (Caledonia Springs, Canada, *W. F. Macrae*), Mackinaw, *C. G. Loring, Jr.*, *Prof. Whitney*, and westward to the Pacific. July. — Sepals, petals, and lip 6'' or 7'' long. — This is the *Aplectrum aphyllum*, Nutt. in herb.; and from the range and the size of the flowers it can hardly be other than *C. striata*, Lindl., but it does not at all agree with the character as to the lip and spur.

15. APLÉCTRUM, Nutt. PUTTY-ROOT. ADAM-AND-EVE.

Perianth neither gibbous, nor with any trace of a spur or sac at the base. Lip free, the palate 3-ridged. Otherwise the flowers and the scape (invested below with 3 greenish sheaths) as in *Corallorhiza*. But, instead of a coral-like

root, a slender naked rootstock produces each year a thick, globular, solid bulb or corm, often 1' in diameter (filled with exceedingly glutinous matter), which sends up late in summer a large, oval, many-nerved and plaited, petioled, green leaf, lasting through the winter, and early in the succeeding summer its scape, a foot or more high, is terminated by a loose raceme of dingy rather large flowers. (Genus too near the last. The name composed of a privative and *πλήκτρον*, a spur, from the total want of the latter.)

1. **A. hyemale**, Nutt. — Woods, in rich mould: rather rare or local. — Each corm lasts 2 or 3 years before it shrivels, so that 3 or 4 are found horizontally connected. Perianth greenish-brown, or the lip whitish, and somewhat speckled with purple, 5" – 6" long.

16. CYPRIPIEDIIUM, L. LADY'S SLIPPER. MOCCASON-FLOWER.

Sepals spreading; all three distinct, or in most cases two of them united into one under the lip. Petals spreading, resembling the sepals but usually narrower. Lip a large inflated sac. Column declined; on each side a fertile stamen, with its short filament bearing a 2-celled anther; the pollen loose and pulpy or powdery-granular; on the upper side a dilated-triangular, petal-like but thickish body, which answers to the fertile stamen of other Orchids, and covers the summit of the style; stigma terminal, broad, obscurely 3-lobed, moist and roughish (not smooth and viscid as in the rest of the order). Pollen in most of our species, especially in No. 6, exposed by the conversion of the face of the anther into a viscid, varnish-like film, which adheres to whatever touches it, carrying away some of the pollen. — Root of many tufted fibres. Leaves large, many-nerved and plaited, sheathing at the base. Flowers solitary or few, large and showy. (Name composed of *Κύπρις*, *Venus*, and *πόδιον*, a sock or buskin, i. e. *Venus's Slipper*.)

§ 1. *The three sepals separate. (Stem leafy, single-flowered.)*

1. **C. arietinum**, R. Brown. (RAM'S-HEAD L.) Upper sepal ovate-lanceolate, pointed; the 2 lower and the petals linear and nearly alike (greenish-brown), rather longer than the red and whitish veiny lip, which is prolonged at the apex into a short conical deflexed point; leaves 3 or 4, elliptical-lanceolate, nearly smooth. (*Cryosánthes*, Raf. *Arietinum*, Beck.) — Cold swamps and damp woods, Maine to New York, Wisconsin, and northward: rare. June. — The smallest species: stem slender, 6' – 10' high: lip only 6' long.

§ 2. *Two of the sepals united into one piece under the lip.*

* *Stem leafy to the top, 1 – 3-flowered: lip slipper-shaped or roundish, much inflated, horizontal, and with a rounded open orifice.*

+ *Sepals and linear wavy-twisted petals brownish, pointed, longer than the lip.*

2. **C. candidum**, Muhl. (SMALL WHITE LADY'S SLIPPER.) *Sepals ovate-lanceolate; lip white, flattish laterally, convex above; sterile stamen lanceolate; leaves lance-oblong, acute.* — Bogs, from central and New York (rare) to Kentucky and Wisconsin. May, June. — Little larger than the foregoing species, slightly pubescent, 1-flowered: petals and sepals greenish, not much exceeding the lip, which is less than 1' long.

3. **C. parviflorum**, Salisb. (SMALLER YELLOW L.) *Sepals ovate or ovate-lanceolate; lip flattish from above, bright yellow (1' or less long); sterile stamen triangular; leaves oval, pointed.* — Bogs and low woods; rather common. May, June. — Stem 1°–2° high. Flower fragrant: sepals and petals more brown-purple than in the next, into which, however, it seems to pass.

4. **C. pubescens**, Willd. (LARGER YELLOW L.) *Sepals elongated-lanceolate; lip flattened laterally, very convex and gibbous above (1½'–2' long, scentless, pale yellow.* — Bogs and low woods: common northward and westward, and southward in the Alleghanies. May, June. — Stem 2° high, pubescent, as are the broadly oval acute leaves.

+ + *Sepals and petals plane, rounded, white, not longer than the lip.*

5. **C. spectabile**, Swartz. (SHOWY L.) *Sepals round-ovate or orbicular, rather longer than the oblong petals; lip much inflated, white, pink purple in front (1½' long); sterile stamen heart-ovate.* — Peat-bogs, Maine and W. New England to Illinois, and southward along the Alleghanies. July. — The most beautiful of the genus, downy, 2° high. Leaves ovate, pointed.

* * *Scape naked, 2-leaved at the base, 1-flowered; sepals and petals greenish, shorter than the drooping lip, which has a closed fissure down its whole length in front.*

6. **C. acaulis**, Ait. (STEMLESS L.) *Sepals oblong-lanceolate, pointed, nearly as long as the linear petals; lip obovoid or oblong, rose-purple (rarely white), nearly 2' long, veiny; sterile stamen rhomboid; leaves oblong.* (*C. humile*, Salisb.) — Dry or moist woods, under evergreens: common, especially northward. May, June. — Plant downy: the scape 8'–12' high, with a green bract at the top.

ORDER 115. AMARYLLIDACEÆ. (AMARYLLIS FAMILY.)

Chiefly bulbous and scape-bearing herbs, not scurfy or woolly, with linear flat root-leaves, and regular (or nearly so) and perfect 6-androus flowers, the tube of the corolline 6-parted perianth coherent with the 3-celled ovary; the lobes imbricated in the bud. — Anthers introrse. Style single. Pod 3-celled, several–many-seeded. Seeds anatropous or nearly so, with a straight embryo in the axis of fleshy albumen. — An order represented in our gardens by the *Narcissus* (N. POETICUS), *Jonquil* (N. JONQUILLA), and *Daffodil* (N. PSEUDO-NARCISSUS), the *Snowdrop* (GALANTHUS NIVALIS) and the *Snowflake* (LEUCOJUM VERNUM), &c., but with very few indigenous representatives in this country. Bulbs acrid. Differs from Liliaceæ chiefly in the inferior ovary.

* Pod 3-valved, loculicidal: anthers versatile: perianth funnel-shaped.

1. **Amaryllis**. Flower naked in the throat: the tube short or none. Bulbs coated.
2. **Pancratium**. Flower with a slender tube and narrow recurved lobes; a cup-shaped crown connecting the stamens. Bulbs coated.
3. **Agave**. Flower equally 6-cleft, persistent; no crown. Fleshy-leaved, not bulbous.

* * Pod indehiscent; anthers sagittate.

4. **Hypoxys**. Perianth 6-parted nearly down to the ovary. Bulb solid.

1. **AMARÝLLIS**, L. § ZEPHYRÁNTHESES, Herb. AMARYLLIS.

Perianth funnel-form, from a tubular base; the 6 divisions petal-like and similar, spreading above; the 6 stamens inserted in its naked throat: anthers versatile. Pod membranaceous, 3-lobed. — Leaves and scape from a coated bulb. Flowers 1 or 2, from a 1-2-leaved spathe. (A poetical name.)

1. **A. Atamásco**, L. (ATAMASCO LILY.) Spathe 2-cleft at the apex; perianth white and pink; stamens and style declined. — Penn. (*Muhl.*) Virginia, and southward. June. — Flower 3' long, on a scape 6' high.

2. **PANCRÀTIUM**, L. PANCRATIUM.

Perianth with a long and slender tube, and an equal 6-parted limb; the lobes long and narrow, recurved; the throat bearing a tubular or cup-shaped corolline delicate crown, which connects the bases of the 6 exerted stamens. Anthers linear, versatile. Pod thin, 2-3-lobed, with a few fleshy seeds, often like bulb-lets. — Scapes and leaves from a coated bulb. Flowers large and showy in an umbel-like head or cluster, leafy-bracted. (Name composed of *πάν*, *all*, and *κράύς*, *powerful*, from fancied medicinal properties.)

1. **P. rotátum**, Ker. Leaves ascending, strap-shaped (1°-2° long); scape few-flowered; the handsome (white and fragrant) flower with a spreading large 12-toothed crown, the alternate teeth bearing the filaments. (*Hymenocállis rotata*, *Herbert.*) — Marshy banks of streams, Kentucky, Virginia, and southward. May. — Flowers opening at night or in cloudy weather.

3. **AGÀVE**, L. AMERICAN ALOE.

Perianth tubular-funnel-form, persistent, 6-parted; the divisions nearly equal, narrow. Stamens 6: anthers linear, versatile. Pod coriaceous, many-seeded. Seeds flattened. — Leaves thick and fleshy, often with cartilaginous or spiny teeth, clustered at the base of the many-flowered scape, from a thick fibrous-rooted crown. (Named altered from *άγανός*, *wonderful*, not inappropriate as applied to *A. AMERICANA*, the CENTURY-PLANT.)

1. **A. Virgínica**, L. (FALSE ALOE.) Herbaceous; leaves entire; scape simple (3°-6° high); the flowers scattered in a loose wand-like spike, greenish-yellow, fragrant. — Dry or rocky banks, Penn.? Virginia to Illinois (*Mr. Lumis*), and southward. Sept.

4. **HYPÓXYS**, L. STAR-GRASS.

Perianth persistent, 6-parted, spreading; the 3 outer divisions a little herbaceous outside. Stamens 6: anthers sagittate, erect. Pod crowned with the withered or closed perianth, not opening by valves. Seeds globular, with a crustaceous coat, ascending, imperfectly anatropous, the rhaps not adherent quite down to the micropyle, the persistent seed-stalk thus forming a sort of lateral beak. Radicle inferior! — Stemless small herbs, with grassy and hairy linear leaves and slender few-flowered scapes, from a solid bulb. (Name composed of *ύπό*, *beneath*, and *όξύς*, *sharp*, it is thought because the pod is acute at the base.)

1. *H. erécta*, L. Leaves linear, grass-like, longer than the umbellately 1-4-flowered scape; divisions of the perianth hairy and greenish outside, yellow within. — Meadows and open woods: common. June-Aug.

ORDER 116. HÆMODORACEÆ. (BLOODWORT FAMILY.)

Herbs, with fibrous roots, usually equitant leaves, and perfect 3-6-androus regular flowers, which are woolly or scurfy outside; the tube of the 6-lobed perianth coherent with the whole surface, or with merely the lower part, of the 3-celled ovary. — Anthers introrse. Style single, sometimes 3-partible; the 3 stigmas alternate with the cells of the ovary. Pod crowned or enclosed by the persistent perianth, 3-celled, loculicidal, 3-many-seeded. Embryo small, in hard or fleshy albumen. A small family; chiefly of the Southern hemisphere.

* Ovary wholly adherent to the calyx-tube: style filiform: seeds peltate, amphitropous.

1. *Lachnanthes*. Stamens 3, exserted: anthers versatile. Leaves equitant.

** Ovary free except at the base: style 3-partible: seeds anatropous.

2. *Lophiola*. Stamens 6, on the base of the woolly 6-cleft perianth. Leaves equitant.

3. *Aletris*. Stamens 6, in the throat of the warty-roughened and tubular 6-toothed perianth. Leaves flat.

1. LACHNÁNTHES, L. RED-ROOT.

Perianth woolly outside, 6-parted down to the adherent ovary. Stamens 3, opposite the 3 larger or inner divisions: filaments long, exserted: anthers linear, fixed by the middle. Style thread-like, exserted, declined. Pod globular. Seeds few on each fleshy placenta, flat and rounded, fixed by the middle. — Herb, with a red fibrous perennial root, equitant sword-shaped leaves, clustered at the base and scattered on the stem, which is hairy at the top, and terminated by a dense compound cyme of dingy yellow and loosely woolly flowers (whence the name, from *λάχνη*, *wool*, and *ἄθος*, *blossom*).

1. *L. tinctoria*, Ell. — Sandy swamps, from Rhode Island and New Jersey southward, near the coast. July-Sept.

2. LOPHIOLA, Ker. LOPHIOLA.

Perianth densely woolly, deeply 6-cleft; the divisions nearly equal, spreading, longer than the 6 stamens, which are inserted at their base. Anthers fixed by the base. Pod ovate, free from the perianth except at the base, pointed with the awl-shaped style, which finally splits into 3 divisions, one terminating each valve. Seeds numerous, oblong, ribbed, anatropous. — A slender perennial herb, with creeping rootstocks and fibrous roots, linear and nearly smooth equitant leaves; the stem leafless and whitened with soft matted wool towards the summit, as also the crowded or paniced cyme. Perianth dingy yellow inside; the lobes naked only towards the tip, each clothed with a woolly tuft towards the base (whence the name from *λοφῆιον*, *a small crest*.)

1. *L. aurea*, Ker. (*Conóstylis Americana*, Pursh.) — Boggy pine barrens, New Jersey to Virginia, and southward. June-Aug.

3. **ÁLETRIS**, L. COLIC-ROOT. STAR-GRASS.

Perianth cylindrical, not woolly, but wrinkled and roughened outside by thickly-set points, which look like scurfy mealiness, the tube cohering below with the base only of the ovary, 6-cleft at the summit. Stamens 6, inserted at the base of the lobes: filaments and anthers short, included. Style awl-shaped, 3-cleft at the apex: stigmas minutely 2-lobed. Pod ovate, enclosed in the roughened perianth; the dehiscence, seeds, &c. nearly as in *Lophiola*. — Perennial and smooth stemless herbs, very bitter, with fibrous roots, and a spreading cluster of thin and flat lanceolate leaves; the small flowers in a wand-like spiked raceme, terminating a naked slender scape (2° – 3° high). Bracts awl-shaped, minute. (*Ἄλετρις*, a female slave who grinds corn; the name applied to these plants in allusion to the apparent mealiness dusted over the blossoms.)

1. **A. farinosa**, L. Flowers oblong-tubular, white; lobes lanceolate-oblong. — Grassy or sandy woods: not rare. July, Aug.

2. **A. aurea**, Walt. Flowers bell-shaped, yellow (fewer and shorter); lobes short-ovate. — Barrens, New Jersey to Virginia, and southward. July.

ORDER 117. **BROMELIACEÆ**. (PINE-APPLE FAMILY.)

Herbs (or scarcely woody plants, nearly all tropical), the greater part epiphytes, with persistent dry or fleshy and channelled crowded leaves, sheathing at the base, usually covered with scurf; 6-androus; the 6-cleft perianth adherent to the ovary in the PINE-APPLE, &c., or free from it in

1. **TILLÁNDZIA**, L. LONG MOSS.

Perianth plainly double, 6-parted; the 3 outer divisions (sepals) membranaceous; the 3 inner (petals) colored; all convolute below into a tube, spreading above, lanceolate. Stamens 6, hypogynous! or the alternate ones cohering with the base of the petals: anthers introrse. Ovary free: style thread-shaped: stigmas 3. Pod cartilaginous, 3-celled, loculicidally 3-valved; the valves splitting into an inner and an outer layer. Seeds several or many in each cell, anatropous, club-shaped, pointed, raised on a long hairy-tufted stalk, like a coma. Embryo small, at the base of copious albumen. — Scurfy-leaved epiphytes. (Named for *Prof. Tillands* of Abo.)

1. **T. usneoides**, L. (COMMON LONG MOSS or BLACK MOSS.) Stems thread-shaped, branching, pendulous; leaves thread-shaped; peduncle short, 1-flowered. — Dismal Swamp, Virginia, and southward; growing on the branches of trees, forming long hanging tufts. A characteristic plant of the Southern States, and barely coming within the limits of this work.

ORDER 118. **IRIDACEÆ**. (IRIS FAMILY.)

Herbs, with equitant 2-ranked leaves, and regular or irregular perfect flowers; the divisions of the 6-cleft petal-like perianth convolute in the bud in 2 sets, the tube coherent with the 3-celled ovary, and 3 distinct or monadelphous stamens, alternate with the inner divisions of the perianth and

opposite the stigmas, with extrorse anthers.—Flowers from a spathe of 2 or more leaves or bracts, usually showy. Style single: stigmas 3, opposite the cells of the ovary. Pod 3-celled, loculicidal, many-seeded. Seeds anatropous: embryo straight in fleshy albumen. Rootstocks, tubers, or corms mostly acrid.—Represented in gardens by CROCUS, GLADIOLUS, TIGRIDIA or TIGER-FLOWER, and by three genera in the wild state.

1. **Iris.** Outer divisions of the perianth recurved; the inner erect: stigmas petal-like.
2. **Pardanthus.** Perianth equally spreading: filaments nearly distinct: stigmas dilated.
3. **Sisyrinchium.** Perianth regular and equally spreading: filaments monadelphous to the top: stigmas thread-like.

1. IRIS, L. FLOWER-DE-LUCE.

Perianth 6-cleft; the tube more or less prolonged beyond the ovary; the 3 outer divisions spreading or reflexed; the 3 inner smaller and erect. Stamens distinct; the oblong or linear anthers sheltered under the overarching petal-like stigmas (or rather branches of the style, bearing the true stigma in the form of a thin lip or plate under their apex): most of the style connate with the tube of the perianth. Pod 3-6-angled, coriaceous. Seeds depressed-flattened.—Perennials, with sword-shaped or grassy leaves, and large showy flowers; ours all with creeping and more or less tuberous rootstocks. (*Ἴρις*, the *rainbow deified*, anciently applied to this genus on account of the bright and varied colors of the blossoms.)

* *Stems leafy and rather tall (1°-3° high), from thickened rootstocks, often branching: tube of the perianth shorter than the divisions, which are beardless and crestless, the erect inner ones (petals) much smaller than the outer.*

+ *Flowers violet-blue, variegated with greenish, yellowish or white, and purple-veined.*

1. **I. versicolor**, L. (LARGER BLUE FLAG.) Stem stout, angled on one side; leaves sword-shaped ($\frac{3}{4}$ ' wide); ovary obtusely triangular with the sides flat; flowers ($2\frac{1}{2}$ '-3' long) short-peduncled, the funnel-form tube shorter than the ovary; pod oblong, turgid, with rounded angles.—Wet places: common. May, June.

2. **I. Virginica**, L. (SLENDER BLUE FLAG.) Stem very slender, terete; leaves narrowly linear (2"-3" wide); flowers slender-peduncled ($1\frac{1}{2}$ '-2' long), the tube extremely short; ovary 3-angled, each side 2-grooved; pod sharply triangular. (*I. prismatica*, Pursh. *I. gracilis*, Bigel.)—Marshes, Maine to Virginia and southward, near the coast. June.

+ + *Flowers yellowish or dull reddish-brown.*

3. **I. cùprea**, Pursh. Stem and leaves as No. 1; tube of the perianth cylindrical, as long as the 6-angled ovary; petal-like branches of the style narrow.—Swamps near Cairo, S. Illinois (*Dr. Vasey*), and southward. May.

* * *Stems low (3'-6' high), from tufted and creeping slender (or here and there tuberous-thickened) rootstocks, 1-3-flowered: tube of the perianth long and slender; the violet-blue divisions nearly equal.*

4. **I. vérna**, L. (DWARF IRIS.) Leaves linear, grass-like, rather glaucous; the thread-like tube of the perianth about the length of the divisions, which

are oblong-obovate and on *slender claws*, the outer ones slightly hairy down the orange-yellow base, *crestless*; pod obtusely triangular. — Wooded hillsides, Virginia, Kentucky, and southward. April.

5. **I. cristata**, Ait. (CRESTED DWARF IRIS.) *Leaves lanceolate* (3' - 5' long when grown); those of the spathe *ovate-lanceolate*, shorter than the *thread-like tube of the perianth*; which is 2' long and *much longer than the light blue obovate short-clawed divisions*, the outer ones *crested* but beardless; pod sharply triangular. — Mountains of Virginia, Kentucky, and southward. May. — Creeping rootstocks pungently acrid.

6. **I. lacustris**, Nutt. (LAKE DWARF IRIS.) *Tube of the perianth rather shorter than the divisions* (yellowish, $\frac{1}{2}$ ' - $\frac{3}{4}$ ' long), *dilated upwards*, not exceeding the spathe: otherwise as in the last, and too near it. — Gravelly shores of Lakes Huron and Michigan. May.

2. PARDÁNTHUS, Ker. BLACKBERRY-LILY.

Perianth 6-parted almost to the ovary; the divisions widely and equally spreading, all nearly alike, oblong with a narrowed base, naked. Stamens monadelphous only at the base: anthers oblong. Style club-shaped, 3-cleft, the narrow divisions tipped with a small dilated stigma. Pod pear-shaped; the valves at length falling away, leaving the central column covered with the globose black and fleshy-coated seeds, imitating a blackberry (whence the popular name). — Perennial, with rootstocks, foliage, &c. of an Iris; the branching stems (3° - 4° high) loosely many-flowered; the orange-yellow perianth mottled above with crimson purple spots (whence the name from *πάρδος*, a leopard, and *ἄνθος*, a flower).

1. **P. chinensis**, Ker. (*Ixia chinensis*, L.) — Sparingly escaped from gardens into waste places. July - Sept. (Adv. from China, &c.)

3. SISYRÍNCIUM, L. BLUE-EYED GRASS.

Perianth 6-parted; the divisions alike, spreading. Stamens monadelphous to the top. Stigmas thread-like. Pod globular, 3-angled. Seeds globular. — Low slender perennials, with fibrous roots, grassy or lanceolate leaves, mostly branching 2-edged or winged stems, and fugacious umbelled-clustered small flowers from a 2-leaved spathe. (Name composed of *σῦς*, a hog, and *ῥύγχος*, snout, from a fancy that the hogs are fond of rooting it up.)

1. **S. Bermudiana**, L. Stem winged, naked, or 1 - 2-leaved; leaves narrow and grass-like; divisions of the perianth obovate, more or less notched at the end, and bristle-pointed from the notch. (Leaves of the spathe almost equal, shorter than the flowers.) — Var. **ANCEPS** (*S. anceps*, Cav.) has a broadly winged scape, and the outer leaf of the very unequal spathe longer than the flowers. — Var. **MUCRONATUM** (*S. mucronatum*, Michx.) has a slender and narrowly winged scape, very narrow leaves, those of the spathe sharp-pointed and unequal, one of them usually longer than the flowers. But there are various intermediate forms. — Moist meadows, &c., among grass; common everywhere. June - Aug. — Flowers small, delicate blue, changing to purplish, rarely whitish; or, in var. **ALBIDUM** (*S. albidum*, Raf.) pure white: Illinois, Kentucky, and westward.

ORDER 119. **DIOSCOREÆÆ.** (YAM FAMILY.)

Plants with twining stems from large tuberous roots or knotted rootstocks, and ribbed and netted-veined petioled leaves, small dioecious 6-androus and regular flowers, with the 6-cleft calyx-like perianth adherent in the fertile plant to the 3-celled ovary. Styles 3, distinct. — Ovules 1 or 2 in each cell, anatropous. Fruit usually a membranaceous 3-angled or winged pod. Seeds with a minute embryo in hard albumen. — Represented chiefly by the genus

1. **DIOSCOREËA**, Plumier. YAM.

Flowers very small, in axillary panicles or racemes. Stamens 6, at the base of the divisions of the 6-parted perianth. Pod 3-celled, 3-winged, loculicidally 3-valved by splitting through the winged angles. Seeds 1 or 2 in each cell, flat, with a membranaceous wing. (Dedicated to the Greek naturalist, *Dioscorides*.)

1. **D. villosa**, L. (WILD YAM-ROOT.) Herbaceous; leaves mostly alternate, sometimes nearly opposite or in fours, more or less downy underneath, heart-shaped, conspicuously pointed, 9–11-ribbed; flowers pale greenish-yellow, the sterile in drooping panicles, the fertile in drooping simple racemes. — Thickets, New England to Wisconsin, and common southward. July. — Stems slender, from knotty and matted rootstocks, twining over bushes. Pods 8"–10" long. — A bad name, for the plant is never villous, and often nearly smooth.

ORDER 120. **SMILACEÆ.** (SMILAX FAMILY.)

Shrubby or rarely herbaceous plants, climbing or supported by a pair of tendrils on the petiole of the ribbed and netted-veined simple leaves; with dioecious small flowers; regular perianth of 6 (rarely more) similar greenish deciduous sepals, free from the ovary; as many stamens as sepals, with introrse 1-celled anthers; ovary with 3 (rarely 1 or 4–6) cells and as many elongated spreading sessile stigmas. Ovules one or a pair in each cell, suspended, orthotropous. Fruit a small berry. Seed-coat close, firmly adherent to the hard horny albumen: embryo minute. — Order, as here limited, represented almost solely by the genus

1. **SMILAX**, Tourn. GREENBRIER. CATBRIER.

Character as above. — Flowers in umbels on axillary peduncles, greenish or yellowish. *Ster. Fl.* Stamens inserted on the very base of the sepals: filaments linear: anthers linear or oblong, fixed by the base, 2-locellate. *Fert. Fl.* Filaments, if present, sterile. Stigmas thick and spreading, almost sessile. (The ancient Greek name, of obscure meaning.)

§ 1. *Stems woody, often prickly: ovules solitary. (Ours all glabrous.)*

* *Leaves ovate or roundish, &c., most of them rounded or heart-shaped at the base, and 5–9-nerved, the three middle nerves or ribs stronger and more conspicuous.*

+ *Peduncles shorter or scarcely longer than the petioles, flattened; leaves thickish, inclining to be evergreen, at least southward, green both sides.*

1. **S. Wálteri**, Pursh. Stem low, somewhat angled, prickly near the base or unarmed; *leaves ovate and somewhat heart-shaped* (3'–4' long); *berries coral-red*. (S. China, Walt.)—New Jersey, and southward. July.

2. **S. rotundifolia**, L. (COMMON GREENBRIER.) Stem armed with scattered prickles, as well as the terete branches; branchlets more or less 4-angular; *leaves ovate or round-ovate*, often broader than long, slightly heart-shaped, abruptly short-pointed (2'–3' long); *berries blue-black*, with a bloom. (S. caduca, L., is only a more deciduous and thin-leaved form.)—Moist thickets: common, especially southward. June, July. — Plant yellowish-green, often high-climbing. — Passes into var. **QUADRANGULARIS**, with the branches, and especially the branchlets, 4-angular, often square. (S. quadrangularis, Muhl.) — Penn. to Illinois, and southward.

+ + *Peduncles longer than, but seldom twice the length of the petiole, flattened: leaves tardily deciduous or partly persistent: berries black, with a bloom.*

3. **S. glauca**, Walt. Terete branches and somewhat 4-angular branchlets armed with scattered stout prickles, or naked; *leaves ovate*, rarely subcordate, *glaucous beneath* and sometimes also above, as well as the branchlets when young (about 2' long), abruptly mucronate, the edges smooth and naked. (S. Sarsaparilla, L., in part, but not as to the syn. of Bauhin, whence the name was taken, (S. caduca, Willd. S. spinulosa, Smith? Torr. fl.) — Dry thickets, S. New York to Kentucky and southward. July.

4. **S. tamnoides**, L. Branches and the angular (often square) branchlets sparsely armed with short rigid prickles; *leaves varying from round-heart-shaped and slightly contracted above the dilated base to fiddle-shaped and halberd-shaped or 3-lobed, green and shining both sides*, cuspidate-pointed, the margins often somewhat bristly-ciliate or spinulose. (S. Bona-nox, L., S. hastata Willd., S. pandurata, Pursh, &c., are all forms of this.) — Thickets, New Jersey to Illinois, and (chiefly) southward. July.

+ + + *Peduncles 2–4 times the length of the petiole: leaves ample (3'–5' long), thin or thinnish, green both sides: berries black: stem terete and branchlets nearly so.*

5. **S. hispida**, Muhl. Rootstock cylindrical, elongated; stem (climbing high) below densely beset with long and weak blackish bristly prickles, the flowering branchlets mostly naked; *leaves ovate and the larger heart-shaped, pointed, slightly rough-margined, membranaceous and deciduous*. — Moist thickets, Penn. and W. New York to Michigan. June. — Peduncles 1½'–2' long. Sepals lanceolate, almost 3'' long.

6. **S. Pseudo-China**, L. Rootstock tuberous; stems and branches unarmed, or with very few weak prickles; *leaves ovate-heart-shaped, or on the branchlets ovate-oblong, cuspidate-pointed, often rough-ciliate, becoming firm in texture; peduncles flat (2'–3' long)*. — Dry or sandy soil, New Jersey to Kentucky, and southward. July.

* * *Leaves varying from oblong-lanceolate to linear, narrowed at the base into a short petiole, 3–5-nerved, shining above, paler or glaucous beneath, many of them*

without tendrils: peduncles short, seldom exceeding the pedicels, terete; the umbels sometimes panicked: branches terete, unarmed.

7. **S. lanceolata**, L. *Leaves thinnish, rather deciduous, ovate-lanceolate or lance-oblong; stigmas 3; berries dull red.* — Southeastern Virginia and southward. June.

8. **S. laurifolia**, L. *Leaves thick and coriaceous, evergreen, varying from oblong-lanceolate to linear (2½' - 5' long); stigma solitary; berries black when ripe, 1-seeded.* — Pine barrens of New Jersey to Virginia and southward. July, Aug.

§ 2. **COPROSMANTHUS**, Torr. *Stem herbaceous, never prickly: flowers carrion-scented: ovules mostly in pairs in each cell: leaves long-petioled, membranaceous, mucronate-tipped: berries bluish-black with a bloom.*

9. **S. herbacea**, L. (CARRION-FLOWER.) *Stem erect and recurving, or climbing; leaves ovate-oblong or rounded, mostly heart-shaped, 7 - 9-nerved, smooth; tendrils sometimes wanting; peduncles elongated (3' - 4' long, or sometimes even 6' - 8', and much longer than the leaves), 20 - 40-flowered.* — Var. **PULVERULENTA** (*S. pulverulenta*, Michx. & *S. peduncularis*, Muhl.) has the leaves more or less soft-downy underneath. A shorter-peduncled state of this appears to be *S. lasioneuron*, Hook. — Moist meadows and river-banks: common. June. — Very variable, 1° - 3°, or even 6° - 8° high: petioles 1' - 3' long. Seeds 6.

10. **S. tamnifolia**, Michx. *Stem upright or climbing; leaves heart-halberd-shaped, 5-nerved, smooth; peduncles longer than the petioles. (S. tamnoides, Pursh, not of L.)* — Pine barrens, New Jersey to Virginia and southward. — Leaves abruptly narrowed above the dilated heart-shaped base, tapering to the apex. Berry 2 - 3-seeded.

ORDER 121. LILIACEÆ. (LILY FAMILY.)

Herbs, or rarely woody plants, with regular and symmetrical almost always 6-androus flowers; the perianth not glumaceous, free from the chiefly 3-celled ovary; the stamens one before each of its divisions or lobes (i. e. 6, in one instance 4), with 2-celled anthers; fruit a few-many-seeded pod or berry; the small embryo enclosed in copious albumen. Seeds anatropous or amphitropous. Flowers not from a spathe, except in *Allium*; the outer and inner ranks of the perianth colored alike (or nearly so) and generally similar, except in *Trillium*. — A large family, as here extended, the principal divisions commonly received as orders, but not well limited. For the present purpose they are best regarded as tribes.

Tribe I. TRILLIDEÆ. *Styles or sessile stigmas 3, separate down to the ovary.*

Fruit a several-many-seeded berry. Divisions or leaves of the perianth distinct, the 3 outer often foliaceous. Simple stem from a rootstock, naked below; leaves all in one or two whorls above, broad, more or less conspicuously netted-veined!

1. **Trillium.** *Perianth of two sorts, 3 foliaceous persistent sepals and 3 colored petals. Anthers adnate, introrse. Leaves (3) in a single whorl. Flower single.*
2. **Medeola.** *Perianth of 6 similar colored divisions, deciduous. Anthers fixed by the middle, extrorse. Leaves in two whorls. Flowers several.*

Tribe II. MELANTHIEÆ. Styles or sessile stigmas 3, separate down to the ovary. Fruit a several-many-seeded pod. Seeds with a soft or loose coat. Anthers extrorse, except in No. 11. Perianth withering-persistent. Leaves parallel-veined and alternate. Flowers often polygamous, sometimes dioecious.

* Anthers heart-shaped or kidney-shaped, confluent 1-celled, shield-shaped after opening: pod 3-horned, and septicidal: seeds flat, membranaceous-margined.

+ Sepals with one or two glands or spots on the upper face near the base.

3. **Melanthium.** Flowers polygamous. Sepals entirely free from the ovary, their long claws bearing the stamens.
4. **Zygadenus.** Flowers perfect or polygamous. Sepals nearly free from the base of the ovary: stamens free.

+ + Sepals destitute of glands, and not raised on claws.

5. **Stenanthium.** Perianth below coherent with the base of the ovary; the sepals lanceolate, pointed, longer than the stamens. Racemes compound-panicled.
6. **Veratrum.** Perianth entirely free; the obovate or oblong sepals longer than the stamens. Flowers panicled, polygamous.
7. **Amianthium.** Perianth entirely free, the oval or obovate sepals shorter than the stamens. Flowers racemed, perfect.

** Anthers 2-celled, extrorse: pod loculicidal. Flowers racemed or spiked.

8. **Xerophyllum.** Flowers perfect. Cells of the globose-3-lobed pod 2-seeded. Leaves very slender. Seeds 2 in each cell.
9. **Helonias.** Flowers perfect. Cells of the globose-3-lobed pod many-seeded. Leaves lanceolate. Scape naked. Seeds numerous.
10. **Chamælrimum.** Flowers dioecious. Pod oblong, many-seeded. Stem leafy.

*** Anthers 2-celled, innate or introrse: pod septicidal, many-seeded.

11. **Tofieldia.** Flowers perfect, spiked or racemed. Leaves equitant.

Tribe III. UVULARIÆ. Style 3-parted; i. e. styles 3 united into one only at the base. Fruit a few-seeded pod. Seed-coat soft or loose. Anthers more or less extrorse. Perianth 6-leaved, deciduous. Stems from a rootstock or fibrous roots, leafy. Leaves alternate, broad, parallel-veined.

12. **Uvularia.** Pod 3-angular or 3-lobed. Anthers linear, adnate, on short filaments.

Tribe IV. ASPARAGINEÆ. Style 1, undivided (i. e. the 3 or rarely 2 styles united to the summit into one). Fruit a few-seeded berry. Seeds with a close coat and horny albumen. Stems from a rootstock. Leaves parallel-veined, chiefly alternate. Pedicels often articulated with the flower or in the middle.

* Plants with true leaves. Coat of the hard seed thin and membranous.

+ Anthers extrorsely attached to the filament. Stamens hypogynous. Sepals distinct.

13. **Prosartes.** Anthers linear-oblong, pointless. Flowers terminating the forked branches, on straight jointless pedicels.
14. **Streptopus.** Anthers arrow-shaped, pointed. Flowers lateral along the forked branches: pedicels bent about the middle.
15. **Clintonia.** Anthers oblong, pointless. Flowers terminating a naked scape.

+ + Anthers introrse. Filaments attached to the perianth. Stem simple.

16. **Convallaria.** Perianth bell-shaped, 6-lobed, bearing the stamens. Leaves all at the base of the naked scape, which bears the flowers in a simple raceme.
17. **Smilacina.** Perianth 4-6-parted, spreading, the stamens borne at the base. Stem leafy. Flowers in a simple or compound terminal raceme.
18. **Polygonatum.** Perianth tubular, 6-cleft, bearing the stamens above the middle. Stem leafy to the top. Flowers axillary.

** Plants with branching stems, their true leaves reduced to scales: leaf-like branchlets serving for foliage. Seed-coat hard and black.

19. **Asparagus.** Perianth 6-parted. The apparent leaves very narrow or thread-like.

Tribe V. LILIEÆ. Style 1, undivided (i. e. the three united throughout into one), or rarely a sessile stigma. Fruit a loculicidal pod. Seeds with a fleshy albumen. Anthers introrse or extrorse. Stem commonly from a coated or scaly bulb.

* Leafy-stemmed from a scaly bulb.

20. **Lilium.** Perianth 6-leaved, deciduous. Pod oblong, many-seeded: seeds horizontal, flat.

** Scape naked or nearly so, from a coated or rarely scaly bulb. Seeds mostly globular.

+ Perianth of 6 separate or nearly separate divisions or sepals.

21. **Erythronium.** Flower single. Style club-shaped. Pod obovate.

22. **Ornithogalum.** Flowers corymbed, never blue or reddish. Style 3-sided.

23. **Scilla.** Flowers racemed, purple or blue. Style thread-like.

24. **Allium.** Flowers umbelled, from a spathe. Sepals 1-nerved.

+ + Perianth globular or ovoid, 6-toothed.

25. **Muscari.** Flowers in a dense raceme, numerous, small, mostly blue.

*** No bulb. Stem or scape several-flowered. Pod many-seeded.

26. **Hemerocallis.** Perianth large, funnel-shaped; the sepals united in a narrow tube below. Stamens and long style declined. Seeds globular, black.

27. **Yucca.** Perianth large, of 6 separate broad divisions. Stigmas sessile. Stem woody and persistent: leaves persistent. Seeds flat, horizontal. Flowers panicked.

28. **Narhecium.** Perianth 6-parted, the divisions narrow, yellowish. Filaments woolly. Style slender. Flowers in a raceme. Leaves equitant. Seeds small, long-tailed at both ends. (Transition to Juncaceæ.)

1. TRILLIUM, L. THREE-LEAVED NIGHTSHADE.

Sepals 3, lanceolate, spreading, herbaceous, persistent. Petals 3, larger, withering in age. Stamens 6: anthers linear, on short filaments, adnate, introrse; the cells opening down the margins. Styles (or rather stigmas) awl-shaped or slender, spreading or recurved above, persistent, stigmatic down the inner side. Ovary 3-6-angled. Berry ovate, 3-celled (purple or red). Seeds horizontal, several in each cell. — Low perennial herbs, with a stout and simple stem rising from a short and præmorse tuber-like rootstock, naked, bearing at the summit a whorl of 3 ample, commonly broadly ovate, more or less ribbed but netted-veined leaves, and a terminal large flower; in spring. (Name from *trilix*, triple; all the parts being in threes.) — Monstrosities are not rare with the calyx and sometimes the petals changed to leaves, or with the parts of the flower increased in number.

§ 1. *Flower sessile in the bosom of the leaves, erect: petals varying from spatulate to lanceolate, 1' - 2' long, little exceeding the sepals, withering-persistent (stems 4' - 12' high).*

1. **T. sessile, L.** *Leaves sessile, ovate or rhomboidal, acute, often blotched or spotted; sessile petals erect-spreading (dark and dull purple, varying to greenish).* — Moist woods, Pennsylvania to Wisconsin, and southward.

2. **T. recurvatum, Beck.** *Leaves contracted at the base into a petiole, ovate, oblong, or obovate; sepals reflexed, petals pointed, the base narrowed into a claw, dark purple.* — Indiana to Wisconsin, and southward.

§ 2. *Flower raised on a peduncle: petals withering away after blossoming.*

* *Peduncle slender, erect or inclined: leaves rhombic-ovate, abruptly taper-pointed, sessile or nearly so by a wedge-shaped or acute base: petals plane.*

3. **T. grandiflorum, Salisb.** (LARGE WHITE T. OR WAKE ROBIN.) *Leaves rhomboid-obovate, longer than broad, barely sessile; petals obovate,*

spreading from an erect base, much larger than the sepals (2' - 2½' long), *white, changing with age to rose-color*. — Rich woods, Vermont to S. Penn., Kentucky, Wisconsin, and northward. June. — Flower erect, on a peduncle 2' - 3' long, very handsome.

4. **T. erectum**, L. (PURPLE T. OR BIRTHROOT.) Leaves dilated-rhomboidal, nearly as broad as long; *petals ovate, acutish, dark dull purple, spreading, little longer than the sepals (1' - 1½' long)*. (T. rhomboideum, var. atropurpureum, Michx.) — Rich woods: common, especially northward. May. — Peduncle 1' - 3' long, usually rather inclined than erect.

Var. **album**, Pursh. *Petals greenish-white, or rarely yellowish; ovary mostly dull-purple*. (T. pendulum, Ait., &c.) — With the purple-flowered form, especially from New York westward.

Var. **declinatum**. *Peduncle (fully half the length of the leaves) horizontal, or soon becoming so, or in fruit almost deflexed; petals white, rarely pink*. — Ohio to L. Superior (where it is the principal Trillium, Dr. Robbins), and northward. — Sometimes confounded with the next, if not passing into it.

* * *Peduncle from the first recurved under the short-petioled or almost sessile leaves, scarcely if at all longer than the wavy recurved-spreading petals.*

5. **T. cernuum**, L. (NODDING T. OR WAKE-ROBIN.) Leaves broadly rhomboid, abruptly pointed; *petals white, oblong-ovate, acute, rather longer than the sepals (6'' - 9'' long); styles distinct*. — Moist woods, especially eastward.

6. **T. stylòsum**, Nutt. *Leaves oblong, tapering to both ends, more distinctly ribbed; petals tinged with rose-color, oblong, much longer and broader than the sepals; styles united below the middle*. — Virginia? and southward.

* * * *Peduncle erect or nearly so: leaves distinctly petioled from a rounded base: petals merely spreading, longer than the sepals.*

7. **T. nivale**, Riddell. (DWARF WHITE T.) Small (2' - 4' high); *leaves oval or ovate, obtuse; petals oblong, obtuse, white, scarcely wavy, spreading from an erect base (as in No. 3), equalling the peduncle; styles long and slender*. — Rich woods, Ohio to Wisconsin. April. — Leaves 1' - 2', and petals 1' long.

8. **T. erythrocarpum**, Michx. (PAINTED T.) *Leaves ovate, taper-pointed; petals ovate or oval-lanceolate, pointed, wavy, widely spreading, white painted with purple stripes at the base, shorter than the peduncle*. (T. pictum, Pursh.) — Cold damp woods and bogs, New England and Penn. to L. Superior and northward, and southward in the higher Alleghanies. May, June.

2. MEDËOLA, Gronov. INDIAN CUCUMBER-ROOT.

Perianth recurved, the 3 sepals and 3 petals oblong and alike (pale greenish-yellow), deciduous. Stamens 6: anthers shorter than the slender filaments, oblong, extrorsely attached above the base, but the line of dehiscence of the closely contiguous parallel cells lateral or slightly introrse. Stigmas, or styles stigmatic down the upper side, recurved-diverging from the globose ovary, long and thread-form, deciduous. Berry globose (dark purple), 3-celled, few-seeded. — A perennial herb, with a simple slender stem (1° - 3° high, clothed with flocculent and deciduous wool), rising from a horizontal and tuberous white

rootstock (which has the taste of cucumber), bearing near the middle a whorl of 5-9 obovate-lanceolate and pointed, sessile, lightly parallel-ribbed and netted-veiny, thin leaves; also another of 3 (rarely 4 or 5) much smaller ovate ones at the top, subtending a sessile umbel of small recurved flowers. (Named after the sorceress *Medea*, from the wholly imaginary notion that it possesses great medicinal virtues.)

1. *M. Virginica*, L. (*Gyròmia*, *Nutt.*) — Rich damp woods. June.

3. MELÁNTHIUM, Grobov., L. MELANTHIUM.

Flowers monœciously polygamous. Perianth of 6 separate and free widely spreading somewhat heart-shaped or oblong and halberd-shaped sepals, raised on slender claws, cream-colored, the base marked with 2 approximate or confluent glands, turning greenish-brown and persistent. Filaments shorter than the sepals, adhering to their claws often to near their summit, persistent. Anthers heart-shaped or kidney-shaped, confluent 1-celled, shield-shaped after opening, extrorse. Styles 3, awl-shaped, diverging, tipped with simple stigmas. Pod ovoid-conical, 3-lobed, of 3 inflated membranaceous carpels united in the axis, separating when ripe, and splitting down the inner edge, several-seeded. Seeds flat, broadly winged. — Stem simple (3°-5° high), from a somewhat bulbous base, roughish-downy above, as well as the open and ample pyramidal panicle (composed chiefly of simple racemes), the terminal part mostly fertile. Leaves lanceolate or linear, grass-like, those from the root broader. (Name composed of *μέλας*, *black*, and *άνθος*, *flower*, from the darker color which the persistent perianth assumes after blossoming; but the name is hardly warranted.)

1. *M. Virginicum*, L. (BUNCH-FLOWER.) (*M. Virginicum* & *racemosum*, *Michx.* *Leimanthium Virginicum*, *Willd.* *L. Virginicum* & *hybridum*, *Roem. & Schult.*, *Gray*, *Melanth.*) — Wet meadows, from Southern New York to Illinois, and common southward. July.

4. ZYGADÈNUS, Michx. ZYGADENE.

Flowers perfect or polygamous. Perianth withering-persistent, spreading; the petal-like oblong or ovate sepals 1-2-glandular next the more or less narrowed, but not unguiculate base, which is either free, or united and coherent with the base of the ovary. Stamens free from the sepals and about their length. Anthers, styles, and pod nearly as in *Melanthium*. Seeds margined or slightly winged. — Very smooth and somewhat glaucous perennials, with simple stems from creeping rootstocks or coated bulbs, linear leaves, and pretty large paniced greenish-white flowers; in summer. (Name composed of *ζυγός*, *a yoke*, and *ἀδήν*, *a gland*, the glands being in pairs.)

* *Glands on the perianth conspicuous.*

1. *Z. glaberrimus*, Michx. Stems 1°-3° high from a *creeping rootstock*; leaves *grass-like*, channelled, conspicuously nerved, elongated, tapering to a point; panicle *pyramidal*, many-flowered; perianth nearly free; the sepals ($\frac{1}{2}$ ' long) ovate, becoming lance-ovate, with a pair of orbicular glands above the short claw-like base. — Grassy low grounds, Virginia (*Pursh*) and southward.

2. **Z. glaucus**, Nutt. Stem 1° – 3° high from a coated bulb; leaves flat; panicle rather simple and few-flowered; base of the perianth coherent with the base of the ovary, the thin ovate or obovate sepals marked with a large *obcordate gland*. (*Anticléa glauca*, Kunth.)—Along the St. Lawrence and Great Lakes (Bergen Swamp, Genesee Co., New York, *G. T. Fish*) to N. Illinois: rare.

* * *Glands of the perianth obscure.* (Here also *Amianthium Nuttallii*, Gray.)

3. **Z. leimanthoides**. Stem 1° – 4° high from a somewhat bulbous base, slender; leaves narrowly linear; flowers small ($4''$ in diameter) and numerous, in a few crowded paniced racemes; only a yellowish spot on the contracted base of the divisions of the free perianth. (*Amianthium leimanthoides*, Gray.)—Low grounds, pine barrens of New Jersey (*Durand*, *Knieskern*) and southward.

5. STENÁNTHIUM, Gray (under Veratrum).

Flowers polygamous. Perianth spreading; the sepals narrowly lanceolate, tapering to a point from the broader base, where they are united and coherent with the base of the ovary, not gland-bearing, persistent, much longer than the short stamens. Anthers, pods, &c. nearly as in Nos. 4 and 6. Seeds nearly wingless.—Smooth, with a wand-like leafy stem from a somewhat bulbous base, long and grass-like conduplicate-keeled leaves, and numerous small flowers in compound racemes, forming a long terminal panicle; in summer. (Name composed of *στενός*, narrow, and *άνθος*, flower, from the slender sepals and panicles.)

1. **S. angustifolium**, Gray. Leaves linear, elongated; flowers ($\frac{1}{4}'$ long), white, very short-pedicelled, in slender racemes; the prolonged terminal one, and often some of the lateral, fertile. (*Veratrum angustifolium*, Pursh. *Helonias graminea*, Bot. Mag.)—Low prairies and meadows, Penn. to Illinois and southward towards the mountains.—Stem slender, 2° – 6° high.

6. VERÀTRUM, Tourn. FALSE HELLEBORE.

Flowers monœciously polygamous. Perianth of 6 spreading and separate obovate-oblong (greenish or brownish) sepals, more or less contracted at the base, entirely free from the ovary, not gland-bearing. Filaments free from the sepals and shorter than they, recurving. Anthers, pistils, fruit, &c. nearly as in *Melanthium*.—Somewhat pubescent perennials, with simple stems from a thickened base producing coarse fibrous roots (very poisonous), 3-ranked leaves, and racemed-paniced dull or dingy flowers; in summer. (Name formed of *vere*, truly, and *ater*, black.)

1. **V. viride**, Ait. (AMERICAN WHITE HELLEBORE. INDIAN POKE.) Stem stout, very leafy to the top (2° – 4° high); leaves broadly oval, pointed, sheath clasping, strongly plaited; panicle pyramidal, the dense spike-like racemes spreading; perianth yellowish-green, moderately spreading.—Swamps and low grounds: common. (Much too near *V. album* of Europe.)

2. **V. parviflorum**, Michx. Stem slender (2° – 5° high), sparingly leafy below, naked above; leaves scarcely plaited, glabrous, contracted into sheathing petioles, varying from oval to lanceolate; panicle very long and loose, the terminal raceme wand-like, the lateral slender and spreading; pedicels as long as the flow-

ers; sepals dingy-green, oblanceolate or spatulate (2''-3' long), those of the sterile flowers on claws, widely spreading. (*Melanthium monoicum*, *Walt.* *Leimanthium monoicum*, *Gray.*)—Mountains of Virginia and southward.

3. **V. Woodii**, Robbins. Leaves lanceolate or oblong-lanceolate; *pedicels* (1½''-3'' long) shorter than the flowers, the oblanceolate spreading *sepals* (3''-4½'' long) dingy green turning brownish purple within: otherwise much as in the last, of which it may be a variety; but the flowers are mostly double the size, and the panicle stouter. (Plant 3°-6° high.)—Woods and hilly barrens, Green Co., Indiana, *Wood.* Augusta, Illinois, *Mead.*

7. AMIÁÑTHIUM, Gray. FLY-POISON.

Flowers perfect. Perianth widely spreading; the distinct and free petal-like (white) sepals oval or obovate, without claws or glands, persistent. Filaments capillary, equalling or exceeding the perianth. Anthers, pods, &c., nearly as in *Melanthium*. Styles thread-like. Seeds wingless, oblong or linear, with a loose coat, 1-4 in each cell.—Glabrous, with simple stems from a bulbous base or coated bulb, scape-like, few-leaved, terminated by a simple dense raceme of handsome flowers, turning greenish with age. Leaves linear, keeled, grass-like. (From *ἀμιανθος*, *unspotted*, and *ἄνθος*, *flower*; a name formed with more regard to euphony than to good construction, alluding to the glandless perianth.)

1. **A. muscætoxicum**, Gray. (FLY-POISON.) Leaves broadly linear, elongated, obtuse (½'-1' wide); raceme simple; pod abruptly 3-horned; seeds oblong, with a fleshy red coat. (*Helonias erythrospërma*, *Michx.*)—Open woods, New Jersey and Penn. to Kentucky and southward. June, July.

8. XEROPHYLLUM, Michx. XEROPHYLLUM.

Flowers perfect. Perianth widely spreading; sepals petal-like (white), oval, distinct, without glands or claws, at length withering, about the length of the awl-shaped filaments. Anthers 2-celled, short, extrorse. Styles thread-like, stigmatic down the inner side. Pod globular, 3-lobed, obtuse (small), loculicidal; the valves bearing the partitions. Seeds 2 in each cell, collateral, 3-angled, not margined.—Herb with the aspect of an *Asphodel*; the stem simple, 1°-4° high, from a bulbous base, bearing a simple compact raceme of showy white flowers, thickly beset with needle-shaped leaves, the upper ones reduced to bristle-like bracts; those from the root very many in a dense tuft, reclined, a foot or more long, 1' wide below, rough on the margin, remarkably dry and rigid (whence the name from *ξηρός*, *arid*, and *φύλλον*, *leaf*).

1. **X. asphodeloides**, Nutt. (*X. tènax*, *Nutt.* *X. setifolium*, *Michx.* *Helonias asphodelioides*, *L.*)—Pine barrens, New Jersey and southward: also far westward. June.

9. HELONIAS, L. HELONIAS.

Flowers perfect. Perianth of 6 spatulate-oblong purple sepals, persistent, turning green, shorter than the thread-like filaments. Anthers 2-celled, roundish-oval, blue, extrorse. Styles revolute, stigmatic down the inner side. Pod

obcordately 3-lobed, loculicidally 3-valved; the valves divergently 2-lobed. Seeds many in each cell, linear, with a tapering appendage at both ends.—A smooth perennial, with many oblong-spatulate or oblanceolate evergreen flat leaves, from a tuberous rootstock, producing in early spring a hollow naked scape (1°–2° high), sheathed with broad bracts at the base, and terminated by a simple and short dense raceme. Bracts obsolete: pedicels shorter than the flowers. (Name probably from *ἔλος*, a *swamp*, the place of growth.)

1. **H. bullata**, L. (*H. latifolia*, *Michx.*)—Wet places, New Jersey and Pennsylvania to Virginia: rare and local.

10. CHAMÆLIRIUM, Willd. DEVIL'S-BIT.

Flowers diœcious. Perianth of 6 spatulate-linear (white) spreading sepals, withering-persistent. Filaments and (yellow) anthers as in *Helonias*: fertile flowers with rudimentary stamens. Styles linear-club-shaped, stigmatic along the inner side. Pod ovoid-oblong, not lobed, of a thin texture, loculicidally 3-valved from the apex, many-seeded. Seeds linear-oblong, winged at each end.—Smooth herb, with a wand-like stem from a (bitter) thick and abrupt tuberous rootstock, terminated by a long wand-like spiked raceme (4'–9' long) of small bractless flowers; fertile plant more leafy than the staminate. Leaves flat, lanceolate, the lowest spatulate, tapering into a petiole. (Name formed of *χαμαί*, on the ground, and *λείριον*, lily; of no obvious application.)

1. **C. luteum**. (BLAZING-STAR.) (*C. Carolinianum*, *Willd.* *Veratrum luteum*, L. *Helonias lutea*, *Ait.* *H. dioica*, *Pursh.*)—Low grounds, W. New England to Illinois and southward. June.

11. TOFIELDIA, Hudson. FALSE ASPHODEL.

Flowers perfect, usually with a little 3-bracted involucre underneath. Perianth more or less spreading; the sepals (white or greenish) concave, oblong or obovate, without claws. Filaments awl-shaped: anthers short, innate or somewhat introrse, 2-celled. Styles awl-shaped: stigmas terminal. Pod 3-angular, 3-partible or septicidal; cells many-seeded. Seeds oblong.—Slender perennials, mostly tufted, with fibrous roots, and simple stems leafy only at the base, bearing small flowers in a close raceme or spike. Leaves 2-ranked, equitant, linear, grass-like. (Named for *Mr. Tofield*, an English botanist of the last century.)

§ 1. *Flowers in a simple spike-like raceme or head: anthers introrse: seeds not appendaged: plant smooth and glabrous.*

1. **T. palustris**, Hudson. Scape leafless or nearly so (3'–6' high), slender, bearing a globular or oblong head or short raceme of whitish flowers; leaves tufted, 1' long.—Isle Royale, &c., Lake Superior, and northward. July.

§ 2. *Flowers racemose, but developing from above downwards: short pedicels in threes from a little involucre of as many bracts: anthers innate: seeds tail-pointed at both ends (as in many species of Juncus.)*

2. **T. glutinosa**, Willd. Stem (6'–16' high) and pedicels very glutinous with dark glands; leaves broadly linear, short.—Moist grounds, Maine, Michigan, Wisconsin, and northward: also southward in the Alleghanies. June.

3. **T. pùbens**, Ait. Stem (1° - 2° high) and pedicels *roughened with minute glands*; leaves longer and narrower. — Pine barrens, New Jersey to Virginia and southward. July.

12. UVULÀRIA, L. BELLWORT.

Perianth nearly bell-shaped, lily-like; the 6 distinct sepals spatulate-lanceolate, with a honey-bearing groove or pit at the erect contracted base, much longer than the stamens, which barely adhere to their base, deciduous. Anthers linear, much longer than the filaments, adnate and extrorse, but the long and narrow cells opening nearly along the margin. Style deeply 3-cleft; the divisions stigmatic along the inner side. Pod triangular or 3-lobed, loculicidally 3-valved from the top. Seeds few in each cell, obovoid, with a tumid or fungous rhaps. — Stems rather low, from a rootstock, naked or scaly at the base, forking above, bearing oblong sessile or clasping flat and membranaceous leaves, and yellowish drooping flowers, in spring, solitary or rarely in pairs, on terminal peduncles which become lateral by the growth of the branches. (Name "from the flowers hanging like the *uvula*, or palate.")

* *Leaves clasping-perfoliate: sepals pointed: pod truncate, 3-lobed: rootstock short.*

1. **U. grandiflora**, Smith. Stems 1° - 2° high; flower $1\frac{1}{2}'$ long, *pale greenish-yellow, the sepals nearly smooth within*; anthers blunt-pointed. — Rich woods, Vermont to Wisconsin and southwestward.

2. **U. perfoliata**, L. Smaller; *sepals granular-roughened within; anthers sharper tipped*; otherwise as No. 1. — Common eastward and southward.

3. **U. flava**, Smith. *Flower bright yellow, 1' long; sepals nearly smooth within*; anthers short-pointed. — New Jersey to Virginia: rare.

** *Leaves sessile: pod triangular: stems low (6' - 12'): rootstock creeping.*

4. **U. sessilifolia**, L. *Leaves lance-oblong, pale, glaucous beneath, sessile or partly clasping by a narrow base; sepals blunt (9" long); anthers pointless; the ovoid and sharply triangular pod stipitate.* — Low woods: common. May.

5. **U. pubérula**, Michx. Slightly *puberulent; leaves bright green both sides and shining, oval, with rough edges; styles separate to near the base, not exceeding the short-pointed anthers; pod ovate, not stipitate; otherwise like the last.* — Mountains, Virginia, and southward.

13. PROSÀRTEs, Don. PROSARTES.

Perianth bell-shaped, the 6 sepals lanceolate or linear, deciduous. Filaments thread-like, much longer than the linear-oblong blunt anthers, which are fixed by a point above the base, and extrorse. Ovary with 2 ovules suspended from the summit of each cell: style one: stigmas short, recurved-spreading, or sometimes united into one! Berry ovoid or oblong, pointed, 3 - 6-seeded, red. — Downy low herbs, divergently branched above, with closely sessile ovate and membranaceous leaves, and greenish-yellow drooping flowers, on slender terminal peduncles, solitary or few in an umbel. (Name from *προσαράω*, to hang from, in allusion to the pendent ovules or flowers.)

1. **P. lanuginosa**, Don. Leaves ovate-oblong, taper-pointed, rounded or slightly heart-shaped at the base, closely sessile, downy underneath; flowers

solitary or in pairs; sepals linear-lanceolate, taper-pointed ($\frac{1}{2}$ ' long), soon spreading, twice the length of the stamens, greenish; style smooth; stigmas 3. (*Streptopus lanuginosus*, Michx.) — Rich woods, Western New York to Virginia, Kentucky, and southward along the Alleghanies. May.

14. STRÉPTOPUS, Michx. TWISTED-STALK.

Perianth recurved-spreading from a bell-shaped base; the 6 distinct sepals lanceolate, acute, the 3 inner keeled, deciduous. Anthers arrow-shaped, extrorse, fixed near the base to the short flattened filaments, tapering above to a slender entire or 2-cleft point. Ovary with many ovules in each cell: style and sometimes the stigmas one. Berry red, roundish-ovoid, many-seeded. — Herbs, with rather stout stems, forking and divergent branches, ovate and taper-pointed rounded-clasping membranaceous leaves, and small (extra-) axillary flowers, either solitary or in pairs, on slender thread-like peduncles, which are abruptly bent or contorted near the middle (whence the name, from *στρεπτός*, *twisted*, and *πούς*, *foot or stalk*.)

1. **S. amplexifolius**, DC. *Leaves very smooth, glaucous underneath, strongly clasping; flower greenish-white on a long peduncle abruptly bent above the middle; anthers tapering to a slender entire point; stigma entire, truncate.* (*S. distortus*, Michx. *Uvularia amplexifolia*, L.) — Cold and moist woods, from Penn. northward. June. — Stem 2°–3° high, rough at the base, otherwise very smooth. Sepals $\frac{1}{2}$ ' long. — In this, as in the next, the peduncles are *opposite the leaves*, rather than truly axillary, and are bent round the clasping base underneath them: they are rarely 2-flowered. (Eu.)

2. **S. roseus**, Michx. *Leaves green both sides, finely ciliate, and the branches sparingly beset with short bristly hairs; flower rose-purple, more than half the length of the slightly bent peduncle; anthers 2-horned; stigma 3-cleft.* — Cold damp woods, northward, and in the Alleghanies southward. May.

15. CLINTONIA, Raf. CLINTONIA.

Perianth of 6 separate sepals, bell-shaped, lily-like, deciduous; the 6 stamens inserted at their base. Filaments long and thread-like: anthers linear or oblong, extrorsely fixed by a point above the base, the cells opening down the margins. Ovary ovoid-oblong, 2–3-celled: style long: stigmas 2 or 3, or in ours united into one. Berry ovoid, blue, few–many-seeded. — Stemless perennials, with slender creeping rootstocks, producing a naked scape sheathed at the base by the stalks of 2–4 large oblong or oval ciliate leaves. Flowers rather large, umbelled, rarely single. (Dedicated to *De Witt Clinton*.)

1. **C. borealis**, Raf. *Umbel few- (2–7-) flowered; ovules 20 or more.* (*Dracæna borealis*, Ait.) — Cold moist woods, Massachusetts to Wisconsin and northward, and southward along the Alleghanies. June. — Scape and leaves 5'–8' long. Perianth over $\frac{1}{2}$ ' long, greenish-yellow, somewhat downy outside.

2. **C. umbellata**, Torr. *Umbel many-flowered; ovules 2 in each cell.* (*C. multiflora*, Beck. *Convallaria umbellulata*, Michx.) — Rich woods, S. W. New York, and southward along the Alleghanies. June, July. — Flowers half the size of the last, white, speckled with green or purplish dots.

16. CONVALLÀRIA, L. (in part). LILY OF THE VALLEY.

Perianth bell-shaped (white), 6-lobed, deciduous; the lobes recurved. Stamens 6, included, inserted on the base of the perianth: anthers introrse. Ovary 3-celled, tapering into a stout style: stigma triangular. Ovules 4-6 in each cell. Berry few-seeded (red).—A low perennial herb, glabrous, stemless, with slender running rootstocks, sending up from a scaly-sheathing bud 2 oblong leaves, with their long sheathing petioles enrolled one within the other so as to appear like a stalk, and an angled scape bearing a one-sided raceme of pretty and sweet-scented nodding flowers. (Altered from *Lilium convallium*, the popular name.)

1. *C. majàlis*, L.—High Alleghanies of Virginia, and southward. May.—Same as the European LILY OF THE VALLEY of the gardens. (Eu.)

17. SMILACÌNA, Desf. FALSE SOLOMON'S SEAL.

Perianth 4-6-parted, spreading, deciduous (white), with as many stamens inserted at the base of the divisions. Filaments slender: anthers short, introrse. Ovary 2-3-celled, with 2 ovules in each cell: style short and thick: stigma obscurely 2-3-lobed. Berry globular, 1-2-seeded.—Perennial herbs, with simple stems from creeping or thickish rootstocks, alternate nerved leaves, and white, sometimes fragrant flowers in a terminal and simple or compound raceme. (Name a diminutive of *Smilax*, to which, however, these plants bear little resemblance.)

§ 1. SMILACINA proper. *Divisions of the perianth (oblong-lanceolate) and stamens 6, the latter longer: ovary 3-celled: ovules collateral: racemes crowded in a compound raceme or close panicle: rootstock stout, fleshy.*

1. *S. racemosa*, Desf. (FALSE SPIKENARD.) Minutely downy (2° high); leaves numerous, oblong or oval-lanceolate, taper-pointed, ciliate, abruptly somewhat petioled.—Moist copses: common. June.—Berries pale red, speckled with purple, aromatic. (*S. ciliàta*, Desf., is a dwarf state of this.)

§ 2. ASTERÁNTHEMUM, Kunth. *Divisions of the perianth 6, oblong-lanceolate, longer than the stamens: ovary 2-3-celled: ovules one above the other: raceme single, 5-20-flowered: rootstock rather slender.*

2. *S. stellàta*, Desf. Nearly glabrous, or the 7-12 oblong-lanceolate leaves minutely downy beneath when young, slightly clasping; berries blackish.—Moist banks: common northward. May, June.—Plant 1°-2° high. (Eu.)

3. *S. trifòlia*, Desf. Glabrous, dwarf (3'-6' high); leaves 3 (sometimes 2 or 4), oblong, tapering to a sheathing base; berries red.—Cold bogs, New England to Penn., Wisconsin, and northward. May.

§ 3. MAIÁNTHEMUM, Desf. *Divisions of the reflexed-spreading perianth (oval) and the stamens 4, of equal length: ovary 2-celled: ovules collateral: raceme single, many-flowered: creeping rootstocks slender.*

4. *S. bifòlia*, Ker. Glabrous, or somewhat pubescent, low (3'-5' high); leaves mostly 2 (sometimes 3), heart-shaped, petioled, or in our plant (the var. CANADÉNSIS) one or both often sessile or nearly so and clasping.—Moist woods: very common, especially northward. May. (Eu.)

18. POLYGONÀTUM. TOURN. SOLOMON'S SEAL.

Perianth cylindrical, 6-lobed at the summit; the 6 stamens inserted on or above the middle of the tube, included: anthers introrse. Ovary 3-celled, with 2-6 ovules in each cell: style slender, deciduous by a joint: stigma obtuse or capitate, obscurely 3-lobed. Berry globular, black or blue; the cells 1-2-seeded. — Perennial herbs, with simple erect or curving stems, rising from creeping thick and knotted rootstocks, naked below, above bearing nearly sessile or half-clasping nerved leaves, and axillary nodding greenish flowers: pedicels jointed with the flower. (The ancient name, composed of *πολύς*, *many*, and *γόνυ*, *knee*, alluding to the numerous joints of the rootstocks and stems.) — Ours are all alternate-leaved species, and with the stem terete or scarcely angled when fresh.

1. **P. biflorum**, Ell. (SMALLER SOLOMON'S SEAL.) Glabrous, except the ovate-oblong or lance-oblong *nearly sessile leaves*, which are commonly *minutely pubescent*, at least on the veins (but sometimes smooth), *as well as pale or glaucous underneath*; stem slender (1°-3° high); *peduncles 1-3- but mostly 2-flowered*; *filaments papillose-roughened*, inserted towards the summit of the cylindrical-oblong perianth. (*Convallaria biflora*, Walt. *C. pubescens*, Willd. *Polygonatum pubescens*, *angustifolium*, & *multiflorum*, Pursh.) — Wooded banks: common. — Perianth $\frac{1}{2}$ ' long, greenish.

2. **P. giganteum**, Dietrich. (GREAT S.) *Glabrous throughout*; stem stout and mostly tall, terete; *leaves ovate, partly clasping* (5'-8' long), or the upper oblong and nearly sessile, many-nerved; *peduncles several- (2-8-) flowered*; *filaments smooth and naked*, or nearly so, inserted on the middle of the tube of the cylindrical-oblong perianth. (*Convallaria canaliculata*, Willd. *Polygonatum canaliculatum*, Pursh. *P. commutatum*, Dietrich.) — River-banks, in alluvial soil, 5°-8° high; in dry or less fertile soil 2°-4°. June. (The stem not being at all channelled in the living plant, it is better to discard the earlier name of *canaliculatum*.) — Pedicels 4''-15'' long: perianth 9'' long. — Perhaps passes into the preceding.

3. **P. latifolium**, Desf. Upper part of the stem (2°-3° high), the 1-5-flowered *peduncles, pedicels*, and lower surface of the ovate or oblong mostly *petioled leaves more or less pubescent*; *filaments glabrous*. (*P. hirtum*, Pursh. *Convallaria hirta*, Poir.) — Pennsylvania, *Muhlenberg!*

(*P. MULTIFLORUM*, with hirsute filaments, I have never seen in this country.)

19. ASPÁRAGUS, L. ASPARAGUS.

Perianth 6-parted, spreading above: the 6 stamens on its base: anthers introrse. Style short: stigma 3-lobed. Berry spherical, 3-celled; the cells 2-seeded. — Perennials, with much-branched stems from thick and matted rootstocks, and small greenish-yellow axillary flowers on jointed pedicels. The narrow, commonly thread-like, so-called leaves are really branchlets, acting as leaves, clustered in the axil of a little scale which is the true leaf. (The ancient Greek name.)

1. **A. OFFICINÀLIS**, L. (GARDEN ASPARAGUS.) Herbaceous, tall, bushy-branched; leaves thread-like. — Sparingly escaped from gardens into waste places on the coast. June. (Adv. from Eu.)

20. LÍLIUM, L. LILY.

Perianth funnel-form or bell-shaped, colored, of 6 distinct sepals, spreading or recurved above, with a honey-bearing furrow at the base, deciduous; the 6 stamens somewhat adhering to their bases. Anthers linear, extrorsely inserted towards the middle to the tapering apex of the long filament, which is at first included, at length versatile; the cells dehiscent by a lateral or slightly introrse line. Style elongated, somewhat club-shaped: stigma 3-lobed. Pod oblong, containing numerous flat and horizontal (depressed) soft-coated seeds densely packed in 2 rows in each cell.—Bulbs scaly, producing simple stems, with numerous alternate-scattered or whorled short and sessile leaves, and from one to several large and showy flowers; in summer. (The classical Latin name, from the Greek *λείριον*.)

* *Flowers erect, bell-shaped, the sepals narrowed below into claws.*

1. **L. Philadelphicum**, L. (WILD ORANGE-RED LILY.) *Leaves linear-lanceolate; the upper chiefly in whorls of 5 to 8; flowers 1–3, open-bell-shaped, reddish-orange spotted with purplish inside; the lanceolate sepals not recurved at the summit.—Dry or sandy ground: common.—Stem 2°–3° high: the flower 2½' long.*

2. **L. Catesbæi**, Walt. (SOUTHERN RED LILY.) *Leaves linear-lanceolate, scattered; flower solitary, open-bell-shaped, the long-clawed sepals wavy on the margin and recurved at the summit, scarlet, spotted with dark purple and yellow inside.—Low sandy soil, Pennsylvania? to Kentucky and southward.*

** *Flowers nodding, bell-shaped, the sessile sepals revolute.*

3. **L. Canadense**, L. (WILD YELLOW LILY.) *Leaves remotely whorled, lanceolate, strongly 3-nerved, the margins and nerves rough, flowers few, long-peduncled, oblong-bell-shaped, the sepals recurved-spreading above the middle, orange-spotted inside with brown.—Moist meadows and bogs, especially northward.—Stem 2°–5° high. Flower 2'–3' long.*

4. **L. superbum**, L. (TURK'S-CAP LILY.) *Lower leaves whorled, lanceolate, pointed, 3-nerved, smooth; flowers often many (3–20 or 40) in a pyramidal raceme; sepals strongly revolute, bright orange, with numerous dark purple spots inside.—Rich low grounds.—Stem 3°–7° high: sepals 3' long. L. Caroliniæ, Michx., apparently belongs to this species, which also probably passes into the preceding.*

21. ERYTHRÒNIUM, L. DOG'S-TOOTH VIOLET.

Perianth lily-like, of 6 distinct lanceolate sepals, recurved or spreading above, deciduous, the 3 inner usually with a callous tooth on each side of the erect base, and a groove in the middle. Filaments 6, awl-shaped: anthers oblong-linear, continuing erect. Style elongated. Pod obovate, contracted at the base, 3-valved, loculicidal. Seeds rather numerous, ovoid, with a loose, membranaceous tip.—Nearly stemless herbs, with two smooth and shining flat leaves tapering into petioles and sheathing the base of the commonly one-flowered scape, rising from a deep solid-scaly bulb. Flowers rather large, nodding, in spring. (Name from *έρυθρός*, red, which is inappropriate as respects the American species.)

1. **E. Americanum**, Smith. (YELLOW ADDER'S-TONGUE.) Leaves elliptical-lanceolate, pale green, mottled and commonly dotted with purplish and whitish; *perianth light yellow*, often spotted near the base; style club-shaped; *stigmas united* into one. — Low copses: common. May. — Scape 6'–9' high: flower 1' or more long. — E. BRACTEATUM, Boott, from the Camel's Rump Mountain, Vermont, is probably only an accidental state, with a bract, such as the Western *E. grandiflorum* often has.

2. **E. álbídum**, Nutt. (WHITE DOG'S-TOOTH VIOLET.) Leaves elliptical-lanceolate, less or not at all spotted, not dotted; *perianth bluish-white*; inner divisions toothless; style more slender except at the apex, bearing 3 *spreading stigmas*. — Low thickets, New York (near Albany), and W. Penn. to Wisconsin and Illinois. — At Lake Superior, Dr. Robbins finds a plant like this but *yellow-flowered*, a transition towards *E. grandiflorum*.

22. ORNITHÓGALUM, Tourn. STAR-OF-BETHLEHEM.

Perianth of 6 colored (white) spreading 3–7-nerved sepals. Filaments 6, flattened-awl-shaped. Style 3-sided: stigma 3-angled. Pod membranous, roundish-angular, with few dark and roundish seeds in each cell, loculicidal. — Scape and linear channelled leaves from a coated bulb. Flowers corymbed, bracted. (An ancient whimsical name from *ὄρνις*, a *bird*, and *γάλα*, *milk*.)

1. **O. UMBELLATUM**, L. Flowers 5–8, on long and spreading pedicels; sepals green in the middle on the outside. — Escaped from gardens into moist meadows, eastward. June. (Nat. from Eu.)

23. SCÍLLA, L. SQUILL.

Perianth of 6 colored (blue or purple) spreading sepals, mostly deciduous; the 6 awl-shaped filaments at their base. Style thread-like. Pod 3-angled, loculicidal, 3-valved, with several black roundish seeds in each cell. — Scape and linear leaves from a coated bulb: the flowers in a simple raceme, mostly bracted. (The ancient name.)

1. **S. Fraseri**. (EASTERN QUAMASH. WILD HYACINTH.) Leaves long and linear, keeled; raceme elongated; bracts solitary, longer than the pedicels; stigma minutely 3-cleft; pod triangular, several-seeded. (*Phalángium esculéntum*, Nutt. *Scilla esculenta*, Ker. *Camássia Fraseri*, Torr.) — Moist prairies and river-banks, Ohio to Wisconsin and southwestward. May. — Bulb onion-like, eaten by the Indians. Scape 1° high. Sepals widely spreading, pale blue, 3-nerved, 6'' long.

24. ÁLLIUM, L. ONION. GARLIC.

Perianth of 6 entirely colored sepals, which are distinct, or united at the very base, 1-nerved, often becoming dry and scarious and more or less persistent: the 6 filaments awl-shaped or dilated at their base. Style persistent, thread-like: stigma simple. Pod lobed, loculicidal, 3-valved, with 1 or few ovoid-kidney-shaped amphitropous or campylotropous black seeds in each cell. — Strong-scented and pungent stemless herbs; the leaves and scape from a

coated bulb flowers in a simple umbel, some of them frequently changed to bulblets; spathe 1-2-valved. (The ancient Latin name of the Garlic.)

* *Ovules and seeds single in each cell: leaves broad and flat, appearing in early spring, and dying before the flowers are developed.*

1. **A. tricóccum**, Ait. (WILD LEEK.) Scape naked (9' high from clustered pointed bulbs, 2' long), bearing an erect many-flowered umbel; leaves lance-oblong (5'-9' long, 1'-2' wide); sepals oblong (white), equalling the simple filaments; pod strongly 3-lobed.—Rich woods, W. New England to Wisconsin, Kentucky, and southward in the Alleghanies. July.

** *Ovules mostly 2 in each cell: ovary crested with 6 teeth: leaves long and narrow.*

+ Umbel bearing only flowers and ripening pods.

2. **A. cernuum**, Roth. (WILD ONION.) Scape naked, angular (1°-2° high), often nodding at the apex, bearing a loose or drooping many-flowered umbel; leaves linear, sharply keeled (1° long); sepals oblong-ovate, acute (rose-color), shorter than the simple slender filaments.—Steep banks, Western New York to Wisconsin and southward. July, Aug.

3. **A. stellatum**, Nutt. Scape terete, slender, bearing an erect umbel; leaves flat; sepals equalling the stamens: otherwise resembling the last, but usually not so tall; the pod more crested. Aug.—Rocky slopes, Illinois (Engelmann), and northwestward.

4. **A. Schœnóprasum**, L. (CHIVES.) Scape naked or leafy at the base (6'-12' high) bearing a globular *capitate umbel* of many rose-purple flowers; sepals lanceolate, pointed, longer than the simple downwardly dilated filaments; leaves awl-shaped, hollow. Var. with recurved tips to the sepals (A. Sibiricum, L.)—Shore of Lakes Huron, Superior, and northward. (Eu.)

+ + Umbel often densely bulb-bearing, with or without flowers.

5. **A. vineale**, L. (FIELD GARLIC.) Scape slender, clothed with the sheathing bases of the leaves below the middle (1°-3° high); leaves terete and hollow, slender, channelled above; filaments much dilated, the alternate ones 3-cleft, the middle division anther-bearing.—Moist meadows and fields: a vile weed eastward. June.—Flowers rose-color and green. (Nat. from Eu.)

6. **A. Canadense**, Kalm. (WILD GARLIC.) Scape leafy only at the base (1° high); leaves narrowly linear, flattish; umbel few-flowered; filaments simple, dilated below.—Moist meadows. May, June.—Flowers pale rose-color, pedicelled; or a head of bulbs in their place.

*** *Ovules several in each cell: leaves long and linear.* (Nothóscordum, Kunth.)

7. **A. striatum**, Jacq. Leaves narrowly linear, often convolute, striate on the back, about the length of the obscurely 3-angled naked scape (6'-12' long); filaments dilated below, shorter than the narrowly oblong sepals (which are white with a reddish keel); ovules 4-7 in each cell.—Prairies and open woods, Virginia to Illinois and southward. May.

25. MUSCARI, Tourn. GRAPE-HYACINTH.

Perianth globular or ovoid, minutely 6-toothed (blue). Stamens 6, included: anthers short, introrse. Style short. Pod loculicidal, with 2 black angular

seeds in each cell. — Leaves and scape (in early spring) from a coated bulb: the small flowers in a dense raceme, sometimes musk-scented (whence the name).

1. **M. BOTRYOÏDES**, Mill. Leaves linear; flowers globular ($1\frac{1}{2}$ "–2" long), deep blue, appearing like minute grapes; whence the popular name. — Escaped from gardens into copses and fence-rows, E. Penn., &c. (Adv. from Eu.)

26. HEMEROCÁLLIS, L. DAY-LILY.

Perianth funnel-form, lily-like; the short tube enclosing the ovary, the spreading limb 6-parted; the 6 stamens inserted on its throat. Anthers as in Lily, more or less extrorse. Filaments and style long and thread-like, declined and ascending: stigma simple. Pod (at first rather fleshy) 3-angled, loculicidally 3-valved, with several black spherical seeds in each cell. — Showy perennials, with fleshy-fibrous roots; the long and linear keeled leaves 2-ranked at the base of the tall scapes, which bear at the summit several bracted and large yellow flowers: these collapse and decay after expanding for a single day (whence the name, from *ἡμέρα*, a day, and *κάλλος*, beauty).

1. **H. FÚLVA**, L. (COMMON DAY-LILY.) Inner divisions (petals) of the tawny orange perianth wavy and obtuse. — Roadsides, escaped from gardens, where it is common (as is also the bright yellow-flowered *H. FLAVA*). July. (Adv. from Eu.)

27. YÚCCA, L. BEAR-GRASS. SPANISH BAYONET.

Perianth of 6 petal-like (white) oval or oblong and acute flat sepals, withering-persistent, the 3 inner broader, longer than the 6 stamens. Stigmas 3, sessile. Pod oblong, somewhat 6-sided, 3-celled, or imperfectly 6-celled by a partition from the back, fleshy, at length loculicidally 3-valved from the apex. Seeds very many in each cell, depressed. — Stems woody, either very short, or rising into thick and columnar palm-like trunks, bearing persistent rigid linear or sword-shaped leaves, and terminated by an ample compound panicle of showy (often polygamous) flowers. (An aboriginal name.)

1. **Y. filamentosa**, L. (ADAM'S NEEDLE.) Trunk (from a running rootstock) rising for a foot or less above the earth, covered with the lanceolate unarmed coriaceous leaves (1° – 2° long), which bear filaments on their margins; scape-like flower-stem 6° – 8° high, erect. — Sandy soil, E. Virginia and southward. July.

28. NARTHÈCIUM, Mœhring. BOG-ASPHODEL.

Sepals 6, linear-lanceolate, yellowish, persistent. Filaments 6, woolly: anthers linear, introrse. Pod cylindrical-oblong, pointed with the undivided style and single stigma, loculicidal, many-seeded. Seeds appendaged at each end with a long bristle-form tail, as in many species of *Juncus*. — Rootstock creeping, bearing linear equitant leaves (as in *Tofieldia*), and a simple stem or scape, terminated by a simple dense raceme. (Name from *ναρθήκιον*, a rod, or a box for fragrant ointments, of obscure application.)

1. **N. ossifragum**, Huds. — Herb with scape about a foot high, longer than the leaves: pedicels mostly bracteolate. (Eu.)

Var. **Americanum**. (*N. Americanum*, *Ker.*) Flowers rather smaller (scarcely 3'' long) and leaves narrower than the European plant, which is limited to the Atlantic side of that continent, as is ours here: viz. in sandy bogs, on this side, where it is very local, in the pine barrens of New Jersey only. June, July.

ORDER 122. JUNCACEÆ. (RUSH FAMILY.)

Grass-like or sedge-like herbs, with small flowers, a regular and hypogynous persistent perianth of 6 similar glumaceous sepals, 6 or rarely 3 stamens with 2-celled anthers, a single short style, 3 filiform hairy stigmas, and an ovary either 3-celled or 1-celled with 3 parietal placentæ, forming a loculicidal 3-valved pod. Seeds anatropous, with a minute embryo enclosed at the base of the fleshy albumen.—Rushes, with the flowers liliaceous in structure, but sedge-like in aspect and texture, mainly represented by only two genera.

1. LÛZULA, DC. WOOD-RUSH.

Pod 1-celled, 3-seeded, one seed to each parietal placenta. — Perennials, often hairy, usually in dry ground, with flat and soft usually hairy leaves, and spiked-crowded or umbelled flowers. (Name said to be altered from the Italian *luciola*, a glow-worm.)

* *Flowers loosely long-peduncled, umbelled or corymbed.*

1. **L. pilosa**, Willd. Leaves lance-linear, hairy; umbel mostly simple; sepals pointed, shorter than the obtuse pod; seeds with a curved appendage. — Woods and banks: common northward. May. — Plant 6'–9' high. (Eu.)

2. **L. parviflora**, Desv., var. **melanocarpa**. Nearly smooth (1°–3° high); leaves broadly linear; corymb decomposed, loose; pedicels drooping; sepals pointed, straw-color, about the length of the minutely pointed and brown pod. (*L. melanocarpa*, *Desv.*) — Mountains, Maine to Northern New York, and northward. July. (Eu.)

* * *Flowers crowded in spikes or close clusters. (Plants 6'–12' high.)*

3. **L. campéstris**, DC. Leaves flat, linear; spikes 4–12, somewhat umbelled, ovoid, straw-color, some of them long-peduncled, others nearly sessile; sepals bristle-pointed, longer than the obtuse pods; seeds with a conical appendage at the base. — Dry fields and woods; common. May. (Eu.)

4. **L. arcuata**, Meyer. Leaves channelled, linear; spikes 3–5, on unequal often recurved peduncles, ovoid, chestnut-brown; bracts ciliate-fringed; sepals taper-pointed, longer than the obtuse pod; seeds not appendaged. — Alpine summits of the White Mountains, New Hampshire, and high northward. (Eu.)

5. **L. spicata**, Desvoux. Leaves channelled, narrowly linear; flowers in sessile clusters, forming a nodding interrupted spiked panicle, brown; sepals bristle-pointed, scarcely as long as the abruptly short-pointed pod; seeds merely with a roundish projection at the base. (Our plant is *L. racemosa*, *Desv.*? according to *Godet.*) With the last, and more common. (Eu.)

2. JÚNCUS, L. RUSH. BOG-RUSH.

Pod many-seeded, 3-celled, or 1-celled by the placentæ not reaching the axis. Stamens when 3 opposite the 3 outer sepals. — Chiefly perennials, and in wet soil or water, with pithy or hollow and simple (rarely branching) stems, and paniced or clustered small (greenish or brownish) flowers, chiefly in summer. Plant never hairy. (The classical name, from *jungo*, to join, alluding to the use of the stems for bands.)

Contributed for this edition by DR. ENGELMANN, who has recently published a monograph of the North American species in the Transactions of the St. Louis Academy of Science.

§ 1. TRUE JUNCUS; with naked and simple scapes from matted running rootstocks, many of them barren, furnished at base with short leafless or rarely leaf-bearing sheaths: leaves, if any, terete, knotless and similar to the scape: flowers single or rarely clustered on the pedicels, in sessile panicles, produced apparently from the side of the scape, the exterior sheath or involucreal leaf being similar to and continuing the scape. Flowering in early summer.

* Flowers single on the pedicels or ultimate ramifications of the panicle.

+ Sheaths at base of the stem leafless.

++ Stamens 3.

1. **J. effusus**, L. (COMMON OR SOFT RUSH.) Scape soft and pliant (2° – 4° high); inner sheaths awned; panicle diffusely much branched, many-flowered; flowers numerous, small ($1\frac{1}{4}$ " long), greenish; sepals lanceolate, very acute, as long as the triangular-obovate retuse and pointless greenish-brown pod; anthers as long as filaments; style very short; seeds small (about $\frac{1}{4}$ " long) with short pale points, delicately ribbed and cross-lined. — Var. **CONGLOMERATUS** (*J. conglomeratus*, L.) has the scape more distinctly striate, the panicle closely crowded, and the pod short-pointed. — Marshy grounds: very common; the var. in sphagnous swamps. (Eu.)

++ ++ Stamens 6.

2. **J. filiformis**, L. Scape very slender (1° – 2° high), pliant; panicle few-flowered, almost simple; flowers ($1\frac{1}{2}$ " long); sepals lanceolate, the inner a little shorter and less acute, longer than the broadly ovate obtuse but short-pointed greenish pod; anthers shorter than the filaments; style very short; seed (less than $\frac{1}{2}$ " long) short-pointed at both ends, indistinctly reticulated. — N. New England and New York to Michigan and northward. (Eu.)

3. **J. Smithii**, Engelm. Scape rather slender (2° – 3° high); panicle few-flowered, nearly simple; flowers brown ($1\frac{1}{4}$ " long); outer sepals lanceolate and acute, inner a little shorter, obtusish, shorter than the broadly ovate rather triangular acute and pointed deep chestnut-brown pod; anthers as long as filaments; style short, seeds large ($\frac{1}{2}$ " or more long) obtuse, with short appendages at both ends, many-ribbed and reticulated. — Sphagnous swamps, on Broad Mountain, Pennsylvania, *C. E. Smith*.

4. **J. Bálticus**, Dethard. Scape rigid (2° – 3° high); panicle loose; flowers larger (2 " long), chestnut-brown with green; sepals ovate-lanceolate, the outer sharp-pointed, the inner obtusish, as long as the elliptical rather triangular obtuse and pointed deep brown pod; anthers much longer than the

broad filaments; style about the length of the ovary; seeds rather large ($\frac{1}{3}$ " or more long), nearly obtuse, delicately ribbed and cross-lined. — Sandy shores of New England, the Great Lakes, and westward; also in swamps, Lancaster Co., Penn., *Prof. Porter*. (Eu.)

+ + *Innermost sheaths leaf-bearing: stamens 6.*

5. **J. setaceus**, Rostkovius. Scape slender (1° – 3° high); panicle loose, rather few-flowered; flowers greenish ($2''$ long); sepals lanceolate, sharp-pointed, especially the 3 shining exterior ones, spreading in fruit, as long as the nearly globose beak-pointed greenish or light-brown pod; anthers as long as filaments; style conspicuous; seeds ($\frac{1}{3}$ " long) almost globose, ribbed and cross-lined. — Pennsylvania (*Rostkovius*), Virginia? and North Carolina, southward near the coast. — Doubtful if in our district.

* * *Flowers in clusters, 6-androus: innermost sheaths at base of stem leaf-bearing.*

6. **J. Roemerianus**, Scheele. Scape stout and rigid (2° – 3° high), its apex as well as the leaves pungent; panicle compound; 3–6 greenish or light-brown flowers ($1\frac{1}{2}$ " long) in a cluster; outer sepals lanceolate, sharp-pointed, longer than the obtusish inner ones, as long as the elliptical rather triangular obtuse short-pointed brown pod; anthers much longer than the broad filaments; styles shorter than the ovary; seeds ($\frac{1}{3}$ " long) oval, obtuse, very delicately ribbed. (*J. maritimus* of Amer. authors.) — Brackish marshes, New Jersey (*Pursh*), Virginia and southward.

§ 2. GRASSY-LEAVED JUNCI; with simple or rarely branched stems, leafy at base or throughout: leaves flat, or somewhat terete or setaceous and channelled, never knotted: panicle terminal.

* *Flowers crowded in heads (produced in late summer).*

+ *Leaves thread-like, hollow: stems simple: heads single or few: seeds large, the loose seed-coat produced into conspicuous tails: stamens 6.*

7. **J. stygius**, L. Stems slender (6'–16' high) from slender branching rootstocks, 1–3-leaved below, naked above; heads 1 or rarely 2, of 3–4-flowers, about the length of the sheathing scarious awl-pointed bract; flowers pale and reddish ($2\frac{1}{2}$ "– $3''$ long); sepals lanceolate, the inner obtusish, $\frac{3}{4}$ the length of the oblong acuminate pod, as long as the slender stamens; filaments many times longer than the oblong anthers; recurved stigmas shorter than the style; seeds oblong, with a very loose slightly striated coat prolonged at both ends ($1\frac{1}{2}$ " long). — Peat-bogs bordering Perch Lake, Jefferson County, New York. N. Maine, *G. L. Goodale*. (New Brunswick. *Mr. Fowler*. North shore of Lake Superior, *Mr. Wheeler*.) — Flowers larger and pod more abruptly pointed than in the European plant. (Eu.)

8. **J. trifidus**, L. Stems densely tufted from matted creeping rootstocks, erect (5'–10' high), sheathed and mostly leafless at the base, 2–3-leaved at the summit, the upper thread-like leaves subtending the sessile head of 1–4 brown ($1\frac{1}{2}$ "– $2''$ long) flowers; sepals ovate-lanceolate, acute, equalling or rather shorter than the ovate beak-pointed deep brown pod; anthers much longer than the filaments; seeds few, oblong, angled ($1''$ long), faintly striate, short-tailed. — Alpine summits of the mountains of New England, and N. New York, and high northward. (Eu.)

+ + *Leaves flat and open, grass-like : stamens 3.*

9. **J. marginatus**, Rostkovius. Stem erect, from a bulbous and stoloniferous base (1° – 3° high), flattened, leafy; leaves long-linear; heads 3–8-flowered, paniced; flowers purplish with green ($1\frac{1}{2}$ " long); sepals oblong, the 3 outer acute and slightly awned, the inner longer and mostly obtuse and pointless, as long as the almost globular scarcely pointed pod; stamens exceeding the outer sepals; purple anthers shorter than filaments; style very short; seeds ($\frac{1}{4}$ "– $\frac{1}{2}$ " long) slender, pointed at both ends and strongly ribbed. (*J. aristulatus*, Michx.) — Moist sandy places, S. New England to Illinois and southward. — Var. **PAUCICAPITATUS** has smaller and less flattened stems, narrower leaves, and few large 8–15-flowered heads. Sandy coast of New Jersey and elsewhere. — Var. **BIFLORUS** has numerous and small 2–3-flowered heads, in much-branched panicles. (*J. biflorus*, Ell.) New Jersey to Illinois and southward.

10. **J. repens**, Michx. Stems flattened, ascending (4'–6' high) from a fibrous annual root, at length creeping or floating; leaves short, linear, those of the stem nearly opposite and fascicled; heads few in a loose leafy panicle, 3–12-flowered; flowers green (3" long); sepals rigid, lance-subulate, slender-pointed, the 3 outer as long as the linear triangular obtuse pod, the inner ones much longer; stamens as long as the outer sepals; filaments many times longer than the oblong anther; seeds small ($\frac{1}{5}$ " long), obovate, slightly pointed, very delicately ribbed and cross-lined. (*Cephaloxys flabellata*, Desv.) — Miry banks, Maryland (*W. M. Canby*) and southward.

* * *Flowers single on the ultimate peduncles : stamens 6.*

+ *Stem branched : root annual.*

11. **J. bufonius**, L. Stems low and slender (3'–9' high) from a fibrous root, leafy, often branched from the base; panicle spreading, mostly with one-sided dichotomous branches; flowers remote, greenish ($2''$ – $3\frac{1}{2}''$ long); sepals linear-lanceolate, awl-pointed, 3 outer ones much longer than the inner and than the oblong obtuse pod; stamens short; filaments scarcely longer than anthers; seeds elliptical, obtuse ($\frac{1}{5}''$ – $\frac{1}{6}''$ long), very delicately ribbed and cross-lined. — Low grounds by roadsides: common, but not everywhere. June–Sept. — Var. **FASCICULATUS**, with flowers crowded at the end of the branchlets, is a southern form, which has been introduced about the Philadelphia Navy Yard. (Eu.)

+ + *Stems slender, simple, tufted, leafy below : root perennial (fl. early in summer).*

12. **J. Gerardi**, Loisel. (BLACK-GRASS.) Stems scarcely flattened, rigid (1° – 2° high); panicle contracted, usually longer than the bracteal leaf; flowers chestnut-brown with green ($1\frac{3}{4}$ " long); sepals oval-oblong, obtuse, incurved, as long as the oval obtuse and mucronate pod; anthers much longer than the short filaments; style as long as the ovary; seeds (nearly $\frac{1}{2}$ " long) obovate, short pointed at both ends, delicately ribbed and cross-lined. — Salt marshes: common along the coast, especially northward; also in saline marshes of W. New York: rare along the Great Lakes. (Eu.) — The closely allied *J. bulbosus*, L. (to which this was referred in a former edition) has not yet been found in this country.

13. **J. tenuis**, Willd. Stem wiry (9' - 18' high); leaves flat or channelled; panicle shorter than the involucreal leaves, loose, or rarely crowded; flowers green (2" or more long); sepals lanceolate, very acute, spreading in fruit, longer than the ovoid retuse scarcely pointed green pod; anthers nearly equal to the filament; style very short; seeds small (about $\frac{1}{4}$ " long), white-pointed at both ends, delicately ribbed and cross-lined. — Low grounds, fields, and roadsides. (Eu.) — Var. *SECUNDUS* is a smaller plant, with smaller one-sided flowers on the forked branches of the panicle. New England to Pennsylvania.

14. **J. dichotomus**, Ell. Stems rigid ($1\frac{1}{2}^{\circ}$ - 2° high) from a tumid base; leaves filiform, nearly terete, slightly grooved on the inner side; panicle loose, often with 1-sided forked branches, mostly longer than the involucreal leaf; flowers greenish (2" or more long); sepals lanceolate, sharp-pointed, spreading in fruit, as long as the globular beaked light mahogany-colored pod; anthers nearly as long as filaments; style short; seeds small ($\frac{1}{8}$ " - $\frac{1}{4}$ " long), white-pointed at both ends, coarsely cross-lined. — Low sandy grounds, New Jersey (*C. F. Parker*), Delaware (*Prof. Leidy, Mr. Commons*), and southward.

15. **J. Greenii**, Oakes & Tuckerm. Stems rigid (1° - 2° high); leaves nearly terete, very deeply channelled (almost involute) on the inner side; panicle usually much shorter than the principal erect involucreal leaf, dense; the numerous crowded flowers often one-sided ($1\frac{3}{4}$ " long); sepals lanceolate, acute, light greenish-brown, appressed, shorter than the ovoid-oblong obtuse greenish-brown pod; anthers as long as filaments; style very short; seeds ovoid, tail-pointed ($\frac{1}{2}$ " - $\frac{2}{5}$ " long), ribbed and delicately cross-lined. — Sandy coast of New England, and on the Great Lakes near Detroit (*Holzer, J. M. Bigelow*).

16. **J. Vaseyi**, Engelm. Stems rigid (1° - $2\frac{1}{2}^{\circ}$ high); leaves nearly terete, very slightly channelled on the inner side; panicle longer than the involucreal leaf, loose; flowers few, often one-sided, greenish or light brown (2" or more long); sepals lanceolate, acute, appressed, shorter than the oblong and retuse green-brown pod; anthers as long as the filaments; style very short; seeds slender, conspicuously tailed at both ends ($\frac{1}{2}$ " - $\frac{3}{8}$ " long), closely ribbed. — Michigan, with the last (near Detroit, *Holzer, Bigelow*) to N. Illinois (*Vasey*), and westward. June.

§ 3. **KNOTTY-LEAVED JUNCIS**; stems leafy, simple or branching: leaves terete or sometimes laterally compressed, knotted by internal cross-partitions: panicle terminal, with flowers chiefly in heads.

* Seeds barely pointed, without tails.

+ Heads composed of 2 or sometimes only single 6-androus flowers.

17. **J. pelocarpus**, E. Meyer. Stems slender and erect from a slender running rootstock (6' - 18' high), bearing few thread-like slightly knotted leaves, branching above into a compound spreading panicle, bearing in the forks and along one side of the branches single flowers or 2-flowered heads; often with the flowers or in place of them are tufts of leaves; flowers small (1' - $1\frac{1}{4}$ ' long), greenish with red; sepals oblong, obtuse, the 3 inner ones longer, but shorter than the oblong taper-beaked 1-celled pod; anthers much longer than the filaments; style slender; seeds ($\frac{1}{4}$ " long) obovate, short-pointed, delicately ribbed-reticulated. (*J. viviparus, Conrad. J. Conradi, Tuckerm.*) — Sandy, wet or swampy places,

Canada to Wisconsin, and from New England southward near the coast. July, Aug. — The proliferous plants are usually sterile and much larger than the fertile ones, with larger and more diffuse panicles. — Var. *SÚTILIS* (*J. fluitans*, *Michx.*), from Canada, is a small and creeping or floating form, mostly with single 2-flowered heads at the ends of the short stems.

+ + *Heads numerous, of 3-12 flowers (rarely more in No. 21), in early summer.*

++ *Stamens 6.*

18. *J. articulátus*, L. Stems ascending or erect (9' - 15' high), tufted from a short creeping rootstock, with 1 or 2 slender leaves; panicle short, spreading; the crowded heads 3-8-flowered; flowers brown, rarely pale (1¼" - 1½" long); sepals lance-oblong, acute or mucronate, or the 3 inner obtuse and a little longer, shorter than the ovate-oblong acute or abruptly mucronate-pointed incompletely 3-celled commonly deep chestnut-brown shining pod; anthers as long as filaments; ovary attenuated into a short style; seeds (less than ¼" long) obovate, attenuate below, abruptly pointed above, ribbed-reticulated. (*J. lam-pocárpus*, *Ehrh.*) — Wet grounds, New England to Western New York and Delaware. (Eu.) — Var. *OBTUSÁTUS*. Panicle crowded, level-topped; heads 5-flowered, green; sepals obtuse, of equal length, the outer mucronate; pod obtuse, mucronate. — Petty's Island, near Philadelphia, *Dr. Diffenbaugh, Mr. Burke.*

19. *J. alpinus*, Villars, var. *insignis*, Fries. Stems erect (9 - 18' high) from a creeping rootstock, with 1 or 2 slender leaves; panicle meagre, with erect branches bearing distant greenish or light-brown heads, each of 3-6 flowers (1¼" - 1½" long); sepals oblong, obtuse, the outer mucronate or cuspidate and usually longer than the rounded inner ones, as long or shorter than the obtuse short-pointed incompletely 3-celled light-brown pod; anthers as long as filaments; style short; seeds (½" or more in length) spindle-shaped, ribbed-reticulated. (*J. pelocárpus*, *Ed. 1.* *J. articulatus*, var. *pelocarpus*, *Ed. 2.*) — Wet sandy banks, from Lake Champlain (*Robbins, Macrae,*) and along the Great Lakes northward and westward. (Eu.)

20. *J. militáris*, Bigel. Stem stout (2° - 4° high) from a thick creeping rootstock, bearing a solitary stout erect leaf (½° - 3½° long) below the middle which overtops the crowded and rather contracted panicle; heads numerous, 5-12- (rarely 25-) flowered; flowers brownish (1½" long); sepals lanceolate, the outer ones awl-pointed, as long as the ovate-oblong triangular taper-beaked 1-celled pod; anthers longer than filaments; ovary attenuated into a slender style, seeds (¼" - ½" long) globose-obovate, obtuse, abruptly pointed, ribbed-reticulated. — In bogs and streams, Maine to Maryland. — Dr. Robbins finds that in flowing water, at Uxbridge, Mass., this bears numberless capillary submersed leaves, 2° - 3° long, from the rootstock.

++ ++ *Stamens 3.*

21. *J. acuminátus*, Michx. Stems tufted, erect, bearing about 2 leaves and a loose spreading panicle; heads few - many-flowered, greenish, at length straw-colored or darker; sepals lance-awl-shaped, sharp-pointed, equal, as long as or shorter than the triangular-prismatic short-pointed 1-celled pod; anthers a little shorter than the filaments; style almost none; seeds small (½" - ¾" long),

acute at both ends, ribbed-reticulated. (*J. palléscens*, *E. Meyer*, as to the N. American plant.) — Earlier than other species which are likely to be confounded with it: May, June. — A very variable plant, the following forms of which have the appearance of distinct species, but are connected by various intermediate states.

Var. **débilis**. Stems slender (9'–18' high); heads green, 3–6-flowered in a loose panicle; flowers smaller ($1\frac{1}{4}$ "– $1\frac{1}{2}$ " long); pod longer than the sepals. (*J. debilis*, *Gray*, in former editions.) — Wet sandy soil, New Jersey to Kentucky and southward. — Stems sometimes decumbent and rooting.

Var. **robústus**. Stems stout, tall, (2° – 4° high), bearing numerous 5–8-flowered light-brown heads in a large much-branched panicle; flowers small ($1''$ – $1\frac{1}{8}''$ long); ovoid pod scarcely longer than the sepals. — Deep swamps in the Mississippi Valley, from Illinois southward.

Var. **legítimus**. Stems slender (1° – 2° long), bearing fewer and larger heads in a very loose spreading panicle; flowers 5–12 and often more in a head ($1\frac{1}{2}''$ – $2''$ long); sepals as long as the straw-colored or light-brown pod. (*J. acuminatus*, *Michx.* *J. paradóxus*, *E. Meyer.* *J. fraternus*, *Kunth.* *J. subverticillátus*, *Muhl.* *J. Pondii*, *Wood.*) — Common in wet places from S. New England southward and westward. — Heads often proliferous in the autumn.

+ + + *Heads few, crowded, of numerous flowers.*

+ + *Stamens 6.*

22. **J. nodósus**, L. Stem erect (6'–15' high), slender from a creeping thread-like and tuber-bearing rootstock, mostly with 2 or 3 slender leaves; heads few or several, rarely single, 8–20-flowered ($3\frac{1}{2}''$ – $4''$ wide), overtopped by the involucre leaf; flowers brown ($1\frac{1}{2}''$ – $2''$ long); sepals lance-linear, awl-pointed (the 3 outer mostly a little shorter), nearly as long as the slender triangular taper-pointed 1-celled pod; anthers oblong, shorter than the filament; style very short; seeds (about $\frac{1}{4}''$ long) obovate, abruptly mucronate. (*J. Rostkovii*, *E. Meyer.*) — Swamps and gravelly banks, from New England and Pennsylvania north and northwestward. July, Aug. — A very tall form (2° high) occurs on the islands above Niagara, *G. W. Clinton.*

Var. **megacéphalus**, Torr. Stem stout (1° – 3° high), with thick leaves, heads few and large (6"–8" wide) 30–80-flowered; flowers pale green ($2\frac{1}{4}''$ – $2\frac{3}{4}''$ long); outer sepals longer than the inner ones; anthers linear, shorter than the filaments. — From Western New York west and southwestward. Aug. — An intermediate form, with numerous green heads in a compound panicle, occurs in Michigan and Northern Illinois.

+ + + *Stamens 3.*

23. **J. brachycárpus**, Engelm. Stem erect (1° – $2\frac{1}{2}^{\circ}$ high), rigid from a thick white horizontal rootstock, bearing about 2 leaves and 2–10 densely flowered spherical heads (4"–5" wide) in a slightly spreading crowded panicle much exceeding the involucre leaf; flowers pale green (2" long); sepals lance-linear, awl-pointed, the 3 outer much longer, and the ovoid pointed 1-celled pod rather shorter, than the inner sepals; anthers much shorter than filaments; style very short; seeds ($\frac{1}{8}''$ long) abruptly apiculate, ribbed-reticulated. — Moist places in open woods and prairies, Ohio to Michigan, Illinois and southward. May, June.

24. **J. scirpoides**, Lam. Stem erect (1° – 3° high), rigid, from a thick white horizontal rootstock, bearing about 2 leaves with wide and open sheaths, and a panicle of few or many densely-flowered pale-green spherical heads much longer than the involucre leaf; sepals rigid, awl-shaped and (especially the outer) bristly pointed, at length pungent, nearly equalling the oblong-triangular taper-pointed 1-celled pod; seeds ovoid, abruptly pointed at each end, ribbed-reticulated. (*J. polycéphalus*, *Michx.*) — The following forms belong here.

Var. **macrostemon**. Rather slender; leaves terete; branches of the panicle erect and often elongated; heads smaller ($3\frac{1}{2}''$ – $4''$ wide), 15–40-flowered; flowers $1\frac{1}{2}''$ – $1\frac{3}{4}''$ long; outer sepals mostly longer than the inner, as long as the stamens; anthers very small; seeds $\frac{1}{2}''$ long. (*J. echinatus*, *Muhl.* *J. macrostemon*, *Gay.*) — Wet sandy soil, from Staten Island (*C. F. Austin*) southward near the coast.

Var. **echinatus**. Stouter; leaves terete; branches of the compact panicle short; heads larger ($5''$ – $6''$ wide), 40–80-flowered; flowers ($1\frac{3}{4}''$ – $2''$ long); sepals narrower and more sharply pointed, the outer a little longer than the inner ones; stamens shorter and anthers longer than in the preceding, and seeds rather smaller and more slender. (*J. echinatus*, *Ell.*) — From Maryland southward.

Var. **polycéphalus**. Much stouter; leaves laterally flattened ($3''$ – $6''$ wide); panicle spreading, branched, bearing many distant heads as large as those of the foregoing form; flowers $2''$ – $2\frac{1}{2}''$ long; the 3 outer sepals the longer; anthers about as long as the filaments; seeds larger ($\frac{1}{2}''$ long). (*J. polycéphalus*, *Ell.*, *Chapm.*) — From North Carolina southward; and may be looked for in Southern Virginia.

* * *Seeds tailed.*

+ *Stamens 3.*

25. **J. Canadensis**, J. Gay. Tufted stems erect, terete, smooth, bearing 2–3 leaves; heads few- or many-flowered, paniculate; sepals lanceolate, the 3 outer shorter than the inner ones, not much longer than the stamens, equal to or shorter than the triangular prismatic almost 1-celled usually short-pointed pod; style mostly short; seeds more or less distinctly tail-pointed, delicately many-ribbed. — Common in most districts. Aug., Sept. — One of the latest flowering species, and thus easily distinguished, even when quite immature, from the similar but early *J. acuminatus*. This very variable species comprises the following forms.

Var. **longicaudatus**. Stem stout and rigid ($1\frac{1}{2}^{\circ}$ – 3° high), bearing in a decumbent somewhat spreading panicle the numerous 5–50-flowered heads; flowers greenish or light brown ($1\frac{1}{2}''$ – $2''$ long); sepals awl-pointed mostly shorter than the abruptly short-pointed pod; seeds slender ($\frac{3}{8}''$ – $1''$ long), conspicuously tail-pointed. (*J. paradoxus*, *Ed. 1 & 2*, *Chapman*, §c.) — From S. New England southward along the coast, and westward to Wisconsin and Northern Illinois.

Var. **subcaudatus**. Stem slender, often decumbent (1° – 2° high), bearing in simpler spreading panicles fewer 8–20-flowered heads; flowers greenish, as large as last; sepals awl-shaped, but not so rigid; pod mostly tapering; seeds

large ($\frac{1}{2}$ "– $\frac{3}{8}$ " long), with short white membranaceous appendages. — From S. New England southward, especially in New Jersey. — Often confounded with forms of No. 21, from which it is readily distinguished by the large, delicately ribbed and not reticulated, appendaged seeds, and by the proportion of the inner and outer sepals.

Var. **brachycéphalus**. Stem slender ($1\frac{1}{2}$ °– $2\frac{1}{2}$ ° high), bearing numerous small and 3–5-flowered heads in a large and spreading panicle; flowers greenish or light brown ($1\frac{1}{4}$ "– $1\frac{1}{2}$ " long); sepals mostly obtuse, shorter than the brown abruptly short-pointed pod; style longer than in other forms; seeds smaller ($\frac{1}{4}$ "– $\frac{1}{3}$ " long), slender, with rather short appendages. — From Pennsylvania northwestward to Illinois and Wisconsin. — Sometimes confounded with No. 18, and with small-flowered forms of No. 21.

Var. **coarctatus**. Stem slender, shorter (9'–18' high), bearing fewer deep-brown 3–5-flowered heads in a somewhat erect contracted panicle; flowers as large as in the last; sepals acute, or rarely obtusish, much shorter than the prismatic abruptly pointed deep-brown pod; seeds as in the last, (*J. acuminatus*, *Ed. 2*, and of most American authors, but not of Michaux). — New England to Wisconsin and northward, southward to the mountains of Pennsylvania.

+ + *Stamens 6.*

26. **J. asper**, Engelm. Stems tufted, erect (2°–3° high), terete, stout, rigid, and with the rigid leaves rough; panicle with rigid slightly spreading branches, bearing scattered few- (2–6-) flowered heads; flowers greenish with brown ($2\frac{1}{2}$ " long); sepals ovate-lanceolate, awl-pointed, rigid and strongly nerved, the outer much shorter than inner ones, these a little shorter than the triangular-ovoid beaked incompletely 3-celled brown pod; ovary tapering into a conspicuous style; seeds large, oblong, delicately many-ribbed, with white or often reddish appendages ($1\frac{1}{4}$ " long). — Sphagnous swamps, New Jersey, *Pickering, C. E. Smith.* August.

ORDER 123. PONTEDERIACEÆ. (PICKEREL-WEED FAMILY.)

Aquatic herbs, with perfect more or less irregular flowers from a spathe; the petal-like 6-merous perianth free from the 3-celled ovary; the 3 or 6 mostly unequal or dissimilar stamens inserted in its throat. — Perianth with the 6 divisions colored alike, *imbricated* in 2 rows in the bud, the whole together sometimes revolute-coiled after flowering, then withering away, or the base thickened-persistent and enclosing the fruit. Anthers introrse. Ovules anatropous. Style 1: stigma 3-lobed or 6-toothed. Fruit a perfectly or incompletely 3-celled many-seeded pod, or a 1-celled 1-seeded utricle. Embryo slender, in floury albumen.

1. **Pontederia**. Perianth 2-lipped, its fleshy persistent base enclosing the 1-seeded utricle. Stamens 6. Spike many-flowered.
2. **Heteranthera**. Perianth salver-shaped, withering-fugacious. Pod many-seeded. Stamens 3, unequal, of 2 forms. Spathe 1–few-flowered.
3. **Schollera**. Perianth salver-shaped, regular. Stamens 3, alike. Spathe 1-flowered.

1. PONTEDERIA, L. PICKEREL-WEED.

Perianth funnel-form, 2-lipped; the 3 upper divisions united to form the 3-lobed upper lip; the 3 lower spreading, and their claws, which form the lower part of the curving tube, more or less separate or separable down to the base: after flowering the tube is revolute-coiled from the apex downwards, and its fleshy-thickened persistent base encloses the fruit. Stamens 6; the 3 anterior exerted on elongated filaments; the 3 posterior (often sterile or imperfect) with very short filaments, unequally inserted lower down: anthers oval, blue. Ovary 3-celled; two of the cells empty, the other with a single suspended ovule. Utricle 1-celled, filled with the single seed. — Stout herbs, growing in shallow water, with thick creeping rootstocks, producing erect long-petioled mostly heart-shaped leaves, and a 1-leaved stem or scape, terminated by a spike of violet-blue ephemeral flowers. Root-leaves with a sheathing stipule within the petiole. (Dedicated to *Pontedera*, Professor at Padua at the beginning of the last century.)

1. *P. cordata*, L. Leaves arrow-heart-shaped, blunt; spike dense, from a spathe-like bract. — Var. *ANGUSTIFOLIA* (*P. angustifolia*, *Pursh*) has triangular-elongated and tapering leaves scarcely heart-shaped at the base. — Common. July – Sept. — Calyx-tube in fruit crested with 6 toothed ridges. Upper lobe of the perianth marked with a pair of small yellow spots.

2. HETERANTHERA, Ruiz & Pav. MUD-PLANTAIN.

Perianth salver-form with a slender tube; the limb somewhat equally 6-parted, ephemeral, soon withering or decaying. Stamens 3; the 2 posterior filaments thickened in the middle and bearing ovate (yellow) anthers; the other longer, bearing a larger oblong or arrow-shaped (greenish) anther. Pod incompletely 3-celled, many-seeded. — Creeping or floating low herbs, in mud or shallow water, with chiefly rounded long-petioled leaves, and a 1 – few-flowered spathe bursting from the sheathing side or base of a petiole. Flowers blue or white, in summer. (Name from *ἐρέπα*, *different*, and *ἀνθηρά*, *anther*.)

1. *H. reniformis*, Ruiz & Pav. Leaves round-kidney-shaped; spathe 3 – 5-flowered; flowers white. — S. New York to Illinois, and southward.

2. *H. limosa*, Vahl. Leaves oblong or lance-oblong, obtuse at both ends; spathe 1-flowered; flowers blue. (*Leptánthus ovalis*, *Michx.*) — W. Virginia to Illinois, and southward.

3. SCHOLLERA, Schreber (1789). WATER STAR-GRASS.

Perianth salver-form, with 6 nearly equal lance-linear divisions on a very long thread-like tube. Stamens 3, with similar oblong-arrow-shaped anthers (or rarely a fourth which is abortive): filaments nearly equal, awl-shaped. Pod oblong, invested by the withered perianth, 1-celled with 3 projecting parietal placentæ, many-seeded. — A grass-like herb, like a Pondweed, growing wholly under water, only the (small pale-yellow) flowers reaching the surface; the slender branching stems clothed with linear translucent sessile leaves, and bearing a terminal 1-flowered spathe: in summer. (Named after one *Scholler*, a German botanist.)

1. *S. graminea*, Willd. (*Leptánthus*, *Michx.*) — In streams: common.

ORDER 124. COMMELYNACEÆ. (SPIDERWORT FAMILY.)

Herbs, with fibrous or sometimes thickened roots, jointed and often branching leafy stems, and chiefly perfect and 6-androus, often irregular flowers, with the perianth free from the 2-3-celled ovary, and having a distinct calyx and corolla, viz. : Sepals 3, persistent, commonly herbaceous. Petals 3, ephemeral, decaying or deciduous. Stamens hypogynous, some of them often sterile : anthers with 2 separated cells. Style 1 : stigma undivided. Pod 2-3-celled, 2-3-valved, loculicidal, 3-several-seeded. Seeds orthotropous. Embryo small, pulley-shaped, partly sunk in a shallow depression at the apex of the albumen. Leaves ovate, lanceolate or linear, parallel-veined, flat, sheathed at the base ; the uppermost often dissimilar and forming a kind of spathe. — A chiefly tropical family, not aquatic, here represented only by two genera.

1. COMMELÿNA, Dill. DAY-FLOWER.

Flowers irregular. Sepals somewhat colored, unequal ; the 2 lateral partly united by their contiguous margins. Two lateral petals rounded or kidney-shaped, on long claws, the odd one smaller. Stamens unequal, 3 of them fertile, one of which is bent inward : 3 of them sterile and smaller, with imperfect cross-shaped anthers : filaments naked. Pod 3-celled, two of the cells 2-seeded, the other 1-seeded or abortive. — Stems branching, often procumbent and rooting at the joints. Leaves contracted at the base into sheathing petioles ; the floral one heart-shaped and clasping, folded together or hooded, forming a spathe enclosing the flowers, which expand for a single morning and are recurved on their pedicel before and afterwards. Petals blue. Flowering all summer. Ours all with perennial roots, or propagating by striking root from the joints. (Dedicated to the early Dutch botanists *J.* and *G. Commelyn.*)

1. *C. erecta*, L. Stem erect, rather stout (2° – 4° high) ; leaves large ($3'$ – $7'$ long, $1'$ – $2'$ wide), oblong-lanceolate, the upper surface and margins rough backwards, *sheaths fringed with rusty bristles ; spathes crowded and nearly sessile, hooded, top-shaped in fruit* ; odd petal like the others but smaller, round-ovate raised on a claw ; pod 3-celled. (*C. Virginica*, *Ed.* 1. A hairy form is *C. hirtella*, *Vahl.*) — Alluvial and shaded river-banks, Penn. to Illinois and southward. — Our largest species, and the only one with a top-shaped spathe.

2. *C. Virginica*, L. Stems slender, erect, or reclined and rooting towards the base ; leaves oblong- or linear-lanceolate ; *spathes mostly solitary or scattered, peduncled, conduplicate, round-heart-shaped when expanded, pointed, in fruit somewhat hood-like* ; odd petal usually inconspicuous and nearly sessile ; pod 2-celled. (*C. Virginica*, *L.*, as to syn. *Pluk.*, which gave the name : Linnæus's detailed description apparently pertains to No. 1, which however must bear the name which he took from Dillenius, the authority for the species. *C. angustifolia*, *Michx.*, is a narrow-leaved form.) — Damp rich woods and banks, S. New York to Michigan, Illinois, and southward.

3. *C. Cayennensis*, Richard. Stems creeping, glabrous ; *leaves ovate-oblong or lance-oblong, obtuse, small (1'-2' long) ; spathes heart-ovate when ex-*

panded, *peduncled, conduplicate, the base not contracted in fruit*, 3-4-flowered; the odd petal round-ovate, nearly sessile. (*C. agraria, Kunth.*)—Alluvial banks, Illinois and southward.—The smallest-leaved and smallest-flowered species.

2. TRADESCANTIA, L. SPIDERWORT.

Flowers regular. Sepals herbaceous. Petals all alike, ovate, sessile. Stamens all fertile: filaments bearded. Pod 2-3-celled, the cells 1-2-seeded.—Perennials. Stems mucilaginous, mostly upright, nearly simple, leafy. Leaves keeled. Flowers ephemeral, in umbelled clusters, axillary and terminal, produced through the summer: floral leaves nearly like the others. (Named for the elder *Tradescant*, gardener to Charles the First.)

* *Umbels sessile, clustered, usually involucrate by 2 leaves.*

1. **T. Virginica**, L. (COMMON SPIDERWORT.) *Leaves lance-linear, elongated, tapering from the sheathing base to the point, ciliate, more or less open; umbels terminal, many-flowered.*—Moist woods, from W. New York to Wisconsin, and southward: also commonly cultivated.—Plant either smooth or hairy; the large flowers blue, in gardens often purple or white.

2. **T. pilosa**, Lehm. *Leaves broadly lanceolate from a narrowed base, pointed, downy-hairy both sides, minutely ciliate; umbels many-flowered, in very dense terminal and axillary clusters; pedicels and calyx glandular-hairy.* (*T. flexuosa, Raf.*)—Ohio, Illinois, Kentucky, and southward.—Stem stout, smooth below, 2°-3° high, often branched, zigzag above, forming a close cluster of small (8''-9'' broad) lilac-blue flowers in all the upper axils.

** *Umbels long-peduncled, naked.*

3. **T. rosea**, Vent. Small, slender (6'-10' high), smooth; leaves linear, grass-like, ciliate at the base; umbel simple, or a pair; flowers (6'' wide) rose-color.—Sandy woods, Pennsylvania (?) to Kentucky, and southward.

ORDER 125. XYRIDACEÆ. (YELLOW-EYED-GRASS FAMILY.)

Rush-like herbs, with equitant leaves sheathing the base of a naked scape, which is terminated by a head of perfect 3-androus flowers, with extrorse anthers, glumaceous calyx, and a regular colored corolla; the 3-valved mostly 1-celled pod containing several or many orthotropous seeds with a minute embryo at the apex of fleshy albumen:—represented by *Xyris*.—But the anomalous genus *Mayaca*, consisting of a few moss-like aquatic plants, intermediate in character between this family and the last, may be introduced here.

1. MAYACA, Aublet. (SYENA, Schreber.)

Flowers single, terminating a naked peduncle. Perianth persistent, of 3 herbaceous lanceolate sepals, and 3 obovate petals. Stamens 3, alternate with the petals. Ovary 1-celled with 3 parietal few-ovuled placentæ: style filiform: stigma simple. Pod 3-valved, several-seeded.—Moss-like low herbs, creeping or floating in shallow water, densely leafy; the leaves narrowly linear, sessile,

1-nerved, pellucid, entire, notched at the apex: the peduncle solitary, sheathed at the base. (An aboriginal name.)

1. **M. Michauxii**, Schott & Endl. Peduncles not much exceeding the leaves, nodding in fruit; petals white. (*Syena fluviatilis*, *Pursh.*) — S. E. Virginia, and southward. July.

2. XÝRIS, L. YELLOW-EYED GRASS.

Flowers single in the axils of coriaceous scale-like bracts, which are densely imbricated in a head. Sepals 3; the 2 lateral glume-like, boat-shaped or keeled and persistent; the anterior one larger and membranaceous, enwrapping the corolla in the bud and deciduous with it. Petals 3, with claws, which cohere more or less. Fertile stamens 3, with linear anthers, inserted on the claws of the petals, alternating with 3 sterile filaments, which are cleft and in our species plumose or bearded at their apex. Style 3-cleft. Pod oblong, free, 1-celled, with 3 parietal more or less projecting placentæ, 3-valved, many-seeded. — Flowers yellow, produced all summer. Ours apparently all perennials. (*Ξυρίς*, an ancient name of some plant with 2-edged leaves, from *ξυρόν*, a razor.)

1. **X. flexuosa**, Muhl., Chapm. Scape slender (10' - 16' high), barely flattened at the summit, often from a bulbous base, very smooth, much longer than the narrowly linear leaves, both commonly twisted with age; head roundish-ovoid (3" - 4" long); *lateral sepals* oblong-lanceolate, *finely ciliate-scarious on the narrow wingless keel*, and usually with a minute bearded tuft at the very apex, shorter than the bract. (*X. Jupacai*, *Michx.* in part. *X. Indica*, *Pursh.* *X. bulbosa*, *Kunth*, & Ed. 2.) — Sandy or peaty bogs, from E. Massachusetts southward near the coast; also Illinois, Wisconsin, and Michigan.

Var. **pusilla**. Small and very slender, seldom twisted, 2' - 9' high, the base not bulbous. (*X. brevifolia*, *Muhl.*, in part, &c., not of *Michx.*) — From New Jersey and Pennsylvania northward to the base of the White Mountains and Lake Superior. — Head 2" - 3" long.

2. **Xyris torta**, Smith. Scape terete and one-edged, slender, 9' - 20' high, from a bulbous base, and with the *linear-filiform rigid leaves* becoming spirally twisted; head ovoid becoming spindle-shaped and acute (5" - 9" long); sepals exceeding the bract; *lateral sepals winged on the keel and fringed above the middle*. — Pine barrens of New Jersey (near Batsto, *D. C. Eaton*) and in the Southern States: in dry sand.

3. **X. Caroliniana**, Walt. Scape flattish, 1-angled below, 2-edged at the summit, smooth, 1° - 2° high, the base hardly bulbous; *leaves linear-sword-shaped, flat*; head globular-ovoid (5" - 7" long); *lateral sepals obscurely lacinate-fringed above on the winged keel*, rather shorter than the bract. (*X. Jupacai*, partly, *Michx.* *X. anceps*, *Muhl.*) — Sandy swamps, &c., Rhode Island to Virginia and southward, near the coast.

4. **X. fimbriata**, Ell. Scape somewhat angled, 2-edged above, rough (2° high), rather longer than the linear-sword-shaped or strap-shaped leaves, the base not bulbous; head oblong (6" - 10" long); *lateral sepals lanceolate-linear, nearly twice the length of the bract, above the middle conspicuously fringed on the wing-margined keel, and even plumose at the summit*. — Pine barrens of New Jersey to Virginia, and southward.

ORDER 126. **ERIOCAULONACEÆ.** (PIPEWORT FAMILY.)

Aquatic or marsh herbs, stemless or short-stemmed, with a tuft of fibrous roots, a cluster of linear and often loosely cellular grass-like leaves, and naked scapes sheathed at the base, bearing dense heads of monœcious or rarely diœcious small 2-3-merous flowers, each in the axil of a scarious bract; the perianth double or rarely simple, chaffy; anthers introrse; the fruit a 2-3-celled 2-3-seeded pod: the ovules, seeds, embryo, &c. as in the preceding order. — Chiefly tropical plants, a few in northern temperate regions.

1. **Eriocaulon.** Perianth double, the inner (corolla) tubular-funnel-form in the staminate flowers; the stamens twice as many as its lobes (4 or 6). Anthers 2-celled.
2. **Præpalanthus.** Perianth as in the last: the stamens only as many as the lobes of the inner series, or corolla (3). Anthers 2-celled.
3. **Lachnocaulon.** Perianth simple, of 3 sepals. Stamens 3, monadelphous below. Anthers 1-celled.

1. **ERIOCAULON,** L. PIPEWORT.

Flowers monœcious and androgynous, i. e. both kinds in the same head, either intermixed, or the central ones sterile and the exterior fertile, rarely diœcious. *Ster. Fl.* Calyx of 2 or 3 keeled or boat-shaped sepals, usually spatulate or dilated upwards. Corolla tubular, 2-3-lobed, each of the lobes bearing a black gland or spot. Stamens twice as many as the lobes of the corolla, one inserted at the base of each lobe and one in each sinus: anthers 2-celled. Pistils rudimentary. *Fert. Fl.* Calyx as in the sterile flowers, often remote from the rest of the flower (therefore perhaps to be viewed as a pair of bractlets). Corolla of 2 or 3 separate narrow petals. Stamens none. Ovary often stalked, 2-3-lobed, 2-3-celled, with a single ovule in each cell: style 1: stigmas 2 or 3, slender. Pod membranaceous, loculicidal. — Leaves mostly smooth, loosely cellular and pellucid, flat or concave above. Scapes or peduncles terminated by a single head, which is involucre by some outer empty bracts. Flowers, also the tips of the bracts, &c., usually white-bearded or woolly. (Name compounded of *ἔριον*, wool, and *καυλός*, a stalk, from the wool at the base of the scape and leaves of the original species. Excepting this and the flowers, our species are wholly glabrous.) — The North American species are all stemless, with a depressed head, and have the parts of the flowers in twos, the stamens 4.

1. **E. decangulære**, L. (*syn. Pluk. &c.*) Leaves obtuse, varying from linear-lanceolate to linear-awl-shaped, rather rigid; scapes 10-12-ribbed (1°-3° high; head hemispherical, becoming globular (2"-7" wide); scales of the involucre acutish, straw-color or light brown; chaff (bracts among the flowers) pointed. (*E. serotinum*, *Walt.*) — Pine-barren swamps (New Jersey?) Virginia, and southward. July - Sept.

2. **E. gnaphalodes**, Michx. Leaves spreading (2'-5' long), grassy-awl-shaped, rigid, or when submersed thin and pellucid, tapering gradually to a sharp point, mostly shorter than the sheath of the 10-ribbed scape; scales of the involucre very obtuse, turning lead-color; chaff obtuse. (*E. decangulare*, L., in

part, viz. as to *pl. Clayt.* *E. compressum*, *Lam.*) — Pine-barren swamps, New Jersey to Virginia, and southward. June — Aug.

3. **E. septanguläre**, Withering. *Leaves short* (1' — 3' long), *awl-shaped, pellucid*, soft and very cellular; *scape 7-striate*, slender, 2' — 6' high, or when submersed becoming 1° — 6° long, according to the depth of the water; *chaff acutish*. (*E. pellucidum*, *Michx.*) — In ponds or along their borders, from New Jersey and Penn. to Michigan, and northward. Aug. — Head 2'' — 3'' broad; the bracts, chaff, &c. lead-color, except the white coarse beard. (Eu. Coast of Ireland, &c. only.)

2. PÆPALÁNTHUS, Mart. (Sp. of ERIOCAULON of authors.)

Stamens as many as the (often involute) lobes of the funnel-form corolla of the sterile flowers, and opposite them, commonly 3, and the flower ternary throughout. Otherwise nearly as in *Eriocaulon*. (Name from *παιπάλη*, *dust* or *flour*, and *ἄνθος*, *flower*, from the meal-like down or scurf of the heads and flowers of many South American species.)

1. **P. flávidus**, Kunth. Tufted, stemless; leaves bristle-awl-shaped (1' long); scapes very slender, simple, minutely pubescent (6' — 12' high), 5-angled; bracts of the involucre oblong, pale straw-color, those among the flowers mostly obsolete; perianth glabrous; sepals and petals of the fertile flowers linear-lanceolate, scarious-white. (*Eriocaulon flavidum*, *Michx.*) — Low pine barrens, S. Virginia and southward.

3. LACHNOCAÛLON, Kunth. HAIRY PIPEWORT.

Flowers monœcious, &c., as in *Eriocaulon*. Calyx of 3 sepals. Corolla none! *Ster. Fl.* Stamens 3: filaments below coalescent into a club-shaped tube around the rudiments of a pistil, above separate and elongated: anthers 1-celled! *Fert. Fl.* Ovary 3-celled, surrounded by 3 tufts of hairs (in place of a corolla). Stigmas 3, two-cleft. — Leaves linear-sword-shaped, tufted. Scape slender, simple, bearing a single head, 2 — 3-angled, hairy (whence the name from *λάχνος*, *wool*, and *καυλός*, *stalk*).

1. **L. Michauxii**, Kunth. (*Eriocaulon villosum*, *Michx.*) — Low pine barrens, Virginia (*Pursh*), and southward.

ORDER 127. CYPERÀCEÆ. (SEDGE FAMILY.)

Grass-like or rush-like herbs, with fibrous roots, mostly solid stems (culms), closed sheaths, and spiked chiefly 3-androus flowers, one in the axil of each of the glume-like imbricated bracts (scales, glumes), destitute of any perianth, or with hypogynous bristles or scales in its place; the 1-celled ovary with a single erect anatropous ovule, in fruit forming an achenium. Style 2-cleft when the fruit is flattened or lenticular, or 3-cleft when it is 3-angular. Embryo minute at the base of the somewhat floury albumen. Stem-leaves when present 3-ranked. — A large, widely diffused family. (See Plates 1 — 6.)

Tribe I. SCIRPINEÆ. Flowers normally perfect and alike, rarely some of them with stamens or pistil abortive: spikes all of one sort.

Subtribe 1. Cyperæ. Scales of the spike strictly 2-ranked, conduplicate and keeled (in all our species). Spikes usually aggregated into clusters, heads, umbels, spikes, &c.

* Flower destitute of any bristles, also of any beak to the achenium.

1. **Cyperus.** Spikes few-many-flowered, usually elongated or slender.
2. **Kyllingia.** Spikes 1-flowered (but of 3 or 4 scales), glomerate in a sessile head.

* Flower furnished with bristles: achenium beaked by the persistent base of the style.

3. **Dulichium.** Spikes 6-10-flowered, slender, clustered on an axillary peduncle.

Subtribe 2. Scirpæ. Scales of the spike or head imbricated all round, convex or open, all or nearly all of them with a perfect flower in the axil.

* Flower with one or more inner scales (either bractlets or perianth).

4. **Fuirena.** Scales of the spike awned below the apex. Flower surrounded by 3 stalked petal-like scales alternating with 3 bristles.
5. **Lipocarpa.** Flower enclosed by 2 inner scales, one next the axis and the other next the awnless scale of the spike. No bristles.
6. **Hemicarpha.** Flower with a single very minute inner and thin scale next the axis of the spike: no bristles.

** Flower without inner scales, either wholly naked or with some bristles (perianth).

← Bristles generally present.

7. **Eleocharis.** Achenium with a tubercle jointed on its apex, consisting of the bulbous persistent base of the style. Spike solitary, terminating the leafless and bractless culm.
8. **Scirpus.** Achenium naked at the apex, or pointed with the continuous simple base of the style. Culms often leafy at the base or apex. Spikes one or more.
9. **Eriophorum.** Achenium, &c., as in *Scirpus*. Bristles after flowering exceedingly lengthened into cottony hairs.

← ← Bristles always none.

6. **Hemicarpha** will be looked for here when the minute inner scale is overlooked.
10. **Fimbristylis.** Style bulbous at the base, or constricted at the junction, deciduous from the achenium (with or rarely without the jointed bulb). Culms leafy at the base, and bracted at the summit, bearing usually several spikes.

Subtribe 3. Rhynchosporæ. Scales of the spike or head imbricated in few or several ranks, some of the lower commonly empty, and of the upper subtending abortive or staminate flowers. No inner scales.

11. **Dichromena.** Spikes crowded into a leafy-involucrate head, laterally flattened, the scales more or less conduplicate and keeled. Achenium crowned with a beak or tubercle formed of the enlarged persistent base of the style. No bristles.
12. **Rhynchospora.** Spikes terete or flattish, the scales convex, and either loosely enwrapping or regularly imbricated. Achenium crowned with a persistent tubercle or beak, and at the base commonly surrounded by bristles.
13. **Cladium.** Spikes terete, few-flowered, the scales, &c. as in the preceding. Achenium destitute of tubercle. No bristles.

Tribe II. SCLERINEÆ. Flowers monœcious; the staminate and pistillate in the same or in different clustered spikes. Achenium naked, bony or crustaceous, supported on a hardened disk.

14. **Scleria.** Spikes few-flowered: lower scales empty. No bristles, &c.

Tribe III. CARICINEÆ. Flowers monoœcious in the same (androgynous) or in separate spikes, or sometimes dioœcious. Achenium enclosed in a sac (*perigynium*), which answers to a single or a pair of inner scales or bractlets.

15. **Carex.** No bristle-form hooked appendage projecting from the sac which encloses the achenium.

1. CYPÈRUS, L. GALINGALE. (Pl. 1.)

Spikes many - few-flowered, mostly flat, variously arranged, mostly in clusters or heads, which are commonly disposed in a simple or compound terminal umbel. Scales 2-ranked, conduplicate and keeled (their decurrent base below often forming margins or wings to the hollow of the joint of the axis next below), deciduous when old. Stamens 1-3. No bristles or inner scales. Style 2-3-cleft, deciduous. Achenium lenticular or triangular, naked at the apex. — Culms mostly triangular, simple, leafy at the base, and with one or more leaves at the summit, forming an involucre to the umbel. Peduncles unequal, sheathed at the base. All flowering in late summer or autumn. (Κύπερος, the ancient name.)

§ 1. PYCRÈUS, Beauv. *Style 2-cleft: achenium flattened: spikes flat, many-flowered: only the lowest scale empty. (Ours all annuals.)*

1. **C. flavescens**, L. *Stamens 3; spikes becoming linear, obtuse, clustered on the 2-4 very short rays (peduncles); scales obtuse, straw-yellow; achenium shining, orbicular.* — Low grounds, mostly near the coast. — Culms 4' - 10' high; spikes 5'' - 8'' long. Involucre 3-leaved, very unequal. (Eu.)

2. **C. diandrus**, Torr. *Stamens 2, or sometimes 3; spikes lance-oblong, scattered or clustered on the 2-5 very short or unequal rays; scales rather obtuse, purple-brown on the margins or nearly all over; achenium dull, oblong-obovate: otherwise much like the last.* — Var. **CASTANEUS**, Torr. (*C. castaneus*, Bigel.) is only a form with browner scales. — Low grounds: common.

3. **C. Nuttállii**, Torr. *Stamens 2; spikes lance-linear, acute and very flat ($\frac{1}{2}$ ' - 1' long), crowded on the few very short (or some of them distinct) rays; scales oblong, yellowish-brown, rather loose; achenium oblong-obovate, very blunt, dull.* — Salt or brackish marshes, Massachusetts to Virginia, and southward. — Culms 4' - 12' high. — *C. minimus?* Nutt. (*C. Cleaveri*, Torr.) is a depauperate form of this, with a 1-leaved involucre, and only one or two spikes!

4. **C. flavicomus**, Michx. *Stamens 3; spikes linear (4'' - 9'' long), spiked and crowded on the whole length of the branches of the several-rayed umbel, spreading; scales oval, very obtuse, yellowish and brownish, broadly scarious-(whitish-) margined; achenium obovate, mucronate, blackish; culm stout (1° - 3° high); leaves of the involucre 3-5, very long.* — Low grounds, Virginia and southward.

§ 2. PAPÏRUS, Thouars. *Style 3-cleft: achenium triangular: stamens 3: spikes many-flowered: the scarious winged margins of the joints of the axis early separating down to the base in the form of a pair of free scales, which are persistent after the proper scale falls away: otherwise as in § 3. (Ours annual.)*

5. **C. erythrorhizos**, Muhl. *Culm obtusely triangular (3' - 3° high); umbel compound, many-rayed; involucre 4-5-leaved, very long; involucels bristle-form; spikes very numerous, crowded in oblong or cylindrical nearly sessile heads or spikes, spreading horizontally, linear, flattish (3'' - 6'' long), bright chestnut-colored; scales lanceolate, mucronulate.* — Alluvial banks, Penn. to Wisconsin? Illinois, and common southward. — Root fibrous, red.

§ 3. **CYPERUS** proper. *Style 3-cleft: achenium triangular: spikes many-flowered, flat or almost terete; only the lowest scale empty; the joints of the axis wing-margined or naked.*

* *Stamen only one: spikes short and small (2''-3'', or becoming 4''-5'' long), collected in globular heads, ovate or linear-oblong, compactly many-flowered: the scales merely ascending on the naked marginless axis.*

+ *Low annuals: involucre 2-3-leaved: heads few: scales pointed.*

6. **C. inflexus**, Muhl. Dwarf (1'-5' high); *spikes oblong becoming linear, 7-13-flowered, in 1-5 ovate heads (either sessile and clustered or short-peduncled); scales nerved, tapering into a long recurved point; achenium obovate, obtuse.* — Sandy wet shores: common. — Sweet-scented in drying.

7. **C. acuminatus**, Torr. Slender (3'-12' high); *spikes ovate, becoming oblong, 16-30-flowered, pale; scales obscurely 3-nerved, short-tipped; achenium oblong, pointed at both ends.* — Low ground, Illinois and southwestward.

+ + *Tall perennial (1°-4° high): heads many, greenish: scales pointless.*

8. **C. virens**, Michx. Culm either sharply or obtusely triangular; leaves and involucre very long, keeled; umbel compound, many-rayed; spikes ovate or oblong; achenium oblong or linear; scales acutish, obscurely 3-nerved. (*C. végetus*, Torr.) — Wet places, Virginia and southward.

* * *Stamens 3: spikes 3-10 in a cluster at the summit of the culm or of each ray of an umbel, flat, oblong or linear, the crowded scales ascending, strongly keeled, ovate, with abruptly sharp-pointed slightly-spreading tips: achenium broadly obovate, sharply triangular.*

9. **C. compréssus**, L. *Low annual (3'-9' high), with a single sessile or a few umbellate clusters of linear green spikes (of 6''-10'' long, 15-30-flowered), the axis naked.* — Sterile fields, Somerset Co., Maryland, *W. M. Canby* (also adventive near Philadelphia Navy Yard), and southward.

10. **C. dentatus**, Torr. *Perennial, with running rootstocks (6'-12' high); clusters several or many in a simple or twice or thrice compound umbel; spikes ovate-oblong and rather few-flowered, or when well-developed linear and 15-30-flowered (3''-5'' long); the scales reddish-brown with green back; joints of the axis wing-margined.* — Sandy swamps, Massachusetts to Northern New York (*J. A. Paine*), and southward. — Spikes often abortive and changed into leafy tufts as in *Juncus*.

* * * *Stamens 3: spikes narrowly linear, flat, scattered along the upper part of the rays of an open simple or somewhat compound umbel: scales appressed, pointless: joints of the spike wing-margined: perennial by slender running rootstocks, which bear small nut-like tubers, by which the plants multiply pestiferously in cultivated grounds.*

11. **C. rotundus**, L., var. **Hydra**. (NUT-GRASS.) Culm slender ($\frac{1}{2}^{\circ}$ - $1\frac{1}{2}^{\circ}$ high), longer than the leaves; umbel simple or slightly compound, about equalling the involucre; the few rays each bearing 4-9 dark chestnut-purple 12-40-flowered acute spikes (5''-9'' long); scales ovate, closely appressed, nerveless except on the keel. (*C. Hydra*, Michx.) — Sandy fields, Virginia and southward: and adventive near Philadelphia, *C. F. Parker*. (Eu.)

12. **C. phymatodes**, Muhl. Culm (1° – $2\frac{1}{2}^{\circ}$ high) equalling the leaves; umbel often compound, 4–7-rayed, much shorter than the long involucre; spikes numerous, light chestnut or straw-color, acutish, 12–30-flowered ($4''$ – $7''$ long); scales oblong, narrowly scarious-margined, nerved, the acutish tips rather loose; achenium oblong. (*C. repens*, Ell.)—Low grounds, along rivers, &c., Vermont to Wisconsin, and common southward.

* * * * *Stamens 3: spikes narrowly linear or slender-awl-shaped, numerous and densely crowded or spiked on the summit of the rays of the open simple or sometimes compound umbel, spreading or sometimes reflexed: scales erect-appressed, conduplicate or keeled, pointless: joints of the axis of the spike with scaly-winged margins partly embracing the achenium: involucre of 3 to several long leaves: annuals with fibrous roots, or sometimes apparently more or less perennial from a tuberos or bulbous thickened base: no running rootstocks.*

+ *Spikes flat, becoming straw-color ($\frac{1}{2}'$ – $1'$ long); the scales strongly conduplicate.*

13. **C. strigosus**, L. Culm mostly stout (1° – 3° high); most of the rays of the umbel elongated ($1'$ – $5'$), their sheaths 2-bristled; spikes 10–25-flowered, scales oblong-lanceolate, several-nerved, much longer than the oblong-linear achenium.—Damp or fertile soil: very common, especially southward. (*C. stenolepis*, Torr., is of this group and nearly related.)

+ + *Spikes slender and rather awl-shaped, almost terete, at least when mature; the scales less conduplicate and more appressed to the axis.*

14. **C. Michauxianus**, Schultes. Culm stout, mostly low ($5'$ – $20'$ high); rays of the umbel mostly all short and crowded; spikes 10–20-flowered, yellowish-brown at maturity ($3''$ – $7''$ long), the short joints of its axis winged with very broad scaly margins which embrace the ovate triangular achenium; the scales ovate, obtusish, imbricately overlapping.—Low grounds and sandy banks: common.—Root truly annual: stem seldom bulbous-thickened at the base.

15. **C. Engelmanni**, Steud. Resembles the foregoing; but the spikes more slender and terete, somewhat remotely 5–15-flowered, the zigzag joints of the axis slender and narrowly winged, and the oblong or oval broadly scarious scales proportionally shorter, so as to expose a part of the axis of each joint, the successive scales not reaching the base of the one above on the same side; achenium oblong-linear, very small. (*C. tenuior*, Engelm.)—Low grounds and sandy banks, Virginia to Wisconsin and southward: also adventive at the Philadelphia Navy Yard.

* * * * *Stamens 3: spikes loosely or somewhat remotely 6–12-flowered, flattish and greenish, several crowded together in one sessile or in a few peduncled heads or dense clusters; their scales ovate, convex on the back, many-nerved, applied to and little longer than the ovate or obovate and sharply triangular achenium: perennials, propagating from the hard clustered corms or bulb-like tubers at the base of the culms.*

16. **C. Schweinitzii**, Torr. Culm rough on the angles (1° – 2° high); leaves linear; umbel simple, 4–8-rayed; spikes crowded along the upper part of the mostly elongated rays, erect, loose ($4''$ – $6''$ long); scales awl-pointed; joints of the axis narrowly winged.—Dry sandy shores and ridges, from Lake Ontario to Ohio, Illinois, and northwestward.—Flowers rather large.

17. **C. Gràyii**, Torr. Culm thread-form, wiry (6'–12' high); leaves almost bristle-shaped, channelled; umbel simple, 4–6-rayed; spikes 5–10 in a loose head, spreading; joints of the axis winged; scales rather obtuse, greenish-chestnut-color. — Barren sands, Plymouth, Mass., to New Jersey, near the coast.

18. **C. filicúlmis**, Vahl. Culm slender, wiry, often reclined (8'–15' high); leaves linear (1"–2" wide); spikes numerous and clustered in one sessile dense head, or in 1–7 additional looser heads on spreading rays of an irregular umbel; joints of the axis naked; scales blunt, greenish. (*C. mariscoides*, Ell.) — Dry sterile soil: common, especially southward.

§ 4. **MARÍSCUS**, Vahl. Style 3-cleft: the achenium triangular: stamens 3: spikes 1–few-flowered, narrow or awl-shaped, with 2 lower scales short and empty, and inclined to persist on the common axis when the rest of the spike disarticulates and falls, crowded in dense heads; otherwise nearly as in the penultimate division of § 3. (*Perennials with clustered small tubers at base of the culms, as in the preceding division: spikes green, merely tawny with age.*)

19. **C. Lancastriénsis**, T. C. Porter, n. sp. Culm (1°–2° high) triangular; leaves rather broadly linear; umbel of 6–9 mostly elongated rays; spikes very numerous in short-oblong or globular dense heads, soon reflexed, 3–6-flowered, linear-awl-shaped; the joints of the axis broadly winged; scales oblong, obtuse, twice the length of the linear-oblong achenium. — Rich soil, banks of the Susquehanna near Lancaster, Penn., *Prof. Porter*. — Most like the Southern *C. Baldwinii*, Torr.; but twice the size; the more numerous spikes 4"–5" long, more linear, less pointed, on a setaceous-bracted axis of 6" or 7" in length, with longer scales and achenium, &c.

20. **C. ovularis**, Torr. Culm sharply triangular (6'–12' high); umbel 1–6-rayed; spikes (50–100) in a globular very dense head, 2–4-flowered, oblong, blunt (1½"–2" long); joints of the axis winged; scales ovate, obtuse, a little longer than the obovate-oblong achenium. — Sandy dry soil, S. New York to Illinois and southward.

21. **C. retrofractus**, Torr. Culm and leaves usually minutely downy and rough on the obtusish angles (1°–3° high); umbel many-rayed; spikes slender-awl-shaped, very numerous in obovate or oblong heads terminating the elongated rays, soon reflexed, 1–2-flowered in the middle (3"–5" long); scales usually 4 or 5, the two lowest ovate and empty, the fertile lanceolate and pointed, the uppermost involute-awl-shaped; achenium linear. (*Scirpus retrofractus*, L.) — Sandy fields, New Jersey to Virginia, and southward.

2. KYLLÍNGIA, Rottböll. KYLLINGIA. (Pl. 1.)

Spikes of 3 or 4 two-ranked scales, 1–1½-flowered; the 2 lower scales minute and empty, as in *Cyperus*, § 4, but style oftener 2-cleft, and achenium lenticular: the spikes densely aggregated in solitary or triple sessile heads. Involucre 3-leaved. (Named after *Peter Kylling*, a Danish botanist of the 17th century.)

1. **K. pumila**, Michx. Head globular or 3-lobed, whitish-green (about 4" broad); spikes strictly 1-flowered; upper scales ovate, pointed, rough on the keel; stamens and styles 2; leaves linear. — Low grounds, Ohio to Illinois and southward. Aug. — Culms 2'–9' high; root annual.

3. DULÍCHIUM, Richard. DULICHUM. (Pl. 1.)

Spikes many- (6 - 10-) flowered, linear, flattened, sessile in 2 ranks on axillary solitary peduncles emerging from the sheaths of the leaves: scales 2-ranked, lanceolate, decurrent, forming flat wing-like margins on the joint below. Perianth of 6 - 9 downwardly barbed bristles. Stamens 3. Style 2-cleft above. — Achenium flattened, linear-oblong, beaked with the long persistent style. — A perennial herb, with a terete simple culm (1° - 2° high), jointed and leafy to the summit; the leaves short and flat, linear, 3-ranked. (The name of a Greek island; its singular application to an American plant unexplained.)

1. **D. spathaceum**, Pers. — Borders of ponds: common. July - Sept.

4. FUIRÈNA, Rottböll. UMBRELLA-GRASS. (Pl. 2.)

Spikes many-flowered, terete, clustered or solitary, axillary and terminal. Scales imbricated in many ranks, awned below the apex, all floriferous. Perianth of 3 ovate or heart-shaped petaloid scales, mostly on claws, and usually with as many alternating small bristles. Stamens 3. Style 3-cleft. Achenium triangular, pointed with the persistent base of the style. Culms from a perennial root, obtusely triangular. (Named for *G. Fuiren*, a Danish botanist.)

1. **F. squarrosa**, Michx. Stem (1° - 2° high) leafy; leaves and sheaths hairy; spikes ovoid-oblong (6" long), clustered in heads, bristly with the spreading awns of the scales; perianth-scales ovate and awn-pointed, the interposed bristles minute. — Var. **PUMILA**, Torr., is a dwarf form, 1' - 6' high, with 2 - 6 spikes; perianth-scales ovate-lanceolate and oblanceolate. — Sandy wet places, Massachusetts to Virginia, and southward; also Michigan: northward mostly the small variety. Aug.

5. LIPOCÁRPHA, R. Br. LIPOCARPHA. (Pl. 2.)

Spikes terete, many-flowered, in a terminal close cluster, which is involucrate by leafy bracts. Scales spatulate, regularly imbricated all round in many ranks, awnless, deciduous, a few of the lowest empty. Inner scales (bractlets) 2 to each flower, thin, one between the scale of the spike and the flower, one between the latter and the axis of the spike. Bristles or other perianth none. Stamens 1 or 2. Style 2 - 3-cleft. Achenium flattish or triangular, naked at the tip. — Culms leafy at the base. (Name formed of *λίπος*, *fat*, and *κάρφος*, *chaff*, from the thickness of the inner scales of some species.)

1. **L. maculata**, Torr. Annual; culm (4' - 8' high) much longer than the linear concave leaves; spikes (1" - 2" long) green and dark-spotted; inner scales delicate; stamen one; achenium oblong with a contracted base. — Petty's Island, Philadelphia, Prof. Leidy, &c. Probably adventive from S. States.

6. HEMICÁRPHA, Nees. HEMICARPHA. (Pl. 2.)

Spike, flowers, &c. just as in *Scirpus*, except that there is a minute translucent scale (readily overlooked) between the flower and the axis of the spike. Stamen only one. Style 2-cleft. Bristles or other perianth none. (Name from *ἥμι*, *half*, and *κάρφος*, *straw* or *chaff*, in allusion to the single inner scalelet.)

1. **H. subsquarrōsa**, Nees. Dwarf or minute annual (1' - 5' high); involucre 1-leaved, as if a continuation of the bristle-like culm, and usually with another minute leaf; spikes 2-3 (barely 2" long); scales brown, tipped with a short, recurved point. (*Scirpus subsquarrosus*, *Muhl.*)—Sandy borders of ponds and rivers; often growing with *Cyperus inflexus*. July-Sept.—Var. **DRUMMÓNDII** (*H. Drummondii*, *Nees.*) is a form with single and paler or greenish heads: Illinois and southward.

7. ELEÓCHARIS, R. Br. SPIKE-RUSH. (Pl. 3.)

Spike single, terminating the naked culm, many-several-flowered. Scales imbricated all round in many (rarely in 2 or 3) ranks. Perianth of 3-12 (commonly 6) bristles, usually rough or barbed downwards, rarely obsolete. Stamens 3. Style 2-3-cleft, its bulbous base persistent as a tubercle, which is jointed with the apex of the lenticular or triangular achenium.—Leafless, chiefly perennial, with tufted culms sheathed at the base, from matted or creeping rootstocks: flowering in summer. (Name from *ἔλος*, a marsh, and *χαίρω*, to delight in; being marsh plants.)

§ 1. *Spike terete, hardly if at all thicker than the spongy-cellular culm; the scales firmly persistent: style mostly 3-cleft: bristles of the perianth 6 (rarely 7), firm or rigid, mostly barbed downwardly, and equalling or surpassing the triangular or double convex achenium.*

* *Spike linear or lanceolate-awl-shaped, few-flowered; the scales (only 3-9) few-ranked, convolute-clasping the long flattened joints of the axis, lanceolate, herbaceous (green) and several-nerved on the back, and with thin scarious margins.*

1. **E. Robbinsii**, Oakes. *Flower-bearing culms exactly triangular*, rather stout, erect (8'-2° high), also producing tufts of capillary abortive stems or fine leaves, which float in the water; sheath obliquely truncate; achenium oblong-obovate, triangular, minutely reticulated, about half the length of the bristles, tipped with a flattened awl-shaped tubercle.—Shallow water, from Pondicherry Pond, New Hampshire (*Robbins*), to New Jersey (*C. E. Smith*, &c.), and southward.—Spike varying from 4" to 10" long, by scarcely over a line wide; the long scales rather remote and sheath-like.

* * *Spike cylindrical and many-flowered, 1'-2' long; the scales regularly imbricated in several ranks, firm-coriaceous with a narrow scarious margin and no midrib, pale, nerveless or faintly striate: culms large and stout (2°-4° high): sheaths at the base often leaf-bearing.* (**LIMNÓCHLOA**, *Nees.*)

2. **E. equisetoides**, Torr. *Culm terete, knotted as if jointed by many cross partitions; achenium smooth, with a conical-beaked tubercle.*—Shallow water, Rhode Island (*Olney*), Michigan (*Houghton*), Delaware, and southward.

3. **E. quadrangulata**, R. Br. *Culm continuous and sharply 4-angled; achenium finely reticulated, with a conical flattened distinct tubercle.*—Shallow water, New York (outlet of Oneida Lake, *A. H. Curtiss*) to Michigan and southward: rare.

§ 2. *Spike terete and turgid-ovate, very much thicker than the slender culm; the scales thin-coriaceous or firm-membranaceous and persistent, ovate: style 3-cleft: bristles*

of the perianth stout, downwardly barbed, as long as the striated and pitted-reticulated triangular achenium and its tubercle; culms tufted from fibrous roots, about 1° high.

4. **E. tuberculosa**, R. Br. Culms flattish, striate; spike 3''–6'' long, many-flowered; tubercle flattish-cap-shaped, as large as the body of the achenium. — Wet sandy soil, Mass. along the coast to Virginia and southward.

5. **E. simplex**, Torr. Culms sharply triangular, capillary, twisting when dry; spike 2''–3'' long, few-flowered; conical-beaked tubercle much smaller than the achenium. (*E. tortilis*, Schultes.) — Eastern shore of Maryland (*W. M. Canby*), and southward.

§ 3. Spike terete, much thicker than the culm, many-flowered; the scales imbricated in many or more than three ranks, thin-membranaceous or scarious in texture, with a thicker midrib, usually brownish or purplish, sometimes deciduous at maturity. (ELEÓGENUS, Nees.)

* Achenium smooth and lenticular, and style 2-cleft, or in No. 6 more commonly 3-cleft: culms slender or thread-form, terete or compressed.

6. **E. obtusa**, Schultes. Culms nearly terete, tufted (8'–14' high) from fibrous roots; spike globose-ovoid and with age oblong, obtuse, rarely becoming acute (dull brown); the scales very obtuse and numerous (80–130), densely crowded in many ranks; style 3- (rarely 2-) cleft; achenium obovate, shining, tumid-margined, about half the length of the 6–8 bristles, crowned with a short and very broad flattened tubercle. — Muddy places: most common.

7. **E. olivacea**, Torr. Culms flattish, grooved, diffusely tufted on slender matted rootstocks (2'–4' high); spike ovate, acutish, 20–30-flowered; scales ovate, obtuse, rather loosely imbricated (purple with a green midrib and slightly scarious margins); achenium obovate, dull, abruptly beaked with a narrow tubercle, shorter than the 6–8 bristles. — Wet sandy soil, Mass. to New Jersey near the coast, and southward: also shore of Lake Ontario, *J. A. Paine*.

8. **E. palustris**, R. Br. Culms nearly terete, striate (1°–2° high), from running rootstocks; spike oblong-lanceolate, pointed, many-flowered; scales ovate-oblong, loosely imbricated in several ranks, reddish-brown with a broad and translucent whitish margin and a greenish keel, the upper acutish, the lowest rounded and often enlarged; achenium obovate, somewhat shining, crowned with a short ovate or ovate-triangular flattened tubercle, shorter than the usually 4 bristles. — Var. *GLAUDESCENS* (*S. glaucescens*, Willd. !): culms slender or filiform; tubercle narrower and acute, beak-like, sometimes half the length of the achenium. — Var. *CÁLVA* (*E. calva*, Torr.): bristles wanting; tubercle short, nearly as in the true *E. palustris*, but rather narrower (*Watertown*, New York, *Dr. Crawe*). — Very common, either in water, when it is pretty stout and tall; or in wet grassy grounds, when it is slender and lower. (Eu.)

9. **E. compressa**, Sullivant. Culms flat, striate, erect (1°–2° high); spike ovate-oblong, or at length lanceolate, 20–30-flowered (4''–7'' long); scales lanceolate-ovate, acute, dark purple with broad white pellucid margins and summit; achenium obovate-pear-shaped, compressed, crowned with a small conical and pointed tubercle; bristles 1–4, very slender, fragile, shorter than or equalling the achenium, sometimes none or a single rudiment.) — Wet places, N. New York

to Ohio and Illinois. — Culms tufted on running rootstocks, strikingly flat, often spirally twisted in drying.

* * *Achenium triangular or turgid: style 3-cleft.*

+ Bristles equalling or surpassing the smooth achenium, downwardly barbed, persistent.

10. **E. rostellata**, Torr. Culms flattened and striate-grooved, wiry, erect (1° – $2\frac{1}{2}^{\circ}$ high), the sterile ones reclining, rooting and proliferous from the apex (1° – 2° high), the sheath transversely truncate; spike spindle-shaped, 12–20-flowered; scales ovate, obtuse (light-brown); achenium obovate-triangular, narrowed into the confluent pyramidal tubercle, which is overtopped by the 4–6 bristles. — Marshes, Rhode Island (*Olney*), E. Massachusetts (*W. Boott*), and Vermont (*Tuckerman*) to New Jersey (*Dr. Allen*), Alexandria, Virginia (*A. H. Curtiss*), Michigan and southward.

11. **E. intermedia**, Schultes. Culms capillary, wiry, striate-grooved, densely tufted from fibrous roots, diffusely spreading or reclining (6'–12' long); spike oblong-ovate, acutish, loosely 10–20-flowered (2''–3'' long); scales oblong, obtuse, green-keeled, the sides purplish-brown; achenium obovoid with a narrowed base, beaked with a slender conical-awl-shaped distinct tubercle, which nearly equals the 6 bristles. (*E. reclinata*, *Kunth*!) — Wet slopes: common from New York and Penn. to Illinois and northward.

12. **E. microcarpa**, Torr., var. **FILICULMIS**, Torr. Like the preceding, but more capillary and heads smaller ($1\frac{1}{2}$ ''–2'' long), sometimes proliferous, the one or more short new culms from the axil of its lowest scale, which persists as an herbaceous bract; achenium very much smaller, with sharper angles and a short conical tubercle, which is hardly equalled by the 3–6 slender bristles. — Wet pine barrens, New Jersey and southward.

+ + Bristles 2–4, shorter than the achenium, slender and fragile, or none.

13. **E. tenuis**, Schultes. Culms almost capillary, erect from running rootstocks, 4-angular and flattish (1° high), the sides concave; spike elliptical, acutish, 20–30-flowered (3'' long); scales ovate, obtuse, chestnut-purple with a broad scarious margin and green keel; achenium obovate, roughish-wrinkled, crowned with a small depressed tubercle, persistent after the fall of the scales; bristles half the length of the achenium, or wanting. (*E. elliptica*, *Kunth*!) — Wet meadows and bogs: common, early-flowering; June.

14. **E. melanocarpa**, Torr. Culms flattened, grooved, wiry, erect (9'–18' high); spike cylindrical-ovoid or oblong, thick, obtuse, densely many-flowered (3''–6'' long); scales closely many-ranked, roundish-ovate, very obtuse, brownish with broad scarious margins; achenium smooth, obovate-top-shaped, obtusely triangular, the broad summit entirely covered like a lid by the flat depressed tubercle, which is raised in the centre into a short abrupt triangular point; bristles shorter than the (soon blackish) achenium, often obsolete. — Wet sand, Plymouth, Massachusetts, to Virginia, and southward along the coast.

15. **E. tricostata**, Torr. Culms flattish (1° – 2° high); spike soon cylindrical, densely many-flowered (6''–9'' long, thickish); scales ovate, very obtuse, rusty brown, with broad scarious margins; achenium obovate, with 3 prominent thickened angles, minutely rough-wrinkled, crowned with a short-conical acute tubercle; bristles none. — Quaker Bridge, New Jersey (*Knieskern*), and southward.

§ 4. *Spike more or less flattened, thicker than the slender or capillary culm, few-many-flowered; the thin membranaceous scales somewhat 2-3-ranked: style 3-cleft: bristles of the perianth 3-6, fragile or fugacious. Small or delicate species, differing from the last subdivision chiefly in the flattish spikes.* (CHÆTOCYPÈRUS, Nees.)

* *Achenium tumid, obscurely triangular, many-ribbed on the sides.*

16. **E. aciculàris**, R. Br. Culms finely capillary (2'-8' long), more or less 4-angular; spike 3-9-flowered; scales ovate-oblong, rather obtuse (greenish with purple sides); achenium obovate-oblong, with 3-ribbed angles and 2-3 times as many smaller intermediate ribs, also transversely striate, longer than the 3-4 very fugacious bristles; tubercle conical-triangular. (S. trichòdes, Muhl., &c.) — Muddy shores: common. (Eu.)

** *Achenium triangular, with smooth and even sides.*

17. **E. pygmæa**, Torr. Culms bristle-like, flattened and grooved (1'-2' high); *spike ovate, 3-8-flowered*; scales ovate (greenish), the upper rather acute; achenium ovoid, acutely triangular, smooth and shining, tipped with a minute tubercle; bristles mostly longer than the fruit, sometimes wanting. (S. pusillus, Vahl?) — Brackish marshes and brackish river-banks.

8. SCÍRPUS, L. BULRUSH OR CLUB-RUSH. (Pl. 3.)

Spikes several-many-flowered, solitary or in a terminal cluster which (except in No. 1) is subtended by a 1-several-leaved involucre (this when simple often appearing like a continuation of the culm), terete, the scales being regularly imbricated all round in many or several ranks, or rarely somewhat compressed, the fewer scales inclining to be 2-ranked. Flowers under all the scales, or all but one or two of the lowest, all perfect. Perianth of 3-6 (straight or rarely tortuous) bristles, or sometimes wanting. Stamens mostly 3. Style 2-3-cleft, simple, not bulbous at the base, wholly deciduous, or sometimes leaving a tip or point to the lenticular or triangular achenium. — Culms sheathed at the base; the sheaths usually leaf-bearing. Perennials, except No. 8-9; flowering in summer. (The Latin name of the Bulrush.)

§ 1. SCÍRPUS proper (including ISÓLEPIS, which simply wants the bristles).

Bristles when present rigid, not elongated and contorted or exerted after flowering, mostly barbed downwards.

* *Spike solitary, few-flowered, small, often flattish: achenium triangular, smooth.*

+ *No involucre leaf or bract and no leaves below: base of style indurated and persistent, but continuous with the achenium.* (Transition to Eleocharis and Blysmus.)

1. **S. pauciflorus**, Lightfoot. Culms striate-angled, very slender (3'-9' high), scarcely tufted, on slender running rootstocks, at the base with a short truncate sheath; scales of the ovate spike evidently 2-ranked, chestnut-brown, pointless, all flower-bearing, the two lower larger; bristles 3-6, downwardly barbed, about the length of the conspicuously beak-pointed triangular achenium. (S. Bæóthryon, Ehrh. Blysmus pauciflorus, Caruel.) — Very wet places, New York (Jefferson Co., Dr. Crave, & Herkimer Co., J. A. Paine) to N. Illinois (Dr. Vasey), Michigan, and northwestward. (Eu.)

+ + *Involucre a short awl-shaped bract; culms tufted (3'–12' high), filiform.*

2. **S. cæspitōsus**, L. *Culms terete, wiry, densely sheathed at the base, in compact turfy tufts; the upper sheath bearing a very short awl-shaped leaf; spike ovoid, rusty-color; involucre a rigid-pointed scale, resembling the lowest proper scale of the spike and scarcely surpassing it; bristles 6, smooth, longer than the abruptly short-pointed achenium.*—Alpine tops of the mountains of Maine, New Hampshire, and N. New York. On Roan Mountain, N. Carolina. Also, away from mountains, Bergen Swamp, Genesee Co., New York, *G. W. Clinton, J. A. Paine*. N. Illinois, *Dr. Vasey*. (Eu.)

3. **S. Clintonii**, Gray. *Culms acutely triangular, almost bristle-like; sheaths at the base bearing a very slender almost bristle-shaped leaf shorter (usually very much shorter) than the culm; involucre a rigid-pointed scale, mostly shorter than the chestnut-colored ovate spike, which has pointless scales: otherwise as the next. (S. planifolius, var. brevifolius, Torr.)*—Rather dry plains, New York, Jefferson Co., *Dr. Crave*; near Buffalo, *G. W. Clinton*. June.

4. **S. planifolius**, Muhl. *Culms triangular, leafy at the base; the leaves linear, flat, as long as the culm, and like it rough-edged; involucre a bristle-tipped scale usually overlapping the ovate or oblong chestnut-colored spike, the green midrib of the scales extended into sharp points; bristles of the perianth upwardly hairy, as long as the pointless achenium.*—Dry or moist ground, New England to W. New York (*G. W. Clinton*), Penn., and Delaware. June.

+ + + *One-leaved involucre more conspicuous, and as if continuing the culm.*

5. **S. subterminalis**, Torr. *Aquatic: culms (1°–3° long, thickish-filiform) partly and the shorter filiform leaves wholly submersed, cellular; the filiform green bract 6"–12" long, much surpassing the oblong spike; scales somewhat pointed; bristles 6, bearded downwards, rather shorter than the abruptly-pointed achenium.*—Slow streams and ponds, New Jersey and New England to Michigan and westward.—Var. **TERRĒSTRIS**, Paine, Cat. Less tall; stem and leaves firmer; fully-fruited spike more turgid: growing chiefly emersed, in a quaking morass, Litchfield, Herkimer Co., New York, *J. A. Paine*.

* * *Spikes clustered (rarely only one), appearing lateral from the one-leaved involucre, which resembles the naked culm, seeming to be a continuation of it.*

+ *Culm sharply triangular, stout, chiefly from running rootstocks: spikes many-flowered, rusty brown, closely sessile in one cluster: sheaths at base more or less leaf-bearing. (Very rarely a second and smaller involucre leaf.)*

6. **S. pungens**, Vahl. *Running rootstocks long and stout; culm sharply 3-angled throughout (1°–4° high) with concave sides; leaves 1–3, elongated (4'–10' long), keeled and channelled; spikes 1–6, capitate, ovoid, usually long overtopped by the pointed involucre leaf; scales ovate, sparingly ciliate, 2-cleft at the apex and awl-pointed from between the acute lobes; anthers tipped with an awl-shaped minutely fringed appendage; style 2-cleft (rarely 3-cleft); bristles 2–6, shorter than the obovate plano-convex and mucronate smooth achenium. (S. triquetèr, Michx., not of L. S. Americanus, Pers.)*—Borders of salt and fresh ponds and streams: very common. (Eu.)

7. **S. Olneyi**, Gray. *Culm 3-wing-angled, with deeply excavated sides, stout (2°–7° high), the upper sheath bearing a short triangular leaf or none; spikes 6–12,*

closely capitate, ovoid, obtuse, overtopped by the short involucre leaf; scales orbicular, smooth, the inconspicuous mucronate point shorter than the scarious apex; *anthers with a very short and blunt minutely bearded tip; style 2-cleft; bristles 6, scarcely equalling the obovate plano-convex and mucronate achenium.* — Salt marshes, Martha's Vineyard (*Oakes*), and Rhode Island (*Olney*), to Delaware, and common southward. — Cross-section of the stem strongly 3-rayed, with the sides parallel. — Much nearer than the last to the European *S. TRIQUÈTER*, which has similar anthers, and an abbreviated or almost abortive leaf; but its culm is wingless, and the cluster of spikes compound, some of them umbellate-stalked.

8. *S. Tórreyi*, Olney. Rootstocks slender if any (so that the plant is readily pulled up from the mud); *culm 3-angled, with concave sides, rather slender (2°-4° high), leafy at the base; leaves 2 or 3, more than half the length of the culm, triangular-channelled, slender; spikes 1-4, oblong or spindle-shaped, acute, distinct, pale chestnut-color, long overtopped by the slender erect involucre leaf; scales ovate, smooth, entire, barely mucronate; style 3-cleft; bristles longer than the unequally triangular obovate very smooth and long-pointed achenium.* (*S. mucronatus, Pursh?, Torr. Fl. N. Y.*) — Borders of ponds, both brackish and fresh, New England to Penn. and Michigan.

+ + *Culm triangular, tall and stout, from slender running rootstocks: spikes many-flowered loosely umbellate or corymbed, involucrellate-bracted.*

9. *S. Cánbyi*, Gray. Culm (3°-5° high) 3-angled, usually sharply so above, obtusely below, the sheath at the base extended into a long and slender triangular and channelled leaf; the involucre leaf similar, continuing the culm; spikes oblong (4'-6' long), single or sometimes proliferously 2 or 3 together, nodding on the apex of the 5-9 long filiform and flattened peduncles or rays of the dichotomous umbel-like corymb, or the central one nearly sessile; scales of the spike loosely imbricated, oblong-ovate, acute, pale, thin and scarious, with a greenish nerved back; bristles of the perianth 6, firm, furnished above with spreading hairs rather than barbs, equalling the slender abrupt beak of the obovate-triangular shining achenium. — In a mill-pond, near Salisbury, Maryland, *A. Commons, W. M. Canby.* — A remarkable species: leaf 2°-4° long: involucre leaf 4'-8' long: rays or peduncles 1½'-3' long, each subtended by a single involucrellate leaf or bract, the lowest like that of the involucre but short, the uppermost reduced to scale-like bracts. Achenium (1½", and its beak ½" long.

+ + + *Culm terete, very tall and stout, from a deep running rootstock, naked; the sheaths at the base bearing a short and imperfect leaf or none: spikes numerous and clustered in a one-sided compound umbel-like panicle, the principal rays of which mostly surpass the involucre leaf: involucrellate bracts small, scale-like and rusty-scarious: scales of the spike rusty or chestnut-brown, scarious, with a salient midrib extended into a mucronate point.*

10. *S. válidus*, Vahl. (GREAT BULRUSH.) Culm 3°-9° high, ½'-1' thick at base; spikes ovate-oblong (3"-4" long); scales mostly a little downy on the back and ciliate; style 2-cleft; *achenium pale and dull, obovate with a narrowed base, plano-convex, mucronate-pointed, usually overtopped by the 4-6 slender*

downwardly barbed bristles. (*S. acutus*, Muhl. *S. lacustris*, of Amer. authors and in former editions.) — Common everywhere in still fresh water. — Achenium ($\frac{3}{8}$ " long, half the size of that of the European *S. lacustris*, also narrower, pale, not shining. — A slender variety with narrower heads, very smooth scales, and shorter or fragile bristles, was sparingly collected by Rev. J. W. Chickering at Havana, New York.

11. **S. riparius**, Presl., which largely takes the place of the preceding southwestward, probably within our limits, is known by the 2-4 rather shorter and linear *plumose scales, rather than bristles of the perianth.*

+ + + + *Culms slender from an annual root, terete, plano-convex or obtusely triangular, naked; the sheaths at the base rarely extended into a short leaf: spikes few or several, sometimes solitary, in a sessile cluster, much overtopped by the involucre leaf: bristles of the perianth often few or wanting.*

12. **S. débilis**, Pursh. Culms obtusely triangular, with somewhat hollowed sides, 1° - 2° high, yellowish-green; spikes 3-12, capitate, ovate-oblong, obtuse (3" - 4" long), chestnut-brown; involucre leaf often horizontal at maturity; scales roundish; stamens 3; style 2-3-cleft; bristles 6, stout, downwardly barbed, equalling or two surpassing the obovate turgidly plano-convex (or bluntly 3-sided) abruptly mucronate-pointed smoothish achenium. (*S. juncoïdes*, Roxburgh.) — Swamps, Mass. to Virginia and southward. Aug., Sept.

13. **S. Smithii**, n. sp. Culms terete, slender, 3' - 12' high, often leaf-bearing from the upper sheath, dull green as are the 1-3 oblong-ovate acute spikes; involucre leaf always erect; scales oblong-oval; style 2-cleft; bristles 1 or 2 minute rudiments or none; achenium somewhat lenticular, smooth, deciduous with the scales. (*S. debilis*, Gray, *Gram. & Cyp.* 135.) — Wet shores, Lake Ontario to Illinois and Delaware Bay (in tidal mud). July. — Named for C. E. Smith, who indicated and insisted on its distinctions.

14. **S. supinus**, L., var. **Hallii**. Culms filiform, 5' - 12' high; upper sheath rarely distinctly leaf-bearing; spikes 1-7 in a sessile or sometimes geminately proliferous cluster, ovate-oblong becoming cylindrical, greenish; scales ovate, strongly keeled, mucronate-pointed; stamens 2; style 2-cleft; *bristles none; achenium obovate-orbicular, mucronate, plano-convex, strongly wrinkled transversely.* (*S. Hallii*, Gray, addend. ed. 2.) — Wet shores, Illinois, E. Hall, &c., and southwestward. — In Texas occurs the normal *S. supinus*, with 3-cleft style and triangular achenium, as in Europe, where it sometimes has short bristles, as in the variety of No. 12. (Eu.)

* * * *Spikes clustered in simple or mostly compound umbellate or cymose-panicled clusters, many-flowered, terete: involucre of mostly several obvious and flat leaves: culm tall, from tufted or running rootstocks; triangular, leafy, sedge-like: style mostly 3-cleft.*

+ *Spikes large, 6" - 15" long: midrib of the scales extended beyond the mostly lacerate or two-cleft apex into a distinct awn.*

15. **S. maritimus**, L. (SEA CLUB-RUSH.) Leaves flat, linear, as long as the stout culm (1° - 3° high), those of the involucre 1-4, very unequal; spikes few-several in a sessile cluster, and often also with 1-4 unequal rays bearing 1-7 ovate or oblong-cylindrical (rusty-brown) spikes; awns of the

scales soon recurved; *achenium* obovate-orbicular, compressed, flat on one side, convex or obtuse-angled on the other, minutely pointed, shining, longer than the 1-6 unequal and deciduous (sometimes obsolete) bristles. — Var. *MACROSTACHYOS*, Michx. (*S. robustus*, Pursh) is a larger form, with very thick oblong-cylindrical heads, becoming 1' - 1½' long, and the longer leaf of the involucre often 1° long. — Salt marshes: common on the coast, and near salt springs in the interior (W. New York, &c.). (Eu.)

16. *S. fluviatilis*, Gray. (RIVER C.) Leaves flat, broadly linear (½' or more wide), tapering gradually to a point, the upper and those of the very long involucre very much exceeding the compound umbel; rays 5-9, elongated, recurved-spreading, each bearing 1-5 ovate or oblong-cylindrical acute heads; *achenium* obovate, sharply and exactly triangular, conspicuously pointed, opaque, scarcely equaling the 6 rigid bristles. (*S. maritimus* var. ? *fluviatilis*, Torr., excl. syn.) — Borders of lakes and large streams, W. Vermont to Penn., Wisconsin and Illinois. — Culm very stout, sharply triangular, 3° - 5° high. Leaves roughish on the margin, like the last; those of the umbel 3-7, the largest 1° - 2° long. Principal rays of the umbel 3' - 4' long, sheathed at the base. Heads paler and duller than in the preceding; the scales less lacerate, and their awns less recurved, the fruit larger and very different.

+ + Spikes very numerous, small, 1' - 3' long; their scales mucronate-pointed or blunt: umbel-like cymose panicle irregular, compound or decomposed: culm 2° - 5° high, unusually leafy; leaves broadly linear, green and rather soft, rough on the edges: bristles of the perianth very slender, often more or less tortuous and naked below: — transition to § *Trichophorum*.

17. *S. sylvaticus*, L. Spikes lead-colored, clustered 3-10 together at the end of the mostly slender ultimate divisions of the open decomposed panicle, ovoid (1½" long); the scales blunt or barely mucronate; bristles 6, downwardly barbed their whole length, straight, scarcely longer than the convex-triangular short-pointed *achenium*. — Base of the White Mountains, New Hampshire (*Oakes*), and northward. (Eu.) (*S. MICROCÁRPUS*, Presl., *S. lenticularis*, Torr., apparently a form of *S. sylvaticus* with a 2-cleft style and flat *achenium*, approaches our northwestern borders.)

18. *S. atrovirens*, Muhl. Leaves somewhat more rigid; spikes dull greenish-brown, densely conglomerate (10-30 together) into close heads, these also usually densely clustered in a less compound panicle; scales pointed; bristles sparsely and strongly downwardly barbed above the middle, naked below, nearly straight, as long as the conspicuously pointed and obovate-oblong triangular *achenium*. (*S. sylvaticus*, var. *atrovirens*, Ed. 2. — Wet meadows and bogs, New England to Kentucky and northward: common.

19. *S. polyphyllus*, Vahl. Culm usually more leafy; spikes yellow-brown, ovate, becoming cylindrical, clustered 3-8 together in small heads on the short ultimate divisions of the open decomposed umbel; scales mucronate; bristles 6, usually twice bent, soft-barbed towards the summit only, about twice the length of the *achenium*. (*S. exaltatus*, Pursh. *S. brúnneus*, Muhl.) — Swamps and shady borders of ponds, W. New England to Illinois, and common southward: rare at the North.

§ 2. **TRICHOPHORUM**, Richard. *Bristles capillary, naked, not barbed, elongating, becoming tortuous and entangled, much longer than the triangular achenium, when old mostly projecting beyond the rusty-colored scales: culm, leaves, &c. as in the preceding subdivision; umbel-like cymose panicle decompound.*

20. **S. lineatus**, Michx. Culm triangular, leafy (1° – 3° high); leaves linear, flat, rather broad, rough on the margins; umbels terminal and sometimes axillary, loose, drooping, the terminal with a 1–3-leaved involucre much shorter than the long and slender rays; spikes oblong, becoming cylindrical ($2''$ – $4''$ long), on thread-like drooping pedicels; bristles at maturity scarcely exceeding the ovate green-keeled and pointed scales; achenium sharp-pointed. — Low grounds, W. New England to Wisconsin, and common southward.

21. **S. Eriophorum**, Michx. (WOOL-GRASS.) Culm nearly terete, very leafy (2° – 5° high); leaves narrowly linear, long, rigid, those of the involucre 3–5, longer than the decompound cymose-panicled umbel, the rays at length drooping; spikes exceedingly numerous, ovate, clustered, or the lateral ones pedicelled, woolly at maturity ($1\frac{1}{2}''$ – $3''$ long); the rusty-colored bristles much longer than the pointless scales; achenium short-pointed. (*Eriophorum cyperinum*, L.) — Var. **CYPERINUS** (*S. cyperinus*, Kunth) is the form with nearly all the spikes conglomerate in small heads. — Var. **LAXUS** (*S. Eriophorum*, Kunth) has the heads scattered, the lateral ones long-pedicelled. Various intermediate forms occur; and the umbel varies greatly in size. — Wet meadows and swamps: common northward and southward.

9. ERIOPHORUM, L. COTTON-GRASS. (Pl. 3.)

Spikes, scales, achenium, &c. as in *Scirpus*. Bristles of the perianth of numerous (in one species few) flat and delicate capillary bristles, which lengthen greatly after flowering, much exceeding the scales, and forming of the capitate spike a (white or reddish) conspicuous cotton-like tuft in fruit. Stamens 1–3. Style 3-cleft. Perennials. (Name composed of *ἔριον*, wool or cotton, and *φορά*, bearing.)

* *Bristles of the flower only 6, crisped, white; spike single: small, involucre none.*

1. **E. alpinum**, L. Culms slender, many in a row from a running root-stock 6'–10' high), scabrous, naked; sheaths at the base awl-tipped. — Cold bogs, New England to Penn., Wisconsin, and northward. June. (Eu.)

* * *Bristles very numerous, not crisped, forming dense cottony heads in fruit.*

+ *Culm bearing a single spike: involucre none.*

2. **E. vaginatum**, L. Culms in close tufts (1° high), leafy only at the base, above with 2 inflated leafless sheaths; root-leaves long and thread-form, triangular-channelled; scales of the ovate spike long-pointed, lead-color at maturity. — Cold and high peat-bogs, New England to mountains of Penn. (*Prof. T. Green*), Wisconsin, and northward; rare. May, June. (Eu.)

(*E. russèolum*, Fries, with copper-colored wool, found in New Brunswick by *Rev. J. Fowler*, may be expected in N. E. Maine.)

+ + *Culm leafy, bearing several umbellate-clustered heads, involucrate.*

3. **E. Virginicum**, L. Culm rigid (2° – 4° high); leaves very narrowly linear, elongated, flat; spikes nearly sessile, crowded in a dense cluster or head;

wool rusty or copper-color, only thrice the length of the scale; stamen 1. — Bogs and low meadows: common. July, Aug.

Var. **álbum**, with the wool white. — Oswego and Jefferson Counties, New York, *Dr. Crawe, A. H. Curtiss*.

4. **E. polystáchyon**, L. Culm rigid (1° – 2° high), obscurely triangular; leaves linear, flat, or barely channelled below, triangular at the point; involucre 2–3-leaved; spikes several (4–12), on nodding peduncles, some of them elongated in fruit; achenium obovate; wool white, very straight (1' long or more).

— Var. **ANGUSTIFÓLIUM** (*E. angustifolium*, *Roth*, and of European botanists, not of American, and the original *E. polystachyon* of *L.*) has smooth peduncles.

— Var. **LATIFÓLIUM** (*E. latifolium*, *Hoppe*, & *E. polystachyon*, *Torr.*) has rough peduncles, and sometimes broader and flatter leaves. — Both are common in bogs, northward; often with the peduncles obscurely scabrous, indicating that the species should be left as Linnæus founded it. June, July. (Eu.)

5. **E. grácile**, Koch. Culm slender (1° – 2° high), rather triangular; leaves slender, channelled-triangular, rough on the angles; involucre short and scale-like, mostly 1-leaved; peduncles rough or roughish-pubescent; achenium elliptical-linear. (*E. triquètrum*, *Hoppe*. *E. angustifolium*, *Torr.*) — Cold bogs, New England to Illinois, and northward. June–Aug. — Spikes 3–7, small, when mature the copious white wool 6"–9" long. Scales brownish, several-nerved; or in our plant, var. **PAUCINÉRVIVM**, *Engelm.*, mostly light chestnut-color, and about 3-nerved. (Eu.)

10. FIMBRÍSTYLIS, Vahl. (Pl. 3.)

Spikes several-many-flowered, terete; the scales all floriferous, regularly imbricated in several ranks. Perianth (bristles, &c.) none. Stamens 1–3. Style 2–3-cleft, often with a dilated or tumid base, which is deciduous (except in No. 4) from the apex of the naked lenticular or triangular achenium. Otherwise as in *Scirpus*. — Culms leafy at the base. Spikes in our species umbelled, and the involucre 2–3-leaved. (Name compounded of *fimbria*, a fringe, and *stylus*, the style, which is fringed with hairs in the genuine species.)

§ 1. **FIMBRISTYLIS** proper. Style 2-cleft, flat and ciliate, falling away, base and all, from the lenticular achenium; scales of the many-flowered spike closely imbricated.

1. **F. spadicea**, Vahl., var. **castánea**. Culms (1° – $2\frac{1}{2}^{\circ}$ high) tufted from a perennial root, rigid, as are the thread-form convolute-channelled leaves, smooth; spikes ovate-oblong becoming cylindrical, dark chestnut-color (2" thick); stamens 2 or 3; achenium very minutely striate and obscurely reticulated. (*F. castanea*, *cylindrica*, &c., *Vahl.*) — Salt marshes along the coast, New York to Virginia, and southward. July–Sept. — Spikes less dark-colored and scales mostly thinner than in the original Jamaica plant.

2. **F. láxa**, Vahl. Culms slender (2'–12' high) from an annual root, weak, grooved and flattish; leaves linear, flat, ciliate-denticulate, glaucous, sometimes hairy; spikes ovate, acute (3" long); stamen 1; achenium conspicuously 6–8-ribbed on each side, and with finer cross lines. (*F. Baldwiniana*, *Torr.* *F. brizoides*, *Nees*, &c.) — Low, mostly clayey soil, Penn. to Illinois and southward. July–Sept. — Scales broader and less pointed than in *F. annua*.

§ 2. **TRICHELÓSTYLIS**, Lestib. *Style 3-cleft and the achenium triangular: otherwise nearly as in § 1: the spikes small and fewer-flowered.*

3. **F. autumnális**, Rœm. & Schult. Annual (3'-16' high), in tufts; culms flat, slender, diffuse or erect; leaves flat, acute; umbel compound; spikes oblong, acute (1"-2" long), single or 2-3 in a cluster; the scales ovate-lanceolate, mucronate; stamens 1-3. (*Scirpus autumnalis*, L.) — Low grounds, Maine to Illinois, and southward. Aug. - Oct.

§ 3. **ONCÓSTYLIS**, Martius. *Style 3-cleft, slender, its thickish base more tardily deciduous from the apex of the triangular achenium.*

4. **F. capilláris**, Gray. Low annual, densely tufted (3'-9' high); culm and leaves nearly capillary, the latter short; umbel compound or paniced; spikes (2" long) ovoid-oblong; stamens 2; achenium minutely wrinkled, very obtuse. (*Scirpus capillaris*, L.) — Sandy fields: common. Aug., Sept.

(**F. CONGÉSTA**, Torr., a diminutive Southern species, with the 2-cleft style not ciliate, has been detected in ballast-sand at Camden, New Jersey, opposite Philadelphia, by *C. F. Parker*: probably only a waif.)

11. **DICHRÒMENA**, Richard. **DICHROMENA.** (Pl. 4.)

Spikes aggregated in a terminal leafy-involucrate head, more or less compressed, few-flowered, all but 3 or 4 of the flowers usually imperfect or abortive. Scales imbricated somewhat in 2 ranks, more or less conduplicate or boat-shaped, keeled, white or whitish. Stamens 3. Style 2-cleft. Perianth, bristles, &c., none. Achenium lenticular, wrinkled transversely, crowned with the persistent and broad tubercled base of the style. — Culms leafy, from creeping perennial rootstocks; the leaves of the involucre mostly white at the base (whence the name, from *δῖς*, *double*, and *χρῶμα*, *color*). — Differs (too little) from the next genus in the involucre bracts and flattened spikes.

1. **D. leucocéphala**, Michx. Culm triangular (1°-2° high); leaves narrow; those of the involucre 4-7; achenium truncate, not margined. — Damp pine barrens of New Jersey to Virginia and southward. Aug., Sept.

2. **D. latifolia**, Baldwin. Culm stouter, nearly terete; leaves broadly linear, those of the involucre 8-9, tapering from base to apex; achenium round-obovate, faintly wrinkled, the tubercle decurrent on its edges. — S. Virginia? and southward.

12. **RHYNCHÓSPORA**, Vahl. **BEAK-RUSH.** (Pl. 4.)

Spikes paniced or variously clustered, ovate, globular, or spindle-shaped, terete, or sometimes flattish; but the scales open or barely concave (not boat-shaped nor keeled); the lower ones commonly loosely imbricated and empty, the uppermost often subtending imperfect flowers. Perianth in the form of (mostly 6) bristles, or occasionally wanting. Achenium lenticular, globular, or flat, crowned with a conspicuous tubercle or beak consisting of the persistent and indurated base or even of the greater part of the style. — Chiefly perennials, with more or less triangular and leafy culms; the spikes in terminal and axillary clusters: flowering in summer. (Name composed of *ῥύγχος*, *a snout*, and *σπορά*, *a seed*, from the beaked achenium.)

§ 1. **PSILOCÁRYA**, Torr. *Spikes ovate, terete, the numerous scales all alike and regularly imbricated; a perfect flower under each: stamens mostly 2: style 2-cleft; its base or the greater part of it enlarging and hardening to form the beak of the lenticular or tumid more or less cross-wrinkled achenium: bristles wholly wanting* (whence the name).

1. **R. scirpoides**. Annual, 4' - 10' high; leaves flat; spikes in broad and open cymes, 20 - 30-flowered; scales oblong-ovate, acute, chestnut-colored; achenium obscurely wrinkled, beaked with the sword-shaped almost wholly persistent style, and somewhat margined. (*Psilocarya scirpoides*, Torr. & Ed. 2.) — Inundated places, Rhode Island and Plymouth, Massachusetts.

(*R. nitens* (*Scirpus nitens*, Vahl. *Psilocarya rhynchosporoides*, Torr.), like this, but with a more wrinkled and short-beaked achenium, takes its place in Southern States.)

§ 2. **EURHYNCHÓSPORA**. *Spikes terete or biconvex, few-many-flowered; some of the lower scales almost always empty: stamens mostly 3: style conspicuously 2-cleft, its base only forming the tubercle or beak of the mostly lenticular achenium: bristles of the perianth usually present, and merely rough or barbed-denticulate* (not plumose).

* *Achenium transversely wrinkled: bristles mostly 6, upwardly denticulate.*

2. **R. cymósa**, Nutt. *Culm triangular; leaves linear* ($\frac{1}{4}$ ' wide); cymes corymbose; the spikes crowded and clustered; achenium round-obovate, twice the length of the bristles, four times the length of the depressed-conical tubercle. — Low grounds, Penn. and New Jersey to Virginia, and southward.

3. **R. Torreyána**, Gray. *Culm nearly terete, slender; leaves bristle-form; cymes paniced, somewhat loose, the spikes mostly pedicelled; achenium oblong-obovate, longer than the bristles, thrice the length of the broad compressed-conical tubercle.* — Swamps; pine barrens of New Jersey, and southward.

4. **R. inexpánsa**, Vahl. *Culm triangular, slender; leaves narrowly linear; spikes spindle-shaped, mostly pedicelled, in drooping panicles; achenium oblong, half the length of the slender bristles, twice the length of the triangular-subulate tubercle.* — Low grounds, Virginia and southward.

* * *Achenium smooth and even.*

+ *Bristles of the perianth 6, long and conspicuous, upwardly denticulate.*

5. **R. fúsca**, Rœm. & Schultes. *Leaves bristle-form, channelled; spikes ovate-oblong, few, clustered in 1 - 3 loose heads* (dark chestnut-color); achenium obovate, half the length of the bristles, equalling the triangular-sword-shaped acute tubercle, which is rough-serrulate on the margins. — Low grounds, New Jersey to Maine and Lake Superior: rare. — Culm 6' - 12' high. (Eu.)

6. **R. gracilénta**, Gray. *Leaves narrowly linear; spikes ovoid, in 2 - 4 small clusters, the lateral long-peduncled; achenium ovoid, rather shorter than the bristles, about the length of the flat awl-shaped tubercle.* — Low grounds, S. New York, New Jersey, and southward. — Culm very slender, 1° - 2° high.

+ + *Bristles none or 1 - 3 and minute: spikes pale, 1-flowered.*

7. **R. pállida**, M. A. Curtis. *Culm* (1° - 2° high) acutely triangular; leaves and spikes as in the next species, but only a terminal dense cluster, which is less white or turns pale reddish-tawny; achenium obovate-lenticular,

tipped with a minute depressed and apiculate tubercle; the delicate bristles four or five times shorter or obsolete. — Bogs in pine barrens of New Jersey (*W. M. Canby, C. F. Parker*), and in N. Carolina.

+ + + *Bristles long, denticulate downwardly, or both ways in No. 12.*

+ *Spikes white or whitish, becoming tawny with age, perfecting only a single flower: stamens usually 2: bristles 9-12, or even 20.*

8. **R. álba**, Vahl. Culm slender (1° - 2° high), triangular above; leaves narrowly linear or almost bristle-form; spikes lanceolate, densely crowded in a head-like terminal corymb and usually one or two lateral ones; achenium oblong-obovate with a narrowed base, scarcely longer than the flattened-awl-shaped tubercle, shorter than the bristles. — Bogs, especially eastward. (Eu.)

+ + *Spikes chestnut-colored or darker in No. 11 and 12, few-several-flowered: stamens 3: bristles usually 6.*

9. **R. capillácea**, Torr. *Leaves bristle-form; spikes 3-6 in a terminal cluster, and commonly 1 or 2 on a remote axillary peduncle, oblong-lanceolate (pale chestnut-color, 3" long); achenium oblong-ovoid, stipitate, very obscurely wrinkled, about half the length of the (6, rarely 12) stout bristles, and twice the length of the lanceolate-beaked tubercle. — Bogs and rocky river-banks, Penn. to N. Vermont, New York, and Michigan. — Culm 6'-9' high, slender.*

10. **R. Knieskérnii**, Carey. *Leaves narrowly linear, short; spikes numerous, crowded in 4-6 distant clusters, oblong-ovate (scarcely 1" long); achenium obovate, narrowed at the base, equalling the bristles, twice the length of the triangular flattened tubercle. — Pine barrens of New Jersey, on bog iron ore exclusively (*Knieskern*), and southward: rare. — Culms 6'-18' high, slender.*

11. **R. glomeráta**, Vahl. *Leaves linear, flat; spikes numerous in distant clusters or heads (which are often in pairs from the same sheath), ovoid-oblong; achenium obovate, margined, narrowed at the base, as long as the lance-awl-shaped flattened tubercle, which equals the (always) downwardly barbed bristles. — Low grounds: common, especially eastward. — Culm 1° - 3° high. — A state with small paniced clusters is *R. paniculata*, Gray.*

12. **R. cephalántha**, Torr. *Leaves narrowly linear, flat, keeled; spikes very numerous, crowded in 2 or 3 or more dense globular heads which are distant (and often in pairs), oblong-lanceolate, dark brown; achenium orbicular-obovate, margined, narrowed at the base, about as long as the awl-shaped beak, half the length of the stout bristles, which are barbed downwards and sometimes also upwards. — Sandy swamps, Long Island to New Jersey, and southward. — Culm stout, 2° - 3° high: the fruit larger than in the last, of which it may be only a marked variety.*

§ 3. **CERATOSCHËNUS**, Nees. *Spikes spindle-shaped or lanceolate, acuminate, in fruit flattish, large, cymose-panicled, producing only one perfect and 1 to 4 staminate flowers; their scales few, the lower mostly empty: stamens 3: bristles of the perianth rigid, either short or slender, minutely scabrous upward: style simple or barely 2-toothed at the apex, filiform and gradually thickened downwards, in fruit almost all of it persistent as a very long, exerted, slender-awl-shaped, upwardly roughened beak, several times longer than the smooth and flat obovate achenium: coarse perennials: spikes in flower 4", in fruit including the pro-*

jecting beak or style about 1' long. (This long beak gives the name, from *κέρας*, a horn, and *σχοίνος*, a rush.)

13. **R. corniculata**, Gray. (HORNED RUSH.) *Cymes* decomposed, *diffuse*; bristles awl-shaped, stout, unequal, shorter than the achenium. — Wet places, Penn. to Illinois, and southward. — Culm 3°–6° high. Leaves about 6" wide.

14. **R. macrostachya**, Torr. *Cymes* decomposed, or in the northern form somewhat simple and smaller, and the spikes usually more clustered; bristles capillary, twice the length of the achenium. — Borders of ponds, Massachusetts, Rhode Island, New Jersey, and southward: rare. — Perhaps it runs into the preceding.

13. CLADIUM, P. Browne. TWIG-RUSH. (Plate 5.)

Spikes ovoid or oblong, of several loosely imbricated scales; the lower ones empty, one or two above bearing a staminate or imperfect flower; the terminal flower perfect and fertile. Perianth none. Stamens 2. Style 2–3-cleft, deciduous. Achenium ovoid or globular, somewhat corky at the summit, or pointed, without any tubercle, in which it differs from *Rhynchospora*. (Name from *κλάδος*, a twig or branch, perhaps on account of the twice branching styles of some species.)

1. **C. mariscoides**, Torr. Perennial; culm obscurely triangular (1°–2° high); leaves narrow, channelled, scarcely rough-margined; cymes small; the spikes clustered in heads 3–8 together on 2 to 4 peduncles; style once 3-cleft. — Bogs, New England to Illinois, and northward. July.

14. SCLÈRIA, L. NUT-RUSH. (Pl. 5.)

Flowers monœcious; the fertile spikes 1-flowered, usually intermixed with clusters of few-flowered staminate spikes. Scales loosely imbricated, the lower ones empty. Stamens 1–3. Style 3-cleft. Achenium globular, stony, bony, or enamel-like in texture. Bristles, &c. none. Perennials, with triangular leafy culms, mostly from creeping rootstocks: flowering in summer: all in low ground or swamps. (Name *σκληρία*, hardness, from the indurated fruit.)

1. **S. triglomerata**, Michx. Culm (2°–3° high) and broadly linear leaves roughish; fascicles of spikes few, terminal and axillary, in triple clusters, the lowermost peduncled; stamens 3; achenium smooth and polished, on an obscure crustaceous ring or disk. — Vermont to Wisconsin, and common southward.

2. **S. reticularis**, Michx. Culms slender (1° high); leaves narrowly linear; clusters loose, axillary and terminal, sessile or the lower on short slender peduncles; stamens 2; achenium globular, regularly pitted-reticulated, not hairy, resting on a double disk, each of three greenish appressed superposed calyx-like lobes, the inner larger. — Eastern Massachusetts to Virginia and southward: rare northward.

3. **S. laxa**, Torr. Culms slender and weak (1°–2° high); leaves linear; clusters loose, the lower mostly long-peduncled and drooping; achenium globular, irregularly pitted-reticulated or pitted-rugose, towards the base minutely hairy on the somewhat spiral wrinkles: otherwise as in the foregoing. — E. Massachusetts to New Jersey, Pennsylvania, and southward.

4. *S. pauciflora*, Muhl. Somewhat downy or smoothish; culms slender (9'–18' high); leaves narrowly linear; clusters few-flowered, the lower lateral ones when present peduncled; bracts ciliate; stamens 3; *achenium globose, papillose-roughened, white*; the disk a narrow ring bearing 3 pairs of minute tubercles. — S. New England to W. New York (rare), and more common southward.

5. *S. verticillata*, Muhl. Smooth; culms simple, slender (4'–12' high), terminated by an interrupted spike of 4–6 rather distant sessile and small clusters; bracts minute; leaves linear; stamens 1 or 2; *achenium rough-wrinkled with short elevated ridges, globular-triangular*; the disk obsolete. — W. New York and Penn. to Michigan and southward: rare. — Plant faintly sweet-scented; achenium small, $\frac{3}{8}$ " long.

15. CAREX, L. SEDGE. (Pl. 5, 6.)

Staminate and pistillate flowers separated (*monœcious*), either borne together in the same spike (*androgynous*), or in separate spikes on the same stem, very rarely on distinct plants (*diœcious*). Scales of the spikes equally imbricated around the axis, each subtending a single staminate or pistillate flower. Stamens 3, rarely 2. Ovary enclosed in an inflated sac (composed of either one or two inner scales (bractlets) united by their margins), forming a rounded or angular bladderly sac (*perigynium*), which encloses the lenticular, plano-convex, or triangular achenium, tipped with more or less of the persistent (rarely jointed) base of the style. Stigmas 2 or 3, long, projecting from the narrow orifice of the perigynium. — Perennial herbs, chiefly flowering in spring and maturing in summer, frequently growing in wet places, often in dense tufts. Culms triangular, bearing the spikes in the axils of green and leaf-like or scale-like bracts, and terminal; commonly with sheaths at the base which enclose more or less of the stalks of the spikes. Leaves grassy, usually rough on the margins and keel. (A classical name, of obscure signification; derived by some from *careo*, to want, the upper spikes being mostly sterile; and by others from *κείρω*, to cut, on account of the sharp leaves.)

Contributed for the first edition of this work, and revised for the second, by JOHN CAREY, Esq.; with some present additions, from recent discoveries, and a few alterations, chiefly from the subsequent investigations of the late DR. FRANCIS BOOTT, published in his magnificent Illustrations of the genus *Carex*, and from notes furnished by WM. BOOTT, Esq.

ABRIDGED SYNOPSIS OF THE SECTIONS.

- A.** Spike solitary and terminal, simple, diœcious or androgynous: bracts small, colored and scale-like. — (This division, retained for the convenience of students, is merely artificial, and combines species having no real natural affinity.) PSYLLOPHORÆ, Loiseleur.
- ‡ 1. Spike diœcious, or with a few staminate flowers at its base. No. 1–2.
2. Spike androgynous, staminate at the summit. No. 3–6.
- Spike androgynous, staminate at the base. No. 36 and 138 may be sought here.
- B.** Spike solitary, single, androgynous, staminate at the summit: bracts and scales of the fertile flowers green and leaf-like. Stigmas 3. PHYLLOSTACHYS, Torr. & Gr. No. 7–9.
- C.** Spikes several or numerous, androgynous (occasionally diœcious in No. 11 and 33), sessile, forming compact or more or less interrupted, sometimes paniculate, compound or decomposed spikes. Stigmas 2. VIGNEA, Beauv.

- § 1. Spikes approximate, with staminate and pistillate flowers variously situated. No. 10-12.
2. Spikes pistillate below, staminate at the summit. No. 13-28.
3. Spikes pistillate above, staminate at the base. No. 29-45.

D. Staminate and pistillate flowers borne in separate (commonly more or less stalked) simple spikes on the same culm; the one or more staminate (sterile) spikes constantly uppermost, having occasionally more or less fertile flowers at base or apex; the lower spikes all pistillate (fertile), or sometimes with staminate flowers at the base or apex. Stigmas 3 (or only 2 in No. 46, 53, 65, 150, &c.). CAREX proper.

* Perigynia with merely a minute or short point, scarcely ever prolonged into a beak.

- § 1. Perigynia not inflated (slightly so in No. 55, 56), smooth, nerved or nerveless, with a minute straight point, glaucous-green, becoming whitish, or more or less spotted or tinged with purple. Scales blackish-purple or brown. Staminate spikes 1-3, or the terminal spike androgynous and staminate at the base, the rest all fertile. No. 46-64.
2. Perigynia slightly inflated, smooth, nerved, obtuse and pointless, or with a straight or oblique point. Scales brown, becoming tawny or white. Staminate spike solitary (except sometimes in No. 71) or androgynous and pistillate above, the rest all fertile. No. 65-81.
3. Perigynia slightly inflated, hairy (in No. 83 smooth at maturity), nerved, with a minute straight point. Terminal spike androgynous, pistillate at the apex, the rest all fertile. No. 82, 83.
4. Perigynia not inflated, smooth, regularly striate, with a short, entire, obliquely bent or recurved point, remaining green at maturity. Staminate spike solitary. Bracts green and leaf-like (except in No. 84). No. 84-91.
5. Perigynia not inflated, smooth or downy, not striate, with a minute, obliquely bent, white and membranaceous point, reddish-brown or olive-colored at maturity. Terminal spike all staminate or with 2-3 fertile flowers at the base; the rest all fertile, or with a few sterile flowers at the apex. Bracts reduced to colored sheaths, or with a short green prolongation. No. 92, 93.
- * Perigynia with a distinct beak, either short and abrupt, or more or less prolonged.
6. Perigynia not inflated, hairy, with a rather abrupt beak, terminating in a membranaceous notched or 2-toothed orifice. Bracts short: culms mostly low and slender; leaves all radical, long and narrow. Staminate spike solitary. No. 94-101.
7. Perigynia slightly inflated, hairy or smooth, with a short beak terminating in an entire or slightly notched orifice. Bracts long and leaf-like: culms tall and leafy. Staminate spike solitary (in No. 102 pistillate at the summit): fertile spikes erect (except in No. 102). No. 102, 103.
8. Perigynia slightly inflated, smooth and shining, green, few-nerved or nerveless, with a straight tapering beak terminating in 2 small membranaceous teeth. Staminate spike solitary (often androgynous in No. 107): fertile spikes all on slender and pedulous stalks. No. 104-110.
9. Perigynia slightly inflated, smooth, nerved, with a tapering somewhat serrulate beak, terminating in 2 distinct membranaceous teeth; becoming tawny or yellow at maturity. Staminate spike solitary. No. 111-115.
10. Perigynia slightly inflated, rough or pubescent, rarely smooth, with an abrupt straight beak. Staminate spikes usually two or more. No. 116-119.
11. Perigynia moderately inflated, smooth or pubescent, conspicuously many-nerved, with a straight beak terminating in 2 rigid more or less spreading teeth. Staminate spikes 1-5. No. 120-127.
12. Perigynia much inflated, smooth, conspicuously many-nerved, with a long tapering 2-toothed beak. Staminate spike solitary. No. 128-136.
13. Perigynia much inflated, obovoid or obconic, smooth, few-nerved, with an extremely abrupt, very long, 2-toothed beak, tawny or straw-colored at maturity, horizontally spreading or deflexed. Terminal spike staminate, or androgynous and fertile at the apex. No. 137-138.
14. Perigynia much inflated, smooth, shining and straw-colored at maturity, with a tapering and more or less elongated 2-toothed beak. Staminate spikes 2-5. No. 139-151.

A. *Spike solitary, simple; its scales or bracts small and scarious or colored (never green or foliaceous).* PSYLLÓPHORA, Loiseleur.

§ 1. *Spike diœcious, or the fertile merely with a few staminate flowers at the base.*

1. **C. gynócrates**, Wormskiold. Culm and bristle-form radical leaves smooth, or minutely rough at the top; sterile spike linear; fertile spike ovoid, loosely flowered; perigynia oblong, short-beaked, with a white membranaceous obtusely 2-toothed apex, narrowed at the base, nerved throughout, smooth, spreading horizontally at maturity, longer than the acute or acutish scale; stigmas 2. — Swamps, Wayne and Genesee Co., New York (*Sartwell, &c.*), Michigan, and northward. (Eu.)

2. **C. scirpoidea**, Michx. Leaves flat; spike narrowly cylindrical; perigynia ovoid, with a minute point, densely hairy, dark purple at maturity, about the length of the pointed ciliate scale; stigmas 3. (*C. Wormskioldiana, Hornem. C. Michauxii, Schw.*) — Alpine summits of the mountains of Maine and New Hampshire, Willoughby Mountain, Vermont (*Wood*), Drummond's Island, Michigan, and northward. (Eu.)

§ 2. *Spike androgynous, staminate at the summit.*

* *Stigmas 2: leaves bristle-form.*

3. **C. capitata**, L. Spike small, roundish-ovoid; perigynia broadly elliptical with a notched membranaceous point, compressed, smooth, spreading, longer than the rather obtuse scale. — Alpine summits of the White Mountains, New Hampshire, *Robbins, Oakes.* (Eu.)

* * *Stigmas 3: leaves very narrow, shorter than the culm.*

4. **C. pauciflora**, Lightfoot. Spike few-flowered; sterile flowers 1 or 2; perigynia awl-shaped, reflexed; scales deciduous. (*C. leucoglòchin, Ehrh.*) — Peat-bogs, from New England and W. New York northward. (Eu.)

5. **C. polytrichoides**, Muhl. Culm capillary; spike very small, few-flowered; perigynia erect, alternate, oblong, compressed-triangular, obtuse, slightly nerved, entire at the apex, green, twice the length of the ovate scale. (*C. leptalea, Wahl. C. microstachya, Michx.*) — Low grounds and bogs: very common.

* * * *Stigmas 3: leaves very (about 1') broad, longer than the naked culm.*

6. **C. Fraseriana**, Sims. Pale or glaucous and glabrous; leaves without midrib, many-nerved, smooth, with minutely crisped cartilaginous margins (9' – 18' long), convolute below around the base of the scape-like culm: spike oblong, the fertile part becoming globular; perigynia ovoid, inflated, mucronately tipped with a minute entire point, longer than the scarious oblong obtuse scale; often enclosing a short appendage at the base of the achenium. — Rich woods, mountains of Pennsylvania? Virginia, and southward: rare, and a most remarkable plant.

B. *Spike solitary, simple, androgynous, staminate at the summit: bracts and scales of the pistillate flowers green, leaf-like, tapering from a broad base, the lowest much longer than the spike, the uppermost equalling the slightly inflated perigynia: style jointed at the base: stigmas 3.* (Leaves long and grassy, much exceeding the short almost radical culms.) PHYLLÓSTACHYS, Torr. & Gr.

7. **C. Willdenovii**, Schk. *Sterile flowers* 4-8, closely imbricated; *perigynia* 6-9, somewhat alternate, *oblong, rough on the angles and tapering beak*; achenium oblong, triangular, finely dotted; *stigmas downy*. — Copses, Mass. to N. Virginia and westward.

8. **C. Steudèlii**, Kunth. *Sterile flowers* 10-15, rather loosely imbricated into a linear (apparently distinct) spike; *perigynia* 2-3, *roundish-obovoid, smooth, with a long and abrupt rough beak*; achenium roundish, obscurely triangular, very minutely dotted; *stigmas downy*. (*C. Jamèsii*, Schw.) — Woody hillsides, N. New York to Illinois and Kentucky.

9. **C. Bäckii**, Boott. *Sterile flowers* 3, *inconspicuous*; *perigynia* 2-4, *loose, globose-ovoid, with a conical beak, smooth throughout*; achenium globose-pyriform, scarcely dotted; *stigmas smooth*. — Rocky hills, W. Massachusetts (Mount Tom, Prof. Whitney), and N. New York to Ohio, Lake Superior, and northward. — Culms generally shorter, and the leafy scales broader and more conspicuous, than in the last two.

C. *Spikes several or numerous, androgynous* (rarely diceious), *sessile, forming a compact or more or less interrupted sometimes paniculate-compound inflorescence*: *stigmas* 2: *achenium lenticular*. VIGNÈA, Beauv.

§ 1. **Spikes approximated, the staminate and pistillate flowers variously situated**: *perigynia plano-convex, nerved, with a rough slightly toothed beak*: bracts light brown, resembling the scales, or with a prolonged point, shorter than the (at maturity) brown and chaffy spikes. — SICCATÆ.

10. **C. bromoides**, Schk. *Spikes* 4-6, *alternate, oblong-lanceolate, some of the central ones wholly fertile*; *perigynia* erect, narrow-lanceolate with a tapering point, solid and spongy at the base, longer than the lanceolate scale; style jointed at the base. — Swamps: common. — Slender, occasionally diceious.

11. **C. siccata**, Dew. *Spikes* 4-8, *ellipsoid, the uppermost and commonly 1-3 of the lowest fertile below, the intermediate ones frequently all staminate*; *perigynia* ovate-lanceolate, compressed, with a long rather abrupt beak, about the length of the scale; style minutely hairy. (*C. pallida*, *C. A. Meyer.*) — Sandy plains, New England to Illinois, and northwestward.

12. **C. disticha**, Huds. *Spikes numerous, short and ovoid, the upper or middle ones frequently almost all staminate, the lower principally or entirely fertile*; *perigynia* ovate-lanceolate, the margins not united to the top, leaving a deep cleft on the outer side; scale ovate, pointed, nearly the length of the perigynium. (*C. intermedia*, Good. *C. Sartwellii*, Dew., and former editions.) — Seneca Co., New York (*Sartwell*) to Illinois, Wisconsin, and northward. (Eu.)

§ 2. *Spikes pistillate below, staminate at the summit.*

* *Perigynia of a thick and corky texture, with a short 2-toothed roughly margined beak, nerved towards the base, dark chestnut-brown and polished at maturity*: spikes decomposed, paniculate: scales light brown, with white membranaceous margins; the bracts at the base resembling them, and with a short bristly prolongation. — PANICULATÆ.

13. **C. teretiuscula**, Good. *Spikes with short appressed branches, crowded in a slender spiked panicle*; *perigynia* ovate, *unequally biconvex, short-stalked, with 3-5 short nerves on the outer side near the broad somewhat heart-shaped*

base; scale acute, rather shorter than the perigynium; *achenium obovoid-pyriform, obtusely triangular*. (*C. paniculata*, var. *teretiuscula*, *Wahl.*) — Swamps, especially northward. (Eu.)

Var. **máior**, Koch. Spikes more paniced; perigynia rather narrower. (*C. Ehrhartiana*, *Hoppe*. *C. prairiea*, *Dew.*) — Bogs and low grounds, New England to Wisconsin, and northward. (Eu.)

14. **C. decompósita**, Muhl. Panicle large, with very numerous densely-crowded spikes on the rather short spreading branches; *perigynia obovate, unequally biconvex, sessile, with a short very abrupt beak, conspicuously nerved on each side*, about the length of the ovate pointed scale. (*C. paniculata*, var. *decomposita*, *Dew.*) — Swamps, W. New York (*Sartwell*) to Pennsylvania, Illinois, and southwestward.

* * *Perigynia small*, compressed, 2-3-nerved, membranaceous, with a short 2-toothed rough beak, yellow or brown at maturity: *spikes decomposed, with numerous small very densely-flowered heads*: scales of the fertile spikes tawny, with the green keel prolonged into a rough point: bracts short and resembling them at the base, or often becoming green and bristle-shaped, and much exceeding the culm. — MULTIFLORÆ.

15. **C. vulpinoidea**, Michx. Spike oblong and dense, or more or less interrupted ($1\frac{1}{2}' - 2\frac{1}{2}'$ long), of 8-10 crowded clusters; perigynia ovate from a broad base, with a more or less abrupt beak, diverging at maturity. (*C. multiflora*, *Muhl.* *C. bracteosa* and *C. polymorpha*, *Schw.* *C. microspërma*, *Wahl.*) — Varies with the perigynium narrower, and the beak tapering and more strongly serrulate. (*C. setacea*, *Dew.*) — Low meadows: everywhere common.

* * * *Perigynia on short stalks*, plano-convex, without a margin, membranaceous, with a thick and spongy base and a long tapering 2-toothed rough beak, distinctly nerved (only obscurely so in No. 19 and 20), widely spreading and yellow at maturity: spikes dense, more or less aggregated, sometimes decomposed: scales of the fertile spikes tawny, with a sharp point: bracts bristle-shaped, shorter than the thick and triangular culms. — VULPINÆ.

16. **C. crus-córvi**, Shuttleworth. Spike very large, decomposed, the lower branches long and distinct, the upper shorter and aggregated; *bracts often 2-toothed at the base; perigynia attenuated from an ovate dilated and truncate base into a very long slightly-winged beak, much exceeding the scale; style tumid at the base*. (*C. sicæformis*, *Boott.* *C. Halei*, *Dew.*) — Swamps, Ohio to Wisconsin, and southwestward. — A conspicuous, very large species, with spikes 4' - 9' long, often somewhat paniculate, and *glaucous leaves 6" wide*.

17. **C. stipáta**, Muhl. Spikes 10-15, aggregated, or the lower mostly distinct and sometimes compound; *perigynia lanceolate, with a long beak tapering from a truncate base, many-nerved, much exceeding the scale; style slightly tumid at the base*. (*C. vulpinoidea*, *Torr.*, *Cyp.*, not of *Michx.*) — Low grounds: common. — Culm flaccid: spikes pale.

18. **C. conjúcta**, Boott. Resembles the preceding; but the spikes (6-12) more simple; *perigynia ovate from a subcordate flat (not corky-tumid) base, short-beaked, fewer-nerved, longer and broader than the pointed scale; style bulbous at the base*. (*C. vulpina* of former editions; — from which it

differs in its flaccid culm, transversely wrinkled sheaths, orbicular achenium, &c.)—Ohio to Illinois and Kentucky.

19. **C. alopecoïdea**, Tuckerman. Head of 8–12 aggregated spikes, oblong, dense; *perigynia compressed, nerveless or very obscurely nerved, ovate from a broad truncate or somewhat heart-shaped base, a little longer than the scale; achenium pyriform; base of the style not tumid.* (C. cephalophora, var. maxima, Dew.)—Woods, W. New York to Penn., Michigan, &c.—Resembles the last, but smaller, with shorter and more compact spikes; easily distinguished by the nearly nerveless *perigynia*, and the different achenium and style.

20. **C. muricata**, L. Spikes 4–6, ovoid, approximate but distinct, the lowermost sometimes a little remote; *perigynia ovate-lanceolate, somewhat compressed, nerveless, or very obscurely nerved towards the base, rather longer than the scale; achenium ovate; base of the style not tumid.*—Fields, Massachusetts (introduced?), Ohio, and Kentucky: rare.—Spikes mostly looser than in the last, the *perigynia* narrower, with a longer and more tapering beak. (Eu.)

* * * * *Perigynia sessile, plano-convex, compressed, more or less margined, membranaceous, with a rather short and rough (or wholly smooth in No. 26) 2-toothed beak, spreading and green at maturity: scales of the fertile spikes tawny or white: bracts bristle-shaped, commonly shorter than the culm.*—
MUHLENBERGIANÆ.

21. **C. sparganioides**, Muhl. Spikes 6–12, ovoid; *the upper ones aggregated, the lower distinct and more or less distant; perigynia broadly-ovate, nerveless, rough on the narrow margin, about twice the length of the ovate-pointed scale; achenium roundish-ovate; style short, merely tumid at the base.*—Var. MINOR, Boott, is merely a reduced form. (C. cephalophora, var. Torr. C. muricata, var. cephaloïdea, Dew. C. cephaloïdea, Dew. in part.)—Low rich grounds.—A robust species, with rather wide pale-green leaves; sometimes with 1–2 short branches of a few spikes each at the base of the compound spike (probably C. divulsa, Pursh, not of Goodenough).

22. **C. cephaloïdea**, Dew. (in part), Boott. Spikes 5 or 6, contiguous; *the broadly ovate perigynia wing-margined, spongy at the base, shorter beaked, equalling or shorter than the cuspidate-tipped scale; style bulbous at the base.*—New York? Illinois (Vasey).—Much resembles and has been confounded with the small form of the foregoing.

23. **C. cephalophora**, Muhl. Spikes 5–6, small, and densely aggregated in a short ovoid head; *perigynia broadly ovate, with 3–4 indistinct nerves on the outer side, scarcely longer than the ovate roughly-pointed scale; achenium and style as in the last.* (C. cephaloïdea, Dew. (in part.))—Var. ANGUSTIFOLIA, Boott, is a narrow-leaved, smaller form. (C. Leavenworthii, Dew.)—Woods and fields: common.

24. **C. Muhlenbergii**, Schk. Spikes 3–9, closely approximate, forming an oblong head; *perigynia orbicular-ovate, with a very short beak, prominently many-nerved on both sides, about the length of the ovate roughly-pointed scale; achenium orbicular, with a very short bulbous style.*—Fields: rather common, especially southward.—Culms 12'–18' high, and with the leaves pale and rigid: commonly with a bract to each spike.

Var. **enérvis**, Boott. Scales sometimes pointless; *perigynia* nearly or wholly nerveless; spikes often bractless. — Hudson River, New York, *J. L. Russell*. — Distinguished from *C. cephaloidea* by its rigid culm, narrower leaves, and firmer perigynium, spongy at the base.

25. **C. rosea**, Schk. Spikes 4–8, the 2 uppermost approximate, the others all distinct, and the lowest often remote; *perigynia* oblong (about 8–10 in each spike), narrow at the base, widely diverging at maturity, twice as long as the broadly ovate obtuse scale. — Var. **MINOR**, Boott, has the 4–6 spikes smaller and more separated, the scales less obtuse and with a rough mucronate point; *perigynia* more erect; leaves narrower. — Var. **RADIATA**, Dew, is still more slender, almost capillary, and has only 3 or 4 remote and 3–4-flowered spikes. (*C. neglécta*, *Tuckerm.*) — Moist woods and meadows: common.

26. **C. retrofléxa**, Muhl. Spikes 3–6, all approximate, the 1–2 lowest distinct but not remote; *perigynia* (about 5–7 in each spike) ovate, or ovate-lanceolate, smooth on the margin and beak, not much exceeding the ovate-lanceolate pointed scale, widely spreading or reflexed at maturity. (*C. rosea*, var. *retrofléxa*, *Torr., Cyp.*) — Copses and moist meadows: less common than the last, from which it is distinguished by the smaller approximate spikes, longer and sharper scales, and especially, from every species in this subsection, by the smooth perigynium.

* * * * * *Perigynia* plano-convex, without a beak, of a thick and leathery texture, prominently nerved, smooth (except on the angles), with a minute and entire or slightly notched white membranaceous point: achenium conformed to the perigynium, crowned with the short thick style: bracts like the scales (brown), the lowest with a prolonged point: rootstock creeping. — **CHORDORHIZÆ.**

27. **C. chordorhiza**, Ehrh. Culms branching from the long creeping rootstock (4'–9' high), smooth and naked above, clothed at the base with short appressed leaves; spikes in an ovoid head; *perigynia* ovate, a little longer than the scale. — Cold bogs, New York to Wisconsin, and northward. (Eu.)

28. **C. tenélla**, Schk. Spikes 2–6, very small, rather remote, or the upper approximate, with 2 or 3, rarely 4, fertile flowers; *perigynia* ovate, twice as long as the scale. (*C. loliacea*, *Schk. suppl.*, not of *L.* *C. dispérma*, *Dew.* *C. gracilis*, *Ed. 1*, not of *Ehrh.*) — Cold swamps, New England to Penn., Wisconsin, and northward. — A slender species, 6'–12' high, with long grassy leaves, growing in loose tufts. (Eu.)

§ 3. Spikes pistillate above, staminate at the base.

* Spikes roundish-ovoid, rather small, more or less distant on the zigzag axis (closely aggregated in No. 30): *perigynia* plano-convex, smooth, pale green, becoming whitish or silvery: scales white and membranaceous; the bracts resembling them, or prolonged and bristle-shaped. — **CANESCÉNTES.**

+ *Perigynia* mostly somewhat thickened and leathery, distinctly nerved, and with a smooth or minutely serrulate short point, entire or slightly notched at the apex.

29. **C. trispérma**, Dew. Spikes 2–3, very small, with about 3 fertile flowers, remote, the lowest with a long bract; *perigynia* oblong, with numerous slender nerves, longer than the scale. — Cold swamps and woods, especially on mountains, New England to Penn., Michigan, and northward. — Resembling the last, but larger spikes and fruit, and weak spreading culms, 1°–2° long.

30. **C. tenuiflora**, Wahl. *Spikes 3, few-flowered, closely approximated; perigynia ovate-oblong, about the length of the broadly ovate scale.* — Cold swamps, N. New England to Wisconsin, and northward. (Eu.)

31. **C. canescens**, L. (in part). *Pale or glaucous; spikes 5-7 (about 12-20-flowered), the upper approximated, the rest distinct, the lowermost remote; perigynia ovate, equalling the pointed scale.* (*C. curta*, Good. *C. Richardi*, Michx.) — Marshes and wet meadows. common, especially northward. (Eu.)

Var. **vitalis** is a more slender and weak form, not glaucous, with smaller and roundish 6-15-flowered spikes, the more pointed perigynia spreading (and often tawny) at maturity: perhaps a good species. (Var. *alpicola* and var. *sphærostachya*, Ed. 1. *C. tenella*, Ehrh. *C. Persoonii*, Steber. *C. vitalis*, Fries. *C. Gebhardi*, Hoppe. *C. sphærostachya* and *C. Buckleyi*, Dew.) — On mountains, and high northward. (Eu.)

32. **C. Norvègica**, Schk. *Pale; stem 1° or less high, angled; spikes 2-5, rather approximate, oblong, short-bracted, with a few staminate flowers below the numerous fertile ones, or the terminal one all staminate; perigynia oval or oblong, lenticular, many-nerved, with a short entire beak, equalling the obtuse scale.* — Salt marsh, Wells, Maine, *Rev. J. Blake*. (Eu.)

← ← *Perigynia thin, spongy-thickened at the base, scarcely nerved, 2-toothed.*

33. **C. Deweyana**, Schw. *Spikes about 4; the 2 uppermost approximate, the others distinct, the lowest long-bracted; perigynia oblong-lanceolate, tapering into a rough serrate-margined beak, rather longer than the sharply pointed or awned scale.* — Copses, New England and New York to Wisconsin, and northward.

* * *Spikes ovoid or obovoid, more or less clustered; perigynia concave-convex, compressed, margined or winged, nerved, with a rough 2-toothed beak, often tawny at maturity: scales tawny or white, awnless: bracts bristle-shaped, usually falling before the maturity of the spikes.*

← *Spikes small: perigynia usually (but not always) becoming spongy-thickened at the base, the margins rigid.* — STELLULATÆ.

↔ *Spikes completely or incompletely diœcious.*

34. **C. exilis**, Dew. *Spike commonly solitary and cylindrical, diœcious or androgynous (staminate) and contracted below, often with 1-6 small additional fertile spikes contiguous to the terminal larger one; perigynia ovate-lanceolate, plano-convex, with a few fine nerves only on the convex side, spreading, turning brownish, longer than ovate acute or obtuse scale; leaves involute-filiform.* — Swamps, E. New England to New Jersey, near the coast: also borders of mountain lakes, Essex County, New York.

35. **C. stérilis**, Willd. *Spikes 4-6, all staminate and rarely all pistillate on some plants, many androgynous, especially the lower spikes, all oblong or the fertile roundish; perigynia ovate from a broad somewhat heart-shaped sharply margined base, flat, diverging or the lower recurved, very acutely 2-toothed at the apex, about equalling the acute or pointed scale; leaves narrow, involute, pale.* (*C. stellulata*, var. *sterilis*, of former ed.) — Swamps and wet meadows: common, especially northward.

++ ++ *Spikes androgynous.*

36. **C. stellulata**, L. Spikes 3-5, the uppermost much contracted at the base by the numerous staminate flowers; perigynia ovate or slightly heart-shaped at the base, which has thickish or obtuse margins, the apex more minutely toothed; scales rather blunt and considerably shorter; leaves flatter and pale: otherwise nearly as in the last. — Lake Superior and northward. (Eu.)

Var. **scirpoides**. Culms slender and weak (9' - 20'); leaves very narrow; spikes contiguous, smaller; perigynia with a rounded or truncate base, plano-convex, almost twice the length of the obtuse scale. (C. scirpoides, Schk.) — Wet places: common.

Var. **angustata**, is remarkable for the narrow lanceolate perigynia more than twice the length of the blunt scale and oblong achenium: otherwise as in var. scirpoides. — Fairfield, New York.

+ + *Spikes rather large: perigynia thickened and spongy on the angles, with a more or less dilated membranaceous margin or wing.* — OVULES.

37. **C. sychnocéphala**, Carey. *Spikes densely clustered, forming a short compound spiked head, subtended by 3 very long and unequal persistent leafy bracts; perigynia tapering from an abruptly contracted ovate base into a long slender beak, somewhat exceeding the lanceolate abruptly mucronate scale.* (C. cyperoides, Dew., not of L.) — Jefferson County (Vasey & Knieskern) and Little Falls, New York, Vasey. — Different in habit from the rest of this section; recognized at once by the ovoid compound spike, subtended by long leafy bracts, by which the lower spikes are partly concealed.

38. **C. arida**, Schw. & Torr. *Spikes 8-10, approximate ($\frac{3}{4}$ ' long), oblong-cylindrical, contracted at each end; perigynia narrowly lanceolate (4-5 lines in length), tapering into a long beak more than twice the length of the ovate-lanceolate scale; achenium sessile, narrowly oblong.* (C. Muskingumensis, Schw.) — Wet meadows, Ohio to Wisconsin and Kentucky. — In characters scarcely distinguished from the next, but strikingly different in appearance; much larger, with long, dry, and chaffy-looking spikes.

39. **C. scoparia**, Schk. *Spikes 5-8, club-shaped, at length ovate, more or less approximate, sometimes forming a dense head; perigynia elliptical-lanceolate, tapering into a long slender beak, longer than the lanceolate pointed scale; achenium distinctly stalked, exactly oval.* — Low meadows: everywhere common. — Spikes brownish or straw-colored when ripe. — Var. **minor**, Boott. Spikes more rusty, smaller, contiguous; perigynia narrowly lanceolate. — Base of White Mountains, New Hampshire, and northward.

40. **C. lagopodioides**, Schk. *Spikes 10-15 or more, approximate, or the lower more separated; perigynia lanceolate, nearly twice the length of the ovate-oblong rather pointed scale; achenium narrowly oval, on a short stalk; leaves very tapering to the apex; their sheaths loose, enlarging upwards, sharp-edged.* — Moist, rather shady places: common.

41. **C. cristata**, Schw. *Spikes smaller, 8-12 closely aggregated, globular, greenish; perigynia oblong or ovate, recurved at maturity; scales obtuse; otherwise as in the last, of which in former editions it was taken for a variety.* — Wet or moist ground: common.

Var. **mirabilis**, Boott, has *broadly ovate perigynia* with a shorter beak, longer than the acute scale. (*C. festucacea*, var. *mirabilis* of former editions. *C. mirabilis*, *Dew.*)—Mass. to Ohio, &c.

42. **C. adústa**, Boott (not of former ed.). *Spikes 4-10, pale or brown, globular*, or the upper club-shaped, the *lower remote* and sometimes compound; *perigynia oval, ovate, or roundish*, gradually tapering to a beak, many-nerved, with the narrow *wing wider above the middle*, turgid at maturity, equalling the scale in length and breadth; achenium large, orbicular, sessile. (*C. argyrántha*, *Tuckerm.*, is a very delicate form of this, found in rocky woods.)—Moist copses, &c., from Rhode Island (*Olney*) and New Jersey (*Kneiskern*), northward and westward: rare.

43. **C. fœnea**, Willd. *Spikes 3-8, pale or silvery green*, finally straw-colored, mostly approximate, *ovoid*, generally acute, the uppermost contracted or club-shaped at the sterile base; *perigynia oval, orbicular or obovate*, short-beaked, broadly winged, *appressed*, transversely wrinkled, a little longer than the ovate or lanceolate white scale; achenium on a short stalk, oval.—Sandy and mostly salt or brackish marshes, &c., along the coast, from New England southward.

Var. ? **ferruginea**, referred here by Dr. Boott, with rusty-colored acute spikes, and longer-beaked perigynia, generally acutish at base and exceeding the acute or mucronate scale (Ohio, *Sullivant*), connects this with *C. straminea*.

Var. ? **sabulónum**, also referred here by Dr. Boott, has 2-10 drooping rather remote spikes, more or less obovate or club-shaped, contracted at base, pale green turning straw-color; perigynia broadly winged at base, slightly exceeding the pointed scale: it is *C. adústa* of former editions, not of Boott.—Sands of the sea-shore from Maine southward.—Leaves narrow, often involute.

44. **C. straminea**, Schk. *Spikes 2-12, pale or tawny*, varying from obovate-globular to club-shaped, contiguous or rather remote; *perigynia orbicular-ovate or oval, often heart-shaped* at base, *very flat*, abruptly contracted into a short or tapering into a longer beak, winged, much wider and commonly longer than the usually acute or pointed scale; achenium nearly sessile, oval.—Open grounds and borders of woods: common, and very variable. The following are the varieties designated by Dr. Boott.—Var. **ΤΥΠΙΚΑ**, with 3-6 roundish spikes; perigynia spreading, broadly winged, rather longer and much broader than the scale.—Var. **ΤÉΝΕΡΑ**, more slender and delicate, with 3-6 smaller oval or obovate spikes, more tawny; perigynia with a short and broad beak, rather longer and broader than the scale. (*C. tenera*, *Dew.* *C. festucacea*, var. *tenera*, of former ed., in part.)—Var. **ΑΡÉΡΤΑ**. Spikes 4-8, tawny, tapering at base, drooping; perigynia long-beaked, thrice the length of the very sharp-pointed scale, loosely spreading in the spike.—Var. **FESTUCÆA**. Spikes 5-8, club-shaped, tawny or greenish; perigynia abruptly short-beaked and mostly narrowly winged, longer than the acute or mucronate scale; plant tall and rather rigid. (*C. festucacea*, *Schk.*, and former ed.)—Var. **ΗΥΑΛÎΝΑ**, a chiefly Western form, approaching the next species, with larger and thick pale spikes, usually 6, all tapering at base; perigynia greenish, with a wide spongy wing, and a long beak, about twice the length of the brown pointless scale; rather tall and stout, with broad leaves. (*C. straminea*, var. *Crawei*, *Boott*; but probably by some mistake said to be collected in Michigan by the late *Dr. Crave*. *C. hya-*

lina, *Boott*, is a small Texan form of it.) — Var. *MEADII* (Illinois, *Dr. Mead*), resembles the last, but has rather smaller and rounder spikes, thinner wings to the perigynia, and long-tapering or rough awn-pointed scales.

45. *C. alata*, Torr. Spikes 3–10, pale, turgid-ovoid, contiguous, mostly large (6''–10'' long); perigynia dilated orbicular or obovate, broadly winged, abruptly short-beaked, either heart-shaped or wedge-shaped at the base, longer and thrice broader than the lanceolate or ovate acute or rough awn-pointed scale; achenia stipitate. — W. New York (*Sartwell*) to Virginia and southward. — All these, from No. 38 to the present, run together variously.

D. *Staminate and pistillate flowers borne in separate (commonly more or less stalked) simple spikes on the same culm*; the one or more staminate (sterile) spikes constantly uppermost, having occasionally more or less fertile flowers intermixed; the lower spikes all pistillate (fertile), or sometimes with staminate flowers at their base or apex: *stigmas 3*: *achenium sharply triangular* (only usually 2 stigmas and the achenium lenticular in No. 46–56, 65, 149, 150). *CAREX* proper.

§ 1. *Perigynia without a beak, or scarcely any, smooth, not inflated* (slightly in No. 55, 56), *terminating in a minute, straight, entire or notched point*, glaucous-green when young, becoming whitish, often spotted or tinged with purple, or occasionally nearly black at maturity: *pistillate scales blackish-purple* (barely brown in No. 55, 56, 64), *giving a dark appearance to the spikes*.

* *Sterile spikes 1–3, stalked*, often with more or less fertile flowers: *pistillate spikes 3–5, frequently with sterile flowers at the apex*: bract of the lowest spike leaf-like, mostly with dark-colored expansions (auricles) at the base, and very minute sheaths or none. (*Culm and leaves more or less glaucous*.)

+ *Stigmas 2* (in No. 46 and 47* sometimes 3): *perigynia lenticular*. — *ACÛTÆ*.

++ *Alpine, saxatile: pistillate scales pointless: leaves flat*.

++ *Scales awnless, mostly obtuse*.

46. *C. rigida*, Good. *Sterile spike solitary; the fertile 2–4, cylindrical, erect, rather loosely flowered*, the lower on short peduncles; lowest bract about the length of the culm, with rounded auricles; *stigmas 2–3*; perigynia elliptical, with an entire *scarcely pointed apex, nerveless*, about as long as the obtuse scale; culm rigid, nearly smooth except towards the top, about the length of the firm erect leaves. (*C. saxatilis*, *Fl. Dan.*, partly of *L.*) (Eu.) — Our plant is the

Var. ? *Bigelovii* (*C. Bigelovii*, Torr. *C. Washingtonia*, *Dew.*), with 3–5 longer and laxer fertile spikes, the lowest long-stalked, spreading, and sometimes remote; the sterile or terminal one often fertile at the top; perigynia more or less nerved: perhaps a distinct species. — Alpine summits of the mountains of New England, New York, and high northward.

++ ++ *Not alpine, paludose: pistillate scales awnless, single-nerved*.

a. *Leaves with involute margins when dry; their sheaths not fibrillose*.

47. *C. vulgaris*, Fries. *Sterile spike 1–3; the fertile 2–4, approximated, oblong, erect, densely-flowered*, occasionally staminate at the apex, the lowest on a very short stalk; lowest bract barely the length of the culm, with *small blackish rounded auricles*; *perigynia ovate-elliptical, stalked, nerved* especially towards the base, with a very short abrupt entire or minutely notched point, *longer than the obtuse appressed black scale*; culm slender, sharply triangular,

nearly smooth, except at the top. (*C. acuta*, var. *vulgaris*, *L.* *C. cæspitosa*, of authors.)—Wet banks, &c. New England to Wisconsin and northward.—Grows in small patches (not in dense tufts like No. 50), and varies in height from 3' to 18', with narrow leaves shorter than the culm. (Eu.) The following may be appended.

47^a. *C. limula*, Fries? Fertile spikes less approximate or rather remote; their bracts surpassing the culm (at least the lowest) and with rougher margins; leaves longer and more numerous at the base of the rougher culm; perigynia nerveless; stigmas often 3: otherwise like *C. vulgaris*.—E. New England, near Boston, *W. Boott*, who rather doubtfully identifies it with the Lapland plant. The specimens in Herb. Suec. Norm. differ in their flat leaves, and narrower, longer, and even pointed scales. (Eu.)

48. *C. aquatilis*, Wahl. Sterile spikes commonly 2–3; the fertile 3–5, cylindrical, inclining to club-shaped, erect, densely-flowered, sessile, or the lower on very short stalks; bracts long, 1–2 lowest exceeding the culm; perigynia obovate-elliptical, stalked, nerveless, with a very short entire point about the length of the lanceolate scale; culm smooth, not much exceeding the pale-green glaucous leaves.—Margins of lakes and rivers, New England to Wisconsin, and northward.—Robust, 2°–3° high; the thick fertile spikes 1'–3' long. (Eu.)

b. *Leaves with more or less revolute margins when dry; sheaths at length fibrillose, i. e. when old splitting up or resolved more or less into slender parallel or loosely reticulated fibres.*

49. *C. torta*, Boott. Sterile spikes 1–2, commonly 1, fertile 3–4, elongated, narrowly-cylindrical or slightly club-shaped, loosely few-flowered at the base, occasionally more or less staminate at the apex, the lower on smooth slender stalks, spreading or drooping; bracts with oblong auricles, or very slightly sheathing, the lowest about the length of the culm, the rest bristle-shaped, shorter than their respective spikes; perigynia elliptical, short-stalked, tapering to a distinct point, with a minutely notched or jagged membranaceous orifice, very smooth, nerveless, the empty tips spreading or obliquely recurved at maturity, scarcely exceeding the narrow obtuse scale; achenium broadly obovate, much shorter than the perigynium; culm very smooth, leaves short, slightly rough on the margin only. (*C. acuta*, var. *sparsiflora*, *Dew.*?)—Rills and wet banks, N. New England, New York, &c., and along the mountains from Penn. southward.—Well marked by its smooth flaccid culm (12'–18' high), soft and short grassy leaves, and the tortuous empty apex of the perigynium.

50. *C. aperta*, Boott. Sterile spikes 1–2, oblong-cylindrical, acute; fertile 2–4, oblong, erect, the uppermost approximate and sessile; the lower distant and short-stalked, staminate at the apex, or often entirely fertile; lowest bract about the length of the culm, with oblong brown auricles, or very slightly sheathing, the upper bristle-shaped, shorter than the spikes; perigynia roundish-ovate, stalked, without nerves, covered with very minute transparent dots, and sometimes very slightly rough at the apex, with an abrupt very short notched orifice, broader and much shorter than the lanceolate pointed brown scale; culm sharply triangular, smooth below, exceeding the rough sharp-pointed leaves.—Wet meadows, Rhode Island and Mass. to Illinois and far westward.—Culm 1°–2° high, with commonly

2 fertile spikes 9''–18'' in length, appearing somewhat bristly from the long and spreading scale. Differs from the next chiefly in the rounder perigynium and nearly smooth culm, and should perhaps be referred to it.

51. **C. stricta**, Lam. (not of Good.) Sterile spikes 1–3; the fertile 2–4, cylindrical, slender, usually barren at the summit, sessile, or the lower on a short stalk; lower bract with rounded or oblong brown auricles, seldom exceeding the culm; perigynia ovate-acuminate or elliptical, nerveless, or very obscurely few-nerved, often minutely rough on the short, entire, or slightly notched point, usually shorter and broader than the narrow reddish-brown scale; culm slender, sharply triangular, rough, longer than the narrow and rigid rough and glaucous leaves; their older sheaths with conspicuous reticulated fibres. (C. acuta, Muhl., &c., not of L. C. Virginiana, Smith in Rees, Cycl. C. acuta, var. erecta, Dew. C. angustata, Boott.)—Var. STRICTIOR has shorter and more densely flowered fertile spikes, and perigynia equalling or somewhat exceeding the scale. (C. strictior, Dew.)—Wet meadows and swamps: very common.—Grows in large and very compact tufts: culms 2°–2½° high. Scales of the fertile spikes very variable; the lower commonly acute, the upper narrower and obtuse. (C. xerocarpa, S. H. Wright, in Dew. Cat., seems to be a mere state of C. stricta.)

52. **C. lenticularis**, Michx. Sterile spike single and mostly fertile at the top; the fertile 2–5, erect, cylindrical (6''–12'' long), sessile, or the lower short-peduncled, densely-flowered; bracts exceeding the culm; perigynia ovate-oval, sessile, more or less nerved, abruptly short-pointed, the point entire, slightly exceeding the oblong and very obtuse scale; culm (6'–20' high) and leaves smooth or nearly so, pale.—Wet gravelly banks and shores, N. Maine (J. Blake, C. E. Smith), N. New York (Torrey, Knieskern, J. A. Paine), Upper Michigan (Prof. Porter, &c.), and northward.

++ ++ ++ Paludose or maritime: pistillate scales awned or pointed from the broad and strong more or less triple-nerved centre or midrib.

53. **C. salina**, Wahl. Sterile spikes 2–3; the fertile 2–4, cylindrical, erect, often sterile at the apex, on more or less included stalks; bracts long, with rounded auricles, the two lowest commonly exceeding the culm; perigynia ovate-elliptical, with a minute entire point, nerveless, rather shorter than the roughly-awned dark-brown scale; culm (1°–2° high) rough at the top, rather exceeding the leaves.—Salt marshes, Massachusetts (Greene, W. Boott), Maine (G. L. Goodale), and far northward. (Eu.)

54. **C. maritima**, Vahl. Sterile and fertile spikes each 2–4 (the latter rarely 5 or 6) (1'–2' long), spreading or drooping on slender peduncles; perigynia nearly orbicular, with a short entire point, much shorter than the long-awned greenish scale; culm (1°–2° high) and the broad flat leaves smooth. (C. paleacea, Wahl.)—Salt marshes, Cambridge and Medford, Mass. (Greene, W. Boott), Wells, Maine, (J. Blake) and northward: rare. (Eu.)

55. **C. crinita**, Lam. Sterile spikes 1–2, often with fertile flowers variously intermixed; the fertile 3–5, long-cylindrical (2'–3' long), densely flowered, often staminate at the apex, on exerted nodding stalks; bracts very long, exceeding the culm; perigynia roundish-obovate, slightly inflated, obscurely nerved, with a short entire point, shorter than the oblong mostly notched roughly-serrate awned

light-brown scale; cum (2° – 4° high) *rough and sharply angled above, leafy below; the pale leaves* ($3''$ – $4''$ wide), rough on the edges, their surface and the *sheaths smooth*. — Varies in size (but usually tall); and with the lower fertile scales often very long-awned, the fruit imperfect and deformed (var. *MÓRBIDA*, *Carey* in *Sill. Jour.* *C. paleacea* of authors). — Wet grounds by streams: common.

56. *C. gynándra*, Schw., Boott. *Sheaths rough* with minute hairiness; fertile spikes rather thicker and looser, and oftener staminate at the apex; perigynia more ovate or oblong and elliptical; the scales longer and less spreading but mostly shorter-awned: otherwise as in the preceding, — to which it is very nearly related. — In similar situations, but less common, from New England to Michigan.

+ + *Stigmas* 3: perigynium obtusely triangular, indistinctly few-nerved, more or less compressed: *pistillate spikes on filiform drooping stalks*. — LIMÓSÆ.

57. *C. Barráttii*, Schw. & Torr. *Sterile spike mostly single, sometimes 2 or even 3, dark purple; fertile mostly 2 or 3, cylindrical, commonly staminate at the top; lower bract usually shorter than the culm; sheaths obsolete or minute; perigynia oval or oval-lanceolate, obliquely divergent, scarcely notched at the point, about the length of the ovate and blunt black-purple scale; culm* (1° – 2° high) sharply triangular, nearly smooth, longer than the glaucous flat leaves; the old sheaths at base splitting into threads. (*C. flácca*, of former ed., and probably a mere geographical variety of that European species.) — Marshes, New Jersey near the coast, *Collins, Knieskern*; and Townsend, Delaware, *W. M. Canby*.

58. *C. limòsa*, L. *Staminate spike solitary; fertile 1–2, oblong, 10–20-flowered, occasionally with staminate flowers at the apex; bracts very narrow, the lowest shorter than the culm; perigynia ovate, with a minute entire point, about equal to the ovate mucronate dull or purplish-brown scale*. — Peat-bogs, New England to Pennsylvania, Wisconsin, and northward. — Culm $6'$ – $12'$ high, erect, sharply triangular, longer than the acute and rigid keeled leaves. (Eu.)

59. *C. rariflòra*, Smith. Resembles the last (of which it was thought to be a variety), but smaller, $4'$ – $9'$ high; culm obtuse-angled; leaves flatter and rather broader; pistillate spikes with only 5–10 less crowded flowers; perigynia very short-pointed or bluntish, rather shorter than and involved in the broadly-ovate black-purple scale. — Mt. Katahdin, Maine (*G. L. Goodale*), and northward. (Eu.)

60. *C. irrigua*, Smith. *Staminate spike solitary; the fertile 2–4, ovoid or oblong, occasionally staminate at the apex, or with a few sterile flowers at the base; lowest bract as wide as the leaves, longer than the culm; perigynia roundish-ovate or obovate, with an entire orifice, much shorter than the tapering and slender-pointed dark purple scale*. (*C. Magellánica*, *Lam.*, according to *Boott.* *C. limòsa*, var. *irrigua*, *Wahl.* *C. paupércula*, *Michx.*) — Peat-bogs, New England to Penn., Wisconsin, and northward. — Taller than No. 58, growing in loose clumps, with weaker and nodding stems, often exceeded by the leaves. (Eu.)

* * *Uppermost spike club-shaped, pistillate above and staminate at the base; the rest all fertile or with a few sterile flowers below: lowest bract leaf-like, scarcely equalling the culm, with minute light-brown auricles and no sheaths: culm and leaves of a pale glaucous-green*. — ATRÁTÆ.

← *Scales of the spike deep-colored, purple or dark brown.*

61. **C. Buxbaumii**, Wahl. *Spikes 3-4, obovoid or oblong, the uppermost short-stalked (rarely altogether staminate), the others nearly sessile, the lowest somewhat remote; perigynia elliptical, obtusely triangular, compressed, obscurely nerved, with a distinctly notched orifice, scarcely equalling the ovate sharp-pointed or short-awned (brown-purple) scale. (C. canescens, L., in part.)* — Peat-bogs: not rare. (Eu.)

62. **C. atrata**, L. *Spikes 3-4, oblong-ovoid, approximate, all on short filiform stalks, at length drooping; perigynia ovoid, with a short notched point, about the length of the ovate acute or dark brown-purple scale. — Alpine summits of the White Mountains, New Hampshire. — About 12'-15' high, with rather rigid leaves, nearly equalling the culm. Fruit at first pale straw-color, often becoming dark purple or nearly black. Stigmas sometimes 2. (Eu.)*

63. **C. alpina**, Swartz. *Spikes 3-4, small, oval or globular, crowded into a head at the summit of the slender naked culm, nearly sessile, mostly overtopped by a foliaceous bract; perigynia orbicular or obovate, pointed with a small short beak, minutely notched at the orifice, roughish, longer than the ovate bluntish black-purple scale. (C. Váhlíi, Schk.) Isle Royale, &c., Lake Superior, Prof. Whitney, C. G. Loring, Jr., and northward. — Culms in ours 1°-2° high, the leaves all at the base: spikes 2"-4" long. (Eu.)*

+ + *Scales and spikes greenish turning straw-color.*

64. **C. Shortiana**, Dew. *Spikes 3-5, cylindrical, erect, more or less distant ($\frac{1}{2}$ '-1 $\frac{1}{2}$ ' long), and the lowest rather remote, all androgynous and densely flowered; the terminal one about half staminate, the rest with only a few barren flowers at the base, the lower on short stalks; perigynia broadly obovate, abruptly contracted at the base into a short stalk, with an extremely minute entire point, little longer than the short-pointed somewhat obovate scale. — Marshes, S. Pennsylvania to Illinois and southward. — Plant 1°-3° high: leaves flat, 3" wide.*

§ 2. *Perigynia without a beak (except in No. 67, &c.), smooth, slightly inflated, bluntly triangular, nerved, with an obtuse and pointless orifice, or a short (and straight or oblique) entire or notched point: bracts leaf-like, sheathing: staminate spike solitary (except sometimes in No. 71), or androgynous and pistillate above; the rest all fertile.*

* *Staminate spike on an elevated stalk (rarely short-stalked or sessile, or with 1-2 small ones at its base): pistillate spikes 1-6, erect, the upper on very short, the lower on more or less elongated exerted stalks (short and included in No. 73): bracts shorter than the culm (except in No. 65 and 72): perigynia with an entire and straight or obliquely bent point, glaucous-green when young, becoming cream-colored or yellow at maturity, sometimes spotted with purple; pistillate scales dark-brown with white margins, fading to tawny. (Leaves mostly radical, more or less glaucous.) — PANICEÆ.*

← *Stigmas mostly 2: perigynium wholly pointless, turgid-obovate.*

65. **C. aurea**, Nutt. *Fertile spikes 3-4, oblong, loosely flowered, the lowest often very remote; perigynia pear-shaped, obtuse, longer than the ovate acute scale; achenium lenticular. (C. pyriformis, Schw.) — Wet grassy banks, especially on limestone, W. New England to Wisconsin, and northward. — A*

slender delicate species, 4' - 8' high, with long grassy leaves, and bracts exceeding the culm. Sterile spike often with some fertile flowers at the apex.

+ + *Stigmas* 3: *perigynium* somewhat pointed and 3-sided.

+ + *Staminate spike* (or the cluster in No. 71) long-stalked.

66. **C. livida**, Willd. *Fertile spikes* 1 - 2, rarely with a third near the base of the culm, 10 - 15-flowered; *perigynia* ovoid-oblong, with faint pellucid nerves, tipped with a straight obtuse point, rather longer than the ovate scale. (*C. limosa*, var. *livida*, Wahl. *C. Grayana*, Dew.) — Peat-bogs and wet pine barrens, New Jersey, Oneida Co., New York, and high northward. — Rarely with a single (sterile) spike, or with an additional fertile one on an erect stalk 4' - 9' long, from the base of the culm. Plant very glaucous, the leaves rigid and finely tapering. (Eu.)

67. **C. vaginata**, Tausch. *Sterile spike* with its stalk commonly bent to a right angle with the culm at flowering time, afterwards erect; fertile 2 or 3, remote, erect, slender-peduncled, loosely flowered; bracts foliaceous, short, with dilated sheaths; *perigynia* short-ovate when mature and with a distinct terete beak or beak-like oblique point, emarginate at the orifice, exceeding the ovate acute scale; culm slender (1° - 2° long), weak and reclining, naked, *stoloniferous*; the long-creeping sterile shoots bearing tufts of flat green leaves (2'' - 3'' wide) almost equalling the fertile culms. (*C. sparsiflora*, Fries. *C. phæostachya*, Smith.) — Moist banks, Bergen swamp, Genesee Co., New York (*J. A. Paine*), Lake Superior (*Robbins and Porter*), and northward. (Eu.)

68. **C. panicea**, L. *Sterile spike* always erect; fertile 1 - 3, mostly 2, erect, remote, oblong or short-cylindrical, rather loosely flowered, only the lower slender-peduncled; sheaths of the short foliaceous bracts shorter and narrower; *perigynia* turgid-ovate at maturity, obscurely nerved, tipped with a short bent entire point (mostly straw-colored), longer than the ovate blunt scale. — Moist grounds, Massachusetts to Delaware (*W. M. Canby*): rare. (Eu.)

69. **C. Meadii**, Dew. Differs from the last only in the denser fertile spikes, the sterile one sometimes longer; and the *perigynia* more triangular, less turgid, paler, less indistinctly nerved, the scales pointed; culms more rigid and roughish: the more slender forms closely approach the next. (*C. panicea*, chiefly of former ed.) — Wet prairies, &c., Ohio to Illinois and Wisconsin.

70. **C. tetanica**, Schk. *Fertile spikes* 1 - 3, commonly 2, oblong-cylindrical, loosely flowered, especially at the tapering base, remote; *perigynia* when young pointed at each end, at maturity obovoid, scarcely inflated, with a slightly bent point, longer than the ovate obtuse and often abruptly mucronate or awn-pointed scale. (*C. conoidea*, Gray, *Gram. & Cyp.*, not of Schk. *C. Woodii*, Dew.) — Margins of lakes and rivers, W. Mass. to Penn., Michigan, and southward.

71. **C. Cráwei**, Dew. *Sterile spikes* often 1 or 2 small ones at the base of the terminal, which is occasionally fertile at the apex; *fertile spikes* 3 - 6, remote, and the lowest near the root, oblong or cylindrical, densely flowered, and sometimes slightly compound at the base; their short peduncles included, or the lowest exserted; *perigynia* ovoid-oblong, obscurely nerved, with a very small straight or slightly recurved point, longer than the ovate obtuse or acute or short-pointed scale. (*C. heterostachya*, Torr.) — Wet places, S. Herkimer and Jefferson

Counties, New York, to Illinois, Michigan, and northwestward. — Variable, 4'–12' high; the taller forms resembling the next; but the perigynium is less round and with fewer and more indistinct nerves, the bracts do not exceed the culm, and the staminate spike is long-peduncled.

++ ++ *Staminate spike nearly sessile: perigynium turgid, many-nerved, greenish.*

72. **C. granulàris**, Muhl. Sterile spike mostly shorter than the approximate uppermost of the 3 or 4 fertile; these cylindrical, dense, the lowest remote and slender peduncled, sometimes from near the root; perigynia globular-ovoid, contracted into a short often slightly bent point, longer than the pointed scale; plant pale, glabrous; bracts long, exceeding the culm. — Wet meadows: common.

73. **C. Torrèyi**, Tuckerman. Sterile spike overtopping the 2 or 3 ovoid contiguous nearly sessile fertile spikes; perigynia obovate, strongly nerved, the very obtuse or retuse summit abruptly tipped with a very short cylindraceous beak-like point, longer than the ovate mostly acute or pointed scale; culm, leaves, and short sheathless bracts downy. (C. abbreviata, Schw., Boott.) — Bethlehem, Pennsylvania, Schweinitz, New York? and high northward. — Probably overlooked from its close external resemblance to the next; but it is very distinct.

* * *Staminate spike sessile, or short-stalked* (except in No. 75): pistillate spikes 2–5, erect, all on more or less exserted stalks; bracts longer than the culm (except in No. 75): perigynia very obtuse, with an abrupt and minute or almost obsolete point, green and somewhat pellucid at maturity: pistillate scales tawny, fading to white. — PALLESCÉNTES.

74. **C. palléscens**, L. Fertile spikes 2–3, ovoid, densely flowered, approximate; perigynia obovoid-oblong, obscurely nerved, about the length of the scale. — Var. **UNDULATA** is a mere state with the lower bract at the base transversely wavy-lined. (C. undulata, Kunze.) — Meadows, New England to Penn. and northward. — Plant 8'–18' high: culm and leaves slightly pubescent. (Eu.)

75. **C. conoidea**, Schk. *Staminate spike on a long stalk; fertile 2–3, oblong, closely flowered*, the lower distant; perigynia oblong-conical, with impressed nerves, slightly oblique at the summit, rather longer (or sometimes shorter) than the sharply pointed or awned scale; bracts not exceeding the culm. (C. tetanica, Schw. & Torr., not of Schk.) — Moist meadows: rather common.

76. **C. grisea**, Wahl. Fertile spikes 3–6, oblong, rather loosely flowered, remote, or the 2 upper contiguous and the lowest distant; perigynia oblong, rather longer than the broadly ovate abruptly strongly awned scale (the awn rough-hispid; style bulbous-thickened; leaves light green, flat, rather broad. (C. laxiflora, Schk., not of Lam.) — Moist grounds: rather common.

77. **C. flaccospérma**, Dew. Differs from the preceding in having the flaccid leaves paler or glaucous; spikes longer, cylindrical (about 1' long), with more numerous flowers, usually more distant, their bracts shorter; perigynia larger (2''–3'' long), laxer, turning brownish, twice or thrice the length of the short-pointed or short-awn-pointed scale; style not thickened. (C. grisea, var. mítica, of former ed. Prof. Dewey now proposes to change the name to *xanthosperma*, because flaccosperma is a hybrid word and “yellow-fruited” was meant. But the fruit is not yellow, and the original name has been adopted by Boott and Steudel.) — Low grounds, Penn. and New Jersey, *Knieskern* (a dubious form), and common southward.

* * * *Uppermost spike fertile-flowered at the apex (rarely all staminate): pistillate spikes 3-5, oblong or cylindrical, loosely flowered, distant, on exerted filiform stalks: bracts equalling or often exceeding the culm: perigynia oblong, with a short and abrupt notched point (obsolete in No. 80), green and membranaceous at maturity: pistillate scales tawny or white.* — GRACILLIMÆ.

+ Fertile spikes nodding or pendulous.

78. **C. Davisii**, Schw. & Torr. *Fertile spikes oblong-cylindrical, rather thick; perigynia somewhat contracted at each end, scarcely longer than the conspicuously awned scale. (C. aristata, Dew., not of R. Br. C. Torreyana, Dew.)* — Wet meadows, Massachusetts to Wisconsin, and southward. — Larger than the next (1°-2° high), and with stouter and longer spikes.

79. **C. formosa**, Dew. *Fertile spikes oblong, short, all commonly with 2 or 3 barren flowers or empty scales at the base; perigynia somewhat contracted at each end, nearly twice as long as the pointed or cuspidate scale.* — Wet meadows, Massachusetts to W. New York.

80. **C. gracillima**, Schw. *Fertile spikes linear, slender; perigynia obtuse and slightly oblique at the orifice, longer than the oblong awned or awnless scale. (C. digitalis, Schw. & Torr., not of Willd.)* — Wet meadows, New England to Kentucky, Wisconsin, and northward. — When the uppermost spike is altogether staminate this resembles *C. arctata*; but is distinguished by the obtuse and sessile perigynium.

+ + Fertile spikes nearly erect, all but the lowest short-peduncled or nearly sessile.

81. **C. æstivalis**, M. A. Curtis. *Spikes slender, loosely flowered; perigynia acutish at both ends, twice the length of the ovate obtuse or mucronate scale; achenium somewhat stipitate; sheaths of the lower leaves pubescent: otherwise nearly as the last, but smaller (1°-1½° high).* — Saddle Mountain, W. Massachusetts (*Dewey*), mountains of Penn., Virginia, and southward.

§ 3. *Perigynia without a beak, hairy (in No. 83 becoming smooth at maturity), slightly inflated, bluntly 3-angled, obtuse, conspicuously nerved, with a minute abrupt straight point: bracts narrow, with very short or obsolete sheaths, the lowest exceeding the culm; pistillate scales tawny or white; spikes 2-4, erect, the uppermost androgynous, fertile-flowered at the apex and club-shaped: the rest all fertile.* — VIRESCENTES.

82. **C. virescens**, Muhl. *Spikes oblong or cylindrical, on short stalks; perigynia ovoid, nearly entire at the orifice, rather longer than the ovate awned scale; leaves and sheaths hairy. (C. costata, Schw.)* — Rocky woods and hillsides, New England to Michigan, and southward. — Culms rough, slender, 1°-2° high: fertile spikes 6''-12'' long.

83. **C. triceps**, Michx. *Spikes ovoid, nearly sessile, closely approximate; perigynia broadly obovoid, entire at the orifice, downy when young, smooth at maturity, rather longer than the pointed scale; sheaths very hairy; leaves more or less so. (C. hirsuta, Willd. C. viridula, Schw. & Torr., not of Michx.)* — Varies with the spikes rather longer and on stalks, and leaves nearly smooth. (*C. hirsuta*, var. *pedunculata*, Schw. & Torr.) — Woods and meadows: the smoother form southward. — Culm 12'-18' high. Spikes 6''-9'' long.

§ 4. *Perigynia* without a beak, smooth, not inflated, 3-angled, regularly striate, terminating in a short, entire, rather obliquely bent or recurved point, remaining green at maturity: pistillate scales membranaceous, mostly tipped with a rough point or awn, brown or spotted, fading to white: staminate spike solitary: pistillate spikes 2-5, more or less remote, the lowest often near the base of the culm.

* Sterile spike club-shaped: fertile spikes (erect, the uppermost commonly near the base of the sterile) all on stalks principally included within sheathing bracts, except sometimes the lowest, and shorter than the spikes or not much exceeding them: perigynia ovoid-triquetrous, narrowed at each end: culms numerous, diffuse and in fruit becoming prostrate: leaves all radical, very broad, finely and closely nerved throughout, with 3 distinct ribs. — PLANTAGINEÆ.

84. **C. plantaginea**, Lam. Fertile spikes commonly 4, oblong, about 5-8-flowered; bracts very short, dark purple, or the lowest greenish. (*C. latifolia*, Schk.) — Shady woods, mostly on hillsides in rich soil, New England to Penn., Wisconsin, and northward; and southward through the Alleghanies.

85. **C. Careyana**, Torr. Fertile spikes 2-3, ovoid or oblong, about 3-5-flowered; bracts green, the upper equal to the spikes, the lower somewhat exceeding them; perigynia large (2"-2½" in length); leaves dark green. — In similar situations with the last, N. New York to Penn. and Ohio: rare.

86. **C. platyphylla**, Carey. Fertile spikes 3, filiform, loosely 3-4-flowered; bracts as in the last; perigynia small; culms slender; leaves pale or whitish-green. — In similar situations with No. 84; and commoner southward.

* * Sterile spike short, club-shaped, pedunculate: fertile spikes 2-4, all on filiform exerted stalks, with long sheathing bracts resembling the leaves; the uppermost, as well as the leaves, exceeding the slender and at length prostrate culms: perigynia as in the last subsection. — DIGITALES.

87. **C. retrocúrva**, Dew. Fertile spikes ovoid or oblong, compactly 3-8-flowered, on long drooping stalks, frequently with one or two staminate flowers at their base; leaves glaucous, 3-4 lines wide, with 3 prominent nerves. — Copses and hillsides, New England to W. New York, Virginia, and southward. — Closely approaching the next; perhaps only a variety of it.

88. **C. digitális**, Willd. Fertile spikes linear-oblong, loosely 6-9-flowered, on long stalks, the lowest sometimes drooping; leaves and bracts narrow, dark green; perigynia smaller than in the last. (*C. oligocárpa*, Schw. & Torr., not of Schk. *C. Vanvléckii*, Schw.) — Copses and hillsides: not rare. — Slender, 6'-12' high, growing in tufts, with numerous culms and long grassy leaves.

* * * Sterile spike linear, either conspicuously peduncled or smaller and nearly sessile in the same species: fertile spikes 2-6, erect; the 1-2 uppermost commonly near the base of the sterile, on an included stalk; the rest on exerted stalks, with long sheathing bracts resembling the leaves; the uppermost exceeding the erect culm: perigynia with obtuse angles, about the length of the scale. — OLIGOCÁRPÆ.

† *Perigynia* distinctly nerved, and with a hyaline orifice: style nearly even: scales of the pistillate spikes awnless or barely awn-pointed.

89. **C. laxiflora**, Lam. Fertile spikes slender and loosely several-many-flowered on a zigzag rhachis, cylindrical, or sometimes rather dense and oblong;

perigynia oval-triangular, narrowed to each end, the point either strongly or sometimes slightly recurved. — A most polymorphous species; very common in open woods, copses, &c. The leading forms were collated by Dr. Boott as follows. — The typical form taken by him (*C. striatula*, Michx.) has the leaves long and narrow (2"–4" wide), spikes about 1' long and loosely flowered; perigynia with a slightly recurved or almost straight point, their scale except the lowest nearly pointless; sterile spike mostly conspicuous and long-peduncled. — Var. *STYLOFLÉXA*, Boott. (*C. styloflexa*, Dew. *C. fusiformis*, Chapman.) Slender, the weak filiform culms 1° – $2\frac{1}{2}^{\circ}$ long; fertile spikes 2–4, short, 5–10-flowered, the lowest on a long setaceous peduncle; bracts mostly shorter than the culm; perigynia more tapering or triangular-fusiform, the point commonly recurved. New Jersey (*C. F. Austin*), Penn. (*Prof. Porter*), and southward. — Var. *PLANTAGÍNEA*, Boott (var. *patulifolia* of former edition. *C. plantaginea*, Schk.), has the root-leaves 5"–7" broad, otherwise as in the typical form. — Var. *INTERMÉDIA*, Boott (*C. anceps*, Willd., Schk.), includes various slender, narrow-leaved forms, with loosely-flowered spikes, but otherwise as the next. — Var. *BLÁNDA* (*C. blanda*, Dew. *C. laxiflora*, var. *striatula*, Ed. 2), includes narrow-leaved forms, 6'–18' high, with the sterile spike usually short or club-shaped and nearly sessile; the fertile oblong and dense, the uppermost approximate; bracts much surpassing the culm; perigynia obovate with a short abruptly bent point; scale usually awn-pointed. — Var. *LATIFÓLIA*, Boott, has a broadly and very sharp-angled culm, and very broad leaves and bracts (8"–15" wide), inconspicuous sterile spike, the fertile ones cylindrical and loosely flowered, but the broad perigynium much longer than the truncate or abruptly short-pointed scale. *Var. v. Scous, 1857.*

+ + *Perigynia densely striate, or as if finely wrinkled rather than nerved, of a firm texture, obscurely triangular, with a callous orifice: style thickened above the base: scale with the somewhat 3-nerved keel extended into a stout rough awn or point.*

90. *C. oligocárpa*, Schk. Fertile spikes small, 3–8-flowered; the point of the perigynium straight or slightly oblique, not recurved; leaves rough only on the edge; sheaths smooth. (*C. Sartwelliana*, Gay.) — Woods, W. New England to Illinois and Kentucky. — Culm slender, 6'–12' long.

91. *C. Hitchcockiana*, Dew. Fertile spikes very loosely 3–5-flowered; apex of the perigynium recurved; sheaths and upper side of the leaves roughly pubescent. — Woods, New England to Illinois and Kentucky. — Culm 1° – 2° high, stouter, and fruit larger than in the last.

§ 5. *Perigynium without a beak, smooth or downy, not inflated, obovoid-triangular, with a minute obliquely bent white and membranaceous point, reddish-brown or olive-colored at maturity: bracts reduced to colored sheaths, or with a short green prolongation: leaves all radical, narrow or bristle-shaped.* — *DIGITATÆ.*

92. *C. ebúrnea*, Boott. Sterile spike solitary; the fertile 3–4, erect, about 5-flowered, approximated and elevated on long stalks above the staminate spike: the lowest sometimes a little remote; perigynia obscurely nerved, smooth and shining, rather longer than the broad and obtuse membranaceous whitish scale. (*C. alba*, var. *setifolia*, Dew.) — Limestone rocks, N. New England to Kentucky, and northward. — A delicate species, with very slender culms, 4'–10' high, and

bristle-shaped leaves, forming dense tufts. The fertile spikes 2''-3'' in length, about 1'' broad.

93. **C. pedunculata**, Muhl. Spikes 3-5, commonly 4, the uppermost sterile with 2-3 fertile flowers at the base, the rest fertile with a few staminate flowers at the apex, all on long stalks, remote, 1-2 of the lowest near the base of the culm; sheaths with green tips much shorter than the stalks; perigynia with a long attenuated base and a minutely notched orifice, somewhat downy, especially on the angles, about the length of the broadly obovate abruptly awned or pointed dark-purple scale. — Dry woods and hillsides, E. New England to Penn., Wisconsin, and northward. — Culms 4'-10' high, prostrate at maturity, in tufts, partly concealed by the very long and narrow grassy leaves.

§ 6. *Perigynia with a straight or slightly bent more or less abrupt beak, hairy, not inflated, terminating in a membranaceous notched or 2-toothed orifice: bracts short, either green and slightly sheathing or auriculate at the base, or small and resembling the scales: scales dark brown or purple with white margins, fading lighter or sometimes turning nearly white: staminate spike solitary; the fertile 2-3, nearly sessile and erect, or the lower on a long radical peduncle. (Culms mostly low and slender: leaves all radical, long and narrow.)* — MONTANÆ.

94. **C. umbellata**, Schk. Culms very short (1'-3', rarely 6' high), in close tufts; staminate spike sometimes with a few pistillate flowers; fertile spikes 4-5, ovoid, few-flowered; the uppermost commonly close to the sterile spike and sessile, the rest on stalks arising from the base of the stem and of about equal height, nearly concealed by the long grassy leaves; perigynia ovoid, 3-angled, with a rather long abrupt beak, about the length of the ovate pointed scale. — Rocky hillsides, New England to Illinois, and northward.

95. **C. Novæ-Angliæ**, Schw. Sterile spike sessile, short and usually inconspicuous; fertile 2-4, greenish-purple, 3-8-flowered, contiguous and sessile, or the lowest rather distant (sometimes even radical) and more or less peduncled; the lower or all the leafy bracts exceeding the culm; perigynia globular-pear-shaped with a much attenuated base and a short conical 2-toothed beak, minutely hairy, longer and broader than the ovate mucronate-pointed purple scale (with green midrib and hyaline margins); achenium apiculate with the very short persistent base of the style; culms very slender (4'-10' long), weak, soon reclined or procumbent. — Saddle Mountain, Massachusetts, Adirondack Mountains, New York, and high northward. — Too near *C. pilulifera*, L., of Europe and the following.

96. **C. Emmonsii**, Cew. Paler, and the spikes greenish, not purple, usually more crowded than in the foregoing, often a long-peduncled one from the base; bracts short, rarely equalling the culm; perigynia oval and more 3-sided, hairy, and with a longer cylindrical beak; base of the style deciduous by an articulation. (*C. Novæ-Angliæ*, var. *Emmonsii*, Ed. 2. *C. Davisii*, Dew., &c.) — Dry woody hills: not rare.

97. **C. Pennsylvanica**, Lam. Sterile spike commonly on a short stalk; fertile 1-3, usually 2, approximate, nearly sessile, ovoid, 4-6-flowered, the lowest commonly with a colored scale-like long-awned bract; perigynia roundish-ovoid, with

a short and abrupt minutely-toothed beak about the length of the ovate pointed chestnut-colored scale. (*C. marginata*, Muhl.) — Dry woods and hills: common, especially northward.

98. **C. varia**, Muhl. Sterile spike sessile; fertile 2-3, mostly 3, distinct, on very short stalks, ovoid, 6-10-flowered; the lowest and sometimes the 2 lower with green leaf-like bracts; perigynia obovoid, with an abrupt distinctly toothed beak, about the length of the ovate pointed light-brown scale. (*C. Pennsylvanica*, var. *Muhlenbergii*, Gray, *Gram. & Cyp.*) — Dry wooded hills: common, especially northward. Closely resembles the last; but has wider, shorter, and more rigid glaucous leaves; also taller (1° - $1\frac{1}{2}^{\circ}$) and more erect than No. 96, broader-leaved, and the spikes scattered. All these seem to run together.

99. **C. PRÆCOX**, Jacq. Sterile spike club-shaped; fertile 2-3, oblong-ovoid, aggregated near the base of the sterile spike, sessile, or the lowest sometimes on a very short stalk, with a leaf-like bract scarcely exceeding the spike; perigynia ovoid-triangular, attenuated at the base, with a short beak and nearly entire orifice, about equal to the ovate pointed dark-brown scale; achenium obovoid, with a prominent ring at the apex surrounding the base of the style; culm 3'-6' high; leaves short, rather rigid. (*C. verna*, Villars, *Dew.*, not of *Schk.*) — Rocky hills, Salem and Ipswich, Massachusetts. (Nat. from Eu.)

100. **C. Richardsonii**, R. Br. Sterile spike peduncled, cylindrical; fertile 1 or 2, sessile or short-stalked, approximate, oblong, longer than the scale-like brownish and mostly short-pointed bracts; perigynia obovoid-triangular, with a tapering base, obtuse, nearly beakless, the short point with an almost entire orifice, rather shorter than the ovate acutish brown or chestnut-colored scale; culm (5'-9' high) and rigid leaves rough. — Dry ground, near Rochester, New York (*Dewey*); prairies of Illinois (*Meud*); Wisconsin (*Sartwell*), and northward. — A well-marked species, in aspect most like No. 97.

101. **C. pubescens**, Muhl. Sterile spike usually sessile; fertile 3-4, oblong or cylindrical, loosely flowered, somewhat approximated, or the lowest a little remote, on a short stalk, with a narrow leaf-like bract about the height of the culm; fruit ovoid and sharply triangular, downy, attenuated at the base, with an abrupt slender beak nearly entire at the orifice, a little longer than the ovate abruptly-pointed white scale; culm and flaccid flat leaves soft-downy. — Moist woods and meadows, New England to Wisconsin and Kentucky. Differs from the other species of this section in its greater size and in aspect, and especially in the sharply angled perigynium.

§ 7. *Perigynia* slightly inflated, with a short beak, terminating in an entire or slightly notched orifice; staminate spike solitary, stalked (in No. 102 usually pistillate at the summit): culms tall and leafy. (Two quite unlike species, artificially combined merely for convenience.) — ANÓMALÆ.

102. **C. miliacea**, Muhl. Staminate spike commonly fertile at the summit; fertile spikes 3, cylindrical, rather slender, loosely flowered at the base, on filiform nodding stalks; bracts exceeding the culm, with short or nearly obsolete sheaths; perigynia ovoid-triangular, very smooth and thin, with an entire or very minutely notched orifice, longer than the ovate short-awned white scale. (*C. prasina*, *Wahl.*) — Rills and wet meadows: rather common.

103. **C. scabrata**, Schw. Fertile spikes 4-5, cylindrical, erect, rather distant, *densely flowered*, the lower on long stalks; bracts *without sheaths*, exceeding the culm; perigynia ovoid, contracted at the base, *prominently few-nerved, rough*, spreading at maturity, with an obliquely notched beak, longer than the ovate slightly ciliate brown scale; culm leaves and bracts *very rough*. — Wet meadows and swamps, E. New England to Penn., Michigan, and northward.

§ 8. *Perigynia slightly inflated, 3-angled, smooth and shining* (minutely pubescent in No. 104 and one form of 108), *green, with a straight tapering beak* (short-pointed in No. 107), terminating mostly in 2 small membranaceous teeth: lower bracts green and sheathing: pistillate scales tawny or white: staminate spike solitary, stalked: *pistillate spikes 3-4, loosely flowered, all* (except in No. 104) *on filiform nodding stalks*.

* Fertile spikes mostly slender, remote; perigynia somewhat nerved: bracts equalling or exceeding the culm. — DÉBILES.

+ Leaves and sheaths more or less soft-pubescent: fertile spikes nearly erect.

104. **C. Sullivantii**, Boott. Fertile spikes 3-5, commonly 4, narrowly cylindrical, erect, rather dense, the upper approximate, the lowest often remote, tapering towards the base and slightly compound, all on rough peduncles; bracts sheathing, not exceeding the culm; perigynia elliptical, hairy, slightly stalked, very obscurely nerved, with an entire or notched orifice, rather longer than the ovate ciliate rough-awned or merely mucronate white scale. — Woods, Columbus, Ohio, *Sullivant*.

105. **C. Knieskérnii**, Dew. Less pubescent; fertile spikes 2 or 3, on longer and somewhat spreading peduncles; perigynia glabrous, more evidently nerved: otherwise like the preceding: too little known. — Copses, Oneida Co., New York, *Dr. Knieskern, Dr. Vasey*.

+ + Glabrous or nearly so: fertile spikes mostly nodding or spreading, loose.

106. **C. arctata**, Boott. Fertile spikes slender-cylindrical, narrowed towards the base; perigynia ovate, short-stalked from a blunt base, short-beaked, longer than the pointed scale. (*C. sylvatica, Dew., not of Hudson.*) — Woods and meadows, New England to Pennsylvania, and northward.

107. **C. glabra**, Boott. Fertile spikes oblong or short-cylindrical, rather dense and the terminal one oftener partly fertile; perigynia elliptical-oblong, not stalked, somewhat contracted at the base, and short-pointed at the apex, but nearly beakless, minutely and sharply 2-toothed at the orifice, prominently nerved, almost twice the length of the blunt brownish-margined scale. — Oneida Co., New York, *Dr. Knieskern*. Near Philadelphia, *C. E. Smith*. Probably not rare, but confounded with the next: also resembling *C. formosa*.

108. **C. débilis**, Michx. Staminate spike occasionally fertile at the apex; fertile spikes slender-cylindrical, with loose alternate flowers on a somewhat zigzag rhachis; perigynia spindle-shaped or oblong-lanceolate, tapering into a slender beak with a hyaline 2-lobed tip, twice as long as the obtuse and pointless scale. (*C. tenuis, Rudge. C. flexuosa, Muhl.*) — Moist meadows: rather common, especially southward. — Perigynium often rusty-dotted: — in var. *PÛBERA* minutely pubescent and passing to *C. venusta, Dew.* of the Southern States. Bear Meadows, Centre Co., Penn., *Prof. Parter*.

* * Fertile spikes short : perigynia nerveless : bracts short. — FLÉXILES.

109. **C. capillaris**, L. Fertile spikes commonly 3, minute, with about 6 alternate flowers ; perigynia oblong-ovoid, contracted at the base, tapering into a long slightly serrulate beak, with an oblique nearly entire orifice, longer than the ovate scale. — Point de Tour, Lake Michigan ; alpine summits of the White Mountains, New Hampshire ; and high northward. — An extremely delicate species, 4' - 6' high, with spikes 3'' - 6'' long, and a line or less in width. (Eu.)

110. **C. fléxilis**, Rudge. Sterile spike short and club-shaped ; fertile spikes oblong or sometimes with a few staminate flowers at the base and becoming club-shaped ; the upper bracts short and scale-like, the lower bristle-shaped, very slightly sheathing ; perigynia ovoid, obscurely nerved, tapering into a beak about the length of the ovate hairy-fringed scale ; leaves pale green and glaucous, and with the bracts soft-hairy. (*C. blepharóphora*, Gray.) — Moist and shady places, Connecticut (near Salisbury), Central and Northern New York, Lake Superior, and Newfoundland : rare.

§ 9. *Perigynia slightly inflated, obtusely 3-angled, nerved, smooth, tapering into a beak, with two mostly distinct teeth, becoming tawny or yellow at maturity : achenium obovate-triquetrous, contracted at the base ; staminate spike usually solitary.* — FLAVÆ.

* *Perigynia erect, slender-beaked : spikes remote ; the staminate one usually long-stalked : bracts not exceeding the culm, and with long sheaths.*

111. **C. lævigata**, Smith. Fertile spikes 3, cylindrical, on exerted nodding stalks ; perigynia ovoid, tapering into a 2-cleft beak, rather longer than the light-brown or purplish pointed and awned scale ; culm smooth. (*C. Greeniana*, Dew.) — Massachusetts (Tewksbury ? *B. D. Greene*). Introduced ? (Eu.)

112. **C. fúlva**, Good. Fertile spikes 2 - 3, oblong or ovoid, erect, the lowest on an exerted stalk ; perigynia ovoid, not much exceeding the dark-brown scarcely pointed awnless scale ; culm rough. (*C. binervis*, Dew., not of Smith.) — Pond at Tewksbury, Massachusetts, *B. D. Greene* : not since found. (Eu.)

* * *Perigynia ascending, short-beaked : spikes approximate or the lower remote ; the staminate one sessile or nearly so ; bracts much surpassing the culm ; the upper nearly without sheaths.*

113. **C. exténsa**, Good. Fertile spikes 2 - 4, oblong, brown-green, very dense (5'' - 10'' long), the upper nearly sessile, the lowest on a short included stalk ; perigynia ovate, the short conical beak sharply 2-toothed, longer than the ovate pointed purple scale ; leaves and bracts long and narrow, involute, erect, rigid ; culms tufted, 8' - 20' high. — Border of salt-marshes, coast of Long Island, New York, *Dr. T. F. Allen*. (Eu.)

* * *Perigynia spreading or reflexed, longer than the scale : spikes mostly approximate or crowded ; the staminate mostly sessile or short-stalked, often fertile at the apex or middle ; the fertile (2 - 4) erect all or all but the lowest short-stalked or sessile : bracts much exceeding the smooth culm, their sheaths very short.*

114. **C. fláva**, L. Fertile spikes 2, roundish-ovoid, dense, the upper approximated, the lowest often remote on a short exerted stalk ; bracts spreading or reflexed ; perigynia tapering from an ovoid contracted base into a slender recurved

beak, widely spreading or reflexed at maturity. — Wet meadows, especially northward. — Whole plant of a yellowish hue, 6'–15' high, with spikes about 6" in length. (Eu.)

115. **C. Cederi**, Ehrh. Fertile spikes *oblong-ovoid, closely aggregated*, or the lowest rather remote, on very short stalks, densely flowered, sometimes staminate at the apex; leaves and bracts *narrow, rigidly erect*; perigynia *ovoid, with a short and rather abrupt minutely notched beak*, scarcely recurved at maturity. (C. *viridula*, Michx., not of Schw. & Torr. C. *irregularis*, Schw.) — Wet rocks and bluffs, coast of New England to Illinois, Lake Superior, and northward. — Resembles the last; but the fertile spikes and perigynia are much smaller, and the beak more abrupt, shorter, and straight. (Eu.)

§ 10. *Perigynia slightly inflated*, ovoid or obtusely 3-angled, with an abrupt straight beak, nerved, *densely pubescent or rough-puberulent*, the pubescence nearly concealing the nerves, except in No. 119: bracts leaf-like, with short sheaths: scales dark-purple or brown.

* *Perigynia densely pubescent of a thick or somewhat leathery texture, ovoid, with 2 short and diverging membranaceous teeth: bracts much exceeding the nearly smooth culm: staminate spikes 2 or 3, the uppermost stalked, the lower short and sessile: fertile spikes 1–4, usually 2, erect, remote, sessile or on very short stalks.* — LANUGINOSÆ.

116. **C. filiformis**, L. Fertile spikes oblong or ovoid; perigynia very short-beaked and with 2 sharp teeth; leaves and bracts *narrow and involute*; culm very slender ($1\frac{1}{2}^{\circ}$ – 3° high). — Peat-bogs, New England to Penn., Wisconsin, and northward. (Eu.)

117. **C. lanuginosa**, Michx. Fertile spikes oblong or cylindrical; perigynia more hispidly pubescent; leaves and bracts *flat, broader and shorter*; culm stouter (1° – 2° high); staminate spikes usually shorter. (C. *pellita*, Muhl.) — Swamps and wet meadows. New England to Kentucky and northward.

* * *Perigynia thin, downy like the last or roughly granulate, or even smooth, ovoid, the beak terminating in a thin and scarious oblique orifice, either entire or slightly notched*; bracts rigidly erect, shorter than the sharply triangular rough culm. — SCARIOSÆ.

118. **C. vestita**, Willd. Sterile spikes 1–2, the uppermost cylindrical, short-stalked; fertile 1–2, approximate, sessile, ovoid or oblong, sometimes staminate at the apex; perigynia *densely pubescent, with a short thick beak*, a little longer than the ovate pointed scale; leaves *flat*, shorter than the stout and rigid culm. — Sandy soils, growing in tufts, New England to Penn. and southward. — Resembling the last in external appearance, but readily distinguished by the membranaceous beak of the fruit, which is reddish at the base and white and transparent at the orifice; and the style is twisted within the perigynium.

119. **C. polymorpha**, Muhl. (in part.) Sterile spikes 1–4, the uppermost on a long stalk; the lower short, often with a few fertile flowers at the base; fertile spike *solitary or rarely 2, remote, cylindrical, sometimes staminate at the apex, erect, on partly exerted stalks*; perigynia *few- (5–10-) nerved, very minutely roughened with granular dots, or smooth, abruptly contracted into a slender*

cylindrical usually purplish beak, with a whitish hyaline entire orifice, longer than the ovate blunt purplish scale. (*C. Halseyana*, *Dew. & Ed. 1.* *C. striata*, *Torr. N. Y. Fl.*, not of *Michx.*) — Varies considerably; in one form with the fertile spikes filiform, and the flowers alternate and very distant on the rhachis. — Upland meadows, Rhode Island and Mass. to Pennsylvania.

§ 11. *Perigynia moderately inflated, conspicuously many-nerved, smooth or pubescent, with a straight beak terminating in 2 rigid more or less spreading teeth: bracts leaf-like, with very short sheathing bases, equalling or exceeding the culm: staminate spikes 1-5.*

* *Perigynia with a short and thick beak, and short teeth.* — LACÚSTRES.

+ *Perigynia hairy, sometimes glabrate, turgid-ovate.*

120. **C. striata**, *Michx.* (not of *Ed. 1.*) Sterile spikes 3, the uppermost slender-stalked; fertile spikes 1-2, oblong, erect, remote, sessile or on short stalks (or the lower rarely on a slender stalk); perigynia minutely hairy or smoothish, or rarely smooth, rather thin, longer than the blunt or pointed scale, the teeth usually scariously lobed at the base; leaves and bracts long and narrow, rather rigid, involute, with slender or setaceous rough extremities. (*C. polymorpha*, *Ed. 1.*) — Wet places, New Jersey to Virginia, and southward.

121. **C. Houghtonii**, *Torr.* Sterile spikes mostly 2; fertile 2-3, oblong or cylindraceous, thicker and less distant, olive-colored; perigynia hairy, thick-walled, longer than the pointed or short-awn-tipped scale; the teeth at the orifice narrow and entire; leaves and bracts flat, shorter and broader, and culm lower (9' - 15') than in the foregoing. — Wisconsin (Lake La Biche), *Dr. Houghton*; shore of Lake Ontario, *Prof. Whitney*; Medford above Bangor, Maine, *J. Blake*; and northwestward.

+ + *Perigynia very smooth, very finely many-nerved.*

122. **C. riparia**, *Curtis.* Sterile spikes 2-5, the uppermost stalked; fertile spikes 2-3, oblong-cylindrical, erect, remote, nearly sessile, or the lowest on a short stalk, large and thick (2' - 3' long, 4" - 6" wide), olive-colored; perigynia lanceolate-conical, coriaceous, tipped with rather slender short teeth, longer than the lance-ovate awned scale. (*C. lacustris*, *Willd.* and former ed.) — Borders of streams, ponds, and swamps: common. — Very robust, 3° - 5° high: leaves 3" - 5" wide, and sheaths nodose-reticulated. (Eu.)

123. **C. paludosa**, *Good.* More slender, with spikes smaller, leaves narrower, perigynia ovate, flattened, and more strongly nerved than the preceding, the orifice merely notched, and hardly exceeding the awned scale. — Border of a salt marsh at Dorchester, Mass., *W. Boott.* (Nat. from Eu.?)

* * *Perigynia with an elongated tapering beak and with long and setaceous or awn-like spreading or divergent teeth.* — ARISTATÆ.

+ *Staminate spikes 2-5 (rarely with some fertile flowers): fertile spikes remote, erect, rather loose, the uppermost almost sessile, without sheaths, the lowest often on an exerted sometimes spreading peduncle: perigynia ascending.*

124. **C. aristata**, *R. Br.* Fertile spikes 2-4, cylindrical; perigynia ovate-lanceolate, smooth, tapering into a slender beak tipped with very slender at length diverging awn-like teeth, longer than the ovate-lanceolate awned and above hispid-ciliate scale; culm smooth; sheaths and under surface of the leaves

hairy. (*C. atherodes*, Spreng. *C. mirata*, Dew.) — Lake shores and river-banks, N. New York to Michigan, and northwestward. — Culm 2°–3° high: leaves 2''–3'' wide. Fertile spikes 2'–3' long, often loosely flowered towards the base. (Eu. *C. orthostachys*.)

125. *C. trichocarpa*, Muhl. More slender; leaves and bracts rough, but not hairy; fertile spikes 2 or 3, fewer-flowered; perigynia more ovate and with shorter and stouter teeth, downy-pubescent, the scale short-awned or awnless. (*C. striata*, Ed. 1, not of Michx.) — In water or wet ground: common, especially northward.

Var. *imberbis*. Perigynia glabrous; sheaths rather rough. Penn-Yan, New York, Sartwell. Illinois, Mead, E. Hall, and northwestward.

+ + *Staminate spikes solitary, with a filiform bract; fertile 3–5, cylindrical, densely-flowered, on long exserted and at length drooping stalks, mostly approximate: perigynia widely spreading, reflexed at maturity.*

126. *C. comosa*, Boott. Fertile spikes thick (1'–3' long, and 6''–7'' wide), the lowest sometimes very remote; perigynia tapering from a stalked ovoid-triangular base into a long deeply 2-forked beak, the sharp elongated teeth widely spreading or somewhat recurved; scales lanceolate, with a long bristle-shaped awn shorter than the mature fruit; culm rough and triquetrous. (*C. furcata*, Ell., not of Lapeyr. *C. Pseudo-Cyperus*, Schw. & Torr., Dew., &c., in part, not of L.) — Wet places: common. — A robust species 2°–3° high, formerly confounded with the next, which it greatly resembles; but it differs especially in the larger fertile spikes, longer beak of the fruit, and the longer, smooth and widely spreading teeth, giving to the spikes a comose or bristly appearance.

127. *C. Pseudo-Cyperus*, L. Fertile spikes narrower and sometimes slightly compound at the base; perigynia as in the last, but with a shorter beak and shorter less spreading teeth; scale about the length of the mature fruit. — Border of lakes and in bogs, New England to Pennsylvania, and northward. (Eu.)

§ 12. *Perigynia much inflated, conspicuously many-nerved, smooth, with a long tapering 2-toothed beak: bracts leaf-like, much exceeding the culm: scales tawny or white: staminate spike stalked, always solitary.* — LUPULINÆ.

* *Bracts with very short or obsolete sheaths.*

128. *C. hystricina*, Willd. Sterile spike often bearing a few fertile flowers at the base or apex; fertile spikes 2–4, oblong-cylindrical, densely flowered, the uppermost on a very short stalk, the others on long stalks and at length nodding, the lowest often very remote; perigynia spreading, tapering from an ovoid base into a long slender beak with sharp smooth teeth, longer than the awned scale. — A variety with shorter ovoid spikes, the lowest very remote on a filiform stalk, 4'–6' long, with rather smaller perigynia not much longer than the awn, is *C. Cooleyi*, Dew. — Wet meadows: common. — Plant pale or yellowish green, with fertile spikes 9'' to 1½' long. Distinguished from the preceding by the more inflated less diverging fruit, its beak longer and teeth shorter; and from the following by the smaller nodding spikes and many-nerved perigynium, with longer and smooth teeth.

129. *C. tentaculata*, Muhl. Fertile spikes 2–3, ovoid, oblong, or cylindrical, densely flowered, approximate and diverging horizontally, the uppermost ses-

sile, the lower on short exerted stalks; perigynia spreading, tapering from an ovoid few- (about 10-) nerved base into a long slender beak with oblique orifice or short minutely serrulate teeth, much longer than the lanceolate awned scale; achenium papillose-roughened. (*C. rostrata*, Muhl., not of Michx.) — Wet meadows: very common. — Var. *GRACILIS*, Boott, is a slender form with 2 much smaller fertile spikes (4"–12" long, 3"–4" broad). — Var. *ALTIOR*, Boott, is tall (2° high), with more scattered large fertile spikes, and the more tapering beak of perigynium with longer teeth; perhaps a hybrid with *C. lupulina*. Penn-Yan, New York, Sartwell, and Amherst, Mass., Tuckerman.

130. *C. intumescens*, Rudge. Fertile spikes 1–3, ovoid, loosely few- (5–8-) flowered, closely approximated, sessile, or the lower on a very short exerted peduncle; perigynia erect-spreading, tapering from an ovoid 15–20-nerved base into a long sometimes rough beak. (*C. folliculata*, Schk., Michx., not of L.) — Wet meadows and swamps: very common. — Culm slender, about 18" high: fertile spikes usually contiguous: perigynia 6"–7" long, very ventricose.

131. *C. Grayii*, Carey. Fertile spikes 2 (sometimes single), globose, densely (15–30-) flowered, separate, on short exerted peduncles; perigynia (8" long) spreading and deflexed, tapering from an ovoid 25–30-nerved base into a long smooth and shining beak. — River bottoms, Oneida Co., New York, to Ohio and Illinois: rather rare. — Culm robust, 3° high; leaves broader; and flowers in July, a month later than the last.

* * Bracts all or all but the uppermost conspicuously sheathing.

+ Fertile spikes approximate, or only the lowest one distant, erect, very large and turgid, many-flowered: perigynia ascending, long-beaked from an ovate-ventricose base: sterile spikes rarely 2.

132. *C. lupulina*, Muhl. Fertile spikes 2–4, cylindraceous or oblong (1'–2' long, 1' thick), the lower on exerted stalks; perigynia (6½"–7" long) often raised on a short stalk-like base, smooth or with the beak rough above, much longer than the lanceolate rough awn-pointed scale; stem (2°–3°) and long broadly linear leaves and bracts smooth, the latter with rough margins (3"–4" wide). (*C. lùrida*, Wahl.) — Wet grounds, common. — *C. Canadensis*, and *C. Bellavilla*, Dew., appear to be depauperate and attenuated states of this, with more distant lax, and fewer-flowered spikes.

133. *C. lupuliformis*, Sartwell. Fertile spikes 4–5, cylindrical (2'–3' long), less approximate; perigynia sessile (7"–8" long); achenium broader, with mamillated angles; scale more awned; otherwise as in the foregoing, of which it is probably a mere variety. (*C. lupulina*, var. *polystachya*, Schw. & Torr.) — Swamps, New York to Delaware, &c.

+ + Fertile spikes distant, few-several-flowered: perigynia lanceolate, ovate-lanceolate or spindle-shaped, loose or widely spreading at maturity: staminate spike small, short-stalked: obtusely angular culms and grassy soft leaves smooth.

134. *C. folliculata*, L. Fertile spikes 3–4, remote, 12–20-flowered, all or the lowest on exerted peduncles, turning yellowish at maturity; perigynia tapering ovate-lanceolate from a broadish base, short-beaked, at length widely spreading, rather exceeding the ovate white rough-awned scale. (*C. xanthophÿsa*, Wahl.) — Peat-bogs, New England to Penn., and northward, and in one form

southward. — Plant 2°–4° high, of yellowish appearance, with long foliaceous bracts, and leaves 4'' wide.

135. *C. rostrata*, Michx. More slender and smaller than the last (10'–15' high), strictly erect, rather rigid; leaves narrow; fertile spikes 1–3, commonly 2; *perigynia scarcely spreading at maturity, lanceolate, obtusely triangular, slender-beaked, about twice the length of the brownish blunt scale.* — Cold bogs, mountains of N. New York, New England, and northward.

136. *C. subulata*, Michx. *Fertile spikes 3–5, very remote, on included peduncles loosely few- (4–8-flowered, commonly with a few staminate flowers at the apex; perigynia awl-shaped, reflexed at maturity; the long slender beak deeply cleft or grooved down one side, tipped with 2 awn-like and at length rigidly deflexed teeth.* (*C. Collinsii, Nutt. C. Michauxii, Dew.*) — Cedar swamps, Canada (*Michaux*) to Rhode Island (*Olney*), and New Jersey near the coast: rare.

§ 13. *Perigynia much inflated, obovoid or obconic, rather few-nerved, smooth, with a long and slender 2-toothed beak abruptly produced from the obtuse or at maturity even retuse summit, tawny-brown or straw-colored at maturity, spreading horizontally, or the lower deflexed, very densely aggregated in thick spikes: these, 1–7 in number, are either all androgynous, staminate at base, or the terminal wholly staminate, the others sometimes wholly pistillate: leaves and bracts flat, the latter much longer than the culm.* — SQUARRŌSÆ.

137. *C. squarrosa*, L. *Spikes often only one, commonly 2–5, globular, ovoid, or cylindraceous (6''–7'' thick), the terminal one with a slender-contracted base from the numerous staminate flowers, the others almost wholly fertile, on short slender peduncles, their bracts scarcely sheathing; perigynia sparingly nerved, longer than and concealing the blunt or short-pointed scales.* (*C. typhina, Michx.*) — Low grounds, E. New England to Illinois and southward. — Remarkable for its dense, short and thick spikes, about 1' long, to which the spreading beaks of the perigynia give a bristly appearance.

138. *C. stenolepis*, Torr. *Spikes 5–7, the terminal one wholly staminate, or sometimes fertile at the top or throughout; the others with a few staminate flowers at the base or sometimes none, cylindrical (1' or more long), the upper approximated, nearly sessile on the zigzag stem, the lower remote on exserted stalks, their bracts sheathing; perigynia shorter than the long awn-like rough scales.* (*C. Frankii, Kunth. C. Shortii, Steud., not of Torr.*) — Marshes, Pennsylvania (*Prof. Porter*) and Virginia to Illinois, and southwestward. — Somewhat resembling the last; but the spikes are narrower and more numerous, and of a still more bristly appearance from the projecting points of the scales.

§ 14. *Perigynia much inflated, smooth and shining, becoming straw-colored at maturity, with a tapering more or less elongated 2-toothed beak: bracts leaf-like, with very short or obsolete sheaths (conspicuously sheathing only in No. 139); scales brown or tawny: staminate spikes 2–5, rarely 1, stalked.* — VESICARIÆ.

* *Perigynia conspicuously nerved: bracts usually much exceeding the culm.*

— *Fertile spikes oblong or cylindrical, many-flowered.*

139. *C. retrorsa*, Schw. Sterile spikes 1–3, the uppermost occasionally with a few fertile flowers, the rest more or less pistillate at the base; *fertile spikes 4–5, oblong-cylindrical, erect, the upper approximate and clustered on short or in-*

cluded stalks, *the lowest remote* on a long exserted stalk, and (with one or more of the others) often bearing 1–2 short branches at the base; *perigynia crowded, spreading and at length reflexed*, strongly *few-nerved*, tapering from an ovoid contracted base into a *conspicuously toothed beak, much longer than the lanceolate scale*. (*C. reversa*, *Spreng.*) — Marshy borders of streams, New England to Penn., Wisconsin, and northward. — Culm nearly smooth ($1\frac{1}{2}^{\circ}$ – $2\frac{1}{2}^{\circ}$ high): leaves and bracts 3''–4'' wide, much exceeding the thick spikes, which are 1'– $1\frac{1}{2}'$ long. — **Var. HARTII** (*C. Hartii*, *Dew.*) is a slender state, with fertile spikes distant, the lower long-peduncled. — Yates Co., New York, *Dr. Hart Wright*.

140. **C. gigantea**, Rudge. Sterile spikes 1–5; sometimes with a few fertile flowers; *fertile spikes* 3–5, *cylindrical, somewhat erect*, or spreading on exserted peduncles, distant or the upper contiguous, *all or most of them staminate at the apex; perigynia ascending, at length horizontal, many-nerved*, abruptly tapering from a broadly or globular-ovate ventricose base into a long and slender sharply 2-toothed beak, much longer than the ovate-lanceolate mostly awn-pointed scale; achenium broader than high, strongly triquetrous, with concave faces. — Swamps, Delaware (*W. M. Canby*), Kentucky (*Short*), and southward. — Allied to the preceding and to No. 133. Culm 2° – 3° high: perigynia about 6'' long.

141. **C. Schweinitzii**, Dew. Sterile spikes commonly 2, the lower often pistillate at the base; *fertile spikes* 3–4, *cylindrical, somewhat drooping, densely flowered, often staminate at the apex*, and occasionally the lower rather compound at the base, on smooth nearly included stalks; *perigynia ascending, oblong-ovoid, rather lightly few-nerved, tapering into a smooth short-toothed beak, a little longer than the lanceolate roughly long-awned scale*. — Wet swamps, New England, New Jersey, W. New York, and northward: not common. — Culm 10'–15' high, smooth: bracts and leaves 2''–3'' wide, smooth except the margins, much exceeding the culm: fertile spikes (2'–3' long, rather narrow) and the whole plant turning straw-color. Perigynia $2\frac{1}{2}''$ –3'' long, thin.

142. **C. utriculata**, Boott. Sterile spikes 3–4; *fertile spikes* 2–5, commonly 3, rather distant, sessile, or the lowest (sometimes loose and attenuated at the base), peduncled, *cylindrical* ($1\frac{1}{2}'$ –4' long) *thick, and densely very many-flowered; perigynia ovate*, either ventricose and abruptly or inclining to elliptical and more gradually contracted into a cylindrical smooth beak, longer than the lanceolate very acute or awn-pointed scale; culm stout and thick, obtusely angular, spongy at base (2° – 3° high); leaves flat (3'–4' broad), pale, nodose-reticulated. (*C. ampullacea*, var. *utriculata* of former ed.) Swamps, New England to Penn., Michigan, and common northward. — Fruit sometimes almost that of *C. vesicaria*, sometimes that of *C. ampullacea*, into which it merges northward: the rough mostly awn-like points of the scale usually distinguish it from both European species.

143. **C. Vaseyi**, Dew. Differs from the last in the *slender culm with acute rough angles*; fertile spikes (2 or 3) looser and fewer-flowered; perigynia more tapering into the beak, and scales less pointed; from *C. vesicaria*, *L.* of Europe (of which it is the nearest representative) in the more pointed scales and fewer-nerved perigynia tapering gradually into a longer beak; from the next (into which it probably passes) by the larger *elongated-ovate perigynia tapering* into a slender beak (the roughness of which, indicated by *Dr. Boott*, is rarely obvious,

the teeth, however, rough-serrulate). (*C. monile*, *Dew.* in 1845, not of *Tuckerm.* *C. vesicaria*, var. *cylindræca*; *Dew.*) — Swamps, Yates Co., New York, *Sartwell*, to Illinois ?

144. **C. monile**, Tuckerman. Sterile spikes 3, rarely 2 or 4; *fertile spikes mostly 2, rarely 3 or solitary, cylindraceous or cylindrical (1' - 2½' long), mostly slender, slightly or the lower when remote longer-peduncled; perigynia globular-ovate, very ventricose, shining, abruptly contracted into a short smooth beak, longer than the ovate-lanceolate acute or scarcely pointed scale; culm slender (1° - 3° high), sharply triangular, rough on the angles; leaves narrow (barely 2" wide).* — Wet places: common, especially northward. — *Perigynia 2" - 2½" long.* (*C. vesicaria*, var. *alpigena*, *Fries*, of N. Eu. appears to be a form of this.)

145. **C. Olneyi**, Boott. Resembles the two preceding and *C. vesicaria*; the *fertile spikes denser and thicker, short-cylindrical (1' - 1½' long); perigynia (2½" - 3" long) turgid-ovate, with the short beak rough-serrate at the apex and on the awl-shaped teeth; the scales acute or bluntish; leaves (1" - 2" broad) rough.* — Wet ground, Providence, Rhode Island, *Olney*.

146. **C. Tuckermanni**, Boott. Sterile spikes about 2; *fertile spikes 2 - 3, commonly 3, oblong or cylindrical, stout, somewhat approximate, on rough stalks, the lowest often nodding; perigynia shining, thin and delicate, much inflated, ovate, ascending, tapering into a rather abrupt slender and cylindrical smooth beak, much longer than the ovate or lanceolate acute or short-pointed scale; the very long narrow bracts and leaves rough, much surpassing the rough-angled culm.* (*C. bullata* of authors, not of *Schk.* *C. cylindrica* of former ed.; but the *cylindrica* of Schweinitz belongs mainly to the next or to some others, and is too much confused for preservation.) — Swamps, W. New England to Penn., Illinois, and northward. — Differs from the next in the more numerous and longer fertile spikes (8" - 2' long), and the larger, more inflated and membranaceous fruit (4" - 5" long), with a smooth beak.

147. **C. bullata**, Schk. Sterile spikes 2 - 3; *fertile spikes most frequently only one, sometimes 2, approximated, oblong or cylindrical, stout, sessile or on short smooth stalks; perigynia spreading, shining, turgid-ovate, tapering into a long cylindrical rough beak, much longer than the lanceolate pointless scale; bracts and leaves narrow, about the length of the smooth or roughish culm.* (*C. cylindrica*, *Schw.*, at least in part, *Tuckerm.*, &c. — Wet swamps, New England to Penn. and southward, chiefly eastward.

+ + *Fertile spikes globular or oblong, few-flowered.*

148. **C. oligosperma**, Michx. Sterile spikes 1 - 2, slender; fertile spikes 1 - 2, densely 6 - 18-flowered, the lower on a very short stalk; perigynia turgid-ovate (2½" long), tapering into a short minutely toothed beak, not much longer than the ovate awnless scale; culm very slender; leaves and bracts involute. (*C. Oakesiana*, *Dew.*) — Wet borders of ponds, &c., E. Mass. to N. New York, Wisconsin, and northward, chiefly in cold or mountainous regions.

* * *Perigynia nerveless or nearly so: bracts mostly shorter than the culm.*

+ *Perigynia pointed with a very short smooth beak, minutely 2-toothed at the apex: stigmas either 2 or 3: fertile spikes 2 or sometimes one, dense, short, usually brown-purple or purplish, the upper almost sessile, the lower short-peduncled: ster-*

ile spikes 1 or 2, often solitary. (The two following are very much alike; and one or the other is doubtless *C. miliaris*, Michx. The first is regarded by Andersson as an extreme form of *C. ampullacea*, the second of *C. vesicaria*.)

149. *C. rotundata*, Wahl. ? *Leaves and bracts involute*, smoothish; fertile spikes oblong or cylindraceous, or the upper ovate or globular; perigynia short-ovate, about the length of the broadly ovate (obtuse, or in our specimens acute or pointed) scale. — Gravel-bars at the outlet of Moosehead Lake, N. Maine, *C. E. & A. H. Smith*. — The specimens are a foot high, with one leaf on the obtusely angular culm, and only the lowest spike leafy-bracted: sterile spike single: the fruit not fully formed. (Eu.)

150. *C. pùlla*, Good. *Leaves and bracts flat*, with a slender triangular apex, rough-margined; culm rather sharply triangular; perigynia inflated-ovate, mostly longer than the ovate scale. (*C. saxatilis*, L., but that was in part *C. rigida*: the name is appropriate for that but not for this species, and is better discarded altogether.) (Eu.) — Var. ? *MILIARIS*. Culm more slender, 1° or more high; fertile spikes paler (1–3), ovoid or oblong, in our specimens staminate at the apex, each with a very rough leafy bract, the lowest often surpassing the culm. (*C. miliaris*, Michx.) — Outlet of Moosehead Lake, with or near the preceding, July 29, *C. E. & A. H. Smith*; and far northward.

+ + *Perigynia abruptly contracted into a very long and 2-toothed beak.*

151. *C. longiróstris*, Torr. Sterile spikes usually 3, at the summit of a long slender stalk; the lower often bearing some fertile flowers; fertile spikes 2–4, cylindrical, more or less distant, on long filiform at length drooping stalks, loosely flowered; perigynia globose-ovoid, smooth and shining, abruptly contracted into a slender cylindrical beak, which is longer than the body, rough on the margin, and 2-cleft at the membranaceous orifice, a little longer than the lanceolate light-colored or white scale. (*C. Sprengèlii*, Dew.) — Shady rocks, N. New England to Wisconsin, and northward.

ORDER 128. GRAMINEÆ. (GRASS FAMILY.)

Grasses, with usually hollow stems (culms) closed at the joints, alternate 2-ranked leaves, their sheaths split or open on the side opposite the blade; the hypogynous flowers umbricated with 2-ranked glumes or bracts: the outer pair (glumes proper) subtending the spikelet of one or several flowers; the inner pair (paleæ or palets) enclosing each particular flower, which is usually furnished with 2 or 3 minute hypogynous scales (squamulæ, Juss., lodiculæ, Beauv.). Stamens 1–6, commonly 3: anthers versatile, 2-celled, the cells distinct. Styles mostly 2 or 2-parted: stigmas hairy or feathery. Ovary 1-celled, 1-ovuled, forming a seed-like grain (caryopsis) in fruit. Embryo small, on the outside and at the base of the floury albumen. — Roots fibrous. Sheath of the leaves usually more or less extended above the base of the blade into a scarios appendage (ligule). Spikelets panicled or spiked. Inner (upper) palet usually 2-nerved or 2-keeled, enclosed or partly covered by the outer (lower) palet. Grain sometimes free from,

sometimes permanently adherent to, the palea. — A vast and most important family, as it furnishes the cereal grains, and the principal food of cattle, &c. (See Plates 7–14.)

Tribe I. POACEÆ. R. Br. Spikelets 1-many-flowered, when more than one-flowered centripetal in development; the lowest flowers first developing, *uppermost*, if any. imperfect or abortive, the rest all alike in the spikelet (perfect, or occasionally monœcious or dioecious); only in a few exceptional cases with the lowest of the several flowers less perfect than the upper (viz. staminate only in *Arrhenatherum* and *Phragmites*, neutral in *Uniola*, *Ctenium*, &c.).

Subtribe 1. Oryzæ. Spikelets 1-flowered, in panicles, the flowers often monœcious. Glumes abortive or wanting! Inner palea 3-nerved! Stamens 1–6.

1. **Leersia.** Flowers perfect, strongly flattened contrary to the awnless conduplicate palea.
2. **Zizania.** Flowers monœcious. Palea convex; the lower one awned in the fertile flowers.

Subtribe 2. Agrostidæ. Spikelets 1-flowered, perfect, and occasionally with the rudiment or abortive pedicel of a second flower above, paniced, or clustered, sometimes so contracted as to form a sort of spike, but the spikelets are not directly borne on the common axis. Stamens 1–3.

* PHLEOIDEÆ. Glumes equal, strongly keeled, laterally flattened, boat-shaped, somewhat herbaceous, as well as the palea. Inflorescence densely spiked!

3. **Alopecurus.** Glumes united at the base. Lower palea bearing an awn on the back: the upper palea wanting.
4. **Pleurum.** Glumes distinct, sharp-pointed, much larger than the two thin and truncate awnless palea.
5. **Crypsis.** Glumes distinct, not longer than the palea; both awnless and pointless.

** AGROSTIDEÆ proper. Glumes and palea both membranaceous, or the latter sometimes very thin and delicate. Inflorescence paniced or glomerate, sometimes rather spike-like, but not contracted into a uniform cylindrical spike. Palea not surrounded by a tuft of hairs, or only with some very minute ones at the base.

+ Flower perfectly sessile in the glumes: lower palea 1-nerved: awns none.

6. **Vilfa.** Fruit a caryopsis (seed adherent to the pericarp, as in most grasses). Panicea spiked or contracted.
 7. **Sporobolus.** Fruit an utricle (seed loose in the thin pericarp). Panicea open or close.
- + + Flower slightly raised in the glumes on a short sometimes stalk-like base (*callus*): lower palea 3–5-nerved; and this or the glumes awned or pointed, except in some species of No. 8.
8. **Agrostis.** Glumes equal, or the lower one rather longer, pointless, exceeding the very thin blunt palea. Lower palea pointless, often awned on the back; the upper sometimes wanting. Panicea open.
 9. **Polypogon.** Glumes nearly equal, long-awned, much longer than the palea, the lower of which is often short-awned below the apex. Stamens 3. Panicea contracted.
 10. **Cinna.** Glumes acute, the lower about equalling and the upper slightly exceeding the similar palea. Stamen 1. Flowers raised on a distinct naked stalk, beardless: lower palea short-awned or bristle-pointed just below the tip; the upper 1-nerved.
 11. **Muhlenbergia.** Lower glume mostly smaller. Palea chiefly hairy-bearded at the base, the tip of the lower one mucronate-pointed or awned. Stamens 3.
 12. **Brachyelytrum.** Lower glume almost obsolete, and the upper minute. Lower palea long-awned from the tip; the upper grooved on the back and bearing a long and slender naked pedicel of an abortive second flower. Stamens 2.

*** CALAMAGROSTIDEÆ. Flower with a copious tuft of hairs at the base of the palea: otherwise as in the foregoing subdivision.

13. **Calamagrostis.** Lower palea mostly awned on the back, shorter than the glumes.

*** STIPEÆ. Paleæ coriaceous, or indurated in fruit, commonly shorter than the membranaceous glumes, usually on a rigid base or *callus*; the lower involute, terete, closely enclosing the upper and the grain, mostly 1-3-awned at the apex. Inflorescence racemose or paniced: spikelets usually large, the flower deciduous from the persistent glumes.

14. **Oryzopsis.** Awn simple, straight, deciduous from the palet or sometimes wanting.

15. **Stipa.** Awn simple, twisted below. Callus conspicuous, pointed at the base.

16. **Aristida.** Awn triple. Upper palet small. Callus conspicuous, pointed at the base.

**** Palets coriaceous or cartilaginous, awnless. Here the following would be sought by the student who overlooked the pair of rudimentary flowers in No. 56, and was not acquainted with the recondite theoretical structure of No. 57 and 58.

56. **Phalaris.** Spikelets laterally flattened. A rudiment at the base of each palet.

57. **Milium.** Spikelets dorsally flattish, not jointed with the pedicels: flowers all alike.

58. **Amphicarpum.** Spikelets of two sorts, the fertile subterranean, those of the panicle separating by a joint without ripening grain.

Subtribe 3. Chloridæ. Spikelets rarely 1-flowered, usually 2-several-flowered, with one or more of the upper flowers imperfect, disposed in one-sided spikes! Glumes persistent, the upper one looking outward. Rhachis (axis) jointless. Spikes usually several and racemed or digitate. Stamens 2 or 3.

* Spikelets strictly 1-flowered.

59. **Paspalum** might be looked for here, having to all appearance merely 1-flowered spikelets.

17. **Spartina.** Spikelets much flattened contrary to the glumes, imbricated in 2 ranks on the triangular rhachis of the straight spike.

** Spikelets with one perfect and two or more imperfect or neutral flowers:

+ The perfect flower intermediate.

18. **Ctenium.** Spikelets closely imbricated on one side of the axis of a single curved spike.

+ + The perfect flower below the one or more neutral or rudimentary ones.

19. **Bouteloua.** Lower palet 3-cleft and pointed or 3-awned at the apex. Spikes dense.

20. **Gymnopogon.** Lower palet and the rudiment 1-awned. Spikes filiform, racemed.

21. **Cynodon.** Flower and the rudiment awnless. Spikes slender, digitate.

*** Spikelets several-flowered; more than one of the lower flowers perfect and fertile.

+ Spikes digitate at the summit of the culm, dense.

22. **Dactyloctenium.** Glumes compressed-keeled; outer one awned: lower palet pointed.

23. **Eleusine.** Glumes and palets both awnless and blunt.

+ + Spikes racemed, slender.

24. **Leptochloa.** Spikelets loosely spiked. Lower palet pointless or awned at the tip.

Subtribe 4. Festucineæ. Spikelets several- (few-many-) flowered, paniced; the uppermost flower often imperfect or abortive. Palets pointless, or the lower sometimes tipped with a straight (not twisted nor deeply dorsal) awn or bristle. Stamens 1-3. Culms sometimes reed-like, but not woody.

* Lowest flowers of the spikelet perfect and fertile.

+ Grain free from the palets.

+ + Joints of the rhachis of the spikelet at the insertion of each flower, or the whole rhachis bearded. Glumes and convex palets membranaceous.

25. **Tricuspis.** Spikelets 3-many-flowered. Lower paleæ hairy-fringed on the 3 nerves, one or all of which project into awns or mucronate tips, mostly from notches or clefts.

26. **Graphophorum.** Spikelets 2-few-flowered. Glumes and palets awnless or pointless.

+ + Rhachis of the spikelet and base of the flower not bearded.

a. Lower palet 1-pointed, awned or acute, the nerves when present running into the point.

27. **Diarrhena.** Glumes (short) and the rigid-pointed lower 3-nerved palet coriaceous, convex-boat-shaped. Stamens 2. Pericarp cartilaginous, large. Panicle loosely few-flowered.

28. **Dactylis.** Glumes (rather long) and lower palet awn-pointed, herbaceous, compressed-keeled. Panicle contracted in one-sided clusters.
29. **Koeleria.** Glumes (nearly as long as the spikelet) and lower palet membranaceous, keeled, acute or mucronate, or rather blunt. Panicle contracted, spike-like.
37. **Festuca,** with grain sometimes free, may be looked for here.

b. Lower palet awnless and pointless, blunt (except one *Glyceria*), the nerves parallel.

1. Glumes extremely dissimilar, $1\frac{1}{2}$ -3-flowered.

30. **Eatonia.** Lower glume linear; the upper broadly obovate and folded round the flowers.
2. Glumes similar in shape, but often unequal in size.
31. **Melica.** Lower palet flattish-convex, many-nerved, membranaceous at the top, hardening on the loose grain. Fertile flowers 1-3, the upper enwrapping deformed sterile ones.
32. **Glyceria.** Lower palet convex or rounded on the back, 5-7-nerved, scarious at the tip. Spikelets many-flowered; the flowers deciduous at maturity by the breaking up of the rhachis into joints.
33. **Brizopyrum.** Lower palet laterally much compressed and often keeled, acute, rigid, rather coriaceous, smooth, faintly many-nerved. Spikelets flat, spiked-clustered.
37. **Festuca.** Lower palet slightly if at all laterally compressed, not keeled, only 1-5-nerved, not cobwebby, and nearly without scarious margins, acutish: otherwise as in *Poa*.
34. **Poa.** Lower palet laterally compressed and mostly keeled, 5-nerved, membranaceous, scarious-margined, the margins or nerves below often cobwebby or pubescent: the upper palet not remaining after the lower falls. Spikelets flattened.
35. **Eragrostis.** Lower palet 3-nerved, keeled, deciduous, leaving the upper persistent on the rhachis. Spikelets flat.

+ + Grain adherent to the upper palet.

36. **Briza.** Lower palea rounded, very obtuse, pointless, many-nerved, flattened parallel to the glumes, soon ventricose, scarious-margined. Spikelets somewhat heart-shaped.
37. **Festuca.** Lower palet convex on the back, acute, pointed, or awned at the tip, few-nerved. Spikelets terete or flattish. Styles terminal.
38. **Bromus.** Lower palet convex or keeled on the back, mostly awned or bristle-bearing below the 2-cleft tip, 5-9-nerved. Styles scarcely terminal.

* Lowest flower of the spikelet neutral or staminate.

39. **Uniola.** Spikelets very flat; the one or more lowest flowers neutral, of a single empty palet. Flowers strongly compressed-keeled, crowded, coriaceous.
40. **Phragmites.** Spikelets strongly silky-bearded on the rhachis, loosely-flowered, the lowest flower staminate or neutral. Palets membranaceous.

Subtribe 5. Bambusæ. Culms woody and commonly arborescent. Spikelets and flowers nearly as in the preceding Subtribe, awnless.

41. **Arundinaria.** Spikelets flattened, loosely 5-14-flowered, in depauperate panicles.

Subtribe 6. Hordeinæ. Spikelets 1-several-flowered, sessile on opposite sides of a zigzag jointed rhachis (which is excavated or channelled on one side of each joint), forming a spike. Glumes sometimes abortive or wanting. Otherwise as in Subtribe 4.

* Spikelets single at each joint of the rhachis.

42. **Lepturus.** Spikelets almost immersed in the excavations of the very slender rhachis, 1-flowered. The filiform spikes usually several.
43. **Lolium.** Spikelets many-flowered, placed edgewise on the rhachis of the solitary spike: glume only one, external.
44. **Triticum.** Spikelets 3-several flowered, placed flatwise on the rhachis of the solitary spike: both glumes present, transverse (right and left).

* * Spikelets 2 or more at each joint of the rhachis: spike solitary.

+ Glumes anterior, forming a sort of involucre for the cluster of spikelets.

45. **Hordeum.** Spikelets 1-flowered, 3 at each joint, but the two lateral usually sterile.
46. **Elymus.** Spikelets 1-several-flowered, all perfect and similar.

+ + Glumes none or 1-2 awn-like rudiments in their place.

47. **Gymnostichum.** Spikelets few-flowered, somewhat pedicelled, 1-3 at each joint.

Subtribe 7. Avenæ. Spikelets 2-several-flowered, panicle; the rachis or base of the flowers often villous-bearded. Glumes mostly equalling or exceeding the flowers. Lower palet bearing a twisted, bent, or straight awn on its back or below its apex; the upper palet 2-nerved. Stamens 3.

* Flowers all perfect, or the uppermost rudimentary.

48. **Danthonia.** Spikelets several-flowered. Lower palet firm and rigid, rounded on the back, several-nerved, the 3 middle nerves running into the flattish twisted awn which proceeds from the cleft at the apex.

49. **Avena.** Spikelets 2-several-flowered. Lower palet roundish on the back, and of firmer texture than the glumes, several-nerved, sharply 2-toothed or 2-cleft at the tip, the bent or twisted awn rising only from the midnerve at or below the cleft.

50. **Trisetum.** Spikelets 2-several-flowered. Lower palet laterally compressed and keeled, sharply 2-toothed or 2-pointed at the apex, the slender awn rising at or near the cleft, from the midnerve only.

51. **Aira.** Spikelets small, 2-flowered, with or without the rudiment of a third flower. Palets thin or scarios, the lower awned from towards the base.

** One of the flowers staminate only.

52. **Arrhenatherum.** Lower flower staminate; the perfect one commonly awnless; the uppermost a rudiment: otherwise as in *Avena*.

53. **Holcus.** Lower flower perfect, awnless; the upper staminate and awned: rudiment none: otherwise resembling *Aira*.

Tribe II. PHALARIDÆ, Trin. (not of Kunth). Spikelets 3-flowered; the uppermost or middle (terminal) flower perfect: the two lower (one on each side) imperfect, either staminate, neutral, or reduced to an inconspicuous rudiment.

Subtribe 1. Anthoxanthæ. Lateral flowers mostly awned, staminate or neutral, of 1 or 2 palets; the upper one awnless and diandrous. Upper palet 1-nerved.

54. **Hierochloa.** Lateral flowers staminate and triandrous, of 2 palets.

55. **Anthoxanthum.** Lateral flowers neutral, each of a single awned and hairy palet.

Subtribe 2. Phalaridæ proper. Lateral flowers reduced to a small neutral rudiment or abortive pedicel on each side of the fertile one; which is awnless and triandrous.

56. **Phalaris.** Glumes boat-shaped, keeled, enclosing the coriaceous fertile flower.

Tribe III. PANICEÆ. Spikelets 2-flowered; the lower flower always imperfect, either staminate or neutral; in the latter case usually reduced to a single empty palet (placed next the lower glume, if that be present); the upper (terminal) flower (placed next the upper or inner glume) only fertile. Embryo and groove (when present) on the outer side of the grain! (next the lower palet of the fertile flower). (Flowers polygamous, or hemigamous (when the lower flower is neutral), or sometimes seemingly simple and perfect, from the suppression both of the lower glume and of the upper palet of the neutral flower, sometimes monocious, or rarely dioecious. Rarely both glumes are wanting.)

Subtribe 1. Paspalæ, Griseb. Glumes and sterile palets herbaceous or membranaceous: palets of the fertile flower of firmer texture, coriaceous or chartaceous, awnless, not keeled, more or less flattened parallel with the glumes.

* Spikelets appearing as if simply 1-flowered, from the suppression of the lower glume, the single neutral palet of the sterile flower apparently occupying its place. (Awnless.)

57. **Milium.** Spikelets not jointed with their pedicels, all alike in a terminal open panicle.

58. **Amphicarpum.** Spikelets jointed with their pedicels, of 2 sorts; one in a terminal panicle; the other subterranean, on radical peduncles.

59. **Paspalum.** Spikelets jointed with their short pedicels, all alike, plano-convex, in one-sided spikes or spiked racemes.

* * Spikelets manifestly $1\frac{1}{2}$ -2-flowered (polygamous, the lower flower staminate or often neutral, of one or both palets), the lower glume being present.

60. **Panicum.** Spikelets not involucrate, nor the peduncles bristle-bearing. Lower glume usually small or minute. Sterile flower either staminate or neutral.

61. **Setaria.** Spikelets spiked or dense-panicled, the peduncles continued into naked solitary bristles or awns: otherwise as in *Panicum*.

62. **Cenchrus.** Spikelets enclosed 1-5 together in a hard and spiny or bristly and globular bur-like involucre.

Subtribe 2. Saccharææ. Fertile palets membranaceous or scarious, always of thinner and more delicate texture than the (often indurated) glumes, frequently awned from the tip. Spikelets usually in pairs or threes, panicled or spiked, some of them entirely sterile or rudimentary.

* Spikelets monoecious, imbedded in the separable joints of the spike.

63. **Tripsacum.** Staminate spikelets above, in pairs at each joint: pistillate spikelets single in each joint: glumes indurated.

* * Fertile spikelets with one perfect and one sterile (staminate or mostly neutral) flower: lower palet of the perfect flower awned.

64. **Erianthus.** Both spikelets at each joint of the rachis alike fertile, and involucrate with a silky tuft: otherwise as in the next.

65. **Andropogon.** Spikelets a pair at each joint of the plumose-hairy spikes, one of them sessile and fertile; the other pedicelled and sterile or rudimentary.

66. **Sorghum.** Spikelets in open panicles, 2-3 together, the lateral ones sterile or sometimes reduced to mere pedicels.

1. LEËRSIA, Solander. WHITE GRASS. (Pl. 7.)

Flowers crowded in one-sided panicled spikes or racemes, perfect, but those in the open panicles usually sterile by the abortion of the ovary, those enclosed in the sheaths of the leaves close-fertilized in the bud and prolific. Spikelets 1-flowered, flat, more or less imbricated over each other, jointed with the short pedicels. Glumes wanting. Palets chartaceous, strongly flattened laterally or conduplicate, awnless, bristly-ciliate on the keels, closed, nearly equal in length, but the lower much broader, enclosing the flat grain. Stamens 1-6. Stigmas feathery, the hairs branching.—Perennial marsh grasses: the flat leaves, sheaths, &c., rough upwards, being clothed with very minute hooked prickles. (Named after *John Daniel Leers*, a German botanist.)

* *Spikelets narrowly oblong, rather loosely crowded.*

1. **L. Virginica**, Willd. (WHITE GRASS.) *Panicle simple; the spikelets closely appressed on the slender branches, around which they are partly curved ($1\frac{1}{2}$ " long); stamens 2 (a third imperfect or wanting); palets sparingly ciliate (greenish-white).—Wet woods. Aug., Sept.*

2. **L. oryzoides**, Swartz. (RICE CUT-GRASS.) *Panicle diffusely branched; spikelets flat, rather spreading ($2\frac{1}{2}$ "-3" long); stamens 3; palets strongly bristly-ciliate (whitish).—Very wet places: common. Aug. (Eu.)*

* * *Spikelets broadly oval, imbricately covering each other ($2\frac{1}{2}$ "-3" long).*

3. **L. lenticularis**, Michx. (FLY-CATCH GRASS.) *Smoothish; panicle simple; palets very flat, strongly bristly ciliate (said to close and catch flies); stamens 2: otherwise like the preceding.—Low grounds, Virginia, Illinois, and southward.*

2. ZIZÀNIA, Gronov. WATER OR INDIAN RICE. (Pl. 7.)

Flowers monœcious; the staminate and pistillate both in 1-flowered spikelets in the same panicle. Glumes wanting, or rudimentary and forming a little cup. Palets herbaceo-membranaceous, convex, awnless in the sterile, the lower one tipped with a straight awn in the fertile spikelets. Stamens 6. Stigmas pencil-form. — Large, often reed-like water-grasses. Spikelets jointed with the club-shaped pedicels, very deciduous. (Adopted from Ζιζάνιον, the ancient name of some wild grain.)

1. **Z. aquática**, L. (INDIAN RICE. WATER OATS.) Annual; *lower branches of the ample pyramidal panicle staminate, spreading; the upper erect, pistillate*; pedicels strongly club-shaped; *lower palet long-awned*, rough; styles distinct; grain linear, slender. (*Z. clavulosa*, Michx.) — Swampy borders of streams and in shallow water: common, especially northwestward. Aug. — Culms 3°–9° high. Leaves flat, 2°–3° long, linear-lanceolate. Grain 6" long; largely gathered for food by the Northwestern Indians.

2. **Z. miliacea**, Michx. Perennial; panicle diffuse, ample, the *staminate and pistillate flowers intermixed*; *awns short*; styles united; grain ovate. Penn.? Ohio, and southward. Aug. — Leaves involute.

3. ALOPECÛRUS, L. FOXTAIL GRASS. (Pl. 7.)

Spikelets 1-flowered. Glumes boat-shaped, strongly compressed and keeled, nearly equal, united at the base, equalling or exceeding the lower palet, which is awned on the back below the middle: upper palet wanting! Stamens 3. Styles mostly united. Stigmas long and feathered. — Clusters contracted into a cylindrical and soft dense spike. Root perennial. (Name from ἀλώπηξ, fox, and οὐρά, tail, the popular appellation, from the shape of the spike.)

1. **A. pratensis**, L. (MEADOW FOXTAIL.) Culm upright, smooth (2° high); *palet equalling the acute glumes*; *awn exerted more than half its length, twisted*; the upper leaf much shorter than its inflated sheath. — Meadows and pastures, eastward. May. (Nat. from Eu.)

2. **A. geniculatus**, L. (FLOATING F.) Culm ascending, bent at the lower joints; *palet rather shorter than the obtuse glumes, the awn from near its base and projecting half its length beyond it*; anthers linear; the upper leaf as long as its sheath. — Moist meadows, eastward. June–Aug. (Nat. from Eu.)

3. **A. aristulatus**, Michx. (WILD F.) *Glaucous*; culm decumbent below, at length bent and ascending; *palet rather longer than the obtuse glumes, scarcely exceeded by the awn which rises from just below its middle*; anthers oblong. (*A. subaristatus*, Pers.) — In water and wet places: common. June–Aug. Spike more slender and paler than in the last. (Eu.)

4. PHLÈUM, L. CAT'S-TAIL GRASS. (Pl. 7.)

Palets both present, shorter than the mucronate or awned glumes; the lower one truncate, usually awnless. Styles distinct. Otherwise much as in Alopecurus. — Perennials. Spike very dense, harsh. (An ancient Greek name.)

1. **P. pratense**, L. (TIMOTHY. HERD'S-GRASS in New England and New York.) Tall; *spike cylindrical, elongated*; glumes ciliate on the back,

tipped with a *short bristle*.—Meadows, commonly cultivated for hay. (Nat. from Eu.)

2. **P. alpinum**, L. Low; *spike ovate-oblong*; glumes strongly ciliate on the back, tipped with a *rough awn about their own length*.—Alpine tops of the White Mountains, New Hampshire, and high northward. (Eu.)

5. CRÝPSIS, Ait. CRYPSIS. (Pl. 7.)

Spikelets 1-flowered, in clusters which are crowded in a dense head or short spike bracted by the uppermost leaves. Glumes, palets, &c. as in the next genus, or rather thinner.—Low and spreading tufted annuals, natives of the East; with short leaves, the sheaths of the upper spathaceous. (Name, *κρύψις*, *concealment*, the spikes at first partly hidden by the subtending sheaths.)

1. **C. schœnoides**, Lam. Leaves rather rigid, tapering to a sharp point; heads or spikes oblong, 7"–20" long, thick. (*C. Virginica*, *Nutt.*, excl. syn.)—Waste places, streets of Philadelphia and vicinity, also of Wilmington, Delaware: becoming very common. (Nat. from Eu.)

6. VÍLFA, Adans., Beauv. RUSH-GRASS. (Pl. 7.)

Spikelets 1-flowered, in a contracted or spiked panicle. Glumes 1-nerved or nerveless, not awned or pointed, the lower smaller. Flower nearly sessile in the glumes. Palets 2, much alike, of the same texture as the glumes (membranaceo-chartaceous) and usually longer than they, naked, awnless and mostly pointless; the lower 1-nerved (rarely somewhat 3-nerved). Stamens chiefly 3. Stigmas simply feathery. Grain (caryopsis) oblong or cylindrical, deciduous.—Culms wiry or rigid. Leaves involute, usually bearded at the throat; their sheaths often enclosing the panicles. (Name unexplained.)

1. **V. áspera**, Beauv. *Root perennial*; culms tufted (2°–4° high); lowest leaves very long, rigid, rough on the edges, tapering to a long involute and thread-like point; the upper short, involute; sheaths partly or at first wholly enclosing the contracted panicle; *palets much longer than the unequal glumes*; grain oval or oblong. (*Agrostis áspera*, *Michx.* *A. clandestina* & *A. involuta*, *Muhl.* *A. longifolia*, *Torr.*)—Sandy fields and dry hills, especially southward. Sept.—Spikelets 2"–3" long. Palets rough above, smooth or hairy below, of greatly varying proportions; the upper one tapering upwards, acute, and one half to twice longer than the lower, or else obtuse and equalled or even considerably exceeded by the lower!

2. **V. vaginæflora**, Torr. *Root annual*; culms slender (6'–12' high), ascending; leaves involute-awl-shaped (1'–4' long); panicles simple and spiked, the lateral and often the terminal concealed in the sheaths; *palets somewhat equal, acute, about the length of the nearly equal glumes*: only one third longer than the linear grain. (*Agrostis Virginica*, *Muhl.*, not of *L.*)—Barren and sandy dry fields: common, especially southward. Sept.

3. **V. cuspidata**, Torr. *Root perennial*; culms and leaves more slender than in the preceding; panicle exserted, very simple and narrow; spikelets smaller, the *glumes very acute, and the lower palet cuspidate*.—Borders of Maine (on the St. John's River, *G. L. Goodale*), and northwestward.

4. **V. Virginica**, Beauv. *Root perennial; culms tufted, slender (5'-12' long), often procumbent, branched; leaves convolute, rigid; palets rather shorter than the nearly equal acute glumes. (Agrostis Virginica, L.) — Sandy seashore, Virginia (Clayton) and southward. — Spikelets much smaller and more numerous than in the others.*

7. SPORÓBOLUS, R. Br. DROP-SEED GRASS. (Pl. 7.)

Spikelets 1- (rarely 2-) flowered, in a contracted or open panicle. Flowers nearly as in Vilfa; the palets longer than the unequal glumes. Stamens 2-3. Grain a globular utricle (hyaline or rarely coriaceous), containing a loose seed, deciduous (whence the name, from *σπορά*, *seed*, and *βάλλω*, *to cast forth*).

* *Glumes very unequal: panicle pyramidal, open: ours perennials, except No. 3.*

1. **S. júnceus**, Kunth. *Leaves involute, narrow, rigid, the lowest elongated; culm (1°-2° high) naked above, bearing a narrow loose panicle; glumes ovate, rather obtuse, the lower one half as long as, the upper equalling, the nearly equal palets. (Agrostis juncea, Michx. Vilfa juncea, Trin.) — Dry soil, Pennsylvania to Wisconsin, and (chiefly) southward. Aug. — Spikelets 1''-2'' long, shining.*

2. **S. heterólepis**, Gray. *Leaves involute-thread-form, rigid, the lowest as long as the culm (1°-2°) which is naked above; panicle very loose; glumes very unequal; the lower awl-shaped (or bristle-pointed from a broad base) and somewhat shorter, the upper ovate-oblong and taper-pointed and longer, than the equal palets. (Vilfa heterolepis, Gray.) — Dry soil, Connecticut, and New York to Illinois and Wisconsin. Aug. — Plant exhaling an unpleasant scent (Sullicant), stouter than the last, the spikelets thrice larger. Utricle spherical (1'' in diameter), shining, thick and coriaceous!*

3. **C. cryptándrus**, Gray. *Leaves flat, pale (2'' wide); the pyramidal panicle bursting from the upper sheath which usually encloses its base, its spreading branches hairy in the axils; upper glume lanceolate, rather acute, twice the length of the lower one, as long as the nearly equal palets; sheaths strongly bearded at the throat; root annual? (Agr. & Vilfa cryptandra, Torr.) — Sandy shores, coast of New England, and of the Great Lakes. Aug., Sept. — Culm 2°-3° high. Panicle lead-color: spikelets 1'' long.*

* * *Glumes almost equal, shorter than the broad palets: panicle racemose-elongated, open, the pedicels capillary: sheaths naked at the throat: spikelets not unfrequently two-flowered: root perennial. (COLPODIUM?)*

4. **S. comprésus**, Kunth. *Very smooth, leafy to the top: culms tufted, stout, very flat: sheaths flattened, much longer than the internodes; leaves erect, narrow, conduplicate-channelled; glumes acutish, about one third shorter than the obtuse palets. (Agrostis compressa, Torr. Vilfa, Trin.) — Bogs in the pine-barrens of New Jersey. Sept. — Forming strong tussocks, 1°-2° high. Panicle 8'-12' long: spikelets 1'' long, purplish.*

5. **S. serótinus**, Gray. *Smooth; culms very slender, flattish (8'-15' high), few-leaved; leaves very slender, channelled; panicle soon much exerted, the diffuse capillary branches scattered; glumes ovate, obtuse, about half the length of the*

palets. (Agr. & *Vilfa scrotina*, Torr. *V. ténera*, Trin. *Poa?* *uniflora*, Muhl. *P. modésta*, Tuckerm.) — Sandy wet places, Maine to New Jersey and Michigan. Sept. — A very delicate grass; the spikelets half a line long.

8. AGRÓSTIS, L. BENT-GRASS. (Pl. 7.)

Spikelets 1-flowered, in an open panicle. Glumes somewhat equal, or the lower rather longer, usually longer than the palets, pointless. Palets very thin, pointless, naked; the lower 3-5-nerved, frequently awned on the back; the upper often minute or none. Stamens chiefly 3. Grain (caryopsis) free. — Culms usually tufted, slender; root commonly perennial. (Name from *ἀγρός*, a field, the place of growth.)

§ 1. TRICHÓDIUM, Michx. — *Upper palet abortive, minute, or none.*

1. **A. elàta**, Trin. *Culms firm or stout* (2° - 3° high); leaves flat ($1''$ - $2''$ wide); upper ligules elongated ($2''$ - $3''$ long); *spikelets crowded on the branches of the spreading panicle above the middle* ($1\frac{1}{2}''$ long); lower palet awnless, slightly shorter than the rather unequal glumes; the upper wanting. (A. Schweinitzii, Trin.? A. altissima, Tuckerm., excl. var. laxa. Trich. elatum, Pursh.) — Swamps, New Jersey and southward. October.

2. **A. perénnans**, Tuckerm. (THIN-GRASS.) *Culms slender, erect from a decumbent base* (1° - 2° high); leaves flat (the upper $4'$ - $6'$ long, $1''$ - $2''$ wide); *panicle at length diffusely spreading, pale green; the branches short, divided and flower-bearing from or below the middle; lower palet awnless* (rarely short-awned), shorter than the unequal glumes; the upper minute or obsolete. (Cornucopiæ perennans, Walt. Trich. perennans, Ell. T. decumbens, Michx. T. scabrum, Muhl. Agr. anómala, Willd.) — Damp shaded places. July, Aug. — Spikelets, &c. as in No. 3, into which it seems to vary.

3. **A. scàbra**, Willd. (HAIR-GRASS.) *Culms very slender, erect* (1° - 2° high); leaves short and narrow, the lower soon involute (the upper $1'$ - $3'$ long, less than $1''$ wide); *panicle very loose and divergent, purplish, the long capillary branches flower-bearing at and near the apex; lower palet awnless or occasionally short-awned* on the back, shorter than the rather unequal very acute glumes; the upper minute or obsolete; root biennial? (A. laxiflora, Richard. A. Michauxii, Trin. partly. Trich. laxiflorum, Michx. T. montanum, Torr.) — Exsiccated places: common. June-Aug. — Remarkable for the long and divergent capillary branches of the extremely loose panicle; these are whorled, rough with very minute bristles (under a lens), as also the keel of the glumes. Spikelets $1'$ long. — A variety? from about the White Mountains, &c. (var. montana, Tuckerm.), has a more or less exerted awn, thus differing from the T. montanum, Torr. (A. oreóphila, Trin.), which is a dwarfed form, growing in tufts in hollows of rocks, &c.

4. **A. canina**, L. (BROWN BENT-GRASS.) *Culms $8'$ - 2° high; root-leaves involute-bristle-form, those of the culm flat and broader; panicle loose; glumes slightly unequal, ovate-lanceolate, very acute; palet exertly awned on the back at or below the middle; spikelets brownish or purplish, rarely pale or greenish* ($1''$ - $1\frac{1}{2}''$ long). — Meadows, sparingly naturalized eastward. A mountain form with shorter and more spreading panicle (A. Pickeringii

& *A. concinna*, *Tuckerm.*, *A. canina*, var. *alpina*, *Oakes*, & Ed. 2., and essentially *A. rubra*, *L. ex Wahl.*, and *A. borealis*, *Hartm.*), is indigenous on mountain-tops, Maine to New York; also, an ampler form in the Alleghanies of Pennsylvania (*J. R. Lowrie*), and southward (*A. rupéstris*, *Chapman*, &c.). July - Aug. (Eu.)

§ 2. AGROSTIS proper. *Upper palet manifest, but shorter than the lower.*

5. *A. vulgáris*, With. (RED-TOP. HERD'S-GRASS of Penn., &c.) Rootstocks creeping; culm mostly upright (1° - 2° high); *panicle oblong*, with spreading slightly rough short branches (*purple*); leaves linear, flat; *ligule very short*, truncate; lower palet nearly equalling the glumes, chiefly awnless, 3-nerved; the upper about one half its length. (*A. polymórpha*, *Huds.*, partly. — Varies with a rougher panicle (*A. hispida*, *Willd.*), and rarely with the flower short-awned. — Low meadows; naturalized from Eu., and apparently also native northward. (Eu.)

6. *A. álba*, L. (FIORIN or WHITE BENT-GRASS.) Rootstocks more stoloniferous, and culms often decumbent at the base, ascending; *ligule elongated*, oblong or linear; *panicle* contracted after flowering, either greenish, purplish or brownish; otherwise as in the preceding, and equally variable; rarely with the lower palet short-awned, or even slender-awned below the tip. (*A. stolonífera*, *L.*, partly.) — Meadows and fields, a valuable grass: nat. from Eu.: also indigenous on river-banks, N. New York and northward. (Eu.)

9. POLYPOGON, Desf. BEARD-GRASS. (Pl. 8.)

Spikelets 1-flowered, in a contracted, mostly spike-like panicle. Glumes nearly equal, long-awned, much longer than the membranaceous palets, the lower of which is commonly short-awned below the apex. Stamens 3. Grain free. (Name composed of *πολύ*, *much*, and *πῶγων*, *beard*; from the awns.)

1. *P. MONSPELIÉNSIS*, Desf. Panicle interrupted; glumes oblong, the awn from a notch at the summit; lower palet awned; root annual. Hampton Beach, New Hampshire (*Robbins*), Virginia? and southward. (Nat. from Eu.)

10. CÍNNA, L. WOOD REED-GRASS. (Pl. 8.)

Spikelets 1-flowered, much flattened, crowded in an open flaccid panicle. Glumes lanceolate, acute, strongly keeled, rough-serrulate on the keel; the lower rather smaller, the upper a little exceeding the palets. Flower manifestly stalked in the glumes, smooth and naked; the palets much like the glumes; the lower longer than the upper, short-awned or mucronate on the back below the pointless apex. Stamen one, opposite the 1-nerved upper palet! Grain linear-oblong, free. — A perennial, rather sweet-scented grass, with simple and upright somewhat reed-like culms (2° - 7° high), bearing an ample compound terminal panicle, its branches in fours or fives; the broadly linear-lanceolate flat leaves (4" - 6" wide) with conspicuous ligules. Spikelets green, often purplish-tinged. (Name unexplained.)

1. *C. arundinácea*, L. — Moist woods and shaded swamps: rather common. July, Aug. — Panicle 6' - 15' long, rather dense; the branches and pedi-

cels spreading in flower, afterwards erect. Spikelets $2\frac{1}{2}''-3''$ long. Awn of the palet either obsolete or manifest.

Var. **péndula**, Gray. Panicle loose and more slender, the branches nearly capillary and drooping in flower; pedicels very rough; glumes and palets thinner, the former less unequal; spikelets $1\frac{1}{2}''-2''$ long; upper palet obtuse. (*C. pendula*, Trin. *C. latifolia*, Griseb. *C. expansa*, Link. *Blyttia suavèolens*, Fries.)—Deep damp woods, N. New England to Lake Superior and northward, and on mountains southward.—A slender variety of the last, as is shown by intermediate specimens, always monandrous. (Eu.)

11. MUHLENBERGIA, Schreber. DRO. SEED G. (Pl. 8.)

Spikelets 1-flowered, in contracted or rarely in open panicles. Glumes mostly acute or bristle-pointed, persistent; the lower rather smaller or minute. Flower very short-stalked or sessile in the glumes; the palets usually minutely bearded at the base, herbaceous, deciduous with the enclosed grain, often equal; the lower 3-nerved, mucronate or awned at the apex. Stamens 3. (Dedicated to the Rev. Dr. Henry Muhlenberg, a distinguished American botanist of the early part of this century.)

§ 1. MUHLENBERGIA proper. *Panicles contracted or glomerate, terminal and axillary: perennials (in our species) with branching rigid culms, from scaly creeping rootstocks: leaves short and narrow.*

* *Lower palet barely mucronate or sharp-pointed.* (Sp. of *Cinna*, Kunth, Trin.)

1. **M. sobolifera**, Trin. Culms ascending ($1^{\circ}-2^{\circ}$ high), rarely branching; the simple contracted panicle very slender or filiform; glumes barely pointed, almost equal, one third shorter than the equal palets; lower palet abruptly short-mucronate. (*Agrostis sobolifera*, Muhl.)—Open rocky woods, Mass. to Michigan, Illinois, and southward. Aug.—Spikelets less than $1''$ long.

2. **M. glomerata**, Trin. Culms upright ($1^{\circ}-3^{\circ}$ high), sparingly branched or simple; panicle oblong-linear, contracted into an interrupted glomerate spike, long-peduncled, the branches sessile; glumes awned, nearly equal, and (with the bristle-like awn) about twice the length of the unequal very acute palets. (*Agr. racemosa*, Michx. *A. setosa*, Muhl. *Polypogon racemosus*, Nutt.)—Bogs: common, especially northward. Aug.—Panicle $2'-3'$ long.

3. **M. Mexicana**, Trin. Culms ascending, much branched ($2^{\circ}-3^{\circ}$ high); panicles lateral and terminal, often included at the base, contracted, the branches densely spiked-clustered, linear (green and purplish); glumes awnless, sharp-pointed, unequal, the upper about the length of the very acute lower palet. (*Agr. Mexicana*, L. *A. lateriflora*, Michx.)—Varies with more slender panicles (*A. filiformis*, Muhl.)—Low grounds: common. Aug.

* * *Lower palet bristle-awned from the tip: flowers short-pedicelled.*

4. **M. sylvatica**, Torr. & Gr. Culms ascending, much branched and diffusely spreading ($2^{\circ}-4^{\circ}$ long); contracted panicles densely many-flowered; glumes almost equal, bristle-pointed, nearly as long as the lower palet, which bears an awn twice or thrice the length of the spikelet. (*Agr. diffusa*, Muhl.)—Low or rocky woods: common. Aug., Sept.—In aspect between No. 3 and No. 5.

5. **M. Willdenovii**, Trin. Culms upright (3° high), slender, simple or sparingly branched; contracted *panicle slender, loosely flowered; glumes slightly unequal, short-pointed, half the length of the lower palet*, which bears an awn 3-4 times the length of the spikelet. (*Agrostis tenuiflora*, Willd.)—Rocky woods: rather common. Aug.

6. **M. diffusa**, Schreber. (DROP-SEED. NIMBLE WILL.) Culms diffusely much branched (8'-18' high); contracted *panicles slender*, rather loosely many-flowered, terminal and lateral; *glumes extremely minute, the lower obsolete*, the upper truncate; awn once or twice longer than the palet. (*Dilepÿrum minutiflorum*, Michx.)—Dry hills and woods, from S. New England to Michigan, Illinois, and southward. Aug., Sept.—Spikelets only 1" long.

§ 2. **TRICHÓCHLOA**, DC. *Panicle very loose and open, the long branches and pedicels capillary: leaves narrow, often convolute-bristle-form.*

7. **M. capillaris**, Kunth. (HAIR-GRASS.) Culm simple, upright (2° high) from a fibrous root; *panicle capillary, expanding (6'-20' long, purple)*; glumes unequal, one third or half the length of the long-awned palets, the lower mostly pointless, the upper more or less bristle-pointed.—Sandy soil, W. New England to New Jersey, Kentucky, and southward. Sept.—Pedicels 1'-2' long, scarcely thicker than the awns, which are about 1' long.

12. BRACHYÉLYTRUM, Beauv. (Pl. 8.)

Spikelets 1-flowered, with a conspicuous filiform pedicel of an abortive second flower about half its length, nearly terete, few, in a simple appressed racemed panicle. Glumes unequal, persistent, usually minute, or the lower one almost obsolete. Palets chartaceo-herbaceous, involute, enclosing the linear-oblong grain, somewhat equal, rough with scattered short bristles; the lower 5-nerved, extended into a long straight awn; the upper 2-pointed; the awn-like sterile pedicel partly lodged in the groove on its back. Stamens 2: anthers and stigmas very long.—Perennial, with simple culms (1°-3° high) from creeping rootstocks, downy sheaths, broad and flat lanceolate pointed leaves, and spikelets $\frac{1}{2}$ ' long without the awn. (Name composed of *βραχύς*, *short*, and *ἔλυτρον*, *husk*, from the minute glumes.)

1. **B. aristatum**, Beauv. (*Muhlenbergia erecta*, Schreb. *Dilepÿrum aristosum*, Michx.)—Rocky woods: common. June.—Var. **ENGELMANNI**, is a Western form, with the upper glume awn-pointed, nearly half the length of the palet.

13. CALAMAGRÓSTIS, Adans. REED BENT-G. (Pl. 8.)

Spikelets 1-flowered, and often with a pedicel or rudiment of a second abortive flower (rarely 2-flowered), in an open or spiked panicle. Glumes keeled or boat-shaped, often acute, commonly nearly equal, and exceeding the flower, which bears at the base copious white bristly hairs. Palets thin; the lower bearing a slender awn on the back or below the tip, or sometimes awnless; the upper mostly shorter. Stamens 3. Grain free.—Perennials, with running rootstocks, and mostly tall and simple rigid culms. (Name compounded of *κάλamos*, *a reed*, and *ἀγρόστις*, *a grass*.)

§ 1. **DEYEÚXIA**, Kunth. Rudiment of a second flower present in the form of a plumose or hairy small pedicel behind the upper palet (very rarely more developed and having palets or even stamens): glumes and palets membranaceous, or the latter thin and delicate as in *Agrostis*; the lower 3-5-nerved and awn-bearing.

* Panicle loose and open, even after flowering: the mostly purple-tinged or lead-colored strigose-scarbrous glumes not closing in fruit: copious hairs surrounding the flower about equalling the hyaline lower palet, not surpassed by those of the rudiment: awn delicate, straight.

1. **C. Canadensis**, Beauv. (BLUE JOINT-GRASS.) Culm tall (3° - 5° high); leaves flat when fresh, glaucous; panicle oblong; glumes ovate-lanceolate, acute, $1\frac{1}{4}''$ - $1\frac{1}{2}''$ long; awn from near the middle of the palet, not exceeding and scarcely stouter than the hairs around the flower. (*Arundo Canadensis*, Michx. C. Mexicana, Nutt.)—Wet grounds: common northward. July.

2. **C. Langsdórrffi**, Trin. Spikelets larger, $2\frac{1}{2}''$ - $3''$ long; glumes lanceolate or oblong-lanceolate and gradually taper-pointed; awn stouter: otherwise like the preceding. — White Mountains, New Hampshire, and northward. (Eu.)

* * Panicle strict, its short branches appressed or erect after flowering, and the glumes mostly closed: lower palet less delicate, roughish, sometimes of as firm texture as the glumes: awn stouter.

+ Leaves narrow, inclined to be involute: awn straight.

3. **C. stricta**, Trin. Panicle glomerate and lobed, strict; glumes $1\frac{1}{2}''$ - $2''$ long, ovate-oblong, not acuminate; hairs scarcely or little shorter than the flower, and as long as those of the rudiment; awn from the middle of the thin palea or lower, and barely exceeding it. — Ledges at Willoughby Lake, Vermont (*W. Boott*), Lake Superior, and northward. (Eu.)

+ + Leaves broader, flat: awn stouter, bent, divergent, or twisted when dry.

4. **C. confinis**, Nutt. Panicle elongated, its rather slender branches spreading at flowering-time, afterwards appressed; glumes lance-oblong, very acute, $2''$ long, pale; hairs of the flower copious, equal, slightly or one third shorter than the thin lower palet and than those of the rudiment; awn borne much below the middle of the palet, somewhat surpassing it; grain glabrous. (*Arundo confinis*, Willd.! *C. inexpansa*, Gray.) — Swamps, N. and W. New York (especially Penn Yan, *Sartwell*) and Pennsylvania. July. — Culm tall.

5. **C. Nuttalliana**, Steud. Culm stout (3° - 5° high); panicle contracted and spike-like; glumes lanceolate and tapering into slender awl-shaped tips, $3''$ long; hairs on the lower side scanty and barely half the length of the firm and keeled lower palet, on the other side longer and equalling the copious tuft on the summit of the rudiment; awn borne half-way between the middle and the tapering tip of the palet, stout, not twisted; grain bearded at the top. (*C. Canadensis*, Nutt. *C. coarctata*, Torr., and of former ed.) — Moist grounds, E. New England to Pennsylvania, Virginia, and southward. Aug.

6. **C. Pórtteri**, Gray. Culm slender (2° - 4° high); a woolly-bearded ring at the junction of the broadly linear leaves with the sheath; panicle long and narrow, with the branches appressed; glumes lanceolate, acute, pale, $2''$ to $2\frac{1}{2}''$ long; hairs of the flower and of the short rudiment scanty, and both reaching about to the middle of the flower behind the upper palet, but very short or none at the

base of the firm-membranaceous lower palet, which bears near its base a twisted awn of its own length.—Dry woods, Pulpit Rocks and vicinity, Huntingdon Co., Pennsylvania, *Prof. T. C. Porter*.

7. **C. Pickeringii**, Gray. Culm $1^{\circ} - 1\frac{1}{2}^{\circ}$ high; leaves short; panicle pyramidal, purplish; glumes ovate-oblong, bluntish or bluntly pointed ($1\frac{1}{2}'' - 2''$ long); hairs both of the flower and of the rudiment very short and scanty, one fourth or fifth the length of the flower, none behind the obtuse lower palet, which bears between its middle and base a short and stout (straight or bent, not twisted) awn.—White Mountains, New Hampshire, in the alpine region of Mt. Washington (*Dr. Pickering, &c.*); and a more luxuriant form with smaller spikelets at Echo Lake, Franconia, *W. Boott*. Sept.

§ 2. **CALAMOVÍLFA**. Glumes and equal palets rather chartaceous, compressed-keeled; the lower glume shorter than the upper and shorter than the palets, of which the lower is 1-nerved and entirely awnless: the upper strongly 2-keeled: rudiment of second flower wanting: panicle open and loose.

8. **C. brevipilis**, Gray. Branches of the diffuse pyramidal panicle capillary (purplish); glumes ovate, mucronate; the upper slightly, the lower nearly one half shorter than the palets, which are above twice the length of the hairs and bristly-bearded along the keels. (*Arundo brevipilis, Torr.*)—Sandy swamps, pine-barrens of New Jersey: rare. Sept.—Culm $3^{\circ} - 4^{\circ}$ high: leaves nearly flat.

9. **C. longifolia**, Hook. Culm ($1^{\circ} - 4^{\circ}$ high) stout, from thick running rootstocks; leaves rigid, elongated, involute above and tapering into a long thread-like point; branches of the pyramidal panicle smooth; glumes lanceolate, the upper as long as the similar palets, the lower one fourth shorter; the copious hairs more than half the length of the naked palets.—Sands, along the upper Great Lakes, from Illinois and Michigan northwestward. Aug.—Spikelets $2\frac{1}{2}''$ long. Sheaths clothed with deciduous wool.

§ 3. **AMMÓPHILA**, Host. Rudiment of second flower present and plumose above: glumes nearly equal and rather longer than the equal similar palets, scarious-chartaceous, lanceolate, compressed-keeled: lower palet 5-nerved, slightly mucronate or obscurely awned near the tip; the upper 2-keeled: squamulæ lanceolate, much longer than the ovary: panicle spiked-contracted: spikelets large ($\frac{1}{2}$ long).

10. **C. arenaria**, Roth. (SEA SAND-REED.) Culm stout and rigid ($2^{\circ} - 3^{\circ}$ high) from firm running rootstocks; leaves long, soon involute; panicle contracted into a dense cylindrical spike ($5' - 9'$ long); hairs only one third of the length of the palet. (*Arundo, L. Psamma, Beauv.*)—Sandy beaches, New Jersey to Maine, and northward, and on the Great Lakes. Aug. (Eu.)

14. **ORYZÓPSIS**, Michx. MOUNTAIN RICE. (Pl. 8.)

Spikelets 1-flowered, nearly terete. Glumes herbaceous or thin-membranaceous, several-nerved, nearly equal, commonly rather longer than the oblong flower, which is deciduous at maturity, and with a very short obtuse callus or scar-like base. Lower palet coriaceous, at length involute so as closely to enclose the upper (of the same length) and the oblong grain; a simple untwisted and deciduous awn jointed off its apex. Stamens 3. Squamulæ 2 or 3, conspicuous. Stigmas plumose.—Perennials, with rigid leaves and a narrow raceme or

panicle. Spikelets greenish, rather large. (Name composed of *ὄρυζα*, rice, and *ὄψις*, likeness, from a fancied resemblance to that grain.)

* *Styles distinct, short: culm leafy to the summit: leaves broad and flat.*

1. **O. melanocárpa**, Muhl. Leaves lanceolate, taper-pointed; sheaths bearded in the throat; panicle simple or sparingly branched; awn thrice the length of the blackish palets (nearly 1' long). (*Milium racemosum*, Smith. *Piptathèrum nigrum*, Torr.)—Rocky woods. Aug.—Culm 2°–3° high.

** *Styles united below, slender: culms tufted, naked: leaves concave or involute.*

2. **O. asperifolia**, Michx. Culms (9'–18' high) with sheaths bearing a mere rudimentary blade, overtopped by the long and rigid linear leaf from the base; very simple panicle or raceme few-flowered; awn 2–3 times the length of the rather hairy whitish palets. (*Uráchne*, Trin.)—Hillsides, &c., in rich woods: common northward. May.—Leaves without keels, rough-edged, pale beneath, lasting through the winter. Squamulæ lanceolate, almost as long as the inner palet!

3. **O. Canadénsis**, Torr. Culms slender (6'–15' high), the lowest sheaths leaf-bearing; leaves involute-thread-shaped; panicle contracted (1'–2' long), the branches usually in pairs; palets pubescent, whitish; awn short and very deciduous, or wanting. (*O. parviflora*, Nutt. *Stipa juncea*, Michx. *S. Canadensis*, Poir. *Milium pungens*, Torr. *Urachne brevicaudata*, Trin.)—Rocky hills and dry plains, W. New England to mountains of Penn., Wisconsin, and northward: rare. May.—Glumes 1''–2'' long, sometimes purplish.

15. STIPA, L. FEATHER-GRASS. (Pl. 8.)

Spikelets 1-flowered, terete; the flower falling away at maturity (with the conspicuous obconical bearded and often sharp-pointed callus) from the membranaceous glumes. Lower palet coriaceous, cylindrical-involute and closely embracing the smaller upper one and the cylindrical grain, having a long and twisted or tortuous simple awn jointed with its apex. Stamens mostly 3. Stigmas plumose.—Perennials, with narrow involute leaves and a loose panicle. (Name from *στύπη*, tow, in allusion to the flaxen appearance of the feathery awns of the original species. In our species the awn is naked.)

* *Callus or base of the flower short and blunt: glumes pointless.*

1. **S. Richardsonii**, Link. Culm (1½°–2° high) and leaves slender; panicle loose (4'–5' long), with slender few-flowered branches; glumes nearly equal, oblong, acutish (2½'' long), about equalling the pubescent linear-oblong lower palet, which bears a tortuous awn 6''–8'' long.—Pleasant Mountain, near Sebago Lake, Maine, *C. J. Sprague*, and northwestward. (Flowers rather smaller than in Richardson's plant, as described.)

** *Callus or base of the flower pungently pointed, at maturity villous-bearded: lower palet slender and minutely bearded at the tip: glumes taper-pointed.*

2. **S. avenácea**, L. (BLACK OAT-GRASS.) Culm slender (1°–2° high); leaves almost bristle-form; panicle open; palets blackish, nearly as long as the glumes (about 4'' long); the awn bent above, twisted below (2'–3' long).—Dry or sandy woods, S. New England to Wisconsin, and southward. July.

3. **S. spártea**, Trin., not of Hook. (PORCUPINE GRASS.) Culm rather

stout ($1\frac{1}{2}^{\circ}$ – 3° high); *panicle contracted*; *palets linear*, $\frac{3}{4}$ '–1' long (including the long callus), pubescent below, *shorter than the lanceolate slender subulate-pointed greenish glumes*; the twisted strong awn ($3\frac{1}{2}$ '–7' long, pubescent below, rough above. (S. juncea, Pursh?) — Plains and prairies, from Illinois and N. Michigan northwestward. May–July.

16. ARÍSTIDA, L. TRIPLE-AWNED GRASS. (Pl. 8.)

Glumes unequal, often bristle-pointed. Lower palet tipped with three awns; the upper much smaller. Otherwise much as in Stipa. — Culms branching; leaves narrow, often involute. Spikelets in simple or paniced racemes or spikes. Grain linear. (Name from *arista*, a beard or awn.) All grow in sterile, dry soil, and all ours have the awns naked and persistent, and flower late.

* *Awns separate to the base, not jointed with the palet.*

+ *Awns very unequal; the much shorter or minute lateral ones erect; the elongated middle one horizontal or turned downwards: glumes equal or the upper one longer: low (5'–18' high) and branching, mostly tufted annuals.*

++ *Spikelets few in loose simple spikes or racemes: glumes 3–5-nerved.*

1. **A. ramosíssima**, Engelm. mss. Culms diffusely much branched; glumes (9''–10'' long) rather shorter than the flower; middle awn 1' long, soon abruptly hooked-recurved, the lateral ones hardly 2'' long; ligule truncate, bearded. — Dry prairies of Illinois (*Engelmann, Vasey*) and Kentucky (*Michaux*). — Var. UNIARISTATA, with lateral awns wanting. Ringwood, Illinois, *Vasey*.

++ ++ *Spikelets more numerous: glumes (3''–4'' long) carinately 1-nerved.*

2. **A. dichótoma**, Michx. (POVERTY GRASS.) Culms low, much branched throughout, ascending; spikelets in short narrow clusters; glumes nearly equal, longer than the flower, fully equalling its *minute lateral awns*, the soon *reflexed middle awn about the length of the palet*. — Dry, sandy or gravelly fields: common, especially southward.

3. **A. grácilis**, Ell. Culms slender, erect (6'–18' high), naked above and terminating in a slender raceme- or spike-like virgate panicle; glumes about the length of the flower, the *exserted lateral awns varying from one third to fully half the length of the horizontally bent middle one*; or in var. DEPAUPERATA, from one fifth to one third its length. — Sandy soil, coast of Mass. and from Illinois southward. — Middle awn 6''–9'' long, in the ordinary forms. In the var. which abounds on micaceous hills near Philadelphia (*C. E. Smith*), the flowers are much smaller, and awns shorter; but it passes into the larger form.

+ + *Awns all diverging and alike, or the lateral ones moderately shorter.*

++ *Glumes equal or the upper one longer.*

4. **A. stricta**, Michx. Culms (2° – 3° high) densely tufted from a *perennial* root, bearing a (1°) long *spiked panicle*; leaves involute-thread-form, long, rigid, sometimes downy: awns about the length of the flower (6'') or the lateral one third shorter. — Virginia and southward.

5. **A. oligántha**, Michx. Culms (6''–20'' high) tufted from an *annual?* root, bearing a *loosely few-flowered raceme*; leaves short, somewhat involute when dry; lower glume 3–5-nerved (nearly 1' long); *awns capillary*, $1\frac{1}{2}$ '–3' long, much exceeding the slender flower. — Virginia to Illinois, and common southwestward.

↔ ↔ *Upper glume shorter than the lower: perennials, simple-stemmed, 2°-4° high.*

6. **A. purpurascens**, Poir. *Glabrous; leaves rather involute; flowers in a (10' - 18') long spiked panicle; awns much longer than the flower, the middle one about 1' long. (A. racemosa, Muhl. A. Geyeriana, Steud.)* — Massachusetts to Michigan, Illinois, and southward: common.

7. **A. lanata**, Poir. *Tall and stout; leaves tardily involute, rough on the upper side, rigid; sheaths woolly; panicle (1°-2° long) spike-like or more compound and open; middle awn (1' long) longer than the flower.* — Salisbury, Maryland, *W. M. Canby*, and southward.

* * *Awns united below into one, jointed with the apex of the palet: root annual.*

8. **A. tuberculosa**, Nutt. *Culm branched below (6' - 18' high), tumid at the joints; panicles rigid, loose; the branches in pairs, one of them short and about 2-flowered, the other elongated and several-flowered; glumes (1' long, including their slender-awned tips longer than the palet; which is tipped with the common stalk (about its own length) of the 3 equal divergently-bent awns (1½' - 2' long) twisting together at the base.* — Sandy soil, E. Massachusetts to New Jersey; also Wisconsin, Illinois, and southward.

17. SPARTINA, Schreber. CORD OR MARSH GRASS. (Pl. 9.)

Spikelets 1-flowered, without a rudiment, very much flattened laterally, spiked in 2 ranks on the outer side of a triangular rhachis. Glumes strongly compressed-keeled, acute, or bristle-pointed, mostly rough-bristly on the keel; the upper one much larger and exceeding the pointless and awnless palets, of which the upper is longest. Squamulæ none. Stamens 3. Styles long, more or less united. — Perennials, with simple and rigid reed-like culms, from extensively creeping scaly rootstocks, racemed spikes, very smooth sheaths, and long and tough leaves (whence the name, from *σπάρτιν*, *a cord*, such as was made from the bark of the *Spartium* or Broom.)

* *Spikelets compactly imbricated, very rough-hispid on the keels: spikes (2' - 4' long) more or less peduncled: culm and elongated leaves rigid.*

1. **S. cynosuroides**, Willd. (FRESH-WATER CORD-GRASS.) *Culm rather slender (2° - 6° high); leaves narrow (2° - 4° long, ½' or less wide below), tapering to a very slender point, keeled, flat, but quickly involute in drying, smooth except the margins; spikes 5 - 20, scattered, spreading; rhachis rough on the margins; glumes awn-pointed, especially the upper, the lower equalling the lower palet, whose strong rough-hispid midrib abruptly terminates below the membranous apex. (Trachynotia cynosuroides, Michx. Limnëtis, Pers.)* — Banks of rivers and lakes, especially northward. Aug. — Glumes strongly serrulate-hispid on the keel; the awn of the upper one about ¾' long. Palets somewhat unequal. — Certainly distinct from the next, to which, in strictness, the Linnæan name belongs.

2. **S. polystachya**, Willd., Muhl. (SALT REED-GRASS.) *Culm tall and stout (4° - 9° high, often 1' in diameter near the base); leaves broad (½' to 1'), roughish underneath, as well as the margins; spikes 20 - 50, forming a dense oblong raceme (purplish); glumes barely mucronate, the lower half the length of the equal palets, of which the rough-hispid midrib of the lower one reaches to the apex.*

(*Trachynotia polystachya*, Michx. *Dactylis cynosuroides*, L. in part, excl. var.) — Salt or brackish marshes, within tide-water, especially southward.

3. **S. júncea**, Willd. (RUSH SALT-GRASS.) Culms low (1° – 2° high) and slender; leaves narrow and rush-like, strongly involute, very smooth; spikes 1–5, on very short peduncles; the rhachis smooth; glumes acute, the lower scarcely half the length of the upper, not half the length of the lower palet. (*Dáctylis patens*, Ait.) — Salt marshes and sea-beaches. Aug. (Eu.)

* * Spikelets loosely imbricated, or somewhat remote and alternate, the keels only slightly hairy or roughish under a lens: spikes sessile and erect, soft: leaves, rhachis, &c. very smooth; culm rather succulent.

4. **S. stricta**, Roth. (SALT MARSH-GRASS.) Culm 1° – 4° high, leafy to the top; leaves soon convolute, narrow; spikes few (2–4), the rhachis slightly projecting at the summit beyond the crowded or imbricated spikelets: glumes acute, very unequal, the larger 1-nerved, a little longer than the palets. — Salt marshes, Pennsylvania, &c. (*Muhl.*) — Odor strong and rancid. (Eu.)

Var. **glábra**. (*S. glabra*, *Muhl.*, partly.) Culm and leaves longer; spikes 5–12 ($2'$ – $3'$ long); spikelets imbricate-crowded. — Common on the coast.

Var. **alterniflora**. (*S. alterniflora*, *Loisel.* *Dactylis cynosuroides*, var., L.) Spikes more slender ($3'$ – $5'$ long), and the spikelets remotish, barely overlapping, the rhachis continued into a more conspicuous bract-like appendage: larger glume indistinctly 5-nerved (not so evidently as in the European and Tropical American plant): otherwise as in the preceding form, into which it passes. — Common with the last: also Onondaga Lake, *J. A. Paine*.

18. CTÉNIUM, Panzer. TOOTHACHE-GRASS. (Pl. 9.)

Spikelets densely imbricated in two rows on one side of the flat curved rhachis of the solitary terminal spike. Glumes persistent: the lower one (interior) much smaller; the other concave below, bearing a stout recurved awn, like a horn, on the middle of the back. Flowers 4–6, all but one neutral; the one or two lower consisting of empty awned palets: the one or two uppermost of empty awnless palets: the perfect flower intermediate in position; its palets membranaceous, the lower awned or mucronate below the apex and densely ciliate towards the base, 3-nerved. Squamulæ 2. Stamens 3. Stigmas plumose. (Name *Κτενίου*, a small comb, from the pectinate appearance of the spike.)

1. **C. Americanum**, Spreng. Culm (3° – 4° high from a perennial root) simple, pubescent or roughish; larger glume warty-glandular outside and conspicuously awned. (*Monócera aromática*, *Ell.*) — Wet pine barrens, S. Virginia and southward. — Taste very pungent.

19. BOUTELOÛA, Lagasca (1305). MUSKFT-GRASS. (Pl. 9.)

Spikelets crowded and closely sessile in 2 rows on one side of a flattened rhachis, comprising one perfect flower below and one or more sterile (mostly neutral) or rudimentary flowers. Glumes convex-keeled, the lower one shorter. Perfect flower with the 3-nerved lower palet 3-toothed or cleft at the apex, the 2-nerved upper palet 2-toothed; the teeth, at least of the former, pointed or sub-

ulate-awned. Stamens 3: anthers orange-colored or red. Rudimentary flowers mostly 1-3-awned. Spikes solitary, racemed or spiked; the rhachis somewhat extended beyond the spikelets. (Named for *Claudius Boutelou*, a Spanish writer upon floriculture and agriculture.)

§ 1. **CHONDROSIUM**, Desv. *Spikes pectinate, of very many spikelets, oblong or linear, very dense, solitary and terminal or few in a raceme: sterile flowers 1-3 on a short pedicel, neutral, consisting of 1-3 scales and awns.*

1. **B. oligostachya**, Torr. Glabrous, perennial (6'-12' high); leaves very narrow; spikes 1-5, the rhachis glabrous; glumes and lower fertile palet sparingly soft-hairy; the lobes awl-pointed; sterile flower copiously villous-tufted at the summit of the naked pedicel, its 3 awns equalling the larger glume. (Atheropogon, Nutt.)—N. W. Wisconsin and westward.—Glumes obscurely if at all papillose along the keel. Middle lobe of the lower palet 2-cleft at the tip. Sterile flowers often 2, the second mostly a large awnless scale, becoming hood-like and coriaceous. (Near *B. gracilis*: perhaps *B. juncifolia*, Lag.)

2. **B. hirsuta**, Lagasca. Tufted, annual? (8'-20' high); leaves flat, lance-linear, papillose-hairy or glabrous; spikes 1-4; upper glume hispid with strong bristles from dark warty glands; lower palet pubescent, 3-cleft into awl-pointed lobes; sterile flower and its pedicel glabrous, the 3 awns longer than the glumes and fertile flower. (Atheropogon papillosus, Engelm. *Chondrosium hirtum*, H. B. K.)—Sandy plains, Wisconsin, Illinois, and southwestward.

§ 2. **ATHEROPOGON**, Muhl. *Spikes short, numerous in a long and virgate one-sided spike or raceme, spreading or reflexed, each of few (4-12) spikelets: sterile flowers neutral, rudimentary.*

3. **B. curtipendula**, Gray. Culms tufted from perennial rootstocks (1°-3° high); sheaths often hairy; leaves narrow; spikes $\frac{1}{2}$ ' or less in length, nearly sessile, 30 to 60 in number in a loose general spike (8'-15' long); flowers scabrous; the lower palet of the fertile with 3 short awl-pointed teeth; sterile flower reduced to a single small awn, or mostly to 3 awns shorter than the fertile flower, and 1 or 2 small or minute scales. (*B. racemosa*, Lagasca. *Chloris curtipendula*, Michx. *Atheropogon apludioides*, Muhl. *Eutriana curtipendula*, Trin.)—Dry hills and plains, S. New York to Wisconsin, and southward. July-Sept.—Passes by transitions into, Var. **ARISTOSA**, with spikes shorter; sterile flower of a large saccate lower palet, awned at the 2-cleft tip and from the lateral nerves, the middle awn exserted, and with a rudiment of an inner palet. (*Eutriana affinis*, J. D. Hook.)—Illinois (*Geyer*), and southward.

20. **GYMNOPOGON**, Beauv. NAKED-BEARD GRASS. (Pl. 9.)

Spikelets of one perfect flower, and the rudiment of a second (consisting of an awn-like pedicel mostly bearing a naked bristle), sessile and remotely alternate on long and filiform rays or spikes, which form a crowded naked raceme. Glumes lance-awl-shaped, keeled, almost equal, rather longer than the somewhat equal membranaceous palets; of which the lower is cylindrical-involute, with the midrib produced from just below the 2-cleft apex into a straight and slender bristle-like awn; the upper with the abortive rudiment at its base.

Stamens 3. Stigmas pencil-form, purple. — Root perennial. Leaves short and flat, thickish, 1' - 3' long. (Name composed of *γυμνός*, *naked*, and *πώγων*, *a beard*, alluding to the reduction of the abortive flower to a bare awn.)

1. **G. racemòsus**, Beauv. Culms clustered from a short rootstock (1° high), wiry, leafy; leaves oblong-lanceolate; *spikes flower-bearing to the base* (5' - 8' long), soon divergent; awn of the abortive flower shorter than its stalk, equalling the *pointed glumes*, not more than half the length of the awn of the fertile flower. (*Anthopògon lepturoìdes*, *Nutt.*) — Sandy pine-barrens, New Jersey to Virginia, and southward. Aug., Sept.

2. **G. brevifòlius**, Trin. Filiform *spikes long-peduncled, i. e. flower-bearing only above the middle*; lower palet ciliate near the base, short-awned; *awn of the abortive flower obsolete or minute*; *glumes acute*. (*Anthopògon brevifolius* & *filifòrmis*, *Nutt.*) — Sussex County, Delaware, and southward.

21. CYNODON, Richard. BERMUDA or SCUTCH-GRASS. (Pl. 9.)

Spikelets 1-flowered, with a mere naked short-pedicelled rudiment of a second flower, imbricate-spiked on one side of a flattish rhachis; the spikes usually digitate at the naked summit of the flowering culms. Glumes keeled, pointless, rather unequal. Palets pointless and awnless; the lower larger, boat-shaped. Stamens 3. — Low diffusely-branched and extensively creeping perennials, with short flattish leaves. (Name composed of *κύων*, *a dog*, and *ὀδοός*, *a tooth*.)

1. **C. DACTYLON**, Pers. Spikes 3 - 5; palets smooth, longer than the blunt rudiment. — Penn. and southward; troublesome in light soil. (Nat. from Eu.)

22. DACTYLOCTENIUM, Willd. EGYPTIAN GRASS. (Pl. 9.)

Spikelets several-flowered, with the uppermost flower imperfect, crowded on one side of a flattened rhachis, forming dense pectinate spikes, 2 - 5 in number, digitate at the summit of the culm. Glumes compressed laterally and keeled, membranaceous, the upper (exterior) one awn-pointed. Lower palet strongly keeled and boat-shaped, pointed. Stamens 3. Pericarp a thin utricle, containing a loose globular and rough-wrinkled seed. — Root annual. Culms diffuse, often creeping at the base. (Name compounded of *δάκτυλος*, *finger*, and *κτενίον*, *a little comb*, alluding to the digitate and pectinate spikes.)

1. **D. ÆGYPTIACUM**, Willd. Spikes 4 - 5; leaves ciliate at the base. (*Chloris mucronàta*, *Michx.*) — Cultivated fields and yards, Virginia, Illinois, and southward. (Adv. from Afr.?)

23. ELEUSINE, Gærtn. CRAB-GRASS. YARD-GRASS. (Pl. 9.)

Spikelets 2 - 6-flowered, with a terminal naked rudiment, closely imbricate-spiked on one side of a flattish rhachis; the spikes digitate. Glumes membranaceous, pointless, shorter than the flowers. Palets awnless and pointless; the lower ovate, keeled, larger than the upper. Stamens 3. Pericarp (utricle) containing a loose oval and wrinkled seed. — Low annuals, with flat leaves, and flowers much as in *Poa*. (Name from *Ἐλευσίη*, the town where *Ceres*, the goddess of harvests, was worshipped.)

1. **E. ÍNDICA**, Gærtn. (DOG'S-TAIL OR WIRE GRASS.) Culms ascending, flattened; spikes 2-5 (about 2' long, greenish).—Yards, &c., chiefly southward. (Nat. from Ind.?)

24. LEPTÓCHLOA, Beauv. (OXYDÈNIA, Nutt.) (Pl. 9.)

Spikelets 3-many-flowered (the uppermost flower imperfect), loosely spiked on one side of a long filiform rhachis: the spikes racemed. Glumes membranaceous, keeled, often awl-pointed, the upper one somewhat larger. Lower palet 3-nerved, with the lateral nerves next the ciliate or hairy margins awnless, or bristle-awned at the entire or 2-toothed tip, larger than the upper. Stamens 2 or 3. Seed sometimes loose in the pericarp.—Ours annuals. Leaves flat. (Name composed of λεπτός, slender, and χλόα, grass, from the long attenuated spikes.)

§ 1. LEPTOCHLOA proper. *Lower palet awnless or simply awned.*

1. **L. mucronàta**, Kunth. Sheaths hairy; spikes numerous (20-40, 2'-4' in length), in a long panicle-like raceme; spikelets small; glumes more or less mucronate, nearly equalling or exceeding the 3-4 awnless flowers.—Fields, Virginia to Illinois, and southward. August.

§ 2. DIPLÁCHNE, Beauv. *Lower palet bristle-awned from the 2-toothed apex; the marginal nerves often excurrent into lateral teeth or points.*

2. **L. fasciculàris**, Gray. Smooth; leaves longer than the geniculate-decumbent and branching culms, the upper sheathing the base of the crowded panicle-like raceme, which is composed of many strict spikes (3'-5' long); spikelets slightly pedicelled, 7-11-flowered, much longer than the lanceolate glumes; palets hairy-margined towards the base; the lower one with 2 small lateral teeth and a short awn in the cleft of the apex. (*Festuca fascicularis*, Lam. *F. polystàchya*, Michx. *Diplachne fascicularis*, Beauv., Torr.)—Brackish meadows, from Rhode Island southward along the coast, and from Illinois southward on the Mississippi. Aug.-Sept.—Makes a direct transition to the next genus.

25. TRICÚSPIS, Beauv. (Pl. 10.)

Spikelets 3-12-flowered, somewhat terete; the terminal flower abortive. Glumes unequal. Rhachis of the spikelet bearded below each flower. Palets membranaceous or somewhat chartaceous; the lower much larger than the 2-toothed upper one, convex, 2-3-toothed or cleft at the apex, conspicuously hairy-bearded or villous on the 3 strong nerves, of which the lateral are marginal or nearly so and usually excurrent, as is the mid-nerve especially, into a short cusp or awn. Stamens 3. Stigmas dark purple, plumose. Grain oblong, nearly gibbous.—Leaves taper-pointed: sheaths bearded at the throat. Panicle simple or compound; the spikelets often racemose, purplish. (Name from the Latin *tricuspis*, three-pointed, alluding to the lower palet.)

§ 1. TRICUSPIS proper. (*Windsoria*, Nutt.) *Glumes shorter than the crowded flowers: lower palet 3-cuspidate by the projection of the nerves, and usually with intermediate membranaceous teeth; the upper palet naked.*

1. **T. seslerioides**, Torr. (TALL RED-TOP.) Perennial; culm upright (3° – 5° high), very smooth, as are the flat leaves; panicle large and compound, the rigid capillary branches spreading, naked below; spikelets very numerous, 5–7-flowered, shining, purple ($4''$ long); the flowers hairy toward the base. (*Poa flava*, L.! *P. seslerioides*, Michx. *P. quinquéfida*, Pursh. *Windsoria*, poæfórmis, Nutt. *Uralespis cùprea*, Kunth.)—Dry or sandy fields, S. New York to Illinois and southward. Aug.—A showy grass, with the spreading panicle sometimes 1° wide. Points of the lower palet almost equal, scarcely exceeding the intermediate teeth, thus appearing 5-toothed.

§ 2. **TRIPLÀSIS**, Beauv. (*Diplòcea*, Raf. *Uralespis*, Nutt.) *Glumes much shorter than the somewhat remote flowers: both palets strongly fringe-bearded; the lower 2-cleft at the summit, its mid-nerve produced into an awn between the truncate or awn-pointed divisions.*

2. **T. purpùrea**, Gray. (SAND-GRASS.) Culms many in a tuft from the same annual root, ascending ($6'$ – $12'$ high), with numerous bearded joints; leaves involute-awl-shaped, mostly short; panicles very simple, bearing few 2–5-flowered spikelets, the terminal one usually exserted, the axillary ones included in the commonly hairy sheaths; *awn much shorter than the palet, seldom exceeding its eroded-truncate or obtuse lateral lobes.* (*Aira purpùrea*, Walt. *Diplòcea barbàta*, Raf. *Uralespis purpùrea* and *U. aristulàta*, Nutt.)—In sand, Massachusetts to Virginia along the coast, and southward: also Lake Erie, near Buffalo, G. W. Clinton. Aug., Sept.—Plant acid to the taste.

(*T. CORNÛTA* (*Uralespis cornuta*, Ell., and *Triplasis Americana*, Beauv.!) may perhaps extend north to the borders of Virginia.)

26. GRAPHÉPHORUM, Desv. (DUPÓNȚIA, R. Br.) (Pl. 10.)

Spikelets 2–5-flowered, rather terete. Glumes membranaceous, mostly nearly equalling the remote flowers. A cluster of villous hairs at the base of each flower. Palets thin and membranaceous or scarious; the lower one convex, scarcely keeled, faintly nerved, entire, pointless and awnless. Stamens 3. Stigmas plumose. Ovary glabrous.—Perennial and northern or arctic grasses, with linear flat leaves, their sheaths closed at the base, the spikelets in a loose panicle. (Genus allied to the *Aveneæ*, but awnless; named from *γραφίς*, a pencil, and *φέρω*, to bear, from the tufts of hair at the base of the flowers.)

1. **G. melicoides**, Beauv. Culm 1° – 2° high; leaves roughish; panicle open; glumes unequal, lanceolate, their midrib and the pedicels rough; joints of the rachis unilaterally bearded between the 2–4 flowers. Grand Detour, Upper Michigan, Prof. Porter, Rivière du Loup, E. Canada, Dr. Pickering, and northward: rare.—Var. MAJOR, Gray (*Dupontia Cooleyi*, of former ed.), is a luxuriant form, 2° – 3° high, with ampler panicle; found on the borders of a swamp, Washington, Macomb County, Michigan, by Dr. Cooley. Aug.

27. DIARRHÈNA, Raf. DIARRHENA. (Pl. 10.)

Spikelets several-flowered, smooth and shining, one or two of the uppermost flowers sterile. Glumes ovate, much shorter than the flowers, coriaceous; the lower one much smaller. Lower palet ovate, convex on the back, rigidly cori-

aceous, its 3 nerves terminating in a strong and abrupt cuspidate or awl-shaped tip. Squamulæ ovate, ciliate. Stamens 2. Grain very large, obliquely ovoid, obtusely pointed, rather longer than the palets, the cartilaginous shining pericarp not adherent to the seed. — A nearly smooth perennial, with running rootstocks, producing simple culms (2°–3° high) with long linear-lanceolate flat leaves towards the base, naked above, bearing a few short-pedicelled spikelets (2''–3'' long) in a very simple panicle. (Name composed of *δῖς*, *two*, and *ἀρρήν*, *man*, from the two stamens.)

1. **D. Americana**, Beauv. (*Festuca diandra*, Michx.) — Shaded riverbanks and woods, Ohio to Illinois and southward. Aug.

28. DÁCTYLIS, L. ORCHARD GRASS. (Pl. 10.)

Spikelets several-flowered, crowded in one-sided clusters, forming a branching dense panicle. Glumes and lower palet herbaceous, keeled, awn-pointed, rough-ciliate on the keel; the 5 nerves of the latter converging into the awn-like point; the upper glume commonly smaller and thinner. Stamens 3. Grain lance-oblong, acute, free. — Perennials: leaves keeled. (Name *δακτυλῖς*, *a finger's breadth*, apparently in allusion to the size of the clusters.)

1. **D. GLOMERATA**, L. Rough, rather glaucous (3° high); leaves broadly linear; branches of the panicle naked at the base; spikelets 3–4-flowered. — Fields and yards, especially in shade. A variety with downy spikelets, Medford, Mass., *W. Boott*. June. — Good for hay. (Nat. from Eu.)

29. KCELÈRIA, Pers. KÆLERIA. (Pl. 10.)

Spikelets 3–7-flowered, crowded in a dense and narrow spike-like panicle. Glumes and lower palet membranaceous, compressed-keeled, obscurely 3-nerved, barely acute, or the latter often mucronate or bristle-pointed; the former moderately unequal, nearly as long as the spikelet. Stamens 3. Grain free. — Tufted Grasses (allied to *Dactylis* and *Poa*), with simple upright culms; the sheaths often downy. (Named for *Prof. G. L. Köhler*, or *Kæler*, an early writer on Grasses.)

1. **K. cristata**, Pers. Panicle narrowly spiked, interrupted or lobed at the base; spikelets 2–4-flowered; lower palet acute or mucronate; leaves flat, the lower sparingly hairy or ciliate. — Var. *GRACILIS*, with a long and narrow spike, the flowers usually barely acute. (*K. nitida*, *Nutt.*) — Dry hills, Penn. to Illinois, thence northward and westward. (Eu.)

30. EATÒNIA, Raf. (REBOÛLEA, Kunth.) (Pl. 10.)

Spikelets usually 2-flowered, and with an abortive rudiment or pedicel, numerous, in a contracted or slender panicle, very smooth. Glumes somewhat equal in length, but very dissimilar, a little shorter than the flowers; the lower narrowly linear, keeled, 1-nerved; the upper broadly obovate, folded round the flowers, 3-nerved on the back, not keeled, scarious-margined. Lower palet oblong, obtuse, compressed-boat-shaped, naked, chartaceous; the upper very thin and hyaline. Stamens 3. Grain linear-oblong, not grooved. — Perennial, slen-

der grasses, with simple and tufted culms, and often sparsely downy sheaths, flat lower leaves, and small greenish (or rarely purplish-tinged) spikelets. (Named for *Prof. Amos Eaton*, author of a popular *Manual of the Botany of the United States*, which was for a long time the only general work available for students in this country, and of other popular treatises.)

1. **E. obtusàta**, Gray. *Panicle dense and contracted, somewhat interrupted, rarely slender; the spikelets crowded on the short erect branches; upper glume rounded-obovate, truncate-obtuse, rough on the back; flowers lance-oblong.* (*Aira obtusàta, Michx.* *A. truncàta, Muhl.* *Kœleria truncata, Torr.* *K. paniculàta, Nutt.* *Reboulea gracilis, Kunth*, in part. *R. obtusata, Ed. 1.* *Eatonia purpuràscens, Raf.?*) — Dry soil, N. Pennsylvania to Wisconsin, and southward. June, July.

2. **E. Pennsylvànica**, Gray. *Panicle long and slender, loose, the racemose branches somewhat elongated; upper glume obtuse or bluntly somewhat pointed; the 2 (rarely 3) flowers lanceolate.* (*Kœleria Pennsylvànica, DC.* *Aira mòllis, Muhl.* *Reboulea Pennsylvànica, Ed. 1.*) — Varies, with a fuller panicle, 6' - 8' long, with the aspect of *Cinna* (var. *MAJOR, Torr.*); and, rarely, with the lower palet minutely mucronate-pointed! — Moist woods and meadows: common.

31. MÉLICA, L. MELIC-GRASS. (Pl. 10.)

Spikelets 2 - 5-flowered; the 1 - 3 upper flowers imperfect and dissimilar, convolute around each other, and enwrapped by the upper fertile flower. Glumes usually large, scarious-margined, convex, obtuse; the upper 7 - 9-nerved. Palets papery-membranaceous, dry and sometimes indurating with age; the lower rounded or flattish on the back, 7 - many-nerved, scarious at the entire blunt summit. Stamens 3. Stigmas branched-plumose. — Perennials with soft and flat leaves. Panicle simple or sparingly branched; the rather large spikelets racemose-one-sided. (An old name, from *μέλι*, *honey*.)

1. **M. mùtica**, Walt. *Panicle simple or branched; glumes unequal, the larger almost equalling the spikelet; fertile flowers 2; lower palet naked, glabrous minutely scabrous on the nerves.* (*M. glabra, Michx.* *M. speciòsa, Muhl.*) — Var. *GLÀBRA* (*M. glabra, Pursh*) has the panicle often few-flowered and rather simple, the lower palet very blunt. — Var. *DIFFÙSA* (*M. diffusa, Pursh*) is taller, 2½° - 4° high, with a more compound and many-flowered panicle; the lower palet commonly more scabrous and its tip narrower. — Rich soil, S. E. Pennsylvania to Wisconsin, and southward. June.

32. GLYCÈRIA, R. Br., Trin. MANNA-GRASS. (Pl. 10.)

Spikelets terete or flattish, several - many-flowered; the flowers mostly early deciduous by the breaking up of the rhachis into joints, leaving the short and unequal 1 - 3-nerved membranaceous glumes behind. Palets naked, of a rather firm texture, nearly equal; the lower rounded on the back, scarious (and sometimes obscurely toothed) at the blunt or rarely acute summit, glabrous, 5 - 7-nerved, the nerves parallel and separate. Stamens 3, or in the first section commonly 2. Stigmas plumose, mostly compound. Ovary smooth. Grain oblong, free. — Perennial, smooth marsh-grasses, mostly with creeping bases or

rootstocks; the spikelets paniced. (Name from *γλυκερός*, *sweet*, in allusion to the taste of the grain.)

§ 1. GLYCERIA proper. *Lower palet conspicuously nerved: styles present: plumes of the stigma branched or toothed: grain grooved on the inner side: leaves flat, the sheaths nearly entire.*

* Spikelets ovate, oblong, or linear-oblong, 1'' - 3'' in length,

+ At length nodding in an open panicle, flattish laterally but turgid.

1. **G. Canadensis**, Trin. (RATTLESNAKE-GRASS.) Panicle oblong-pyramidal, at length drooping; spikelets ovate, at length very broad and tumid, Briza-like, 2'' long, pale, with purplish glumes; lower palet acute or blunt-pointed, firm, with not very prominent nerves, longer than the rounded upper one; culm stout, 2° - 3° high; leaves long, roughish. (*Briza Canadensis*, Michx. — Bogs and wet places: common from Penn. northward. July.

+ + Erect in a narrow contracted panicle, somewhat flattened and turgid.

2. **G. obtusa**, Trin. Panicle narrowly oblong, dense; (3' - 5' long); spikelets 6 - 7-flowered; 2'' - 3'' long; lower palet obtuse; culm stout, 1° - 2° high, very leafy; leaves long, smooth. (*Poa obtusa*, Muhl.) — Bogs, E. New England to Penn., near the coast.

3. **G. elongata**, Trin. Panicle narrowly racemose, elongated (1° long), recurving; the branches and 3 - 4-flowered spikelets appressed; lower palet obtuse; leaves very long (1° or more), rough. (*Poa elongata*, Torr.) — Wet woods, New England to Michigan, and northward. July - Aug.

+ + + Diffuse: lower palet truncate-obtuse, prominently 7-nerved; upper 2-toothed.

4. **G. nervata**, Trin. (FOWL-MEADOW GRASS, in part.) Branches of the loose panicle capillary, at length drooping, the very numerous small spikelets ovate-oblong, 3 - 7-flowered; leaves rather long. (*Poa nervata*, Willd. *P. striata*, Michx. *P. parviflora*, Pursh.) — Moist meadows: very common. June. — Culm erect, 1° - 3° high. Spikelets 1'' - 2'' long, commonly purplish.

5. **G. pallida**, Trin. Branches of the rather simple panicle slender, erect-spreading, rough; the spikelets usually few, somewhat appressed, oblong-linear, 5 - 9-flowered (pale, 2'' - 3'' long); lower palet minutely 5-toothed; the upper lanceolate, conspicuously 2-toothed; leaves short, sharp-pointed, pale. (*Windsoria pallida* & *Poa dentata*, Torr.) — Shallow water: common, especially northward. July. — Culms slender, 1° - 3° long, ascending from a creeping base.

6. **G. aquatica**, Smith. (REED MEADOW-GRASS.) Panicle much branched, ample (8' - 15' long); the numerous branches ascending, spreading with age; spikelets oblong or linear-oblong, 5 - 9-flowered (usually purplish, 2'' - 3'' long); lower palet entire; leaves large (1° - 2° long, $\frac{1}{2}$ ' to $\frac{3}{4}$ ' wide. — Wet grounds: common northward. July. — Culm stout, upright, 3° - 5° high. (Eu.)

** Spikelets linear ($\frac{1}{2}$ ' - 1' long), pale, appressed on the branches of the long and narrow racemose panicle, terete except during anthesis: palets minutely roughish, the upper 2-toothed: squamulae unilateral or united: ligule long: culms flattened (1° - 5° high), ascending from a rooting base. (*Glyceria*, R. Br.)

7. **G. fluitans**, R. Br. Spikelets 7 - 13-flowered; lower palet oblong, obtuse, or the scarious tip acutish, entire or obscurely 3-lobed, usually rather longer

than the blunt upper one. (*G. plicata*, *Fries.*) — Shallow water. common. June — Aug. — Leaves short and rather broad, very smooth. Panicle 1° long: the simple branches appressed, finally spreading below. (Eu.)

8. *G. acutiflora*, Torr. Spikelets 5 — 12-flowered, few and scattered; lower palet oblong-lanceolate, acute, shorter than the long tapering point of the upper one. — Wet places, Penn. to Maine: rather rare. June. — Resembles the last; but the erect leaves smaller, the separate flowers twice the length (4'' long), and less nerved.

§ 2. HELEÓCHLOA, *Fries.* (*Scleròchloa*, *Ed.* 1.) Lower palet inconspicuously or obsoletely 5-nerved: stigmas nearly sessile and simply plumose: grain hardly grooved: saline species: panicle contracted with age.

9. *G. maritima*, Wahl. (SEA SPEAR-GRASS.) Sterile shoots procumbent, runner-like; flowering culms erect (1° — 1½° high); branches of the panicle solitary or in pairs; spikelets oblong or linear, 4 — 8-flowered; lower palet rounded at the summit, slightly pubescent towards the base; leaves somewhat involute; ligule elongated. (*Poa maritima*, *Huds.*) — Sea coast: not rare. (Eu.)

10. *G. distans*, Wahl. Culms geniculate at the base, ascending, destitute of running shoots; branches of the panicle 3 — 5 in a half whorl, spreading; spikelets 3 — 6-flowered; lower palet truncate-obtuse; leaves mostly flat; ligule short. (*P. fasciculata*, *Torr.* — Salt marshes along the coast. — Too like the last. (Eu.)

32. BRIZOPÿRUM, Link. SPIKE-GRASS. (Pl. 10.)

Spikelets and numerous flowers compressed, crowded in a densely spiked or capitate panicle. Glumes herbaceous or membranaceous; the lower faintly many-nerved. Lower palet rather coriaceous, laterally much flattened, indistinctly many-nerved, acute. Ovary stalked. — Flowers diœcious, pretty large. Leaves crowded on the culms, involute, commonly rigid. (Name compounded of *Briza*, the Quaking-Grass, and *πυρός*, *wheat*.)

1. *B. spicatum*, Hook. Culms tufted from creeping rootstocks (9' — 18' high); spike oblong, flattened (1' long); spikelets ovate or oblong, 5 — 10-flowered; flowers smooth and naked; grain pointed. (*Uniola spicata*, *L.* *Poa Michauxii*, *Kunth.*) — Salt marshes and shores. Aug. — Pistillate flowers more rigid and almost keeled, with very long plumose stigmas; the sterile smaller and somewhat rounded on the back.

34. PÒA, L. MEADOW-GRASS. SPEAR-GRASS. (Pl. 10.)

Spikelets ovate or lance-ovate, laterally compressed, several- (2 — 10-) flowered, in an open panicle. Glumes mostly shorter than the flowers; the lower smaller. Lower palet membranaceo-herbaceous, with a delicate scarious margin, compressed-keeled, pointless, 5-nerved (the intermediate nerves more obscure or obsolete), the principal nerves commonly clothed at and towards the base with soft hairs: upper palet membranaceous, 2-toothed: base of the flower often cobwebby. Stamens 2 or 3. Stigmas simply plumose. Grain oblong, free. — Culms tufted, from perennial roots, except No. 1. Leaves smooth, usually flat and soft. (*Πόα*, an ancient Greek name for grass or fodder.)

* *Low and spreading (3' - 6' high) from an annual or biennial root, flaccid: branches of the short panicle single or in pairs.*

1. **P. ánnua**, L. (LOW SPEAR-GRASS.) Culms flattened; panicle often 1-sided; spikelets crowded, very short-pedicelled, 3-7-flowered. — Cultivated and waste grounds, everywhere: but doubtful if indigenous. April - Oct. (Eu.)

** *Low; the culms (6' - 20' long) geniculate-ascending from a running rootstock, rigid, very much flattened: panicle simple and contracted.*

2. **P. compréssa**, L. (WIRE-GRASS.) Pale, as if glaucous; leaves short; panicle dense and narrow, somewhat one-sided (1' - 3' long), the short branches mostly in pairs; spikelets almost sessile, 3-10-flowered, flat. — Dry, mostly sterile soil, in waste places; rarely in woods: probably introduced with other and more valuable grasses: apparently indigenous northward. (Eu.)

*** *Low alpine or alpestrine species, erect, in perennial tufts.*

+ *Soft and flaccid, smooth or nearly so, even to the branches of the panicle: leaves short and flat, short-pointed; ligule elongated.*

3. **P. alpína**, L. Culms rather stout (8' - 14' high); leaves broadly linear, especially those of the culm (1½' - 2' long, 1½'' - 3'' wide); panicle short and broad; spikelets broadly ovate, 3-9-flowered (about 3'' long); lower palet villous on the midrib and margins. — Isle Royale, Lake Superior, *C. G. Loring, Jr., Prof. Porter*, N. Maine? and northward. (Eu.)

4. **P. laxa**, Hænke. Culms slender (4' - 9' high); leaves narrow; panicle somewhat raceme-like, narrow, often one-sided and nodding; spikelets 2-4-flowered, one half smaller. — Alpine mountain-tops of Maine, New Hampshire, and N. New York, and high northward. (Eu.)

+ + *More strict and rigid, roughish, especially the panicle: ligule short.*

5. **P. cæsia**, Smith. Culms 6' - 20° high; leaves narrow, short, soon involute; branches of the panicle 2-5 together, very scabrous; spikelets purplish (or sometimes pale), 2-5-flowered; glumes ovate-lanceolate and taper-pointed; flowers lanceolate, somewhat webby at the base; the lower palet villous on the keel and margins below the middle, its nerves obscure. (*P. áspera*, *Gaudin.*) — N. Wisconsin, *I. A. Lapham*; a form with loose open panicle (*P. nemoralis*, *Ed. 2.*) — Var. STRICTIOR, is 6' - 12' high, with a contracted grayish-purple panicle, of smaller flowers. N. shore of Lake Superior, *C. G. Loring, Jr.*, especially Isle Royale, *Prof. Whitney, &c.*, and northward. (Eu.)

**** *Taller (1° - 3°), meadow or woodland grasses: panicle open.*

+ *Spikelets mostly very numerous and crowded on the rather short rough branches (usually in fives) of the oblong or pyramidal panicle, green, or sometimes violet-tinged: flowers acute, crowded, more or less webbed at base.*

6. **P. serótina**, Ehrhart. (FALSE RED-TOP. FOWL MEADOW-GRASS.) Culms tufted without running rootstocks; leaves narrowly linear, soft and smooth; ligules elongated; spikelets 2-4- (rarely 5-) flowered (1'' - 2'' long), all short-pedicelled in an elongated panicle, often tinged with dull purple; flowers and glumes narrow; lower palet very obscurely nerved. (*P. nemoralis*, *Pursh.* *P. crocàta*, *Michx.*) — Wet meadows and low banks of streams: common, especially northward. July, Aug. — A good grass for moist meadows. (Eu.)

7. **P. pratensis**, L. (GREEN OR COMMON MEADOW-GRASS. KENTUCKY BLUE-GRASS.) Culms sending off copious *running rootstocks* from the base, and the *sheaths smooth*; *ligule short and blunt*; panicle short-pyramidal; *spikelets 3-5-flowered, crowded*, and most of them *almost sessile* on the branches, ovate-lanceolate or ovate; *lower palet 5-nerved, hairy along the margins* as well as the keel. — Common in dry soil: imported for pastures and meadows. Indigenous in mountain regions from N. Penn. northward. May - July. (Eu.)

8. **P. trivialis**, L. (ROUGHISH MEADOW-GRASS.) Culms erect from a somewhat decumbent base, but no distinct running rootstocks; *sheaths and leaves more or less rough*; *ligule oblong, acute*; panicle longer or with the branches more distant; spikelets mostly 3-flowered, broader upwards; *lower palet prominently 5-nerved, naked at the margins*: otherwise nearly as in the preceding. — Moist meadows, &c., July. (Nat. from Eu.)

+ + *Spikelets fewer and more scattered, on slender pedicels: plants soft and smooth, flowering early. (No running rootstocks, except in No. 13.)*

++ *Spikelets small (1" - 2" long), pale green, rather loosely 2-4-flowered: flowers oblong, obtuse: lower palet scarcely scarios-tipped: culm-leaves lance-linear, acute, 1' - 3' long.*

9. **P. sylvéstris**, Gray. *Culm flattish, erect*; branches of the oblong-pyramidal panicle short, numerous, in fives or more; *lower palet villous on the keel for its whole length, and on the margins below the middle, sparingly webbed at the base*. — Rocky woods and meadows, W. New York, Penn. and Virginia to Wisconsin, Kentucky, and southward. June.

10. **P. débilis**, Torr. *Culms terete, weak*; branches of the small panicle few and slender (the lower $1\frac{1}{2}' - 2'$ long to the few spikelets), in pairs and threes; *flowers very obtuse, smooth and glabrous*, except a sparing web at their base. — Rocky woodlands, Rhode Island and N. New York to Wisconsin. May.

++ ++ *Spikelets 2" long, light green: oblong-lanceolate flowers and both glumes acute.*

11. **P. alsódes**, Gray. Leaves rather narrowly linear, acute, the uppermost ($2\frac{1}{2}' - 4'$ long) often sheathing the base of the narrow and loose panicle, the capillary branches of which are appressed when young, and mostly in threes or fours; lower palet very obscurely nerved, villous on the keel below, and with a narrow cobwebby tuft at its base, otherwise glabrous. (*P. nemoralis*, Torr. & Ed. 1: but wholly different from the European species of that name.) — Woods, on hillsides, New England to Penn. and Wisconsin. May, June.

++ ++ ++ *Spikelets larger (3" - 4" long), pale green, rarely purple-tinged, few and scattered at the extremity of the long and capillary branches (mostly in pairs or threes) of the very diffuse panicle: flowers 3-6, loose, oblong and obtuse, as is the larger glume: lower palet conspicuously scarios at the apex, villous below the middle on the keel and margins: culms flattish, smooth.*

12. **P. flexuosa**, Muhl. (not of Wahl.) Culms $1^{\circ} - 3^{\circ}$ high, tufted; its leaves all linear ($2' - 5'$ long) and gradually taper-pointed; panicle very effuse (its branches $2' - 4'$ long to the 4-6-flowered spikelets or first ramification); lower palet prominently nerved, no web at the base. (*P. autumnalis*, Muhl. in Ell. *P. cämpyle*, Schult.) — Dry woods, Virginia, Kentucky, and southward. Feb. - May. — Wrongly confounded with the last, but near it. *P. autumnalis* is an

inappropriate name, and there is now no obstacle to restoring to this species the earlier and unobjectionable (but not descriptive) name of *P. flexuosa*.

13. **P. brevifolia**, Muhl. Culms 1° – $1\frac{1}{2}^{\circ}$ high from *running rootstocks*, 2–3-leaved, the *upper leaves very short* ($\frac{1}{2}'$ – $2'$ long), lanceolate, all abruptly cuspidate-tipped; branches of the short panicle mostly in pairs; spikelets 3–4-flowered; lower palet rather obscurely nerved, cobwebby at the base. (*P. pungens*, Nutt., excl. syn. *Ell.* *P. cuspidata*, Barton. The older and more appropriate name is here restored.)—Rocky or hilly woodlands, Pennsylvania, Virginia, and sparingly westward. April, May.—Culm scarcely surpassing the long root-leaves.

35. ERAGRÓSTIS, Beauv. ERAGROSTIS. (Pl. 10.

Spikelets 2–70-flowered, nearly as in *Poa*, except that the lower palet is but 3- (rarely 1-) nerved, not webby-haired at the base, and deciduous; and the upper one persistent on the rachis after the rest of the flower has fallen.—Culms often branching. Leaves linear, frequently involute, and the ligule or throat of the sheath bearded with long villous hairs. Panicle various. (An early name, probably from *ἔρα*, earth, and *Agrostis*, in allusion to the procumbent habit of the original species.)

* *Prostrate and creeping, much-branched: root annual: spikelets flat, imperfectly dioecious, clustered, almost sessile, in the more fertile plant almost capitate.*

1. **E. réptans**, Nees. Spikelets linear-lanceolate, 10–30-flowered; flowers lance-ovate, acute; leaves short, almost awl-shaped. (*Poa reptans*, Michx.)—Gravelly river-borders: common. Aug.—Flower-branches 2'–5' high.

** *Diffusely spreading, or the flowering culms ascending, low (6'–15' high): spikelets large, densely-flowered, flat, forming a narrow crowded panicle.*

2. **E. POËOÏDES**, Beauv. Lower sheaths often hairy; leaves flat, smooth; spikelets short-pedicelled, lance-linear or oblong-linear, 8–20-flowered, lead-colored (2''–5'' long); flowers ovate, obtuse, the lateral nerves evident. (*Poa Eragrostis*, L.)—Sandy waste places, eastward: scarce. (Nat. from Eu.)

Var. **MEGASTACHYA**. Sheaths mostly glabrous; spikelets larger (3''–10'' long), becoming linear, whitish when old, 10–50-flowered. (*E. megastachya*, Link. *Briza Eragrostis*, L.)—Similar situations, and more common. Aug.—Emits a sharp, unpleasant odor. (Nat. from Eu.)

*** *Erect, or in No. 3–5 diffusely spreading and ascending: panicle open, its branches capillary; the spikelets proportionally small, sometimes minute. (Number of flowers in the spikelet very variable, according to age, &c.)*

+ *Culms slender, branching and decumbent or spreading at the base, from an annual root: leaves narrow, flat, soft: branches of the narrow panicle rather short and thickly-flowered, not bearded in the axils, except sometimes the lowest sparingly.*

3. **E. PILÓSA**, Beauv. Panicle elongated-oblong, with rather erect branches (except at flowering-time); spikelets 5–12-flowered (2''–4'' long, purplish-lead-color), becoming linear, about equalling their pedicels; glumes (small) and lower palet obtuse, the latter broadly ovate, 1-nerved (lateral nerves obsolete). (*P. pilosa*, L. *P. Linkii*, Kunth.)—Sandy or gravelly waste places, S. New England to Illinois, and southward. Aug.—Plant 6'–12' high. (Nat. from Eu.)

4. **E. Fránkii**, Meyer. Much branched, diffuse (3'-8' high); panicle ovate-oblong, rather dense, spreading; spikelets 2-5-flowered (1"-1½" long), on slender pedicels; glumes very acute; lower palet ovate, acute, rather obscurely 3-nerved. (*E. erythrogona*, Nees, from the joints of the culm being mostly reddish.)—Low or sandy ground, S. Pennsylvania to Illinois, and southward. Aug.

5. **E. Púrshii**, (Bernh.?) Schrader. Sparingly branched at the decumbent base, then erect (½°-2° high); panicle elongated, the branches widely spreading, very loose; spikelets 5-18-flowered, oblong-lanceolate, at length linear (2"-4½" long), mostly much shorter than their capillary pedicels; glumes and lower palet ovate and acute, or the latter acutish, 3-nerved. (*Poa tenella*? Pursh. *P. Caroliniána*, Spreng. *P. pectinacea* of authors, not of Michx.)—Sandy or sterile open grounds, New Jersey to Virginia, and southward.

← ← Culms simple or branching only at the very base, firm, erect from an annual or perennial root, mostly forming thick tufts: leaves very long: panicle very large, compound, often longer than the culm, with elongated and loosely-flowered branches, their axils often bearded. (Doubtful perennials, or No. 7 annual.)

6. **E. ténuis**, Gray. Panicle virgately elongated (1°-2½° long), very loose, the spreading branches bearded in some of the lower axils, their remote divisions and long diverging pedicels capillary; spikelets 2-6- (sometimes 7-12-) flowered, pale or greenish; glumes lanceolate or awl-shaped, very acute (1½"-2" long), membranaceous, as are the oblong-lanceolate acute flowers; lower palet distinctly 3-nerved; the upper ciliate-scabrous. (*Poa tenuis*, Ell. *P. capillaris*, Michx. *P. trichòdes*, Nutt. *E. Géyeri*, Steud.)—Sandy soil, Illinois, Virginia? and southward. Aug.-Oct.—Leaves rather rigid, 1½°-2° long, glabrous or sparingly hairy: the sheaths hairy or glabrous; the throat strongly bearded. Flowers much larger than in the next, fully 1½" long.

7. **E. capillàris**, Nees. Panicle widely expanding, usually much longer than the culm, its spreading branches (mostly naked in the axils) and long diverging pedicels capillary; spikelets rather terete, very small, 2-4-flowered, greenish or purplish; glumes and flowers ovate, acute (less than 1" long); lower palet obscurely 3-nerved, scarcely keeled; the upper rough-ciliate. (*Poa capillaris*, L. *P. hirsuta*, Michx.)—Sandy dry soil and fields: common, especially southward. Aug., Sept.—Leaves and sheaths very hairy, or nearly glabrous; the former about 1° long, not rigid. Panicle 1°-2° long, soon diffuse.

8. **E. pectinàcea**, Gray. Panicle widely diffuse, its rigid divergent main branches bearded in the axils; the capillary pedicels more or less appressed on the secondary branches; spikelets flat, 5-15-flowered, becoming linear, purple or purplish; glumes and flowers ovate or oblong-ovate, acutish; lower palet strongly 3-nerved; the upper hirsute-ciliate. (*Poa pectinacea*, Michx., ex char. *P. hirsuta*, of Amer. authors. *E. Uniònis* & cognata Steud.?)—Leaves long, rigid, mostly hairy, the sheaths especially so.—Var. **SPECTABILIS**. Leaves and sheaths mostly glabrous; branches of the panicle (the lower reflexed with age) and pedicels shorter; spikelets rather larger. (*E. spectabilis*, Ed. 1. *Poa spectabilis*, Pursh.)—Sandy dry ground, from E. Mass. near the coast, and from Ohio and Illinois southward. Aug.-Oct.—Plant 1°-3° high. Spikelets 2"-3" long, 1" wide, closely flowered.

36. BRIZA, L. QUAKING GRASS. (Pl. 10.)

Spikelets many-flowered, ovate or heart-shaped, flattish-tumid; the flowers closely imbricated. Glumes roundish, unequal (purple). Lower palet roundish and entire, flattened parallel with the glumes, ventricose on the back, heart-shaped at the base, papery-membranaceous and becoming dry, scarious-margined, obscurely many-nerved; the upper palet much smaller, ovate, flat. Stamens 3. Stigmas branched-plumose. Grain flattened parallel with the palets, adhering to the upper one. — Leaves flat. Panicle loose, diffuse, with the large and showy spikelets often drooping on delicate pedicels (whence the name, an ancient Greek appellation for some kind of grain, from *βρίζω*, to *slumber* (Linn.), or *βρίθω*, to *bend downwards*.)

1. **B. MÈDIA**, L. Panicle erect, the branches spreading; spikelets 5–9-flowered (3" long); glumes shorter than the lower flowers; root perennial. — Pastures: sparingly eastward. June. (Adv. from Eu.)

37. FESTÛCA, L. FESCUE-GRASS. (Pl. 10.)

Spikelets 3–many-flowered, paniced or racemose; the flowers not webby at the base. Glumes unequal, mostly keeled. Palets chartaceous or almost coriaceous, roundish (not keeled) on the back, more or less 3–5-nerved, acute, pointed, or often bristle-awned from the tip, rarely blunt; the upper mostly adhering at maturity to the enclosed grain. Stamens 1–3. — Flowers, and often the leaves, rather dry and harsh. (An ancient Latin name.)

* *Flowers awl-shaped, bristle-pointed or awned from the tip: panicle contracted.*

+ *Annuals or biennials, slender, 5'–18' high: leaves convolute-bristle-form.*

1. **F. MYÛRUS**, L. Panicle spike-like, one-sided; spikelets about 5-flowered; glumes very unequal; *awn much longer than the palet*, fully 6" in length; stamen 1. — Dry fields, New Jersey, S. Penn., and southward. July. (Nat. from Eu.)

2. **F. tenella**, Willd. Panicle spike-like, one-sided, or more compound and open; spikelets 7–13-flowered; *awn 1"–3" long, shorter than or equalling the palet*; stamens 2. — Dry, sterile soil, especially southward. June, July.

+ + *Perennial, tufted, 6'–24' high: stamens 3.*

3. **F. ovina**, L. (SHEEP'S FESCUE.) Panicle somewhat one-sided, short, usually more or less compound, open in flowering; spikelets 3–8-flowered; awn not more than half the length of the flower, often much shorter or almost wanting. — Indigenous in Northern New England, Lake Superior, and northward: naturalized farther south as a pasture grass. June. — Varies greatly. — Var. **VIVIPARA** (which with us has running rootstocks), a state with the spikelets partially converted into leafy shoots, is found on the alpine summits of the White Mountains of New Hampshire, and high northward. — Var. **DURIÛSCULA**, (*F. duriuscula*, L.) is a tall form, with spikelets rather larger, usually in a more compound panicle; culm-leaves often flat or less convolute, and the lower with their sheaths either smooth or hairy. New England to Virginia; nat., and indigenous northward. — Var. **RÛBRA** (*F. rubra*, L.) has running rootstocks and forms looser tufts; the leaves often reddish and pubescent above. Naturalized eastward: wild, Lake Superior, *Dr. Robbins*, and northward. (Eu.)

* * *Flowers oblong or lanceolate, awnless or nearly so* ($1\frac{1}{2}''$ – $4''$ long); *grain often free!* (*Root perennial: culms mostly tall: leaves flat.*)

4. **F. ELÀTIOR**, L. (TALLER OR MEADOW FESCUE.) *Panicle narrow, contracted before and after flowering, erect, with short branches; spikelets crowded, 5–10-flowered; the flowers rather remote, oblong-lanceolate; lower palet 5-nerved, scarious-margined, blunt, acute, or rarely with a distinct but very short awn.*—The type is large, 3° – 4° high; spikelets about $6''$ long, in an ample and compound panicle. Rich grass-land.—Var. **PRATÈNSIS**. (*F. pratensis*, *Huds.*) is lower (1° – 3° high), with a simpler or close panicle, of smaller or narrower spikelets; and abounds in grass-lands. June–Aug. (*Nat. from Eu.*)

5. **F. NŪTANS**, Willd. *Panicle of several long and slender spreading branches, mostly in pairs, drooping when old, rough, naked below, bearing near their extremity a few ovate 3–5-flowered spikelets* ($3''$ long) on pretty long pedicels; *flowers ovate-oblong, rather obtuse, close together, coriaceous, smooth, very obscurely 5-nerved.*—Rocky woods and copses. July.—Culm 2° – 4° high, naked above: leaves broadly linear, taper-pointed, dark green, often rather hairy.

38. BRŌMUS, L. BROME-GRASS. (Pl. 10.)

Spikelets 5–many-flowered, paniced. Glumes unequal, membranaceous; the lower 1–5-, the upper 3–9-nerved. Lower palet either convex on the back or compressed-keeled, 5–9-nerved, awned or bristle-pointed from below the mostly 2-cleft tip: upper palet at length adhering to the groove of the oblong or linear grain. Stamens 3. Styles attached below the apex of the ovary.—Coarse Grasses, with large spikelets, at length drooping, on pedicels thickened at the apex. (*An ancient name for the Oat, from βρόμος, food.*)

§ 1. *Lower palet convex on the back; the flowers imbricated over one another before expansion: lower glume 3–5-nerved, the upper 5–9-nerved.*

* *Annuals or biennials, weeds of cultivation, introduced into grain-fields, or rarely in waste grounds, probably all derived from the European B. arvensis, L.*

1. **B. SECÁLINUS**, L. (CHEAT OR CHESS.) *Panicle spreading, even in fruit, the drooping peduncles little branched; spikelets oblong-ovate, turgid, smooth, of 8–10 rather distant flowers; lower palet rather longer than the upper, short-awned or awnless; sheaths nearly glabrous.*—Too common in wheat-fields. June, July. (*Adv. from Eu.*)

2. **B. RACEMŌSUS**, L. (UPRIGHT CHESS.) *Panicle erect, simple, rather narrow, contracted in fruit; flowers closer, more imbricated; lower palet decidedly exceeding the upper, bearing an awn of its own length; culm more slender; sheaths sometimes hairy: otherwise nearly as in the last, for which it is often mistaken in this country.* (*Adv. from Eu.*)

3. **B. MÓLLIS**, L. (SOFT CHESS.) *Panicle erect, closely contracted in fruit; spikelets conical-ovate, somewhat flattened; the flowers closely imbricated, downy (as also the leaves, &c.); lower palet acute, long-awned.*—Wheat-fields, New York to Virginia: scarce. June. (*Adv. from Eu.*)

* * *Perennial: indigenous.* (*Lower glume strongly 3-nerved, the upper 5-nerved.*)

4. **B. KÁLMII**, Gray. (WILD CHESS.) *Panicle simple, small* ($3'$ – $4'$ long); spikelets drooping on capillary peduncles, closely 7–12-flowered, densely silky

all over; awn only one third the length of the lance-oblong flower; lower palet 7-9-nerved, much longer and larger than the upper; culm slender ($1\frac{1}{2}^{\circ}$ - 3° high); leaves and sheaths conspicuously or sparingly hairy. (*B. ciliatus*, *Muhl.* *B. purgans*, *Torr. Fl. N. Y.*)—Dry ground: common northward. June, July.—This is in the herbarium of Linnæus under the name of *B. ciliatus*, but is not the plant he described; thence has arisen much confusion.

§ 2. *SCHEDÓNORUS*, Beauv., Fries. *Lower palet somewhat convex, but keeled on the back and laterally more or less compressed, at least above: flowers soon separating from each other: lower glume 1-nerved, the upper 3-nerved, or with an obscure additional pair.*

* *Perennial, tall (3°-5° high): flowers oblong or lanceolate.*

5. *B. ciliatus*, L. Panicle compound, very loose, the elongated branches at length divergent, drooping; spikelets 7-12-flowered; lower palet tipped with an awn half to three fourths its length, silky with appressed hairs near the margins, at least below (or rarely naked), smooth or smoothish on the back (*B. Canadensis*, *Michx.* *B. pubescens*, *Muhl.*);—or, in var. *PURGANS* (*B. purgans*, *L.*!), clothed all over with short and fine appressed hairs.—River-banks and moist woodlands: common. July, Aug.—Culm and large leaves (3"-6" wide) smooth or somewhat hairy; the sheaths in the larger forms often hairy or densely downy near the top. Variable, comprising several forms, including both the Linnæan species; for which the present name is preferable to the inapplicable *purgans*, taken from Feuillée's South American species.

6. *B. ASPER*, L. Culm slender and panicle smaller; spikelets 5-9-flowered; lower palet linear-lanceolate, scarcely keeled, hairy near the margins, rather longer than the awn; sheaths and lower leaves hairy or downy.—Bethel, Maine, in fields along the river-bank, *W. Boott.* (Nat. from Eu.)

** *Annual or biennial (10'-30' high): flowers slender.*

7. *B. STÉRILIS*, L. Panicle open; spikelets of 5-9 rather distant and 7-nerved roughish linear-awl-shaped long-awned flowers (awn 1' long); leaves rather downy.—Waste places and river-banks, E. Massachusetts, New York, and Pennsylvania: rare. June. (Nat. from Eu.)

39. UNIOLA, L. SPIKE-GRASS. (Pl. 11.)

Spikelets closely many-flowered, very flat and 2-edged; one or more of the lowest flowers sterile (neutral) and consisting of a single palet. Glumes lanceolate, compressed-keeled. Lower palet coriaceous-membranaceous, strongly laterally compressed and keeled, striate-nerved, usually acute or pointed, entire, enclosing the much smaller compressed 2-keeled upper one and the free laterally flattened smooth grain. Stamen 1 (or in *U. paniculata* 3).—Upright smooth perennials, growing in tufts from strong creeping rootstocks, with broad leaves and large spikelets in an open or spiked panicle. (Ancient name of some plant, a diminutive of *unio*, unity.)

* *Spikelets large ($\frac{1}{2}$ '-2' long), ovate or oblong, 9-30-flowered: panicle open.*

1. *U. paniculata*, L. *Leaves narrow, when dry convolute; spikelets ovate, short-pedicelled; flowers glabrous, bluntish, several of the lower sterile; the fer-*

tile with 3 stamens; culm and panicle elongated (4°-8° high). — Sand-hills on the sea-shore, S. Virginia and southward.

2. *U. latifolia*, Michx. *Leaves broad and flat (nearly 1' wide); spikelets at length oblong, hanging on long pedicels; flowers acute, ciliate on the keel, all but the lowest perfect and monandrous.* — Shaded rich hillsides, S. Pennsylvania to Illinois and southward. Aug. — Culm 2°-4° high: panicle loose.

* * *Spikelets small: panicle contracted and wand-like: perfect flowers long-pointed.*

3. *U. grácilis*, Michx. *Spikelets short-pedicelled (2"-3" long), broadly wedge-shaped, acute at the base, 4-8-flowered; the flowers ovate and divergently beaked, long, the lowest one neutral.* — Sandy soil, from Long Island to Virginia, near the coast, and southward. Aug. — Culm 3° high, slender.

40. PHRAGMITES, Trin. REED. (Pl. 11.)

Spikelets 3-7-flowered; the flowers rather distant, silky-villous at their base, and with a conspicuous silky-bearded rhachis, all perfect and 3-androus, except the lowest, which is either neutral or with 1-3 stamens, and naked. Glumes membranaceous, shorter than the flowers, lanceolate, keeled, sharp-pointed, very unequal. Palets membranaceous, slender; the lower narrowly awl-shaped, thrice the length of the upper. Squamulæ 2, large. Styles long. Grain free. — Tall and stout perennials, with numerous broad leaves, and a large terminal panicle. (*φραμίτες*, growing in hedges, which this aquatic Grass does not.)

1. *P. communis*, Trin. Panicle loose, nodding; spikelets 3-5-flowered; flowers equalling the wool. (*Arundo*, *L.*) — Edges of ponds. Sept. — Looks like Broom-Corn at a distance, 5°-12° high: leaves 2' wide. (Eu.)

41. ARUNDINÀRIA, Michx. CANE. (Pl. 11.)

Spikelets flattened, 5-14-flowered; the flowers somewhat separated on the jointed rhachis. Glumes very small, membranaceous, the upper one larger. Palets herbaceous or somewhat membranaceous; the lower convex on the back, many-nerved, tapering into a mucronate point or bristle. Squamulæ 3, longer than the ovary. Stamens 3. Grain oblong, free. — Arborescent or shrubby Grasses, simple or with fasciated branches, and with large spikelets in panicles or racemes; the flowers polygamous. (Name from *arundo*, a reed.)

1. *A. macrosperma*, Michx. (LARGE CANE.) Culms arborescent, 10°-20° high, rigid, simple the first year, branching the second, afterwards at indefinite periods fruiting, and soon after decaying; leaves lanceolate (1'-2' wide), smoothish; panicle lateral, composed of few simple racemes; spikelets purple, erect; lower palet lance-ovate, pubescent, fringed (8" long). *Chapman*, under *A. gigantea*. — River-banks, S. Virginia? Kentucky? and southward, forming canebrakes.

2. *A. tecta*, Muhl. (SMALL CANE.) Culms slender, 2°-10° high, branching; leaves linear-lanceolate (9"-1' wide), roughish, the sheaths bearded at the throat; spikelets solitary or in a simple raceme at the summit of the branches, or frequently on leafless radical culms; lower palet (6" long) ovate-lanceolate, smooth, fringed on the margins. *Chapman*. (*Arundo tecta*, *Walt.*) — Swamps and moist soil, Virginia, Illinois, and southward. April.

42. LEPTÛRUS, R. Br. LEPTURUS. (Pl. 11.)

Spikelets solitary on each joint of the filiform rhachis, and partly immersed in the excavation, 1-2-flowered. Glumes 1-2, including the 2 thin pointless paleas. Stamens 3. Grain free, oblong-linear, cylindrical. — Low and branching, often procumbent Grasses, chiefly annuals, with narrow leaves and slender spikes (whence the name, from λεπτός, *slender*, and οὐρά, *tail*.)

1. **L. paniculatus**, Nutt. Stem slender (6'-20' long), naked and curved above, bearing 3-9 racemously disposed thread-like and triangular spikes; glumes 2, transverse. — Open grounds and salt licks, Illinois (*Mead*), and westward. Aug.

43. LÓLIUM, L. DARNEL. (Pl. 11.)

Spikelets many-flowered, solitary on each joint of the continuous rhachis, placed edgewise; the glume, except in the terminal spikelet, only one (the upper) and external: — otherwise nearly as in Triticum. (Ancient Latin name.)

1. **L. PERÉNNE**, L. (COMMON DARNEL, RAY-OR RYE-GRASS.) Root perennial, *glume shorter than the spikelet; flowers 8-15, awnless or sometimes short-awned.* — Fields and lots: eastward. June. — A pretty good pasture-grass. (Nat. from Eu.)

2. **L. TEMULÉNTUM**, L. (BEARDED DARNEL.) Root annual; culm taller, *glume fully equalling the 5-7-flowered spikelet; awn longer than the flower (½ long).* — Grain-fields: rare. — Grain noxious; almost the only instance of the kind among Grasses. (Adv. from Eu.)

44. TRÍTICUM, L. WHEAT. (Pl. 11.)

Spikelets 3-several-flowered, single at each joint, and placed with the side against the rhachis. Glumes transverse (i. e. right and left), nearly equal and opposite, herbaceous, nerved. Lower palet very like the glumes, convex on the back, pointed or awned from the tip: the upper flattened, bristly-ciliate on the nerves, free, or adherent to the groove of the grain. Stamens 3. (The classical name, probably from *tritius*, beaten, because the grain is threshed out of the spikes.) — The true species are annuals, with the glumes ovate-oblong, turgid and boat-shaped, as in common *Wheat* (*T. vulgare*). Others are perennial, with nearly lanceolate glumes, and 2-ranked spikes, never furnishing bread-corn (§ AGROPÿRON, Gærtn.); to which the following belong. Flowering in summer.

* *Multiplying by long running rootstocks: awn shorter than the flower or none.*

1. **T. répens**, L. (COUCH-, QUITCH, OR QUICK-GRASS.) *Spikelets 4-8-flowered, glabrous or nearly so; glumes 5-7-nerved; rhachis glabrous, but rough on the edges; awns when present straight; leaves flat and often roughish or pubescent above.* — Nat. in cultivated grounds, fields, &c., and very troublesome; indigenous northwestward. — Varies greatly. A tall form, rather bright green (Var. *NEMORALE*, Andersson) bears awns nearly as long as the palet. A lower form is glaucous and mostly awnless or nearly so. A maritime form (nearly Var. *INTERMÉDIUM*, Fries, and approaching *T. laxum*, *Fries*, and *T. acutum*, *DC.*), collected by *Prof. Tuckerman* on the coast of Maine, is glaucous, rigid, with

at length somewhat involute leaves, crowded spikelets, blunt and rigid glumes, and pointless or mucronate-tipped palets; the rhachis not disarticulating as in *T. junceum*, *L.* (Eu.)

2. **T. dasystachyum**, Gray. Perhaps a remarkable variety of the last, but the glaucous leaves are narrow and often involute, and the 5-9-flowered spikelets densely downy-hairy all over. (*T. repens*, var. *dasystachyum*, *Hook.*)—Sandy shores of Lakes Huron and Superior, and northward. Aug.

* * *No obvious running rootstocks, glabrous, or the flat and roughish leaves sometimes hairy above: glumes as well as flowers mostly awned or awn-pointed.*

3. **T. violaceum**, Hornemann. *Spike dense, strict and rigid*, usually tinged with violet or purple; spikelets 3-5-flowered; *glumes conspicuously 5-nerved; awns straight*, varying from half to nearly twice the length of the pallet. (*T. Richardsonii*, *Schrader.* *T. pauciflorum*, *Schweinitz.*)—Mountains of Penn. (*Porter*); in the alpine region of the White Mountains (*Tuckerman*), Wisconsin (*Lapham*), and northward.—Intermediate in character between the last and the next. (Eu.)

4. **T. caninum**, *L.* (AWNED WHEAT-GRASS.) *Spike usually more or less nodding*, at least in fruit, rather dense (3'-6' long); spikelets 3-5-flowered; *glumes 3-5-nerved; awns mostly somewhat bent or spreading, fully twice the length of the pallet.*—Sparingly naturalized in cult. ground and meadows. Indigenous along our northern borders, in a slender form, which approaches the northwestern *T. divergens*, *Nes.* or *T. ægilopoides*, *Turczaninow*, which has narrow and convolute leaves, 5-7-nerved glumes, and widely diverging awns. (Eu.)

• 45. HÓRDEUM, L. BARLEY. (Pl. 11.)

Spikelets 1-flowered, with an awl-shaped rudiment on the inner side, three at each joint of the rhachis; but the lateral ones usually imperfect or abortive, and short-stalked. Glumes side by side in front of the spikelets, 6 in number, forming a kind of involucre, slender and awn-pointed or bristle-form. Palets herbaceous, the lower (anterior) convex, long-awned from the apex. Stamens 3. Grain oblong, commonly adhering to the palets. Spike often separating into joints. Ours are annuals or biennials, or scarcely perennial. (The ancient Latin name.)

1. **H. jubatum**, *L.* (SQUIRREL-TAIL GRASS.) Low; lateral flowers abortive, on a short pedicel, short-awned; the perfect flower bearing a capillary awn (2' long) about the length of the similar capillary glumes, all spreading.—Sandy sea-shore, upper Great Lakes, and westward. June.

2. **H. pratense**, *Huds.* Low (6'-18' high); lateral flowers imperfect or neutral, awnless or merely pointed; perfect flower with awn as long as those of the glumes (3''-6''); spike linear, 1'-2' long. (*H. pusillum*, *Nutt.* and former ed.)—Plains, especially in saline soil, Ohio to Illinois and westward: also sparingly introduced, Virginia and southward along the coast. May, June.

46. ELYMUS, L. LYME-GRASS. WILD RYE. (Pl. 11.)

Spikelets 2-4 at each joint of the rhachis, all fertile and alike, sessile, each 1-7-flowered. Glumes conspicuous, nearly side by side in front of the spikelets,

2 for each spikelet, forming an involucre to the cluster. Palets coriaceous; the lower rounded on the back, acute or awned at the apex. Grain adherent to the involving palets (whence the name, an ancient one for some grain, from ἐλάω, to roll up).

* *Glumes and flowers firm or rigid, both or only the latter awned: spikelets 1-5-flowered: perennials, with slender culms and rather harsh and broad flat leaves.*

← *Spike large and stout.*

1. **E. Virgīnicus**, L. *Spike rigidly upright, dense (2'-3' long, 6" thick), on a short peduncle usually included in the sheath; spikelets 2-3 together, 2-3-flowered, smooth, rather short-awned, about the length of the thickened strongly-nerved and bristle-pointed lanceolate glumes.*—River-banks: common. Aug.—Culm stout, 2°-3° high.

2. **E. Canadēnsis**, L. *Spike soon nodding (5'-9' long), on an exerted peduncle; spikelets mostly in pairs, of 3-5 long-awned rough or rough-hairy flowers; the awl-shaped glumes tipped with shorter awns.* (E. Philadelphicus, L.!)—Var. GLAUCIFOLIUS (E. glaucifolius, Muhl.) is pale or glaucous throughout, the flowers with more spreading awns (1½' long).—River-banks: common.

← *Spike more slender, as also the culm.*

3. **E. Sibiricus**, L. *Glabrous; spike wand-like (2'-6' long, about 3" thick), often somewhat nodding; spikelets in pairs, 3-6-flowered; glumes linear-lanceolate, 3-5-nerved, short-awned, shorter than the flowers, which are rather short-awned.*—South shore of Lake Superior (Porter), and northwestward. (Eu.)

4. **E. striātus**, Willd. *More or less pubescent; spike dense and thickish (2'-4' long), upright or slightly nodding; spikelets mostly in pairs, 1-2- (or rarely 3-) flowered, minutely bristly-hairy; glumes awl-shaped, bristle-awned, 1-3-nerved, about thrice the length of the flowers exclusive of the capillary awn (which is 1' long).*—Var. VILLOSUS (E. villosus, Muhl.!) has very hairy flowers and glumes, and villous sheaths.—Rocky woods and banks. July, Aug.—Palets only 3" long.

* * *Glumes and palets awnless and soft in texture: reed-like perennials.*

5. **E. mollis**, Trin. (not of R. Br.) *Culm (3° high) velvety at top; spike thick, erect (8' long); spikelets 2 or 3 at each joint, 5-8-flowered; the lanceolate pointed 5-7-nerved glumes (1' long) and the pointed palets soft-villous; rhachis of the spikelets separating into joints.*—Shore of the Great Lakes, Maine, and northward. (Near E. arenarius.)

47. GYMNOSTICHUM, Schreb. BOTTLE-BRUSH GRASS. (Pl. 11.)

Spikelets 2-3 or sometimes solitary on each joint of the rhachis, raised on a very short callous pedicel, loosely 2-4-flowered (when solitary flatwise on the rhachis). Glumes none! or small awn-like deciduous rudiments (whence the name, from γυμνός, naked, and στῖχος, rank). Otherwise nearly as in Elymus.

1. **G. Hystrix**, Schreb. *Spike loose (3'-6' long); the spreading spikelets 2-3 together, early deciduous; flowers smoothish or often rough-hairy, tipped with an awn thrice their length (1' long); leaves and sheaths smoothish; culm 3°-4° high; root perennial.* (Elymus Hystrix, L.)—Moist woodlands. July, Aug.

48. **DANTHŌNIA**, DC. WILD OAT-GRASS. (Pl. 12.)

Lower palet (oblong or ovate, rounded-cylindrical, 7-9-nerved) bearing between the sharp-pointed or awn-like teeth of the tip an awn composed of the 3 middle nerves, which is flattish and spirally twisting at the base: otherwise nearly as in *Avena*. Glumes longer than the imbricated flowers. Ours perennials, 1°-2° high, with narrow and soon involute leaves, hairy sheaths bearded at the throat, and a small simple panicle or raceme of about 7-flowered spikelets. (Named for *Danthoine*, a French botanist.)

1. **D. spicata**, Beauv. Culms tufted, low; leaves short, very narrow; spikelets few, 3"-5" long; *lower palet loosely hairy, its teeth short and pointless*. — Dry and sterile or rocky soil. June-Aug.

2. **D. sericea**, Nutt. Taller and not tufted (1°-3° high); leaves larger; spikelets more numerous and paniced, 6"-9" long; *lower palet very silky-villous, tipped with slender awn-pointed teeth*. — Dry or moist sandy soil, Southern Massachusetts (*Dr. Robbins*), New Jersey (*C. E. Smith, C. F. Parker*), and southward: rare. June.

49. **AVÈNA**, L. OAT. (Pl. 12.)

Spikelets 2-many-flowered, paniced; the flowers herbaceo-chartaceous, or becoming harder, of firmer texture than the large and mostly unequal glumes; the uppermost imperfect. Lower palet rounded on the back, mostly 5-11-nerved, bearing a long usually bent or twisted awn on the back or below the acutely 2-cleft tip proceeding from the midnerve only. Stamens 3. Grain oblong-linear, grooved on one side, usually hairy at least at the top, free, but invested by the upper palet. (The classical Latin name.) — The COMMON OAT (*A. SATIVA*, L.) represents the large-flowered annual species of the Old World. The following are smaller-flowered, indigenous perennials.

1. **A. striata**, Michx. *Glabrous and smooth throughout, slender* (1°-2° high); leaves narrow; ligule short, truncate; panicle simple, loose; spikelets (6" long) on capillary pedicels, 3-6-flowered, much exceeding the scarious-margined purple acute glumes; the *lower glume 1-, upper 3-nerved*; rhachis smooth; *flowers short-bearded at the base*; lower palet 7-nerved, much longer than the ciliate-fringed upper one (4" long), mostly shorter than its soon bent or divergent awn, which rises just below the tapering very sharply cuspidate 2-cleft tip. (*Trisetum purpurascens, Torr.*) — Rocky, shaded hills, N. New England, New York, and northwestward. June.

2. **A. Smithii**, T. C. Porter, n. sp. Taller (2½°-4½° high), rather stout; leaves broadly linear (3"-6" wide) and taper-pointed, flat, and with the sheaths and culm *retroscely scabrous*; ligule elongated, acute; panicle larger (6'-12' long), the few branches at length spreading; glumes slightly purplish, scabrous on the *nerves*, of which there are 3 in the lower and 5 in the upper; rhachis minutely hispid; *flowers* (3-5) *not hairy-tufted* at the base; awn one third or half the length of the 7-nerved palet, straight. — Isle Royale, Keweenaw Point, Lake Superior, *Robbins*. Woods near Sault Ste Marie, *C. E. Smith*, for whom the species is named. April, May. — To be compared with the Siberian *A. callosa, Turczaninow*, which was referred to *A. striata* by Trinius.

50. TRISÈTUM, Persoon. TRISETUM. (Pl. 12.)

Spikelets 2-several-flowered, often in a contracted panicle; the lower palet compressed-keeled, of about the same membranaceous texture as the glumes, bearing a bent or flexuous (rarely twisted) awn below the sharply 2-toothed or 2-pointed apex (whence the name, from *tris*, three, and *seta*, a bristle): otherwise nearly as in *Avena*. Ours are perennials.

1. **T. subspicatum**, Beauv., var. **molle**. *Minutely soft-downy; panicle dense, much contracted, oblong or linear (2' - 3' long); glumes about the length of the 2-3 smooth flowers; awn diverging, much exerted. (Avena mollis, Michx.)*—Mountains and rocky river-banks, N. New England to Wisconsin, and northward. July. — About 1° high: leaves flat, short. (Eu.)

2. **T. palustre**, Torr. *Smooth; panicle rather long and narrow (5' long), loose, the branches capillary; spikelets flat (3" long); glumes shorter than the two smooth lanceolate flowers, of which the upper is on a slightly naked joint of the rachis, and bears a slender spreading or bent awn next the short 2-pointed tip, while the lower one is commonly awnless or only mucronate-pointed. (Avena palustris, Michx. Aira pallens, Muhl.)*—Low grounds, Southern New York to Illinois, and southward. June. — Culm slender, 2°-3° high: leaves flat, short. Spikelets yellowish-white, tinged with green.

51. AIRA, L. HAIR-GRASS. (Pl. 12.)

Spikelets small, in an open diffuse panicle, of 2 perfect flowers and often with the pedicel or rudiment of a third, all usually shorter than the membranaceous keeled glumes, and hairy at the base; the upper remotish. Lower palet thin and scarious, 2-cleft or else truncate and mostly denticulate or eroded at the summit, bearing a slender bent or straight awn on its back, commonly near its base. Stamens 3. Styles plumose to the base. Ovary glabrous. Grain oblong. (An ancient Greek name for Darnel.)

§ 1. **DESCHAMPSIA**, Beauv. *Lower palet delicately 3-5-nerved, eroded or toothed at the truncate summit; the awn attached mostly a little above the base: grain not grooved, free: glumes about equalling the flowers: root perennial.*

1. **A. flexuosa**, L. (COMMON HAIR-GRASS.) Culms slender, nearly naked (1°-2° high) above the small tufts of *involute bristle-form root-leaves* (1'-6' long); branches of the small spreading panicle capillary; *awn longer than the palet, at length bent and twisted.* — Dry places: common. June. (Eu.)

2. **A. cæspitosa**, L. Culm tufted (2°-4° high); *leaves flat, linear; panicle pyramidal or oblong (6' long); awn straight, barely equalling the palet.* — Shores of lakes and streams: common northward. June, July. (Eu.)

§ 2. **AIROPSIS**, Desv. *Spikelets very small, of 2 closely approximate flowers, and with no rudiment of a third: lower palet of firmer texture, obscurely nerved, acutely 2-cleft at the apex: grain grooved, adnate: low annuals, with short and setaceous leaves.*

3. **A. PRÆCOX**, L. Culms tufted, 3'-4' high; branches of the *small and dense panicle* appressed; awn from below the middle of the flower. (*Avena præcox, Beauv.*) — Sandy fields, New Jersey to Virginia: rare. (Nat. from Eu.)

4. **A. CARYOPHYLLEA**, L. Culms 5'–10' high, bearing a *very diffuse panicle* of purplish and at length *silvery scarious spikelets*. — Dry fields, Nantucket: also Newcastle, Delaware, *W. M. Canby*. (Nat. from Eu.)

§ 3. **VAHLÓDEA**, Fries. *Glumes boat-shaped, longer than the flowers: lower palet almost coriaceous, nerveless, its truncate-obtuse tip mostly entire; the awn borne at or above the middle: grain grooved, flattish, free: alpine perennial.*

5. **A. atropurpúrea**, Wahl. Culms 8'–15' high, weak; leaves flat, rather wide; panicle of few spreading branches; awn stout, twice the length of the palet. — Alpine tops of the White Mountains, and those of Northern New York. Aug. (Eu.)

52. **ARRHENATHERUM**, Beauv. OAT-GRASS. (Pl. 12.)

Spikelets open-panicled, 2-flowered, with the rudiment of a third flower; the middle flower perfect, its lower palet barely bristle-pointed from near the tip; the lowest flower staminate only, bearing a long bent awn below the middle of the back (whence the name, from ἄρρην, *masculine*, and ἀθήρ, *awn*): — otherwise as in *Avena*, of which it is only a peculiar modification.

1. **A. AVENACEUM**, Beauv. Root perennial; culm 2°–4° high; leaves broad, flat; panicle elongated; glumes scarious, very unequal. (*Avena elatior*, L.) — Meadows and lots: absurdly called *Grass of the Andes*. May–July. (Nat. from Eu.)

53. **HÓLCUS**, L. (partly). MEADOW SOFT-GRASS. (Pl. 13.)

Spikelets crowded in an open panicle, 2-flowered, jointed with the pedicels; the boat-shaped membranaceous glumes enclosing and much exceeding the remotish flowers. Lower flower perfect, its papery or thin-coriaceous lower palet awnless and pointless; the upper flower staminate, otherwise similar, but bearing a stout bent awn below the apex. Stamens 3. Styles plumose to the base. Grain free. (An ancient name, from ἄλκος, *attractive*, of obscure application.)

1. **H. LANATUS**, L. (VELVET-GRASS.) Perennial, soft-downy and pale; panicle oblong; upper glume mucronate-awned under the apex; awn of the staminate flower curved. — Moist meadows. June. (Nat. from Eu.)

54. **HIERÓCHLOA**, Gmelin. HOLY GRASS. (Pl. 13.)

Spikelets 3-flowered, open-panicled; the flowers all with 2 palets; the two lower (lateral) flowers staminate only, 3-androus, sessile, often awned on the middle of the back or near the tip; the uppermost (middle) one perfect, short-pedicelled, scarcely as long as the others, 2-androus, awnless. Glumes equalling or exceeding the spikelet, scarious: palets chartaceous. — Perennials: leaves flat. (Name composed of ἱερός, *sacred*, and χλόα, *grass*; these sweet-scented Grasses being strewn before the church-doors on saints' days, in the North of Europe.)

1. **H. boreális**, Roem. & Schultes. (VANILLA OR SENECA GRASS.) Panicle somewhat one-sided, pyramidal (2'–5' long); peduncles smooth; staminate flowers with the lower palet mucronate or bristle-pointed at or near the tip;

rootstock creeping. (*Holcus odoratus*, L.) — Moist meadows, chiefly northward near the coast and along the Great Lakes. May. — Culm 1°-2° high, with short, lanceolate leaves. Spikelets chestnut-color; the staminate flowers strongly hairy-fringed on the margins, and the fertile one at the tip. (Eu.)

2. *H. alpina*, Roem. & Schultes. Panicle contracted (1'-2' long); one of the staminate flowers barely pointed or short-awned near the tip, the other long-awned from below the middle; lowest leaves very narrow. — Alpine mountain-tops, New England, New York, and northward. July. (Eu.)

55. ANTHOXÁNTHUM, L. SWEET VERNAL-GRASS. (Pl. 13.)

Spikelets spiked-panicled, really 3-flowered; but the lateral flowers neutral, consisting merely of one palet which is hairy on the outside and awned on the back: the central (terminal) flower perfect, small, of 2 awnless chartaceous palets, 2-androus. Glumes very thin, acute, keeled; the upper about as long as the flowers, twice the length of the lower. Squamulæ none. Grain ovate, adherent. (Name compounded of *ἄνθος*, *flower*, and *ἄνωον*, *of flowers*. L.)

1. *A. odoratum*, L. Spikelets (brownish or tinged with green) spreading at flowering-time; one of the neutral flowers bearing a bent awn from near its base, the other short-awned below the tip. — Meadows, pastures, &c. Perennial: very sweet-scented in drying. May-July. (Nat. from Eu.)

56. PHÁLARIS, L. CANARY-GRASS. (Pl. 13.)

Spikelets crowded in a clustered or spiked panicle, with 2 neutral mere rudiments (a scale or a pedicel) in place of lateral flowers, one on each side, at the base of the perfect one, which is flattish, awnless, of 2 shining palets, shorter than the equal boat-shaped and keeled glumes, finally coriaceous or cartilaginous, and closely enclosing the flattened free and smooth grain. — Stamens 3. — Leaves broad, mostly flat. (The ancient name, from *φαλός*, *shining*, alluding either to the palets or the grain.)

§ 1. PHÁLARIS proper. *Panicle very dense, spike-like: glumes wing-keeled.*

1. *P. canariensis*, L. (CANARY-GRASS.) Annual, 1°-2° high; spike oval; rudimentary flower a small lanceolate scale. — Waste places and roadsides, Massachusetts to Pennsylvania: rare. July-Sept. (Adv. from Eu.)

§ 2. DÍGRAPHIS, Trin. *Panicle branched, the clusters open in anthesis: glumes not winged on the back.*

1. *P. arundinacea*, L. (REED C.) Perennial, 2°-4° high; leaves flat (3''-5'' wide; glumes open at flowering, 3-nerved, thrice the length of the fertile flower; rudimentary flowers reduced to a minute hairy scale or pedicel. — Wet grounds: common, especially northward. June, July. — Var. *pfcta*, the leaves striped with white, is the familiar RIBBON-GRASS of the gardens. (Eu.)

57 MÍLIUM, MILLET-GRASS. (Pl. 13.)

Spikelets diffusely paniced, not jointed with their pedicels, apparently consisting of 2 equal membranaceous convex and awnless glumes, including a single coriaceous awnless flower: but theoretically the lower glume is wanting, while

an empty single palet of the lower (neutral) flower, resembling the upper glume, fulfils its office, and stands opposite the narrow upper palet of the terete fertile flower. Stamens 3. Stigmas branched-plumose. Grain not grooved, enclosed in the palets, all deciduous together. (The ancient Latin name of the Millet (which however belongs to a different genus), probably from *mille*, a thousand, because of its fertility.)

1. *M. effusum*, L. Smooth perennial, 3°–6° high; leaves broad and flat, thin; panicle spreading (6'–9' long); flower ovoid-oblong.—Cold and damp woods, New England to Illinois and northward. June. (Eu.)

58. AMPHICÁRPUM, Kunth. (Pl. 13.)

Spikelets jointed with the apex of the pedicels, apparently 1-flowered, of two kinds; one kind in a terminal panicle, like those of *Milium*, except that the rudiment of the lowest glume is ordinarily discernible, and deciduous from the joint without ripening fruit, although the flower is perfect: the other kind solitary at the extremity of slender runner-like radical peduncles (which are more or less sheathed towards the base), much larger than the others, perfect and fertile, subterranean, fertilized in the bud; the enwrapping glume and similar empty palet many-nerved. Flower oblong or ovoid, pointed. Stamens 3 (small in the radical flowers). Stigmas plumose, deep purple. Grain not grooved, in the radical flowers very large, the embryo next the lower palet. Neutral palet somewhat exceeding the glume and the fertile flower. (Name from ἀμφίκαρπος, doubly fruit-bearing.)

1. *A. Púrshii*, Kunth. Annual or biennial? erect, 1°–4° high; leaves lanceolate, copious on the lower part of the culm, hispid, especially on the sheaths; panicle strict, naked; grain ovoid or oblong (2''–3'' long), terete. (*Milium amphicárpon*, *Pursh*.)—Moist sandy pine-barrens, New Jersey, and in the Southern States. Sept.

59. PÁSPALUM, L. PASPALUM. (Pl. 13.)

Spikelets spiked or somewhat racemed in 2–4 rows on one side of a flattened or filiform continuous rhachis, jointed with their very short pedicels, plano-convex, awnless, apparently only one-flowered, as in *Milium*; but, on the other hand, differing from *Panicum* merely in the want of the lower glume; which, however, is occasionally present in some species, as a small scale. Glume and empty palet few-nerved. Flower coriaceous, mostly orbicular or ovate, flat on the inner side, convex on the outer. Stamens 3. (Said to have been a Greek name for Millet.)

* *Spikes with a (1'') broad and thin membranaceous or foliaceous and keeled rhachis, the incurved margins partly enclosing the small two-rowed spikelets. (Smooth, aquatic, or nearly so, decumbent or floating.)*

1. *P. fluitans*, Kunth. Annual; leaves lanceolate, flat (3''–8'') broad; spikes numerous in a raceme; the rhachis somewhat projecting beyond the minute and slightly pubescent spikelets into a tapering point, scabrous on the back. (*Cerèsia fluitans*, *Ell.*)—River-swamps, Virginia, S. Ohio, Illinois, and southward. Sept., Oct.

2. **P. Walterianum**, Schultes. Perennial; leaves linear, short; spikes 3-7, the lowest partly included in the sheath of the uppermost leaf, the rachis blunt; spikelets glabrous. (*P. vaginatum*, Ell.)—Low or wet grounds, New Jersey (Cape May, Nuttall), Delaware (Tatnall, Canby), and southward.

* * Spikes with a narrow wingless rachis: perennials, or mostly so.

+ Spikelets very obtuse, orbicular: spikes one terminal and often 1-5 lateral.

3. **P. setaceum**, Michx. Culm ascending or decumbent (1° - 2° long), slender; leaves ($2''$ wide, flat) and sheaths clothed with soft spreading hairs; spikes very slender ($2'$ - $4'$ long), smooth, mostly solitary on a long peduncle, and usually one from the sheaths of each of the upper leaves on short peduncles or included; spikelets ($\frac{1}{2}''$ wide) narrowly 2-rowed. (Also *P. débile* and *P. ciliatifolium*, Michx.)—Sandy fields: common from E. Mass. to Illinois, southward. Aug.

4. **P. læve**, Michx. Culm upright, rather stout (1° - 3° high); the pretty large and long leaves with their flattened sheaths smooth or somewhat hairy; spikes 2-6, the lateral ones somewhat approximated near the summit of an elongated naked peduncle, spreading ($2'$ - $4'$ long), smooth, except a bearded tuft at their base; spikelets broadly 2-rowed (over $1''$ wide).—Moist soil, S. New England to Kentucky, and southward. Aug.—Either glabrous or sometimes the lower sheaths, &c. very hairy. As here received this perhaps comprises two or more species.

+ + Spikelets acute: spikes always a pair at the summit of the naked peduncle.

5. **P. distichum**, L. (JOINT-GRASS.) Nearly glabrous, rather glaucous; culms ascending (about 1° high) from a long creeping base; leaves linear-lanceolate ($2'$ - $3'$ long); spikes short and closely-flowered ($9''$ - $2'$ long), one short-peduncled, the other sessile; rachis flat on the back; spikelets ovate, slightly pointed (barely $1\frac{1}{2}''$ long).—Wet fields, Virginia and southward. July-Sept.

6. **P. Digitaria**, Poir. Culms ascending (1° - $2\frac{1}{2}^{\circ}$ high) from a creeping base; leaves lanceolate ($3'$ - $6'$ long, $4''$ - $6''$ wide); spikes slender, rather sparsely flowered ($1'$ - $4'$ long), both sessile at the apex of the slender peduncle; spikelets ovate-lanceolate ($2''$ long). *Milium paspalodes*, Ell.)—Virginia and southward.

60. PANICUM, L. PANIC-GRASS. (Pl. 13.)

Spikelets paniced, racemed, or sometimes spiked, not involucrate, $1\frac{1}{2}$ -2-flowered. Glumes 2, but the lower one usually short or minute (rarely even wanting), membranaceo-herbaceous; the upper as long as the fertile flower. Lower flower either neutral or staminate, of one palet which closely resembles the upper glume, and sometimes with a second thin one. Upper flower perfect, closed, coriaceous or cartilaginous, usually flattish parallel with the glumes, awnless (except in § 3), enclosing the free and grooveless grain. Stamens 3. Stigmas plumose, usually purple. (An ancient Latin name of the Italian Millet, *P. Italicum* (now *Setaria Italica*), thought to come from *panis*, bread; some species furnishing a kind of bread-corn.)

§ 1. DIGITARIA, Scop. Spikelets crowded 2-3 together in simple and mostly 1-sided clustered spikes or spike-like racemes, wholly awnless and pointless: lower flower neutral, of a single palet: lower glume minute, sometimes obsolete or wanting: root annual: plant often purplish.

* *Spikes erect; its rhachis filiform and nearly terete.*

1. **P. filiforme**, L. Culms very slender (1° – 2° high), upright; lower sheaths hairy; *spikes* 2–8, alternate, approximated, filiform; spikelets oblong, acute ($\frac{1}{2}$ " long); lower glume almost wanting. — Dry sandy soil, Massachusetts to New Jersey along the coast, Illinois, and southward. Aug.

* * *Spikes spreading; its rhachis flat and thin.*

2. **P. GLABRUM**, Gaudin. Culms spreading, prostrate, or sometimes erect ($5'$ – $12'$ long), glabrous; *spikes* 2–6, widely diverging, nearly digitate; spikelets ovoid (about $1''$ long); *upper glume equalling the flower, the lower one almost wanting.* — Cultivated grounds and waste places: common, especially southward: in some places appearing as if indigenous. Aug., Sept. (Nat. from Eu.)

3. **P. SANGUINALE**, L. (COMMON CRAB-OR FINGER-GRASS.) Culms erect or spreading (1° – 2° high); leaves and sheaths glabrous or hairy; *spikes* 4–15, spreading, digitate; spikelets oblong ($1\frac{1}{2}$ " long); *upper glume half the length of the flower, the lower one small.* — Cultivated and waste grounds. Aug.–Oct. (Nat. from Eu.)

§ 2. PANICUM proper. *Spikelets scattered, in panicles, awnless.*

* *Panicle elongated and racemose, wand-like or pyramidal; the numerous and usually pointed spikelets short-pedicelled, excepting No. 7 and 8.*

+ *Sterile flower neutral and of 2 palets, fully twice the length of the lower glume: spikelets small ($1''$ or $1\frac{1}{2}$ " long): root perennial.*

4. **P. anceps**, Michx. Culms flat, upright (2° – 4° high); leaves rather broadly linear (1° – 2 long, $4''$ – $5''$ wide), smooth; panicle contracted-pyramidal; *spikelets ovate-lanceolate, pointed, a little curved; upper glume 5–7-nerved; neutral flower one third longer than the perfect one.* — Wet sandy soil, New Jersey and Penn. to Virginia, and southward. Aug. — Too near the next: spikelets and branches of the panicle longer.

5. **P. agrostoides**, Spreng. Culms flattened, upright (2° high); leaves long, and with the sheaths smooth; panicles terminal and often lateral, pyramidal ($4'$ – $8'$ long); the *spikelets* racemose, crowded and one-sided on the spreading branches, *ovate-oblong, acute* (purplish); *upper glume 5-nerved, longer than the neutral flower which exceeds the perfect one; perfect flower bearded at the apex.* (*P. agrostidiforme*, Lam. ? *P. multiflorum*, Poir.) — Wet meadows and shores, E. Massachusetts and New York (Oneida Lake, A. H. Curtiss) to Illinois, and common southward. Aug.

+ + *Sterile flower neutral and of a single palet, much longer than the lower glume; spikelets $\frac{1}{2}''$ – $1\frac{1}{2}''$ long; annuals except No. 8: leaves flat; sheaths flattened.*

+ + *Glabrous and smooth throughout; spikelets crowded, appressed, short-pedicelled.*

6. **P. proliferum**, Lam. Culms usually thickish and rather succulent, branched, geniculate and ascending from a procumbent base; sheaths flattened; ligule ciliate; panicles terminal and lateral, compound, pyramidal, the slender primary branches at length spreading; spikelets pale green, rarely purplish; lower glume broad, $\frac{1}{3}$ to $\frac{1}{2}$ the length of the upper; neutral flower little longer than the perfect one. — Marshy river-banks and shores, especially when brackish, but also in the interior, from Mass. and Illinois southward. Aug.

++ ++ *Hispid or hairy on the sheaths, at least the lower: spikelets mostly scattered on slender or capillary pedicels in an ample, loose, at length very effuse panicle: culms mostly branched from the base, erect or ascending (10' - 20' high).*

7. **P. capillare**, L. (OLD-WITCH GRASS.) All the sheaths and usually the leaves copiously hairy or hispid; panicle mostly very compound, the branches divaricate when old; spikelets varying from ovoid to narrowly oblong, pointed; lower glume half the length of the neutral palea which is longer than the elliptical obtuse perfect flower. — Sandy soil and cultivated fields everywhere. Aug. - Oct. — Varies extremely in size and appearance: in depauperate forms the spikelets only $\frac{3}{4}$ " in the larger forms $1\frac{1}{2}$ " in length.

8. **P. autumnale**, Bosc! Root perennial? lower sheaths and margins of the small narrow leaves more or less hairy, otherwise glabrous, except some bristly hairs in the main axils of the very effuse capillary panicle, its much elongated divisions sparingly branched, or even simple and terminated with solitary spindle-shaped spikelets; lower glume minute; perfect flower lanceolate-oblong and pointed, nearly equalling the lance-oblong obtusish upper glume and the neutral palea. (*P. nudum*, Walt.? *P. dichotomiflorum*, Michx. *P. divergens*, Muhl., not of *H. B. K.* *P. fragile*, Kunth.) — Sand-hills, Mason County, Illinois (*Mead, E. Hall*), and southward.

+ + + *Sterile flower staminate, of 2 paleas: lower glume more than half the length of the upper: spikelets large (2" - 2½" long), ovate, pointed, as are the glumes, &c.: perennials, glabrous, with tall or stout and rigid upright culms.*

9. **P. virgatum**, L. Tall (3° - 5° high); leaves very long, flat; ligule silky-bearded; branches of the compound loose and large panicle (9' - 2° long) at length spreading or drooping; spikelets scattered, usually purplish. — Moist sandy soil: common, especially southward. Aug.

10. **P. amarum**, Ell. Culms ($1\frac{1}{2}$ ° high) sheathed to the top; leaves involute, glaucous, coriaceous, the uppermost exceeding the contracted panicle, the simple racemose branches of which are appressed; spikelets pale. — Sandy shores, Connecticut (*Barratt, Robbins*), Virginia, and southward. Aug., Sept.

* * *Panicle loosely spreading or diffuse, short or small: perennials.*

+ *Lower (sterile) flower neutral, or in No. 11, and sometimes in No. 14, staminate, formed of 2 paleas, the upper one scarious and sometimes small and inconspicuous.*

+ *Culm-leaves broadly lanceolate or wider, with 9 - 15 principal nerves.*

11. **P. latifolium**, L. Culm (1° - 2° high) smooth; the joints and the orifice of the throat or margins of the otherwise smooth sheaths often bearded with soft woolly hairs; leaves broadly oblong-lanceolate from a heart-clasping base (often 1' wide), taper-pointed, 11 - 15-nerved, smooth, or sparingly downy-hairy; panicle more or less exserted (2' - 3' long), usually long-peduncled, the branches spreading; spikelets obovate, $1\frac{1}{2}$ " long, downy; lower glume ovate, not half the length of the many-nerved upper one; sterile flower often (not always) with 3 stamens. (*P. Walteri*, Poir.) — Moist thickets: common. June - Aug.

12. **P. clandestinum**, L. Culm rigid (1° - 3° high), very leafy to the top, at length producing appressed branches, the joints naked; sheaths rough with papillæ bearing very stiff and spreading bristly hairs; leaves oblong-lanceolate from a heart-clasping base, very taper-pointed; lateral and usually also the ter-

minal panicle more or less enclosed in the sheaths, or with the terminal one at length long-peduncled (*P. pedunculatum*, Torr.):—otherwise resembling No. 11; but the spikelets more ovoid, often smooth; the lower flower (always?) neutral.—Low thickets and river-banks: common. June—Sept.

13. **P. microcarpon**, Muhl. Culm and sheaths as in No. 11; the broadly lanceolate leaves nearly similar, but longer in proportion and less pointed, not dilated at the rounded bristly-ciliate base, very rough-margined, the upper surface roughish; panicle soon exerted on a slender peduncle, very many-flowered, narrowly oblong (3'–7' long); spikelets only about $\frac{1}{2}$ " long, ovoid, smooth or smoothish; lower glume orbicular and very small. (*P. multiflorum*, Ell.? not of Poir.)—Dry or moist thickets, Pennsylvania and Michigan to Illinois, and southward. July—Sept.

14. **P. xanthophysum**, Gray. Culm simple, or at length branched near the base (9'–15' high); sheaths hairy; leaves lanceolate, very acute (4'–6' long by $\frac{1}{2}$ ' wide), not dilated at the ciliate-bearded clasping base, smooth except the margins, strongly 9–11-nerved; panicle long-peduncled, very simple, the appressed branches bearing a few roundish-obovate spikelets (about $1\frac{1}{2}$ " long); lower glume ovate, acutish, one third or half the length of the 9-nerved upper one.—Dry sandy soil, Maine to Wisconsin, and northward: rare. June.—Plant yellowish-green: spikelets minutely downy: sterile flower sometimes staminate.

15. **P. viscidum**, Ell. Culms upright or ascending, at length much branched, leafy to the top, densely velvety-downy all over, as also the sheaths, with reflexed soft and often clammy hairs, except a ring below each joint; leaves likewise velvety, lanceolate ($\frac{1}{2}$ ' wide), 11–13-nerved; panicle spreading, the lateral ones included; spikelets obovate 1" or $1\frac{1}{2}$ " long, downy; the roundish lower glume scarcely one fourth the length of the 7-nerved upper one.—Damp soil, New Jersey to Virginia, and southward. Aug.

16. **P. pauciflorum**, Ell.? Culms upright, at length much branched and reclining (1°–2° long), roughish; leaves lanceolate (3'–5' long by $\frac{1}{2}$ '– $\frac{1}{2}$ ' wide), rather faintly 9-nerved, hairy or smooth, fringed on the whole margin or next the base with long and stiff spreading hairs, the sheaths bristly throughout with similar hairs; panicle open, nearly simple, bearing few tumid-obovate hairy or smoothish spikelets about $1\frac{1}{2}$ " long; lower glume roundish, about half or a quarter of the length of the upper one.—Wet meadows and copses, E. Massachusetts to Wisconsin, and southward. June, July.—Distinguished by its much larger spikelets, more nerved leaves, and coarser aspect, from any form of the next. It has probably been described under several names, some of them earlier than Elliott's.

++ ++ Leaves linear or lanceolate, with few or indistinct primary nerves.

17. **P. dichotomum**, L.! Culms (8'–20' high) at first mostly simple, bearing a more or less exerted spreading compound panicle (1'–3' long), and lanceolate flat leaves (those tufted at the root usually ovate-lanceolate and very short, thickish); but commonly branching later in the season, the branches often clustered, and bearing nearly simple and included small panicles; spikelets $\frac{1}{2}$ " to about 1" long; oblong-obovate, downy or smooth; lower glume roundish, one third or a quarter the length of the 5–7-nerved upper one.—Founded on an autumnal state of the species, much forked and with densely clustered lateral branchlets

and panicles. (*P. nodiflorum*, Lam.) — Exhibits an interminable diversity of forms; of which a shaggy-hairy and larger-flowered variety is *P. pubescens*, Lam.; and one with smaller spikelets is *P. laxiflorum*, Lam.; while the varied smooth or smoothish states with shining leaves are *P. nitidum*, Lam., and (the more slender forms) *P. barbulatum* and *P. ramulosum*, Michx., &c. Some of these may be good species. — Dry or low grounds: common. June–Aug.

18. **P. depauperatum**, Muhl. Culms simple or branched from the base, forming close tufts (6'–12' high), terminated by a simple and few-flowered contracted panicle, often much overtopped by the narrowly linear and elongated (4'–7') upper leaves; spikelets $\frac{3}{4}$ "–1 $\frac{1}{2}$ " long, oval-obovate, commonly pointed when young; the ovate lower glume one third the length of the 7–9-nerved upper one. (*P. strictum*, Pursh. *P. rectum*, Rœm. & Schult.) — Varies, with the leaves involute, at least when dry (*P. involutum*, Torr.), and with the sheaths either beset with long hairs or nearly smooth: the panicle either partly included, or on a long and slender peduncle. — Dry woods and hills: rather common. June.

+ + Lower sterile flower of a single palet, and neutral.

19. **P. verrucosum**, Muhl. Smooth; culms branching and spreading, very slender (1°–2° long), naked above; leaves linear-lanceolate (2"–3" wide), shining; branches of the diffuse panicle capillary, few-flowered; spikelets warty-roughened (dark green), oval, acute, $\frac{3}{4}$ " long; the lower glume one fourth the length of the obscurely nerved upper one. — Sandy swamps, New England to Virginia, near the coast, and southward. Aug.

§ 3. **ECHINÓCHLOA**, Beauv. Spikelets imbricated-spiked on the branches of the simple or compound raceme or panicle, usually rough with appressed stiff hairs: lower palet of the sterile flower awl-pointed or awned.

20. **P. CRUS-GÁLLI**, L. (BARNYARD-GRASS.) Root annual; culms stout, branching from the base (1°–4° high); leaves lanceolate ($\frac{1}{2}$ ' or more wide), rough-margined, otherwise with the sheaths smooth; spikes alternate (1'–3' long), crowded in a dense panicle; glumes ovate, abruptly pointed; lower palet of the neutral flower bearing a rough awn of variable length. — Varies greatly; sometimes awnless or nearly so; sometimes long-awned, especially so in the var. *HISPIDUM* (*P. hispidum*, Muhl., *P. longisetum*, Torr.), a very large and coarse form of the species with the sheaths of the leaves very bristly. — Moist, chiefly manured soil: the variety in ditches, especially of brackish water; possibly indigenous. Aug.–Oct. (Nat. from Eu.)

61. SETÀRIA, Beauv. BRISTLY FOX-TAIL GRASS. (Pl. 13.)

Spikelets altogether as in *Panicum* proper, and awnless, but with the short peduncles produced beyond them into solitary or clustered bristles resembling awns (but not forming an involucre). Inflorescence a dense spiked panicle, or apparently a cylindrical spike. — Annuals, in cultivated or manured grounds, with linear or lanceolate flat leaves: properly to be regarded as merely a subgenus of *Panicum*. (Name from *seta*, a bristle.)

* Bristles single or in pairs, roughened or barbed downwards.

1. **S. VERTICILLATA**, Beauv. Spike cylindrical (2'–3' long, pale green), composed of apparently whorled short clusters; bristles short, adhesive. (*Panicum verticillatum*, L.) — Near dwellings. (Adv. from Eu.)

* * *Bristles in clusters, roughened or barbed upwards.*

2. **S. GLAÛCA**, Beauv. (FOXTAIL.) *Spike cylindrical, dense, tawny yellow (2' - 4' long); bristles 6 - 11 in a cluster, much longer than the spikelets; perfect flower transversely wrinkled.* — Very common, in stubble, &c. (Adv. from Eu.)

3. **S. VÍRIDIS**, Beauv. (GREEN FOXTAIL. BOTTLE-GRASS.) *Spike nearly cylindrical, more or less compound, green; bristles few, longer than the spikelets; perfect flower striate lengthwise and dotted.* — Cultivated grounds. (Adv. from Eu.)

4. **S. ITÁLICA**, Kunth. *Spike compound, interrupted at the base, thick, nodding (6' - 9' long, yellowish or purplish); bristles 2 or 3 in a cluster, either much longer or else shorter than the spikelets.* (S. Germánica, Beauv.) — Sometimes cultivated under the name of MILLET, or BENGAL GRASS: rarely spontaneous. (Adv. from Eu.)

62. CÉNCHRUS, L. HEDGEHOG- or BUR-GRASS. (Pl. 14.)

Spikelets as in *Panicum*, awnless, but enclosed 1 to 5 together in a globular and bristly or spiny involucre, which becomes coriaceous and forms a deciduous hard and rigid bur: the involucre sessile in a terminal spike. Styles united below. (An ancient Greek name of *Setaria Italica*.)

1. **C. tribuloides**, L. Culms branched and ascending (1° - 2° high) from an annual root; leaves flat; spike oblong, of 8 - 20 spherical heads; involucre prickly all over with spreading and barbed short spines, more or less downy, enclosing 2 or 3 spikelets. — Sandy soil, on the coast, the Great Lakes, and the larger rivers. Aug. — A vile weed.

63. TRÍPSACUM, L. GAMA-GRASS. SESAME-GRASS. (Pl. 14.)

Spikelets monœcious, in jointed spikes, which are staminate above and fertile below. Staminate spikelets 2, sessile at each triangular joint of the narrow rhachis, forming a 1-sided and 2-ranked spike longer than the joints, both alike, 2-flowered: glumes coriaceous, the lower (outer) one nerved, the inner one boat-shaped: palets very thin and membranaceous, awnless: anthers (turning orange or reddish-brown) opening by 2 pores at the apex. Pistillate spikelets single and deeply imbedded in each oblong joint of the cartilaginous thickened rhachis, occupying a boat-shaped recess which is closed by the polished and cartilaginous ovate outer glume, the inner glume much thinner and pointed, 2-flowered; the lower flower neutral, palets very thin and scarious, pointless. Styles united: stigmas very long (purple), hispid. Grain ovoid, free. Culms stout and tall, solid, from very thick creeping rootstocks. Leaves broad and flat. Spikes axillary and terminal, separating spontaneously into joints at maturity. (Name from *τρίβω*, to rub, perhaps in allusion to the polished fertile spike.)

1. **T. dactyloides**, L. Spikes (4' - 8' long) 2 - 3 together at the summit (when their contiguous sides are more or less flattened), and also solitary from some of the upper sheaths (when the fertile part is cylindrical); sometimes, var. **MONOSTACHYUM**, the terminal spike also solitary. — Moist soil, from Connecticut to Pennsylvania, near the coast, and from Illinois southward. Aug. — Culm 4° - 7° high: the leaves like those of Indian Corn. — This is one of our largest and most remarkable Grasses: sometimes used for fodder at the South, where better is not to be had.

64. ERIANTHUS, Michx. WOOLLY BEARD-GRASS. (Pl. 14.)

Spikelets spiked in pairs upon each joint of the slender rhachis; one of them sessile, the other pedicelled; otherwise both alike; with the lower flower neutral, of one membranaceous palet; the upper perfect, of 2 hyaline palets, which are thinner and shorter than the nearly equal membranaceous glumes, the lower palet awned from the tip. Stamens 1-3. Grain free. — Tall and stout reed-like perennials, with the spikes crowded in a panicle, and clothed with long silky hairs, especially in a tuft around the base of each spikelet (whence the name, from *ἔριον*, wool, and *ἄνθος*, flower).

1. **E. alopecuroides**, Ell. Culm (4°-6° high) woolly-bearded at the joints; *panicle contracted*; the *silky hairs longer than the spikelets*, shorter than the awn; stamens 2. — Wet pine barrens, from New Jersey and Illinois southward: rare. Sept., Oct.

2. **E. brevibarbis**, Michx. Culm (2°-5° high), somewhat bearded at the upper joints; *panicle rather open*; silky hairs shorter than the spikelets. — Low grounds, Virginia and southward.

65. ANDROPÖGON, L. BEARD-GRASS. (Pl. 14.)

Spikelets in pairs upon each joint of the slender rhachis, spiked or racemed; one of them pedicelled and sterile, often a mere vestige; the other sessile, with the lower flower neutral and of a single palet; the upper perfect and fertile, of 2 thin and hyaline palets shorter than the herbaceous or chartaceous glumes, the lower awned from the tip. Stamens 1-3. Grain free. — Coarse, mostly rigid perennials, mostly in sterile or sandy soil; with lateral or terminal spikes commonly clustered or digitate; the rhachis hairy or plumose-bearded, and often the sterile or staminate flowers also (whence the name, composed of *ἀνήρ*, *ἄνδρως*, man, and *πώγων*, beard).

* *Spikes digitate, thickish, short-bearded, the sterile spikelet staminate: stamens 3.*

1. **A. furcatus**, Muhl. Tall, 3°-4° high, rigid, the naked summit of the culm (and usually some lateral branches) terminated by 2-5 rigid spikes; spikelets approximated, appressed; hairs at the base of the fertile spikelet, on the rhachis and on the stout pedicel of the awnless staminate spikelet short and rather sparse; awn of fertile flower long and bent; leaves flat, roughish, the lower ones long. — Common in dry sterile soil. Aug. - Oct.

* * *Spikes with slender often zigzag rhachis, silky-villous,*

+ *Single and scattered along the branches, with the silky hairs shorter than the flowers: sterile spikelet conspicuous but mostly neutral; the fertile triandrous.*

2. **A. scoparius**, Michx. Culms slender (1°-3° high), with numerous paniculate branches; lower sheaths and narrow leaves hairy; spikes slender, scattered, mostly peduncled (1'-2' long), very loose, often purplish, silky with lax dull-white hairs; sterile spikelet awn-pointed or awnless; the fertile about half the length of its twisted or bent awn. — Dry ground. July - Sept.

+ + *In pairs or clustered; the copious soft-silky hairs much longer than the flowers: sterile spikelet a small neutral rudiment (in No. 3), or altogether wanting on the summit of the very plumose-hairy pedicel: fertile flower monandrous, its awn capillary: leaves narrow, the lower or their sheaths often rather hairy.*

3. **A. argenteus**, Ell. Culms rather slender (1° – 3° high); *spikes* in pairs (rarely in fours) on short mostly exserted and loosely paniculate peduncles, densely flowered ($1'$ – $2'$ long), very silky with long bright white hairs. (*A. argyræus*, Schultes. *A. Elliottii*, Chapm.)—Delaware (*W. M. Canby*), Virginia, near the coast, and southward. Sept., Oct.

4. **A. Virginicus**, L. Culm flattish below, slender (2° – 3° high), sparingly short-branched above, sheaths smooth; *spikes* 2 or 3 together in distant appressed clusters, shorter than their sheathing bracts, weak ($1'$ long), the spikelets loose on the filiform rhachis, the soft hairs dull white. (*A. vaginatus*, Ell., a form with larger and inflated sheaths.)—Sandy soil, E. Massachusetts to Virginia, Illinois, and southward. Sept., Oct.

5. **A. macrourus**, Michx. Culm stout (2° – 3° high), bushy-branched at the summit, loaded with very numerous *spikes forming thick leafy clusters*; sheaths rough, the uppermost hairy; flowers nearly as in the preceding; the sterile spikelet of each pair wholly wanting, its pedicel slender and very plumose.—Low and sandy grounds, New York to Virginia, near the coast, and southward. Sept., Oct.

66. SÓRGHUM, Pers. BROOM CORN. (Pl. 14.)

Spikelets 2–3 together on the ramifications of an open panicle, the lateral ones sterile or often reduced merely to their pedicels; only the middle or terminal one fertile, its glumes coriaceous or indurated, sometimes awnless: otherwise nearly as in *Andropogon*. Stamens 3. (The Asiatic name of *S. vulgare*, the INDIAN MILLET, to which species belongs GUINEA-CORN, BROOM-CORN, the SWEET SORGHUM, and other cultivated races.)

1. **S. nutans**, Gray. (INDIAN GRASS. WOOD-GRASS.) Root perennial; culm simple (3° – 5° high), terete; leaves linear-lanceolate, glaucous; sheaths smooth; panicle narrowly oblong, crowded or loose ($6'$ – $12'$ long); the perfect spikelets at length drooping (yellowish or russet-brown and shining), clothed, especially towards the base, with fawn-colored hairs, lanceolate, shorter than the twisted awn; the sterile spikelets small and imperfect, deciduous, or reduced to a mere plumose-hairy pedicel. (*Andropogon nutans*, L.)—Dry soil: common, especially southward, where it runs into several marked varieties or perhaps species (*S. avenaceum*, *nutans*, and *secundum*, Chapman). Aug. – Oct.

SERIES II.

CRYPTOGAMOUS OR FLOWERLESS PLANTS.

VEGETABLES destitute of proper flowers (i. e. no stamens and pistils), producing, in place of seeds, minute bodies of homogeneous structure (called *spores*), in which there is no embryo, or plantlet anterior to germination.

CLASS III. ACROGENS.

Cryptogamous plants with a distinct axis (stem and branches), growing from the apex only, containing woody fibre and vessels (especially ducts), and usually with some kind of foliage.

ORDER 129. EQUISETACEÆ. (HORSETAIL FAMILY.)

Leafless plants, with rush-like hollow and jointed stems, arising from running rootstocks, terminated by the fructification in the form of a cone or spike, which is composed of shield-shaped stalked scales bearing the spore-cases underneath. — Comprises solely the genus

1. EQUISETUM, L. HORSETAIL. SCOURING RUSH. (Pl. 15.)

Spore-cases (*sporangia, thecæ*) 6 or 7, adhering to the under side of the angled shield-shaped scales of the spike, 1-celled, opening down the inner side and discharging the numerous loose spores. To the base of each spore are attached 4 thread-like and club-shaped elastic filaments, which roll up closely around the spore when moist, and uncoil when dry. — Stems mostly from running rootstocks, striate-grooved (in many the hard cuticle abounding in silex), hollow, and also with an outer circle of smaller air-cavities corresponding with the grooves; the joints closed and solid, each bearing instead of leaves a sheath, which surrounds the base of the internode above, and is split into teeth corresponding in number and position with the principal ridges of the stem: the stomata in the grooves. Branches, when present, in whorls from the base of the sheath, like the stem, but without the central air-cavity. (The ancient name, from *equus*, horse, and *seta*, bristle.)

§ 1. *Annual-stemmed, not surviving the winter. (Stomata scattered.)*

* *Fruiting in spring from soft and rather succulent pale or brownish fertile stems, the sterile stems or branches appearing later, herbaceous and very different.*

+ *Fertile stems remaining simple, soon perishing; the sterile producing copious branches.*

1. **E. Telmatèia**, Ehrh. (GREAT HORSETAIL.) Stems stout (as thick as the finger); the sheaths of the fertile ones ($1' - 1\frac{1}{2}'$ long) enlarging upwards, deeply 20-30-toothed; sterile stem white, 20-30-furrowed; its branches simple, rough, usually 4-angled and again grooved on the angles. (*E. eburneum*, Roth., Schreber, and Ed. 2. *E. fluviatile*, Smith.)—Shore of the upper Great Lakes, and northwestward: rare. April, May.—Fertile stems $10' - 15'$, the sterile $2^\circ - 5^\circ$ high. (Eu.)

2. **E. arvensè**, L. (COMMON H.) Fertile stems ($4' - 10'$ high) with loose and usually distant about 8-12-toothed sheaths; the sterile slender (at length $1^\circ - 2^\circ$ high), 10-14-furrowed, producing long and simple or sparingly branched 4-angular branches; their teeth 4, herbaceous, lanceolate.—Moist, especially gravelly soil: very common. March-May.—Rootstocks occasionally bearing copious little tubers like small peas (Illinois, *S. A. Collier*).—The var. **SEROTINUM**, Meyer, an accidental state, in which the sterile plant produces a spike of fruit from its summit, is found in New Jersey by *C. F. Austin*.

+ + *Fertile stems when older producing herbaceous 3-sided branches, and lasting through the summer, except the naked top which perishes after fructification.*

3. **E. pratense**, Ehrh. Sterile and finally also the fertile stems producing simple straight branches; sheaths of the stem with ovate-lanceolate short teeth, those of the branches 3-toothed: stems more slender and the branches shorter than in the last. (*E. umbròsum*, Willd. *E. Drummóndii*, Hook.)—Michigan (*Cooley, &c.*), Wisconsin, and northward. April, May. (Eu.)

4. **E. sylvaticum**, L. Sterile and fertile stems (about 12-furrowed) producing compound racemed branches; sheaths loose, with 8-14 rather blunt teeth, those of the branches bearing 4 or 5, of the branchlets 3, lance-pointed divergent teeth.—Wet shady places: common northward. May. (Eu.)

* * *Fruiting in summer; the stems all of one kind, or the fertile contemporaneous with and like the sterile, equally herbaceous, producing mostly simple branches, or sometimes nearly naked.*

5. **E. palùstre**, L. Stems ($10' - 18'$ high) slender, very deeply 5-9-grooved, the grooves separated by narrow wing-like ridges, roughish, the lance-awl-shaped teeth whitish-margined; branches rather few in a whorl.—Wet places, Niagara River (*G. W. Clinton*), Wisconsin (*C. F. Austin*), and northward. June. (Eu.)

6. **E. limòsum**, L. Stems ($2^\circ - 3^\circ$ high) slightly many-furrowed, smooth, usually producing upright branches after fructification; sheaths appressed, with 10-22 (commonly about 18) dark-brown and acute rigid short teeth. (*E. uliginòsum*, Muhl.)—In shallow water: rather common.—Air-cavities none under the grooves, but small ones under the ridges. June, July. (Eu.)

(*E. LITTORALE*, Kùhlewein, a species of N. Europe which stands between No. 5 and No. 6, is said by Milde to grow in Canada, on the authority of a specimen in herb. Hook., and may be looked for northward.)

§ 2. *Evergreen or perennial-stemmed, surviving the winter, mostly rough (the cuticle abounding in silex): fruiting in summer: spike tipped with a rigid little point. (Stomata in regular rows, in our species 1-rowed on each side of the groove.*

* *Stems tall and stout ($1\frac{1}{2}^{\circ}$ - 4° or even 6° high), simple or casually branched, evenly many- (15-40-) grooved: sheaths appressed. (Probably all forms of the Common Scouring Rush.)*

7. **E. lævigatum**, Braun. Stems $1\frac{1}{2}^{\circ}$ - 4° high, sometimes with numerous branches; the ridges convex, obtuse, smooth or minutely roughish with minute tubercles; sheaths elongated, with a narrow black limb and about 22 linear-awl-shaped caducous teeth, 1-keeled below. — Dryish clay soil, Illinois and southward.

8. **E. robustum**, Braun. Stems 3° - 6° high; the ridges narrow, rough with one line of tubercles: sheaths short, with a black girdle above the base, rarely with a black limb, and about 40 deciduous 3-keeled teeth with ovate-awl-shaped points. — River-banks, Ohio to Illinois, and southward. — Passes by var. **AFFINE**, Engelm. (smaller, with 20-25 awl-pointed more persistent teeth) into the next.

9. **E. hyemale**, L. (SCOURING-RUSH. SHAVE-GRASS.) Stems $1\frac{1}{2}^{\circ}$ - 4° high; the ridges roughened by 2 more or less distinct lines of tubercles; sheaths elongated, with a black girdle above the base and a black limb, of about 20 (17-26) narrowly linear teeth, 1-keeled at the base and with awl-shaped deciduous points. — Wet banks: common northward. Used for scouring. (Eu.)

* * *Stems slender, in tufts, 5-10-grooved: sheaths looser.*

10. **E. variegatum**, Schleicher. Stems ascending (6' - 18' long), usually simple from a branched base, 5-10-grooved; sheaths green variegated with black above; the 5-10-teeth tipped with a deciduous bristle. — Shores or river-banks, New Hampshire (Bellows Falls, *Carey*) and Niagara to Wisconsin and northward: rare. (Eu.)

11. **E. scirpoides**, Michx. Stems very numerous in a tuft, filiform (3' - 6' high), flexuous and curving, mostly 6-grooved, with acute ridges; the sheaths 3-toothed, the bristle-pointed teeth more persistent; central air-cavity wanting. — Wooded hillsides, New England to Pennsylvania, Michigan, and northward. (Eu.)

ORDER 130. FILICES. (FERNS.)

Leafy plants, with the leaves (fronds) usually raised on a stalk or petiole (stipe), rising from a root, or mostly from a prostrate or assurgent or even erect rootstock, separately rolled up (circinate) in the bud (except in Suborder IV.) and bearing on the under surface, commonly on the veins or along the margins, the simple fructification, which consists of 1-celled spore-cases (sporangia), opening in various ways, and discharging the numerous minute spores. (Antheridia and pistillidia formed and fertilization effected on the seeding plantlet!) — Of the eight well-marked Suborders, into which the Ferns are divided, four are represented in the Northern United States.

The whole order has been elaborated for this edition by Prof. DANIEL C. EATON, of Yale College.

SUBORDER I. **POLYPODIACEÆ.** THE TRUE FERNS.

Sporangia collected in dots, lines or variously shaped clusters (*sori* or *fruit-dots*) on the back or margins of the frond or its divisions, stalked, cellular-reticulated, the stalk running into a vertical incomplete many-jointed ring, which by straightening at maturity ruptures the sporangium transversely on the inner side, discharging the spores. Fruit-dots often covered (at least when young) by a membrane called the *indusium* (or less properly the *involute*), growing either from the back or the margin of the frond. (Plates 15-18.)

Tribe I. POLYPODIEÆ. Fructification on the back of the frond, in roundish or elongated fruit-dots (*sori*) placed on the veins or at the ends of the veins, without indusium of any kind. Stipes articulated to the rootstock, leaving a distinct scar when separated. Veins free (not reticulated) in our species.

1. **Polypodium.** Sori round, in one or more rows each side of the midrib or of the segments of the frond.

Tribe II. PTERIDEÆ. Fructification marginal or intramarginal, provided with a general indusium formed of the (either altered or unchanged) margin of the frond. Stipes not articulated to the rootstock. Veins free in all our species.

* Sporangia at the ends of the veins, borne on a reflexed portion of the margin of the frond.

2. **Adiantum.** Midrib of the pinnules near the lower margin or none. Stipe black and polished.

** Sporangia borne on a continuous marginal vein-like receptacle, which connects the apices of the veins, and is covered by a delicate whitish indusium formed of the reflexed margin of the pinnule.

3. **Pteris.** Midrib of the pinnules central. Stipe light-colored.

*** Sporangia at or near the ends of the unconnected veins, borne on the under surface of the frond: indusium various.

4. **Chellanthus.** Sori minute, at the ends of the veins, indusium continuous or interrupted. Fronds mostly chaffy, woolly or pulverulent.

5. **Pellaea.** Sori on the upper part of the veins, distinct, or mostly forming a confluent sub-marginal band of sporangia. Indusium membranaceous, continuous, rarely altogether wanting. Sterile and fertile fronds not very unlike: stipes dark-colored: fronds smooth.

6. **Allosorus.** Sori roundish or elongated and extending far down the free veins, at first covered by the reflexed herbaceous margin of the segment, at length exposed and confluent. Sterile and fertile fronds very different: stipes light-colored: fronds smooth.

Tribe III. ASPLENIEÆ. Sori more or less elongated, occupying one or both sides (but not the back) of the veins, covered by a special indusium which is attached by one side to the fertile vein, and is free on the other. Stipes not articulated.

* Sori linear or oblong, borne on cross veins parallel to the midrib.

7. **Woodwardia.** Sori forming a chain-like row each side of the midrib or central vein. Veins reticulated.

** Sori linear or oblong, borne on veins oblique to the midrib.

8. **Asplenium.** Sori on the upper side of the fertile veinlet, rarely on both sides of a veinlet. Veins free in all our species.

9. **Scolopendrium.** Sori linear, confluent in pairs, each pair appearing like a single sorus with a double indusium opening down the middle. Veins free.

** Sori borne partly on veins parallel to the midrib, partly on veins oblique to the midrib.

10. **Camptosorus.** Sori oblong or linear, the outer ones often approximating in pairs. Veins reticulated.

Tribe IV. ASPIDIEÆ. Sori round or roundish, on the back or rarely at the apex of the fertile vein, provided with a special indusium, rarely naked. Stipes not articulated to the rootstock.

* Indusium obsolete or none.

11. **Phegopteris.** Sori round, rather small. Veins free in our species.

** Indusium evident, round or roundish, covering the sporangia, at least when young. Sterile and fertile fronds not very unlike. Veins free in our species.

12. **Aspidium.** Indusium flat, orbicular or round-reniform, fixed by the centre, opening all round the margin.

13. **Cystopteris.** Indusium convex, fixed by a broad base partly under the sorus, commonly reflexed as the sporangia ripen.

*** Indusium obscure, irregularly semicircular. Fertile fronds much contracted and very unlike the sterile ones.

14. **Struthiopteris.** Fertile frond simply pinnate. Sterile frond with free veins.

15. **Onclea.** Fertile frond twice pinnate. Sterile frond with reticulated veins.

*** Indusium roundish or stellate, placed beneath the sporangia, sometimes enclosing them and then bursting open from the top.

16. **Woodsia.** Indusium very delicate, cleft into irregular lobes, or divided into a ciliate fringe. Veins free.

Tribe V. DAVALLIEÆ. Sori roundish or transversely elongated, borne at the ends of the veins or on marginal cross-veinlets, with an indusium attached at the base or base and sides and opening towards the margin of the segment.

17. **Dicksonia.** Sori marginal, very small, the indusium cup-shaped, somewhat 2-valved, the under portion confluent with a lobule of the frond. Veins free.

SUBORDER II. SCHIZÆACEÆ.

Sporangia ovate, sessile, having a complete transverse articulated ring at the apex, and opening by a longitudinal slit. (Pl. 19.)

18. **Schizæa.** Sporangia naked, fixed in a double row to the midrib of the narrow fertile segments. Sterile fronds rigid, simple or dichotomously branched.

19. **Lygodium.** Sporangia borne in a double row on narrow fertile segments, each sporangium seated on a separate veinlet, and provided with a special scale-like indusium. Fronds leafy, climbing.

SUBORDER III. OSMUNDACEÆ.

Sporangia naked, globose, pedicelled, reticulated, having a very incomplete transverse ring around the apex, opening into two valves by a longitudinal slit. (Pl. 19.)

20. **Osmunda.** Fertile pinnae or fronds very much contracted, bearing the abundant and large sporangia upon the margins of the very narrow segments. Veins free.

SUBORDER IV. OPHIOGLOSSACEÆ.

Sporangia spiked, destitute of a ring, naked, coriaceous and opaque, not reticulated, opening by a transverse slit into two valves, discharging very copious powdery spores. Fronds straight, never rolled up in the bud! (Pl. 19.)

21. **Botrychium.** Sporangia in pinnate or compound spikes, distinct. Veins free.

22. **Ophioglossum.** Sporangia cohering in a simple spike. Veins reticulated.

1. **POLYPÓDIUM**, L. POLYPODY. (Pl. 15.)

Fruit-dots round, naked, arranged on the back of the frond in one or more rows each side of the midrib or central vein, or irregularly scattered, each borne in our species on the end of a free veinlet. Rootstocks creeping, branched, often covered with chaffy scales, bearing scattered roundish knobs, to which the stipes are attached by a distinct articulation. (Name from *πολύ*, *many*, and *πούς*, *foot*, alluding to the branching rootstock.)

1. **P. vulgare**, L. Fronds evergreen, oblong, *smooth both sides*, 4' - 10' high, simply and deeply pinnatifid; the divisions linear-oblong, obtuse or somewhat acute, remotely and obscurely toothed; veins once or twice forked; *fruit-dots large, midway between the midrib and the margin*. — Rocks: common. July. (Eu.)

2. **P. incanum**, Swartz. Fronds evergreen and coriaceous, oblong, 2' - 6' high, *grayish and very scurfy underneath with peltate scales*, simply pinnatifid; the divisions oblong-linear, obtuse; *fruit-dots rather small, near the margin*; veins forking, free in the N. American plant! — Rocks and trunks of trees, Virginia and Ohio to Illinois, and southward. Aug.

2. **ADIANTUM**, L. MAIDENHAIR. (Pl. 16.)

Fruit-dots marginal, short; borne on the under side of a transversely oblong, crescent-shaped or roundish, more or less altered margin or summit of a lobe or tooth of the frond reflexed to form an indusium: the sporangia attached to the approximated tips of the free forking veins. — Main rib (*costa*) of the pinnules none, or at the lower margin. Stipes black and polished. (The ancient name, from *a* privative and *διαίνω*, meaning *unwetted*, the smooth foliage repelling rain-drops.)

1. **A. pedatum**, L. Frond forked at the summit of the upright slender stalk (9' - 15' high), the recurved branches bearing on one side several slender spreading divisions, which bear numerous triangular-oblong and oblique short-stalked pinnules; these are as if halved, being entire on the lower margin, from which the veins all proceed, and cleft and fruit-bearing on the other. — Rich, moist woods. July. — A delicate and most graceful Fern.

3. **PTERIS**, L. BRAKE OR BRACKEN. (Pl. 16.)

Sporangia in a continuous slender line of fructification, occupying the entire margins of the fertile frond, and covered by its reflexed narrow edge, which forms a continuous membranaceous indusium, attached to an uninterrupted transverse vein-like receptacle which connects the tips of the forked and free veins. — Fronds once to thrice pinnate or decompound. (The ancient Greek name of Ferns, from *πτερόν*, *a wing*, on account of the prevalent pinnate or feathery fronds.)

1. **P. aquilina**, L. (COMMON BRAKE.) Frond dull green (2° - 3° wide), ternate at the summit of an erect stout stalk (1° - 2° high), the widely spreading branches twice pinnate; pinnules oblong-lanceolate; the upper undivided; the lower more or less pinnatifid, with oblong obtuse lobes, margined all round with the indusium. — Thickets and hills: common. Aug. (Eu.)

4. CHEILÁNTHES, Swartz. LIP-FERN. (Pl. 16.)

Sporangia borne on the thickened ends of free veinlets, forming small and roundish distinct or nearly contiguous marginal fruit-dots, covered by a mostly whitish and membranaceous, sometimes herbaceous, common indusium, formed of the reflexed margin of separate lobes or of the whole pinnule. — Low, mostly with 2–3-pinnate and hairy or chaffy fronds, the sterile and fertile nearly alike, the divisions not halved, the principal vein central. Some species with continuous indusium connect this genus very closely with the next. (Name composed of *χείλος*, a lip, and *ἄνθος*, flower, from the shape of the involucre.)

1. *C. vestita*, Swartz. Fronds (6'–15' high), lanceolate-oblong, hirsute, as are the brown and shining stipes, with *straightish prominently articulated rusty hairs*, twice pinnate; pinnæ rather distant, triangular-ovate; pinnules oblong, crowded (2"–4" long), more or less incised, *the ends of the roundish or oblong lobes reflexed and forming separate herbaceous involucre*s, which are pushed back by the ripened sporangia. (*Nephrodium lanosum*, Michx.!) — Clefts of rocks, Island of New York (*W. W. Denslow*), and New Jersey to Illinois, and southward.

2. *C. tomentosa*, Link. Fronds (12'–20' high) lanceolate-oblong, densely tomentose with *slender and entangled whitish obscurely articulated hairs*, thrice pinnate; primary and secondary pinnæ oblong or ovate-oblong; pinnules distinct, minute ($\frac{1}{2}$ "–1" long), roundish-obovate, sessile or adnate-decurrent, the upper surface less woolly, *the reflexed narrow margin forming a continuous somewhat membranaceous involucre*. — Mountains of Virginia? and Kentucky; thence westward and southward. — Stipe and rachis rather stout, brown, covered with narrow chaffy scales and whitish hairs.

3. *C. lanuginosa*, Nutt. (in herb. Hook. & Sp. Fil. 1851). Stipes slender, at first hairy, black or brown, shining; fronds (3'–6' high) ovate-lanceolate, woolly with *soft whitish distinctly articulated flattened hairs*, becoming smoother above, twice or thrice pinnate; pinnæ (5"–6" long) ovate, the lowest distant, the others contiguous; *pinnules crenately pinnatifid*, or mostly divided into minute and roundish densely crowded segments ($\frac{1}{2}$ "–1" long), *the herbaceous margin recurved forming an almost continuous involucre*. (*C. vestita*, Hook. *C. lanosa*, Eaton, Moore, excl. syn. *C. gracilis*, Mettenius, 1859. *Myriopteris gracilis*, Fée.) — In dense tufts, on dry rocks and cliffs, Wisconsin (*T. J. Hale*), Iowa (*Vasey*), Missouri (*Nuttall, Engelmann*), thence westward and southward.

5. PELLÆA, Link. CLIFF-BRAKE. (Pl. 15.)

Sporangia in roundish or elongated clusters on the upper part of the free veins, distinct or confluent laterally so as to imitate the marginal continuous line of fructification of *Pteris*, commonly covered by a broad membranaceous and continuous (rarely interrupted) general indusium, which consists of the reflexed and altered margin of the fertile pinnule or division. Small Ferns, with 1–3-pinnate fronds, the fertile ones with narrower divisions than the sterile, but otherwise similar. Stipes generally dark-colored, smooth and shining. (Name from *πελλός*, dusky, alluding to the stipe.)

1. *P. gracilis*, Hook. Fronds smooth (3'–6' high), *delicately membranaceous and slender*, of few pinnæ, the lower ones once or twice pinnately parted

into 3-5 decurrent divisions, those of the fertile frond oblong or linear-oblong, entire or sparingly incised; of the sterile ovate or obovate, crenate or incised; veins of the fertile fronds mostly only once forked. (*Pteris gracilis*, *Michx.* *Allosorus gracilis*, *Presl*, and former ed. — Shaded calcareous rocks, Vermont and Northern and Central New York, to Wisconsin and northward: rare. July. — Rootstock very slender, creeping: stipes polished, brownish, darker and sparingly chaffy at the base.

2. **P. atropurpurea**, Link. Smooth, except some bristly-chaffy hairs on the midribs and especially on the *dark-purple and polished stalk* and rhachis, 6'-15' high; *frond coriaceous*, pale, once or below twice pinnate; the divisions broadly linear or oblong, or the sterile sometimes oval, chiefly entire, somewhat heart-shaped or else truncate at the stalked base; veins about twice forked. (*Pteris atropurpurea*, *L.* *Allosorus atropurpureus*, *Kunze*, and former ed.) — Dry calcareous rocks: not common, but of wide range. July. — Rootstock short and stout: stipes clustered.

6. ALLOSORUS, Bernhardt, Link. ROCK-BRAKE.

Fruit-dots roundish or elongated and extending far down on the free forking veins. True involucre or indusium none, the herbaceous margins of the fertile segments at first reflexed and meeting at the midrib, at length opening out flat and exposing the confluent sporangia. — Low, with smooth, 2-3-pinnate fronds, the fertile ones taller than the sterile, and with much narrower divisions. (Name from *ἄλλος*, *various*, and *σῶρος*, *a heap*.)

1. **A. acrostichoides**, Sprengel. Stipes densely tufted, straw-colored; fronds 2-3-pinnate (6'-10' high); fertile segments stalked, linear or linear-oblong (3''-5'' long), the sporangia in lines extending down the veins almost to the midrib, confluent when ripe and covering the under surface of the now fully opened segments; sterile fronds on much shorter stipes, with ovate or obovate decurrent and crenately toothed or incised segments. (*Cryptogramme acrostichoides*, *R. Br.*) — Isle Royale, Lake Superior, thence westward and northward.

7. WOODWARDIA, Smith. CHAIN-FERN. (Pl. 16.)

Fruit-dots oblong or linear, arranged in one or more chain-like rows on transverse anastomosing veinlets parallel and near to the midrib. Indusium fixed by its outer margin to the fruitful veinlet, free and opening on the side next the midrib. Veins more or less reticulated, free towards the margin of the frond. — Large Ferns, with pinnatifid or pinnate fronds. (Named for *Thomas J. Woodward*, an English botanist.)

§ 1. **ANCHISTEA**, Presl. *Sterile and fertile fronds alike: veins forming only one row of meshes (areoles).*

1. **W. Virginica**, Smith. Fronds (2°-3° high) pinnate, with numerous lanceolate pinnatifid pinnae; segments oblong; veins forming a row of narrow areoles along the midrib both of the pinnae and of the lobes, the outer veinlets free; fruit-dots oblong, one to each areole, confluent when ripe. — Wet swamps, Maine to Virginia, and southward. July.

§ 2. LORINSERIA, Presl. *Sterile and fertile fronds unlike: veins of the sterile fronds forming many rows of meshes.*

2. **W. angustifolia**, Smith. Fronds pinnatifid; sterile ones (12'–18' high) with lanceolate serrulate divisions united by a broad wing; fertile fronds taller, with narrowly linear almost disconnected divisions, the areoles and fruit-dots (4''–5'' long) in a single row each side of the secondary midribs. (*W. onocleoides*, Willd. *W. areolata*, Moore.)—Bogs, Massachusetts, near the coast, to Virginia, and southward: rare. Aug., Sept.

8. ASPLENIUM, L. SPLEENWORT. (Pl. 17.)

Fruit-dots oblong or linear, oblique, separate; the straight, or rarely curved, indusium fixed lengthwise by one edge to the upper (inner) side of the fertile vein:—in some species a part of the fruit-dots are double; the fertile vein bearing two indusia placed back to back. Veins free in all our species. (Named, from a privative and *σπλήν*, the spleen, for supposed remedial properties.)

§ 1. ASPLENIUM proper. *Indusium straight or slightly curved, attached to the upper side of the vein, rarely double.*

* *Indusium flat, or flattish, thin. (Fronds evergreen.)*

+ *Fronds pinnately lobed or parted, or simply pinnate.*

1. **A. pinnatifidum**, Nutt. *Fronds (3'–6' long) lanceolate, pinnatifid, or pinnate below, tapering above into a slender prolongation, "the apex sometimes rooting"; lobes roundish-ovate, obtuse, or the lowest pair long-acuminate; fruit-dots irregular, those next the midrib often double, even the slender prolongation fertile.*—Cliffs on the Schuylkill and Wissahickon, near Philadelphia, and southward along the Alleghanies; also sparingly westward: rare. July.—Resembles the Walking-Leaf (*Camptosorus*), but the veins are free. *Stipes brownish, becoming green higher up, and so passing into the broad pale-green midrib.*

2. **A. ebenoides**, R. R. Scott. *Fronds (4'–9' long) broadly lanceolate, pinnatifid, below pinnate, the apex prolonged and slender; divisions lanceolate from a broad base, the lower ones shorter, often proliferous, as is the apex of the frond; fruit-dots much as in the last; stipes black and polished, as is the lower part of the midrib, especially beneath.*—Limestone cliffs on the Schuylkill, near Philadelphia, R. R. Scott. A single plant was found, growing with *Camptosorus* and *Aspl. ebeneum*, of which *Rev. M. G. Berkeley* (*Journ. Royal Horticult. Soc.* July, 1866) considers it a probable hybrid.

3. **A. Trichómanes**, L. *Fronds (3'–8' long) in dense spreading tufts, linear in outline, pinnate; pinnæ numerous, roundish-oblong or oval (3''–4'' long), unequal-sided, obliquely wedge-truncate at the base, attached by a narrow point, the midvein forking and evanescent; the thread-like stipe and rhachis purple-brown and shining. (A. melanocaulon, Willd.)*—Shaded cliffs: common. July. (Eu.)

4. **A. ebeneum**, Ait. *Fronds upright (8'–16' high) pinnate, lance-linear in outline; pinnæ ($\frac{1}{2}'$ –1' long) many, lanceolate, or the lower oblong, slightly scythe-shaped, finely serrate, sessile, the dilated base auricled on the upper or both sides; fruit-dots numerous on both sides of the elongated midvein; stipe and rhachis blackish-purple and shining.*—Rocky, open woods: rather common.

← ← *Fronde once or twice pinnate: pinnæ incised.*

5. **A. montanum**, Willd. *Fronde* (2'–5' high), *ovate-lanceolate, pinnate; the ovate pinnæ 3–7-parted* (or the upper barely cleft) *and cut-toothed*; fruit-dots very short, the basal ones sometimes double. — Cliffs in the Alleghenies, Pennsylvania (*Mr. Lea, Prof. Porter*), to Virginia and southward. July. — Rhachis green, broad and flat: stipe brown at the base.

6. **A. Ruta-muraria**, L. *Fronde* (2'–4' long) *ovate in outline, 2–3-pinnate below, simply pinnate above; the few divisions rhombic-wedge-shaped, toothed or incised at the apex*; veins forking, diverging from the base; fruit-dots few, elongated, soon confluent. — Limestone cliffs, Vermont to Michigan, Virginia, and southward along the mountains: scarce. July. (Eu.)

* * *Indusium slightly curved, strongly convex, thickish: fruit-dots very numerous and crowded. (Fronde tall, simply pinnate, decaying in autumn.)*

7. **A. angustifolium**, Michx. *Fronde* (2°–3° high) thin, *simply pinnate; pinnæ numerous, short-stalked, linear-lanceolate, acuminate, entire or crenulate* (3'–4' long), *those of the fertile frond narrower*; fruit-dots linear, 20–40 each side the midvein. — Rich woods, W. New England to Wisconsin, and southward along the mountains. Sept.

* * * *Lower fruit-dots single, those towards the ends of pinnæ double: indusium straight, slightly convex, thinnish. (Fronde tall and ample, decaying in autumn.)*

8. **A. thelypteroides**, Michx. *Fronde* (2°–3° high) *pinnate; pinnæ deeply pinnatifid, linear-lanceolate* (3'–5' long); the lobes oblong, obtuse, minutely toothed, crowded, each bearing 3–6 pairs of *oblong fruit-dots*. — Rich woods: not rare. July–Sept.

§ 2. **ATHYRIUM**, Roth. *Indusium delicate, curved, often crossing the vein, and attached to both sides of it, thus becoming reniform, or shaped like a horseshoe.*

9. **A. Filix-fœmina**, Bernh. *Fronde* (1°–3° high) *ovate-oblong or broadly lanceolate, twice pinnate; pinnæ lanceolate, numerous; pinnules confluent on the secondary rhachis by a narrow margin, oblong and doubly serrate, or elongated and pinnately incised with cut-toothed segments; fruit-dots short, variously curved, at length confluent. (Aspidium Filix-fœmina & A. asplenoides, Swartz.)* — A narrow form is *Aspidium angustum, Willd.* — Moist woods: common, and very variable. July. (Eu.)

9. **SCOLOPÉNDRIUM**, Smith. HART'S-TONGUE. (Pl. 17.)

Fruit-dots linear, elongated, almost at right angles to the midrib, contiguous by twos, one on the upper side of one veinlet, and the next on the lower side of the next superior veinlet, thus appearing to have a double indusium opening along the middle. (The ancient Greek name, so called because the numerous parallel lines of fruit resemble the feet of the centipede, or *Scolopendra*.)

1. **S. vulgare**, Smith. Frond oblong-lanceolate from an auricled-heart-shaped base, entire or wavy-margined (7'–18' long, 1'–2' wide), bright green. (*S. officinarum, Swartz*, a later name.) — Shaded ravines and under limestone cliffs, Chittenango Falls, and near Jamesville, &c., Onondaga Co., New York, *Pursh, W. Cooper, Lewis Foote, J. A. Paine*: also in Canada West. (For a full account of stations, see *Silliman's Journal* for May and September, 1866. (Eu.)

10. CAMPTOSORUS, Link. WALKING-LEAF. (Pl. 17.)

Fruit-dots oblong or linear, as in *Asplenium*, but irregularly scattered on either side of the *reticulated veins* of the simple frond, those next the midrib single, the outer ones inclined to approximate in pairs (so that their two indusia open face to face), or to become confluent at their ends, thus forming crooked lines (whence the name, from *καμπτός*, *bent*, and *σώρος*, for *fruit-dot*.)

1. **C. rhizophýllus**, Link. (*Asplenium rhizophyllum*, *L.* *Antigrámma*, *J. Smith*, *Torr.* Also *C. rumicifolius*, *Link.*) — Shaded calcareous rocks, W. New England to Wisconsin, and southward: rare or local. July. — Fronds evergreen, growing in tufts, spreading or procumbent (4'–9' long), lanceolate from an auricled-heart-shaped or often hastate base, tapering above into a slender prolongation like a runner, which often roots at the apex and gives rise to new fronds, and these in turn to others; hence the popular name. — A singular form is found at Mount Joy, Penn., by *Mr. Stauffer*, having roundish fruit-dots and inconspicuous veins.

11. PHEGÓPTERIS, Fée. (Sp. of POLYPODIUM, L.) BEECH-FERN.

Fruit-dots small, round, naked (no indusium), borne on the back of the veins below the apex. Stipe continuous with the rootstock. — Our species have free veins and bright-green membranaceous fronds, decaying in early autumn. (Name composed of *φήγός*, an *oak* or *beech*, and *πτερίς*, *fern*.)

* *Fronds twice pinnatifid: pinnæ all sessile, adnate to the winged rhachis.*

1. **P. polypodioides**, Fée. Fronds triangular, longer than broad (4'–9' long), hairy on the veins, especially beneath; pinnæ linear-lanceolate, the lowest pair deflexed and standing forwards; their divisions oblong, obtuse, entire, the basal ones decurrent and adnate to the main rhachis; fruit-dots all near the margin. (*Ph. vulgaris*, *Mettenius*. *Polypodium Phegopteris*, *L.*, and former ed. *P. connéctile*, *Michx.*) — Damp woods: common northward. July. — Rootstock slender, creeping, bearing a few distant slender stalks, rather longer than the fronds. (Eu.)

2. **P. hexagonóptera**, Fée. Fronds triangular, usually broader than long (7'–12' broad), slightly pubescent and often finely glandular beneath; pinnæ lanceolate; upper segments oblong, obtuse, toothed or entire, those of the very large lowest pinnæ elongated and pinnately lobed, basal ones very much decurrent and forming a continuous many-angled wing along the main rhachis; fruit-dots near the margin; some also between the sinus and the midrib. (*Polypodium hexagonopterum*, *Michx.*) — Rather open woods: common, especially southward. July. — Larger and broader than the last, which it often closely resembles.

* * *Fronds ternate, the three divisions petioled: rhachis wingless.*

3. **P. Dryópterus**, Fée. Fronds smooth, broadly triangular (4'–6' wide); the 3 triangular primary divisions widely spreading, 1–2-pinnate; segments oblong, obtuse, entire or toothed; fruit-dots near the margin. (*Polypodium Dryopteris*, *L.*) — Rocky woods: common northward. July. (Eu.) (*P. CALCAREA*, which is more rigid, and minutely glandular-mealy, especially on the rhachis and midrib, is to be sought for northward.)

12. **ASPIDIUM**, Swartz. SHIELD-FERN. WOOD-FERN. (Pl. 18.)

Fruit-dots round, borne on the back or rarely at the apex of the veins. Indusium covering the sporangia, flat or flattish, scarious, orbicular and peltate at the centre, or round-kidney-shaped and fixed either centrally or by the sinus, opening all round the margin. Stipe continuous (not articulated) with the rootstock. — Our species have free veins, and 1-3-pinnate fronds. (Name, *ἀσπίδιον*, a small shield, from the shape of the indusium.)

§ 1. **DRYÓPTERIS**, Adanson. (Nephrodium, *Rich.*, in part, *Hook.* Lástrea, *Bory.*) Indusium round-kidney-shaped, or orbicular with a narrow sinus.

* *Veins simple or simply forked and straight: fronds annual, decaying in autumn, the stalks and slender creeping rootstocks nearly naked.* (Thelypteris, *Schott.*)

1. **A. Thelypteris**, Swartz. Fronds pinnate, lanceolate in outline; pinnæ horizontal or slightly recurved, linear-lanceolate, deeply pinnatifid, the lowest pairs scarcely smaller; lobes oblong, entire, obtuse or appearing acute when in fruit from the strongly revolute margins; veins mostly forked, bearing the (soon confluent) fruit-dots near their middle; indusium minute, smooth and naked. (Polypodium Thelypteris, *L.*) — Marshes: common. Aug. — Stalk 1° long or more, usually longer than the frond, which is of thicker texture than the next, and slightly downy. (Eu.)

2. **A. Noveboracense**, Swartz. Fronds pinnate, lanceolate in outline, tapering both ways from the middle; pinnæ lanceolate, the lowest 2 or more pairs gradually shorter and deflexed; lobes flat, oblong, basal ones often enlarged and incised; veins simple, or forked in the basal lobes; fruit-dots distinct, near the margin; indusium minute, the margin glanduliferous. (Polypodium Noveboracense, *L.* A. thelypteroïdes, *Swartz.*) — Swamps and moist thickets: common. July. — Frond pale-green, delicate and membranaceous, hairy beneath along the midribs and veins.

* * *Veins, at least the lowest ones, more than once forked or somewhat pinnately branching; the fruit-bearing veinlets often obscure or vanishing above the fruit-dot: fronds, at least the sterile ones, often remaining green through the winter: stalks and apex of the thickened rootstock scaly or chaffy, and often the main rhachis also.*

+ *Fronds small, pinnate: pinnæ pinnatifid: indusia very large, persistent.*

3. **A. fragrans**, Swartz. Fronds (4' - 12' high) glandular and aromatic, narrowly lanceolate, with linear-oblong pinnately-parted pinnæ; their crowded divisions (2'' long) oblong, obtuse, toothed or nearly entire, nearly covered beneath with the very large thin imbricated indusia, which are orbicular with a narrow sinus, the margin sparingly glanduliferous and often ragged. — Shaded trap-rocks, Falls of the St. Croix, Wisconsin, *Dr. Parry*; Wisconsin River, *Lapham*, Lake Superior, Canada, and high northward. — Rootstock stout, nearly erect, densely chaffy, as are the crowded stipes and rhachis.

+ + *Large (1° - 2½° high), the fronds mostly twice pinnate with variously toothed and incised pinnules: indusia rather small, shrivelled in age, or deciduous.*

4. **A. spinulosum**, Swartz. Stipes with a few pale-brown deciduous scales; frond ovate-lanceolate, twice pinnate; pinnæ oblique to the rhachis, elongated-trian-

gular, the lower pairs broadly triangular; pinnules set obliquely on the midribs, connected by a very narrow wing, oblong, acute, incisedly serrate or pinnatifid with spinulose-toothed lobes; *indusium* smooth and *without marginal glands*. (*A. spinulosum, elevatum, Braun. Lastrea spinulosa, Presl., Moore.*) — In damp woods, Penn Yan, New York, *Sartwell*; Vermont, *Frost*; and probably northward. July.—The common European type, rare in North America. (Eu.)

Var. **intermedium**. Scales of the stipe few, brown with a darker centre; frond broadly oblong-ovate, twice or often thrice pinnate; *pinnæ spreading, oblong-lanceolate*, the lower ones unequally triangular-ovate; pinnules crowded, ovate-oblong, spreading, pinnately divided; the oblong lobes spinulose-toothed at the apex; *margin of the indusium denticulate and beset with minute stalked glands*. (*A. intermedium, Willd. Dryopteris intermedia, Ed. 1.*) — Woods, everywhere.

Var. **dilatatum**. Scales of the stipe large, brown with a dark centre; frond broader, ovate or triangular-ovate in outline, oftenest thrice pinnate; pinnules lance-oblong, the lowest ones often much elongated; *indusium (in the North American plant) smooth and naked*. (*A. dilatatum, Swartz. A. campylópterum, Kunze.*) — A dwarf state, fruiting when only 5'–8' high, answers to var. *dumetorum*. — N. New England to Wisconsin, chiefly in mountain woods, and northward. (Eu.)

Var. **Boottii**. Scales of the stipe pale-brown; frond elongated-oblong or elongated-lanceolate in outline; pinnules broadly oblong, very obtuse, the lower pinnatifid, the upper and smaller merely serrate; *indusium minutely glandular*. (*A. Boottii, Tuckerm. Dryopteris rigida, Ed. 1; not A. rigidum, Swartz.*) — E. Massachusetts and Connecticut to New York, and northward. — The least dissected form, closely allied to the European *A. remotum, Braun*, and intermediate in appearance between *A. spinulosum* and *A. cristatum*, but passing into the former.

+ + + *Large (2°–4° high): fronds once pinnate, and the pinnæ deeply pinnatifid, or nearly twice pinnate: fruit-dots not very near the margin; the indusia large, thinnish and flat, persistent.*

5. **A. cristatum, Swartz.** Frond linear-oblong or lanceolate in outline (1°–2° long); *pinnæ short (2'–3' long), triangular-oblong*, or the lowest nearly triangular-ovate, from a somewhat heart-shaped base, acute, deeply pinnatifid; the divisions (6–10 pairs) oblong, very obtuse, finely serrate or cut-toothed, the lowest pinnatifid-lobed; *fruit-dots as near the midvein as the margin*; *indusium round-reniform, the sinus mostly shallow, smooth and naked*. (*A. Lancastriense, Spreng.*) — Swamps, &c.: common. July.—Stipes and the stout creeping rootstock bearing broad and deciduous chaffy scales. (Eu.)

Var. **Clintonianum**. Frond in every way much larger (2½°–4° long); *pinnæ oblong-lanceolate, broadest at base (4'–6' long, 1'–2' broad)*, deeply pinnatifid; the divisions (8–16 pairs) crowded or distant, linear-oblong, obtuse, obscurely serrate or cut-toothed, the basal ones sometimes pinnately lobed; veins pinnately forking, the lowest anterior veinlets bearing the *fruit-dots near the midvein*; *indusium orbicular with a shallow sinus, smooth and naked*. — Swampy woods, New England to New Jersey, New York (*G. W. Clinton, &c.*), and westward. July.—Rootstock stout, creeping, chaffy (like the stipes) with large bright-brown scales. A showy Fern, unlike any European form of *A. cristatum*, and often mistaken for *A. Goldianum*.

6. **A. Goldianum**, Hook. Frond broadly ovate, or the fertile ovate-oblong in outline (2° – 3° long); pinnæ (6'–9' long) oblong-lanceolate, broadest in the middle, pinnately parted; the divisions (about 20 pairs) oblong-linear, slightly scythe-shaped (9''–15'' long), serrate with appressed teeth; veins pinnately forking and bearing the fruit-dots very near the midvein; indusium very large, orbicular with a deep narrow sinus, smooth and without marginal glands. — Rich and moist woods, from Connecticut to Kentucky, and northward. July. — A stately Fern, often 4° high, the fronds growing in a circle from a stout ascending chaffy root-stock, and decaying in autumn. Indusium with the sides of the sinus often overlapping, thus appearing to be round and entire as in *Polystichum*.

+ + + + Large (1° – 3° high): stipes very chaffy at the base: fronds twice pinnate, but the upper pinnules confluent, some of the lower pinnatifid-toothed: fruit-dots rather large: the indusium convex, without marginal glands, persistent.

7. **A. Filix-mas**, Swartz. Frond lanceolate in outline (1° – 3° high); pinnæ linear-lanceolate, tapering from base to apex; pinnules oblong, very obtuse, serrate at the apex, and obscurely so at the sides, the basal ones incisely lobed, distinct, the upper confluent; fruit-dots nearer the midvein than the margin, and usually confined to the lower half of each fertile pinnule. — Rocky woods, Keweenaw Peninsula, Lake Superior, Dr. Robbins, and westward. — Frond thickish but not surviving the winter. (Eu.)

8. **A. marginale**, Swartz. Frond evergreen, smooth, thickish and almost coriaceous, ovate-oblong in outline (1° – 2° long); pinnæ lanceolate, broadest above the base; pinnules oblong or oblong-scythe-shaped, crowded, obtuse, entire or crenately-toothed; fruit-dots close to the margin. — Rocky hillsides in rich woods: common, especially northward. Aug.

§ 2. **POLYSTICHUM**, Roth. (*Aspidium*, Hook.) *Indusium orbicular and entire, peltate, fixed by the depressed centre: fronds rigid and coriaceous, evergreen, very chaffy on the rhachis, &c.: pinnæ or pinnules auricled at the base on the upper side, crowded, the teeth or lobes bristle-tipped.*

* Fronds simply pinnate.

9. **A. acrostichoides**, Swartz. Frond lanceolate (1° – $2\frac{1}{2}^{\circ}$ high), stalked; pinnæ linear-lanceolate, somewhat scythe-shaped, half-halberd-shaped at the slightly stalked base, serrulate with appressed bristly teeth; the fertile (upper) ones contracted and smaller, bearing contiguous fruit-dots near the midrib, which are confluent with age, covering the surface. (*Nephrodium acrostichoides*, Michx.) — Var. **INCISUM** (*A. Schweinitzii*, Beck) is a state with cut-lobed pinnæ, a not unfrequent case in the sterile fronds; sometimes the tips of almost all of them fertile more or less. — Hillsides and ravines in woods: common northward, and southward along the Alleghanies. July.

10. **A. Lonchitis**, Swartz. Frond linear-lanceolate (9'–20' high), scarcely stalked, very rigid; pinnæ broadly lanceolate-scythe-shaped, or the lowest triangular, strongly auricled on the upper side and wedge-truncate on the lower, densely spinulose-toothed (1' or less in length), copiously fruit-bearing; fruit-dots contiguous and near the margins. — Woods, southern shore of Lake Superior, and northward. (Eu.)

* *Fronds bipinnate.*

11. **A. aculeatum**, Swartz, var. **Braunii**, Koch. *Frond spreading* ($1\frac{1}{2}^{\circ}$ – 2° long), oblong-lanceolate in outline, with a tapering base, the lower of the many pairs of oblong-lanceolate pinnæ gradually reduced in size and obtuse; pinnules ovate or oblong, obtuse, truncate and almost rectangular at the base, short-stalked, or the upper confluent, sharply toothed, beset with long and soft as well as chaffy hairs. (A. Braunii, *Spenner.*)—Deep woods, mountains of New Hampshire, Vermont, N. New York, and northward. (Eu.)

13. **CYSTÓPTERIS**, Bernhardi. **BLADDER-FERN.** (Pl. 18.)

Fruit-dots roundish, borne on the back of a straight fork of the free veins; the delicate indusium hood-like or arched, attached by a broad base on the inner side (towards the midrib) partly under the fruit-dot, early opening free at the other side, which looks toward the apex of the lobe, and is somewhat jagged, soon thrown back or withering away. — Tufted Ferns with slender and delicate twice or thrice pinnate fronds; the lobes cut-toothed. (Name composed of *κύστις*, a bladder, and *πτερίς*, fern, from the inflated indusium.)

1. **C. bulbifera**, Bernh. *Frond lanceolate, elongated* (1° – 2° long), 2-pinnate; the pinnæ lanceolate-oblong, pointed, horizontal ($1'$ – $2'$ long); the *rhachis* and pinnæ often bearing *bubbles underneath, wingless*; pinnules crowded, oblong, obtuse, toothed or pinnatifid; indusium short, truncate on the free side. (*Aspidium bulbiferum*, Swartz. A. atomarium, *Muhl.!*)—Shaded ravines, &c.: common. July.

2. **C. fragilis**, Bernh. *Frond oblong-lanceolate* ($4'$ – $8'$ long, besides the stalk which is fully as long), 2–3-pinnate; the pinnæ and pinnules ovate or lanceolate in outline, irregularly pinnatifid or cut-toothed, mostly acute, *decurrent on the margined or winged rhachis*; indusium tapering or acute at the free end. — Var. *DENTATA*, Hook., is narrower and less divided, barely twice pinnate, with ovate obtuse and bluntly-toothed pinnules. (*Aspidium tenue*, Swartz.)—Shaded cliffs: common, and very variable. July. (Eu.)

14. **STRUTHIÓPTERIS**, Willd. **OSTRICH-FERN.** (Pl. 15.)

Fruit-dots round, on the pinnæ of a separate contracted and rigid frond, the margins of which are rolled backward so as to form a somewhat necklace-shaped or continuous hollow body enclosing the fruit: there are 3–5 pinnate free veinlets from each primary vein, each bearing a fruit-dot on its middle: fruit-dots crowded and confluent; the sporangia borne on an elevated receptacle which is half-encircled at its base by a very delicate semicircular and ragged evanescent indusium. — Sterile fronds large, very much exceeding the fertile, pinnate, the pinnæ pinnatifid, all growing in a close circular tuft from thick and scaly matted rootstocks. Stipes stout, angular. Pinnate veins free and simple. (Name compounded of *στρουθός*, an ostrich, and *πτερίς*, a fern, from the plume-like arrangement of the divisions of the fertile frond.)

1. **S. Germanica**, Willd. Sterile fronds smooth, broadly lanceolate, the lowest pinnæ gradually much smaller; pinnæ very numerous, narrowly lanceolate, deeply pinnatifid; the lobes oblong, obtuse: fertile frond with somewhat

necklace-shaped pinnæ, the lowest ones much smaller. (*S. Pennsylvànica*, Willd. *Onoclea Struthiopteris*, L., Hook.) — Alluvial soil: not rare northward. — Fronds intermediate between the sterile and fertile condition (bearing a few fruit-dots on contracted, but still herbaceous pinnæ) are sometimes found; a condition analogous to the var. *obtusiloba* of *Onoclea sensibilis*. (Eu.)

15. ONOCLÉA, L. SENSITIVE FERN. (Pl. 18.)

Fertile frond twice pinnate, much contracted; the pinnules short and revolute, usually so rolled up as to be converted into berry-shaped closed involucre, filled with sporangia, and forming a one-sided spike or raceme. Fruit-dots one on the middle of each strong and simple primary vein (with or without sterile cross-veins), round, soon all confluent. Indusium very thin, hood-like, lateral, fixed by its lower side, free on the upper (towards the apex of the pinnule). — Sterile fronds rising separately from the naked extensively creeping rootstock, long-stalked, broadly triangular in outline, deeply pinnatifid into lance-oblong pinnæ, which are entire or wavy-toothed, or the lowest pair sinuate-pinnatifid (decaying in autumn); veins reticulated with fine meshes. (Name apparently from *ὄνος*, a vessel, and *κλείω*, to close, from the singularly rolled up fructification.)

1. *O. sensibilis*, L. — Moist or wet places, along streams: common. July. — A rare abnormal state, in which the pinnæ of some of the sterile fronds, becoming again pinnatifid and more or less contracted, bear some fruit-dots without being much revolute or losing their foliaceous character, is the var. *obtusilobata*, Torr., *N. Y. State Fl.* (Connecticut, New York, &c) This explains the long-lost *O. obtusilobata*, Schkuhr (from Pennsylvania), which, as figured, has the sterile fronds thus 2-pinnately divided. (*Ragiopteris*, Presl., is founded on a young fertile frond of this species with the sterile frond of some *Aspidium*.)

16. WOODSIA, R. Brown. WOODSIA. (Pl. 18.)

Fruit-dots round, borne on the back of simply-forked free veins; the very thin and often evanescent indusium attached by its base all around the receptacle, under the sporangia, either small and open, or else early bursting at the top into irregular pieces or lobes. — Small and tufted pinnately-divided Ferns. (Dedicated to *Joseph Woods*, an English botanist.)

§ 1. *HYPOPÉLTIS*, Torr. *Indusium conspicuous, at first enclosing the sporangia, but early opening at the top, and splitting into several spreading jagged lobes.*

1. *W. obtusa*, Torr. Frond broadly lanceolate, minutely glandular-hairy (6'–12' high), pinnate, or nearly twice-pinnate; pinnæ rather remote, triangular-ovate or oblong (1'–2' long), bluntish, pinnately parted; segments oblong, obtuse, crenately toothed, the lower ones pinnatifid with toothed lobes; veins forked, and bearing the fruit-dots on or below the minutely toothed lobes. *W. Perriniàna*, Hook. & Grev. *Aspidium obtusum*, Weber & Mohr., Willd.) — Rocky banks and cliffs: common, especially westward. July.

§ 2. *WOODSIA* proper. *Indusium minute or evanescent, open and flat from an early stage, and concealed under the fruit-dot, its margin cleft into slender hairs or cilia.*

* *Stipes obscurely jointed near the base; the withered fronds falling away at the joint: cilia of the indusium long and inflexed over the sporangia.*

2. **W. Ilvénsis**, R. Brown. *Fronde oblong-lanceolate (2'-6' long by 1' wide), smoothish and green above, thickly clothed underneath as well as the stalk with rusty bristle-like chaff, pinnate; the pinnæ crowded, oblong, obtuse, sessile, pinnately parted, the numerous crowded segments oblong, obtuse, obscurely crenate, almost coriaceous; the fruit-dots near the margin, somewhat confluent when old. (Nephrodium rufidulum, Michx.)* — Exposed rocks: common, especially northward, and southward in the Alleghanies. June. (Eu.)

3. **W. glabélla**, R. Brown. *Smooth and naked throughout; frond linear (2'-5' high), pinnate; pinnæ roundish or ovate, the lower ones rather remote, (2''-4'' long), cut into 3-7 rounded or somewhat wedge-shaped lobes.* — Rocks, Little Falls, New York (*Vasey*); Willoughby Mountain, Vermont (*Wood, C. C. Frost*); and high northward.

* * *Stipes not jointed: cilia of the indusium very short and hidden by the sporangia.*

4. **W. Oregána**, Eaton. *Smooth throughout; fronds (2'-8' high, 8''-12'' wide) elliptical-lanceolate, pinnate, the fertile ones tallest; pinnæ triangular-oblong, obtuse, pinnatifid; segments oblong or ovate, obtuse, finely toothed, and in larger fronds incised; fruit-dots near the margin; indusium very small, divided almost to the centre into a few necklace-like-jointed cilia.* — Crevices of rocks, south shore of Lake Superior (*Robbins*), and westward.

17. DICKSONIA, L'Her. DICKSONIA. (Pl. 17.)

Fruit-dots small, globular, marginal, each placed on the apex of a free vein or fork; the sporangia borne on an elevated globular receptacle, enclosed in a membranaceous cup-shaped indusium which is open at the top, and on the outer side partly adherent to a reflexed toothlet of the frond. (Named for *James Dickson*, an English Cryptogamic botanist.)

1. **D. punctilóbula**, Kunze. *Minutely glandular and hairy (2°-3° high); fronds ovate-lanceolate and pointed in outline, pale green, very thin, with strong chaffless stalks rising from slender extensively creeping naked rootstocks, mostly bipinnate; primary pinnæ lanceolate, pointed, the secondary pinnatifid into oblong and obtuse cut-toothed lobes; fruit-dots minute, each on a recurved toothlet, usually one at the upper margin of each lobe. (D. pilosiuscula, Willd. Nephrodium punctilobulum, Michx. Dennstædtia, Moore. Sitolóbium, J. Smith.)* — Moist, rather shady places: very common: pleasantly odorous. July.

18. SCHIZÆA, Smith. SCHIZÆA. (Pl. 19.)

Sporangia large, ovoid, striate-rayed at the apex, opening by a longitudinal cleft, naked, vertically sessile in a double row along the single vein of the narrow divisions of the pinnate (or radiate) fertile appendages to the slender and simply linear, or (in foreign species) fan-shaped or dichotomously many-cleft fronds (whence the name, from $\sigma\chi\acute{\iota}\zeta\omega$, to split).

1. **S. pusilla**, Pursh. *Sterile fronds linear, very slender, flattened and tortuous; the fertile ones equally slender ($\frac{1}{4}$ '' wide), but taller (3' - 4' high), and bearing at the top the fertile appendage, consisting of about 5 pairs of crowded*

pinnae (each 1" - 1½" long). — Low grounds, pine barrens of New Jersey: very local. Sept.

19. **LYGÒDIUM**, Swartz. CLIMBING FERN. (Pl. 19.)

Fronds twining or climbing, bearing stalked and variously lobed (or compound) divisions in pairs, with mostly free veins; the fructification on separate contracted divisions or spike-like lobes, one side of which is covered with a double row of imbricated hooded scale-like indusia, fixed by a broad base to short oblique veinlets. Sporangia much as in *Schizæa*, but oblique, fixed to the veinlet by the inner side next the base, one or rarely two covered by each indusium. (Name from *λυγώδης*, *flexible*.)

1. **L. palmatum**, Swartz. Very smooth; stalks slender, flexile and twining (1° - 3° long), from slender running rootstocks; the short alternate branches or petioles 2-forked; each fork bearing a round-heart-shaped palmately 4 - 7-lobed frondlet; fertile frondlets above, contracted and several times forked, forming a terminal panicle. (*Hydroglóssum*, *Willd.*) — Shaded or moist grassy places, Massachusetts to Virginia, Kentucky, and sparingly southward: rare. Sept.

20. **OSMÚNDA**, L. FLOWERING FERN. (Pl. 19.)

Fertile fronds or fertile portions of the frond very much contracted, and bearing on the margins of the narrow rhachis-like divisions short-pedicelled and naked sporangia: these are globular, thin and reticulated, large, opening by a longitudinal cleft into two valves, and bearing near the apex a few parallel striæ, the rudiment of a transverse ring. — Fronds tall and upright, from thickened rootstocks, once or twice pinnate; veins forking and free. Spores green. (*Osmunder*, a Saxon name of the Celtic divinity, Thor.)

* *Fronds twice pinnate, fertile at the top.*

1. **O. regalis**, L. (FLOWERING FERN.) Very smooth, pale green (2° - 5° high); sterile pinnules 13 - 25, varying from oblong-oval to lance-oblong, finely serrulate, especially towards the apex, otherwise entire, or crenately lobed towards the rounded, oblique and truncate, or even cordate and semi-auriculate base, sessile or short-stalked (1' - 2' long); the fertile racemose-panicled at the summit of the frond. (*O. spectabilis*, *Willd.* *O. glaucescens*, *Link*, *Mettenius*.) — Swamps and wet woods: common. The cordate pinnules are commoner in Europe, but are sometimes found here. May, June. (Eu.)

* * *Sterile fronds once pinnate: pinnae deeply pinnatifid; the lobes entire.*

2. **O. Claytoniana**, L. Clothed with loose wool when unfolding, soon perfectly smooth (2° - 3° high); *pinnae oblong-lanceolate*, with oblong obtuse divisions; *some (2 - 5 pairs) of the middle pinnae fertile*, these entirely pinnate; sporangia greenish turning brown. (*O. interrupta*, *Michx.*, &c.) — Low grounds: common. May. — Fruiting as it unfolds. — This, being Clayton's plant (as ascertained in 1839, both from the Claytonian and Linnæan herbaria), must bear the original Linnæan name, though wrongly described from young specimens in which the fructification was thought to be terminal.

3. **O. cinnamomea**, L. (CINNAMON-FERN.) Clothed with rusty wool when young; sterile fronds smooth when full grown, the lanceolate pinnae pin-

natifid into broadly oblong obtuse divisions; *fertile fronds separate*, from the same rootstock, contracted, twice pinnate, covered with the cinnamon-colored sporangia. — Var. *FRONDOSA* is a rare occasional state, in which some of the fronds are sterile below and more sparsely fertile at their summit. (O. Claytoniana, Conrad, not of L.) — Rarely such fronds are fertile in the middle. — Swamps and low copses, everywhere. May. — Growing in large bunches; the fertile fronds in the centre, perfecting fruit as they unfold, 1°-2° long, decaying before the sterile fronds (at length 4°-5° high) get their growth.

21. BOTRYCHIUM, Swartz. MOONWORT. (Pl. 19.)

Rootstock very short, erect, with clustered fleshy roots (which are full of starch, in very minute, irregular granules!); the base of the naked stalk containing the bud for the next year's frond: frond with an anterior fertile and a posterior sterile segment; the former mostly 1-3-pinnate, the contracted divisions bearing a double row of sessile naked sporangia; these are distinct, rather coriaceous, not reticulated, globular, without a ring, and open transversely into two valves. Sterile segment of the frond ternately or pinnately divided or compound; veins all free. Spores copious, sulphur-color. (Name a diminutive of *βότρυς*, a cluster of grapes, from the appearance of the fructification.)

* *Sterile portion of the frond sessile or nearly so on the upper part of the common stalk.*

1. **B. Lunaria**, Swartz. *Sterile segment* nearly sessile, borne near the middle of the common stalk, *oblong, simply pinnate with 5-15 lunate or fan-shaped* very obtuse crenate, incised or nearly entire, *fleshy divisions*, more or less excised at the base on the lower or on both sides, the veins radiating from the base and repeatedly forking; fertile segment paniced, 2-3-pinnate. — Lake Superior (*Lesquereux*), and sparingly northward. — Plant 4'-10' high, very fleshy throughout. (Eu.)

2. **B. simplex**, Hitchcock. *Fronde small (2'-4', rarely 5'-6' high)*, the sterile segment *short-petioled from the middle or upper part of the common stalk, thickish and fleshy, simple and roundish, or pinnately 3-7-lobed*; the lobes roundish-obovate, nearly entire, decurrent on the broad and flat indeterminate rhachis; the veins all forking from the base; fertile segment simple or 1-2-pinnate. — Maine to New York, and northward: rare.

3. **B. lanceolatum**, Angström. *Fronde small (3'-8' high)*; the sterile segment *closely sessile near the top of the long and slender common stalk*, somewhat fleshy, in the smallest ones 3-lobed, in larger ones broadly triangular, ternately twice pinnatifid; the divisions lanceolate, incised or toothed; veins forking from a midvein; fertile part 2-3-pinnate. (B. simplex of former ed.) — New England and New Jersey (*C. F. Austin*) to Ohio and northward: rare. July.

4. **B. Virginicum**, Swartz. *Fronde tall and ample*; sterile segment sessile above the middle of the common stalk, broadly triangular, thin and membranaceous, ternate; the short-stalked primary divisions once or twice pinnate, and then once or twice pinnatifid; the oblong lobes cut-toothed towards the apex; veins forking from a midvein; fertile part 2-3-pinnate. — Plant 1°-2° high, or often reduced to a few inches, when it is *B. gracile, Pursh.* — Rich woods: common. June, July.

* * Sterile portion of the frond long-petioled from near the base of the common stalk.

5. **B. lunarioides**, Swartz. Sterile portion of the frond petioled from near the base, 2-3-ternate, or the ultimate divisions often pinnate or pinnately parted, broadly triangular in general outline; the lobes or divisions obovate, somewhat kidney-shaped, roundish or oblong, somewhat crenate; fertile stalk 3'-6' high; the fertile part mostly 2-pinnate. (*Botrypus lunarioides*, Michx. *Botrychium fumarioides*, Willd.) — Dry and rich woods, especially southward. July. — A state from Hingham, Massachusetts (*C. J. Sprague*), has the two lateral primary divisions of the sterile segment changed into long-stalked fertile fronds.

Var. **obliquum** (*B. obliquum*, Muhl.) is mostly larger (6'-17' high); the fructification more compound; the sterile segment with oblong or lanceolate divisions, either obtuse or oblique at the base, nearly entire, toothed, or irregularly pinnatifid. — New England to Wisconsin, and southward: rather scarce.

Var. **disséctum** (*B. dissectum*, Muhl.) has the divisions of the sterile segment compoundly and laciniately cut into narrow small lobes and teeth: otherwise as the last, into which it passes, and with which it grows.

22. OPHIOGLÓSSUM, L. ADDER'S-TONGUE. (Pl. 19.)

Mode of growth much as in *Botrychium*; but the coriaceous sporangia connate and coherent in two ranks on the edges of a simple spike, which in our species is single and placed on a stalk rising from the base of the simple and reticulated-veined sterile segment. Spores copious, sulphur-color. (Name compounded of *ὄφις*, a serpent, and *γλῶσσα*, tongue.)

1. **O. vulgátum**, L. Sterile segment ovate or elliptical-oblong (2'-3' long) rather fleshy, obtuse, narrowed at the base, and sessile near the middle of the stalk of the fertile spike. — Bogs and meadows: not common. July. — Stalk 6'-12' high, rising from a short oblique rootstock, the bud not enclosed in the base of the stalk. (Eu.)

ORDER 131. LYCOPODIACEÆ. (CLUB-MOSS FAMILY.)

Low plants, usually of Moss-like aspect; with pretty large spore-cases (sporangia or sporocarps) sessile in the axil of simple and sessile, persistent, mostly awl-shaped or lanceolate leaves: — of three genera, including the aquatic and peculiar Isoetes.

* Terrestrial, with erect or creeping stems: spore-cases free in the axils of the leaves.

1. **Lycopodium**. Spore-cases all of one sort, coriaceous, mostly kidney-shaped, 2-valved, filled with innumerable minute spores.
2. **Selaginella**. Spore-cases of two sorts; one very small, oblong or globular, 2-valved, filled with innumerable minute spores; the other larger, 3-4-valved, containing 3 or 4 large spores.

* * Aquatic, with a corm in place of stem, covered above with the dilated bases of the tufted long and rush-like leaves, to which the spore-cases adhere.

3. **Isoetes**. Spore-cases of two sorts; one containing numerous large, the other numerous small spores.

1. **LYCOPÒDIUM**, L., Spring. CLUB-MOSS. (Pl. 20.)

Spore-cases all of one kind (much like those of *Ophioglossum*, only larger), coriaceous, flattened, usually kidney-shaped, 1-celled, 2-valved, mostly by a transverse line round the margin, discharging the subtile spores in the form of a copious sulphur-colored inflammable powder. — Perennials, with evergreen one-nerved leaves, imbricated or crowded in 4–16 ranks. (Name compounded of *λύκος*, a wolf, and *πούς*, foot, from no obvious resemblance.)

§ 1. *Spore-cases in the axils of the ordinary and uniform (dark-green and shining, rigid, lanceolate, spreading, about 8-ranked) leaves.*

1. **L. lucidulum**, Michx. Stems thick, 2 or 3 times forked, the branches ascending (6'–12' high); *leaves widely spreading or reflexed, acute, minutely toothed.* — Cold, damp woods: common northward. Aug. — Little bulblets form in the axils of the leaves of young shoots (*Austin, Rothrock*).

2. **L. Selàgo**, L. Stems thick and rigid, erect, fork-branched, forming a level topped cluster (3'–6' high): *leaves pointed, entire.* — Tops of high mountains, Maine to New York, on the Alleghanies southward, shore of Lake Superior, and northward: rare: both the variety with more erect, and that with widely spreading leaves. (Eu.)

§ 2. *Spore-cases only in the axils of the upper (bracteal) leaves, thus forming a spike.*

* *Leaves of the creeping sterile and of the upright fertile stems or branches, and those of the simple spike all alike, many-ranked (sporangia opening near the base).*

3. **L. inundatum**, L. Dwarf; creeping sterile stems forking, flaccid; the fertile solitary (1'–4' high), bearing a short thick spike; *leaves lanceolate or lance-awl-shaped, acute, soft, spreading, naked, or sometimes bearing a few minute spiny teeth.* — Leaves (curving upwards on the prostrate shoots) narrower in the American than in the European plant (perhaps a distinct species), and passing into the var. *BIGELÒVII*, Tuckerm., which has fertile stems 5'–7' high, its leaves more awl-shaped and pointed, sparser and more upright, often somewhat teeth-bearing. (L. *Carolinianum*, *Bigel.*, not of L.) — Sandy bogs, northward: rare: the var. E. New England to New Jersey and southward. Aug. (Eu.)

4. **L. alopecuroides**, L. Stems stout, very densely leafy throughout; the sterile branches recurved-procumbent and creeping; the fertile of the same thickness, 6'–20' high; *leaves narrowly linear-awl-shaped, spinulose-pointed, spreading, conspicuously bristle-toothed below the middle; those of the cylindrical spike with long setaceous tips.* — Pine-barren swamps, New Jersey to Virginia, and southward. Aug., Sept. — Stems, including the dense leaves, $\frac{1}{2}$ ' thick; the comose spike, with its longer spreading leaves, $\frac{3}{4}$ ' to 1' thick.

* * *Leaves (bracts) of the catkin-like spike scale-like, imbricated, yellowish, ovate or heart-shaped, very different from those of the sterile stems and branches.*

+ *Spikes sessile (i. e. branches equally leafy to the top), single.*

5. **L. annòtinum**, L. Much branched; stems prostrate and creeping (1°–4° long); *the ascending branches similar (5'–8' high), sparingly forked, the sterile ones making yearly growths from the summit; leaves equal, spreading, in about 5 ranks, rigid, lanceolate, pointed, minutely serrulate (pale green); spike soli-*

tary, oblong-cylindrical, thick. — Var. *PUNGENS*, Spring, is a reduced sub-alpine or mountain form, with shorter and more rigid-pointed erectish leaves. (Var. *montanum*, *Tuckerm.*) — Woods: common northward: the var. on the White Mountains, with intermediate forms around the base. July. (Eu.)

6. *L. dendroideum*, Michx. (GROUND-PINE.) *Stems upright (6'–9' high) from a subterranean creeping rootstock, simple below, and clothed with homogeneous lanceolate-linear acute entire leaves appressed-erect in 4–6 rows, bushy-branched at the summit; the crowded branches spreading, fan-like, with the lower row of leaves shorter and the lateral spreading: — in var. OBSCURUM appearing flat, from the leaves of the upper side being also shorter and appressed. (L. obscurum, L.) — Moist woods. Aug. — Remarkable for its tree-like growth. Spike cylindrical, 4–10 on each plant.*

+ + *Spikes peduncled: i. e. the leaves minute on the fertile branches.*

+ *Leaves homogeneous and equal, many-ranked; stems terete.*

7. *L. clavatum*, L. (COMMON CLUB-MOSS.) *Stems creeping extensively, with similar ascending short and very leafy branches; the fertile terminated by a slender peduncle (4'–6' long), bearing about 2–3 (rarely 1 or 4) linear-cylindrical spikes; leaves linear-awl-shaped, incurved-spreading (light green), tipped, as also the bracts, with a fine bristle. — Dry woods: common, especially northward. July. (Eu.)*

+ + *Leaves of two forms, few-ranked: stems or branches flattened.*

8. *L. Carolinianum*, L. *Sterile stems and their few short branches entirely creeping (leafless and rooting on the under side), thickly clothed with broadly lanceolate acute and somewhat oblique 1-nerved lateral leaves widely spreading in 2 ranks, and a shorter intermediate row appressed on the upper side; also sending up a slender simple peduncle (2'–4' high, clothed merely with small bract-like and appressed awl-shaped leaves), bearing a single cylindrical spike. — Wet pine-barrens, New Jersey to Virginia, and southward. July.*

9. *L. complanatum*, L. *Stems extensively creeping (often subterranean), the erect or ascending branches several times forked above; bushy branchlets crowded, flattened, all clothed with minute imbricated-appressed awl-shaped leaves in 4 ranks, with decurrent-united bases, the lateral rows with somewhat spreading tooth-like tips, those of the upper and under rows smaller, narrower, wholly appressed; peduncle slender, bearing 2–4 cylindrical spikes. — Woods and thickets: common: the typical form with spreading fan-like branches, abundant southward; — while northward, especially far northward, it passes gradually into var. *SABINÆFOLIUM* (*L. sabinæfolium*, *Willd.* *L. Chamæcyparissus*, *Braun*), with more erect and fasciated branches. (Eu.)*

2. SELAGINÉLLA, Beauv., Spring. (Pl. 20.)

Fructification of two kinds, namely, of spore-cases like those of *Lycopodium*, but minute and oblong or globular, containing reddish or orange-colored powdery spores; and of 3–4-valved tumid larger ones, filled by 3 or 4 (rarely 1–6) much larger globose-angular spores (*oophoridia*); the latter either intermixed with the former in the same axils, or solitary (and larger) in the lower axils of the leafy 4-ranked sessile spike. (Name a diminutive of *Selago*, an ancient name of a *Lycopodium*, from which this genus is separated.)

* *Leaves all alike and uniformly imbricated; those of the spike similar.*

1. **S. selaginoides**, Gray. *Sterile stems prostrate or creeping, small and slender; the fertile thicker, ascending, simple (1'-3' high); leaves lanceolate, acute, spreading, sparsely spinulose-ciliate. (S. spinòsa, Beauv. S. spinulòsa, Braun.)* — Wet places, New Hampshire (*Pursh*), Michigan, Lake Superior, and northward: rare. — Leaves larger on the fertile stems, yellowish-green. (Eu.)

2. **S. rupéstris**, Spring. *Much branched in close tufts (1'-3' high); leaves densely appressed-imbricated, linear-lanceolate, convex and with a grooved keel, minutely ciliate, bristle-tipped; those of the strongly quad-angular spike rather broader; the two sorts of fructification in the same axils. (Lycopodium rupestre, L.)* — Dry and exposed rocks: very common. — Grayish-green in aspect, resembling a rigid Moss.

* * *Leaves shorter above and below, resembling stipules: the lateral larger, 2-ranked.*

3. **S. ápus**, Spring. *Stems tufted and prostrate, creeping, much branched, flaccid; leaves pellucid-membranaceous; the larger spreading horizontally, ovate, oblique, mostly obtuse; the smaller appressed, taper-pointed; those of the short spikes nearly similar; larger spore-cases copious at the lower part of the spike. (Lycopodium ápodum, L.)* — Low, shady places: not rare, especially southward. — A delicate little plant, resembling a Moss or *Jungermannia*.

3. ISÔETES, L. QUILLWORT. (Pl. 20.)

Stem or trunk a fleshy more or less depressed corm, rooting just above its 2-lobed (or in many foreign species 3-lobed) base, above covered with the dilated and imbricated bases of the awl-shaped or linear somewhat quadrangular leaves, which include 4 air-tubes, intercepted by cross partitions. Sporangia, or *sporocarps*, pretty large, orbicular or ovoid, plano-convex, very thin, sessile in the axils of the leaves, and united at the back with their excavated bases (the thin edges of the excavation folding round partly cover them, forming the *velum*), traversed internally by transverse threads; those of the outer leaves filled with large spherical spores (*macrospores*), their whitish crustaceous integument marked by one circular, and on the upper surface by 3 radiating elevated lines (circumscribing a lower hemisphere, and three upper segments which open valve-like in germination): those of the inner leaves filled with very minute and powdery grayish spores (*microspores*); these are always obliquely oblong and triangular. — Mostly small aquatics, grass-like or rush-like in aspect, some always submerged, others amphibious, a few living in merely moist soil, maturing their fruit in late summer and early autumn, except No. 7, and some forms of No. 6.

Genus much investigated of late by *Prof. Durieu* and the late *J. Gay* of France, and by *Prof. Braun* of Berlin, newly elaborated for this edition by Dr. GEORGE ENGELMANN.

* *Growing under water, only accidentally or in very dry seasons out of water: leaves without stomata (except in some forms of No. 3), and without peripheral bast-bundles.*

1. **I. lacústris**, L. *Leaves (10-25 in number, 2'-6' long) dark green, rigid; sporocarps ovoid or circular, the upper third or less covered by the velum, free part pale and unspotted; both kind of spores the largest of our spe-*

cies; macrospores ($0''.32-0''.38$ wide) covered with short and twisted crested ridges, which often anastomose; microspores ($0''.017-0''.020$ long) smooth. — Mountain lakes, Penn., New York, and New England to Lake Superior, and northward often with No. 3. (Eu.) — The American is distinguished from the European plant by the larger macrospores, therefore *I. macrospora*, *Durieu*.

2. *I. Tuckermàni*, Braun, n. sp. Leaves (10–30, 2'–3' long) very slender, awl-shaped, olive-green, the outer ones recurved; sporocarps ovoid or circular, the upper third covered by the velum, the free part sometimes brownish-spotted; macrospores ($0''.22-0''.28$ wide) on the upper segments covered with parallel and anastomosing ridges, the lower half reticulated; microspores ($0''.013-0''.015$ long) smooth or very delicately papillose. — Mystic and other ponds near Boston, together with the next, *Tuckerman*, *W. Boott*.

3. *I. echinóspora*, *Durieu*. Leaves slender, awl-shaped; sporocarps ovoid or circular; macrospores ($0''.20-0''.25$ wide) beset all over with small entire and obtuse or slightly forked spinules. (Eu.) — In the European form, which has not yet been found in America, the leaves are very slenderly attenuated (3'–4' long), the upper margin of the sporocarp only is covered with the narrow velum, the free part is unspotted, and the slightly papillose microspores are larger ($0''.015-0''.016$ long). The following are the American forms of this species.

Var. *Braunii*. Leaves (15–30 in number, 3'–6' long) dark and often olive-green, straight or commonly recurved, half or two thirds of the sporocarp covered by the velum, the free part often with light brown spots; macrospores as in the species; microspores smaller ($0''.013-0''.014$ long), smooth (*I. Braunii*, *Durieu*.) — Ponds and lakes, New England to Western New York and northward, often with the two preceding. — Often with a few stomata, especially in Niagara specimens.

Var. *muricàta*. Leaves (15–30, 6'–10' long) straight or flaccid, bright green; about one half of the almost circular sporocarp covered by the velum, unspotted; macrospores ($0''.22-0''.27$ wide) with shorter and blunter spinules; microspores as in the last variety, or rarely spinulose. (*I. muricata*, *Durieu*.) — In some ponds north of Boston, *W. Boott*.

Var. *Boóttii*. Leaves (12–20, 4'–5' high) awl-shaped, stiffly erect, bright green, with stomata; sporocarp as in last; macrospores as in the species, but a little smaller and with very slender spinules. (*I. Boottii*, *Braun*, in litt.) — Pond in Woburn, near Boston, partly out of water, *W. Boott*.

* * Growing partly out of water, either by the pond drying up, or by the receding of the ebb tide; leaves with stomata, and in 6 and 7 with four or more peripheral bast-bundles.

4. *I. saccharàta*, Engelm., n. sp. Leaves (10–15, 2'–3' long) slender, olive-green, curved; sporocarps small, ovoid, only the upper edge covered by the velum, nearly unspotted; macrospores ($0''.20-0''.22$ wide) minutely tuberculate; microspores ($0''.012$ long) papillose. — On Wicomico River, eastern shore of Maryland, between high and low tide, *W. M. Canby*.

5. *I. ripària*, Engelm. Leaves (15–30, 4'–8' long) slender, deep green, erect; sporocarps mostly oblong, upper margin to one third covered by the

velum, the free part spotted; macrospores very variable in size ($0''.22-0''.30$ wide), the upper segments covered by short crested ridges, which on the lower hemisphere run together forming a network; microspores larger than in any other species except No. 1 ($0''.013-0''.016$ long), mostly somewhat tuberculated. — Gravelly banks of the Delaware, from above Philadelphia to Wilmington, between flood and ebb tide; margins of ponds, New England (*Robbins, &c.*), and northward. — Distinguished from the nearly allied *I. lacustris* by the stomata of the leaves, the spotted sporocarp, the smaller size of the spores and their reticulation on the lower half.

6. **I. Engelmanni**, Braun. Leaves long (25–100, 9'–20' long), light green, erect or at last prostrate, flat on the upper side; sporocarps mostly oblong, unspotted, the velum very narrow; macrospores ($0''.19-0''.24$ wide) covered all over with a coarse honey-comb-like network; microspores ($0''.012-0''.014$ long) mostly smooth. — Shallow ponds and ditches, from Massachusetts (near Boston, *W. Boott, H. Mann*) to Pennsylvania and Delaware and (probably through the Middle States) to Missouri. — By far the largest of our species, often mature in July.

Var. **gracilis**. Leaves few (8–12 only, 9'–12' long) and very slender; both kinds of spores nearly as in the species. — Southern New England, apparently in deep water.

Var. **válida**. Trunk large and stout (often 1'–2' wide); leaves (50–100, even 200, 18'–25' long) with an elevated ridge on the upper side; sporocarps oblong or linear-oblong ($4''-9''$ long), one third or one half or more covered by the velum; both kinds of spores very small, macrospores $0''.16-0''.22$ wide, microspores $0''.011-0''.013$ long, spinulose. — Delaware (*W. M. Canby*), and Pennsylvania (*Prof. Porter*). Sept.

7. **I. melanópoda**, J. Gay. Leaves (15–50, 6'–10' long) very slender, keeled on the back, straight, bright green, usually with dark brown or black shining bases; sporocarps mostly oblong, with a very narrow velum, brown or spotted; macrospores very small ($0''.14-0''.18$ wide), smoothish, or with faint tubercles or ridges; microspores ($0''.010-0''.012$ long) spinulose. — Shallow ponds, and wet prairies and fields, Central and Northern Illinois (*E. Hall, Vasey*), and westward. June, and sometimes again in November. — Trunk more spherical and more deeply 2-lobed, and both kind of spores smaller than in any other of our species; leaves disappearing during the summer heat. Closely approaching the completely terrestrial species of the Mediterranean region.

Two other species, *I. FLACCIDA*, Shuttleworth, and *I. CHAPMANI*, Engelm., are found in Florida; and three more, *I. NUTTALLII*, Braun, *I. CALIFORNICA*, Engelm., and *I. PYGMÆA*, Engelm., in the Pacific States.

ORDER 132. HYDROPTÉRIDES. (MARSILIACEÆ, R. Br.)

Aquatic cryptogamous plants, of diverse habit, with the fructification borne on submerged branches, consisting of two sorts of organs, contained in irregularly bursting involucre (sporocarps):—here represented by only two genera.

1. **MARSÍLIA**, L. MARSILIA.

Submersed or emersed aquatic plants, with slender creeping rootstocks, sending up elongated petioles, which bear at their apex a whorl of 4 nervose-veined leaflets, and at or near their base, or sometimes on the rootstock, one or more globular but somewhat eccentric sporocarps. These sporocarps or fruit are 2-celled vertically, and with many transverse partitions, and split or burst into 2 lobes at maturity. On the partitions are inserted numerous short-stalked sporangia, of two sorts intermixed; the larger ones containing a single oval or oblong spore, the smaller containing many very minute spores. (Named for *Aloysius Marsili*, an early Italian naturalist, — therefore to be written Marsilia, not Marsilea.)

1. **M. quadrifolia**, L. Leaflets broadly obovate-cuneate, glabrous; sporocarps usually 2 or 3 on a short peduncle from near the base of the petioles, pedicelled, glabrous or somewhat hairy. — In water, the leaflets commonly floating on the surface; Bantam Lake, Litchfield, Connecticut, *Dr. T. F. Allen*. The only known habitat in America. (Eu.)

2. **M. uncinata**, Braun., with hairy leaflets, and villous short-stalked or sessile sporocarps, solitary at the base of each petiole, will doubtless be found in the northwestern part of Wisconsin. It has been confounded with the very similar *M. vestita*, Hook and Grev., of the Southwest.

2. **AZÓLLA**, Lam. AZOLLA. (Pl. 20.)

Plant floating free, pinnately branched, clothed with minute imbricated leaves, appearing like a small *Jungermannia*; fructification sessile on the under side of the branches, of 2 sorts. Sporocarps covered at first with an indusium of a single diaphanous membrane, ovoid: the smaller kind opening transversely all round, containing several roundish *antheridia*? peltately attached to the sides of a central erect column: the large or fertile kind bursting irregularly, filled with numerous spherical sporangia rising from the base on slender stalks, each containing a few globular spores. (Name said to come from *ἄζω*, *to dry*, and *ἄλλω*, *to kill*, being destroyed by dryness.)

1. **A. Caroliniána**, Willd. Leaves ovate-oblong, obtuse, spreading, reddish underneath, beset with a few bristles. — Still water, New York to Illinois and southward. Plant forming little mats on the water, 6" — 12" broad.

SALVÍNIA NATANS, L., said by Pursh to grow floating on the surface of small lakes in Western New York, has not been found by any other person, and probably does not occur in this country. It is therefore omitted.

A D D E N D A .

THE FOLLOWING PLANTS HAVE BEEN DETECTED WITHIN OUR BORDERS TOO LATE FOR INSERTION IN THEIR PROPER PLACES.

Thermopsis mollis, M. A. Curtis. Genus like *Baptisia*, p. 142, but with a long, narrow, flat pod: plant 1°-2° high, minutely soft-downy; leaflets wedge-obovate varying to oblong; raceme reclining; flowers golden yellow; pod long and linear. (*Baptisia mollis*, DC.)—Franklin, S. Virginia (*W. M. Canby*), and southward. May.

Leucanthemum vulgare, Lam. . to p. 266, line 2, add—Var. **TUBULIFLORUM**, S. Tenney. A curious abnormal state, with the rays changed into tubular and palmately or bilabiate 5-lobed (rarely 3-4-lobed) corollas.—Fields, Poughkeepsie, New York, *Prof. Tenney*.

Vaccinium tenellum, Ait. Most like *V. Pennsylvanicum*, No. 11, p. 291; but taller (1°-3° high), with firmer and obscurely serrulate leaves, and a narrower cylindraceous corolla.—Pine-barren swamps, Franklin, S. Virginia (*W. M. Canby*), and southward. April, May.

Potamogeton. The synonymes cited under No. 3 and No. 8, pp. 485, 486, are names given by Chamisso merely as provisional appellations.

INDEX.

*** The names of the Classes, Subclasses, and the Latin names of Orders, are in full capitals ; of the Suborders, Tribes, &c., in small capitals ; of the Genera, &c., as well as popular names and synonymes, in common type.

| | | | | | |
|--------------------|----------|---------------------|-----|-------------------------|-----|
| Abele-tree | 467 | Alkanet | 363 | ANDROMEDEÆ | 287 |
| Abies | 471 | Alligator Pear | 422 | Andropogon | 651 |
| ABUETINEÆ | 468 | Allium | 533 | Androsace | 314 |
| Abutilon | 101 | Allosorus | 660 | Anemone | 36 |
| Acacia | 131, 145 | Allspice, Wild | 423 | Angelica | 192 |
| Acalypha | 436 | Almond Family | 146 | Angelica-tree | 199 |
| Acanthus Family | 338 | Alnus | 460 | Angelico | 194 |
| ACANTHACEÆ | 338 | Aloe | 513 | ANGIOSPERMÆ | 33 |
| Acer | 118 | Alopecurus | 608 | Anise Hyssop | 353 |
| Acerates | 397 | Alpine Azalea | 301 | Anisophyllum | 431 |
| ACERINEÆ | 117 | Alpine Bistort | 415 | ANONACEÆ | 50 |
| Achillea | 265 | Alsine | 91 | Antennaria | 269 |
| Acnida | 413 | ALSINEÆ | 87 | ANTHEMIDEÆ | 220 |
| Aconitum (Aconite) | 46 | Alum-root | 169 | ANTHEMIS | 265 |
| Acorus | 478 | Althæa | 99 | ANTHOXANTHÆ | 606 |
| ACROGENS | 653 | ALYSSINEÆ | 63 | ANTHOXANTHUM | 643 |
| Actæa | 47 | Alyssum | 72 | Antigramma | 663 |
| Actinella | 263 | Amaranth | 411 | ANTIRRHINEÆ | 324 |
| Actinomeris | 258 | Amaranth Family | 411 | ANTIRRHINIDEÆ | 324 |
| Adam-and-Eve | 510 | Amarantus | 411 | Antirrhinum | 326 |
| Adam's Needle | 535 | AMARANTACEÆ | 411 | Anychia | 96 |
| Adder's-mouth | 509 | Amarelloides | 387 | Apalante | 496 |
| Adder's-tongue | 533, 672 | AMARYLLIDACEÆ | 512 | Aphyllon | 323 |
| Adelia | 402 | Amaryllis | 513 | Apetalous Exogenous Pl. | 403 |
| Adenocaulon | 227 | Amaryllis Family | 512 | Apios | 140 |
| Adenorhachis | 161 | Ambrina | 407 | Aplectrum | 510 |
| Adiantum | 658 | Ambrosia | 250 | APOCYNACEÆ | 392 |
| Adice | 445 | Amelanchier | 162 | Apocynum | 393 |
| Adlumia | 60 | American Aloe | 513 | Apple | 161 |
| Adonis | 48 | American Centaury | 384 | Apple of Peru | 382 |
| Æschynomneme | 134 | American Columbo | 386 | AQUIFOLIACEÆ | 305 |
| Æsculus | 117 | American Cowslip | 314 | Aquifolium | 306 |
| Æthusa | 193 | American Frog's-bit | 495 | Aquilegia | 45 |
| Agathophyton | 408 | American Ipecac | 150 | ARABIDEÆ | 63 |
| Agave | 513 | American Ivy | 113 | Arabis | 67 |
| Agrimonia | 151 | Amanthium | 525 | ARACEÆ | 475 |
| Agrimony | 151 | Ammadenia | 92 | Aralia | 198 |
| Agropyron | 637 | Ammannia | 182 | ARALIACEÆ | 198 |
| Agrostemma | 90 | Ammophila | 616 | Arbor Vitæ | 472 |
| AGROSTIDEÆ | 603 | Amorpha | 130 | ARBUTEÆ | 287 |
| Agrostis | 611 | Ampelopsis | 113 | Arbutus | 292 |
| Ailanthus | 110 | Amphicarpæa | 141 | Arbutus (Trailing) | 293 |
| Aira | 641 | Amphicarpum | 644 | Archangelica | 193 |
| Airopsis | 641 | Amsonia | 393 | Archemora | 192 |
| AJUGOIDEÆ | 342 | AMYGDALÆ | 146 | Arctostaphylos | 292 |
| Alder | 460 | Anacharis | 495 | Arenaria | 90 |
| Alder-Buckthorn | 115 | ANACARDIACEÆ | 111 | Arethusa | 506 |
| Alechmilla | 151 | ANAGALLIDEÆ | 313 | ARTHUSEÆ | 497 |
| Aletris | 515 | Anagallis | 316 | Argemone | 59 |
| Alisma | 491 | Anantherix | 398 | Arisæma | 475 |
| ALISMACEÆ | 490 | Anchistea | 660 | Aristida | 613 |
| ALISMEÆ | 490 | Andromeda | 295 | Aristolochia | 404 |

| | | | | | |
|----------------------|----------|--------------------|----------|-----------------------|----------|
| ARISTOLOCHIACEÆ | 403 | Basswood | 103 | Bladderwort Family | 317 |
| Armeria | 312 | Bastard Pennyroyal | 344 | Blazing-Star | 223, 527 |
| Arnica | 271 | Bastard Toad-flax | 425 | Blephilia | 352 |
| Aromatic Wintergreen | 293 | Batodendron | 290 | Blessed Thistle | 272 |
| Arrhenantherum | 642 | Batrachium | 40 | Bletia | 509 |
| Arrow Arum | 476 | Batschia | 363 | Blitum (Blite) | 408 |
| Arrow-Grass | 491 | Bayberry | 457 | Blood-root | 60 |
| Arrow-Grass Family | 490 | Beach Pea | 139 | Bloodwort Family | 514 |
| Arrow-head | 492 | Beak-Rush | 567 | Blue Beech | 457 |
| Arrow-wood | 206 | Bean | 140 | Blueberry | 289 |
| Artemisia | 266 | Bearberry | 292 | Bluebottle | 272 |
| ARTOCARPEÆ | 441 | Beard Grass | 612, 651 | Blue Cohosh | 53 |
| Aruncus | 150 | Bear-Grass | 535 | Blue Curis | 343 |
| Arundinaria | 636 | Beard-tongue | 327 | Blue Flag | 516 |
| Arundo | 636 | Beaver-Poison | 196 | Blue-Hearts | 333 |
| Asarum | 403 | Bech | 455 | Buets | 213 |
| Asarabacca | 403 | Beech-drops | 322 | Blue-eyed Grass | 517 |
| ASCLEPIADACEÆ | 394 | Beech-Fern | 663 | Blue Grass | 600 |
| ASCLEPIADEÆ | 394 | Beet | 405 | Blue Joint-Grass | 615 |
| Asclepias | 394 | Bedstraw | 208 | Blue Lettuce | 281 |
| Ascyrum | 83 | Beggar's Lice | 366 | Blue Tangle | 289 |
| Ash | 401 | Beggar-ticks | 261 | Blue-weed | 361 |
| Ash-leaved Maple | 119 | Bellflower | 285 | Blysmus | 560 |
| Asimina | 50 | Bellis | 239 | Blyttia | 613 |
| ASPARAGINEÆ | 521 | Bellwort | 628 | Boehmeria | 445 |
| Asparagus | 531 | Bengal Grass | 650 | Bog-Asphodel | 535 |
| Aspen | 466 | Benjamin-bush | 423 | Bog-Rush | 637 |
| ASPIDIÆ | 657 | Bent-Grass | 611 | Boitonia | 238 |
| Aspidium | 664 | Benzoïn | 423 | Bonamia | 376 |
| ASPLENIEÆ | 656 | BERBERIDACEÆ | 52 | Boneset | 226 |
| Asplenium | 661 | Berberis | 52 | BORRAGEÆ | 360 |
| Aster | 228 | Berchemia | 114 | BORAGE Family | 360 |
| Asteranthemum | 530 | Bergamot | 351 | BORRAGINACEÆ | 360 |
| ASTERINEÆ | 218 | Bermuda Grass | 622 | Borrchia | 253 |
| ASTEROIDEÆ | 218 | Berula | 196 | Bottlebrush-Grass | 639 |
| Astilbe | 167 | Betonica (Betony) | 358 | Bottle-Grass | 650 |
| ASTRAGALEÆ | 124 | Betony (Wood) | 337 | Botrychium | 671 |
| Astragalus | 132 | Betula | 458 | Botryois | 407 |
| Atamasco Lily | 513 | BETULACEÆ | 458 | Bouncing Bet | 88 |
| Atheropogon | 621 | Bidens | 261 | Bouteloua | 620 |
| Athyrium | 662 | Bigelovia | 245 | Bowman's root | 150 |
| Atragene | 35 | Bignonia | 321 | Boxberry | 293 |
| Atriplex | 409 | BIGNONIACEÆ | 320 | Box-Elder | 119 |
| Aurantiaceæ | 110 | Bignonia Family | 320 | Box Family | 430 |
| Avena | 640 | Bilberry | 289 | Boykinia | 169 |
| AVENEÆ | 606 | Bilsted | 174 | Brachychæta | 239 |
| Avens | 151 | Bindweed | 375 | Brachyelytrum | 614 |
| Avicularia | 417 | Biotia | 228 | Bracted Bindweed | 375 |
| Awlwort | 73 | Birch Family | 458 | Brake, Bracken | 658 |
| Awned Wheat-Grass | 618 | Birthroot | 623 | Bramble | 156 |
| Azalea | 299 | Birthwort | 404 | Brasenia | 55 |
| Azolla | 677 | Birthwort Family | 403 | Brasiletto Family | 125 |
| | | Bishop's-Cap | 170 | Brassica | 70 |
| BACCHARIDEÆ | 219 | Bistorta | 415 | BRASSICEÆ | 63 |
| Baccharis | 247 | Bitter-Cress | 66 | Bread-fruit Family | 441 |
| Bald Cypress | 473 | Bittersweet | 116, 380 | Bristly Foxtail-Grass | 649 |
| Baldwinia | 284 | Bitter-weed | 251 | Briza | 633 |
| Ballota | 359 | Black Alder | 307 | Brizopyrum | 628 |
| Balm | 350 | Blackberry | 167 | Brome-Grass | 634 |
| Balm of Gilead | 467 | Blackberry-Lily | 617 | BROMELIACEÆ | 515 |
| Balmory | 327 | Black Bindweed | 418 | Brooklime | 332 |
| Balsam | 108 | Black-Gum-tree | 201 | Brook-weed | 317 |
| Balsam Family | 106 | Black-Grass | 539 | Broom-Corn | 652 |
| BALSAMIFLUE | 173 | Black Haw | 206 | Broom-Crowberry | 440 |
| BALSAMINEÆ | 106 | Black Horehound | 360 | Broom-rape | 323 |
| BAMBUSEÆ | 605 | Black-Jack | 453 | Broom-rape Family | 322 |
| Baneberry | 47 | Black Moss | 515 | Bromus | 634 |
| Baptisia | 142 | Black-Mustard | 71 | Brunella | 355 |
| Barbarea | 69 | Black Oat-Grass | 617 | Brunnichia | 422 |
| Barberry | 52 | Black Snakeroot | 190 | Buchnera | 333 |
| Barberry Family | 52 | Black Thorn | 148, 160 | BUCHNEREÆ | 325 |
| Barley | 638 | Bladder-Fern | 667 | Buckbean | 390 |
| Barnyard-Grass | 649 | Bladder-Ketmia | 102 | Buckthorn | 114 |
| Barren Strawberry | 153 | Bladder-nut | 117 | Buckthorn Family | 113 |
| Bartonia | 184, 389 | Bladder-nut Family | 117 | Buckeye | 117 |
| Basil | 346, 349 | Bladder-pod | 73 | Buckwheat | 419 |
| Basil Thyme | 349 | Bladderwort | 318 | Buckwheat Family | 414 |

| | | | | | |
|--------------------|----------|-------------------|----------|-----------------------|----------|
| Buffalo-berry | 425 | Cancer-root | 322, 323 | Chelone | 327 |
| Buffalo-nut | 426 | Cane | 636 | CHELONEÆ | 324 |
| Bugbane | 48 | CANNABINEÆ | 442 | CHENOPODIACEÆ | 405 |
| Bagle-weed | 345 | Cannabis | 446 | Chenopodia | 410 |
| Bugloss | 362 | Caper Family | 75 | Chenopodium | 406 |
| Bug-seed | 409 | CAPPARIDACEÆ | 75 | Cherry | 147 |
| Bulrush | 560 | CAPRARIA | 329 | Chervil | 197 |
| Bumelia | 308 | CAPRIFOLIACEÆ | 202 | Chess | 331 |
| Bunch-berry | 200 | Caprifolium | 203 | Chestnut | 454 |
| Bunch-flower | 524 | Capsella | 73 | CHICORACEÆ | 221, 275 |
| Bupleurum | 195 | Carex | 571 | Chickweed | 92 |
| Burdock | 275 | Cardamine | 66 | Chickweed-Wintergreen | 314 |
| Bur-Grass | 650 | Cardinal Flower | 283 | Chimaphila | 303 |
| Burmanna | 496 | Carduus | 274 | CHINCHONEÆ | 208 |
| BURMANNIACEÆ | 496 | CARICINEÆ | 551 | Chinquapin | 455 |
| Bur-Marigold | 261 | Carnation | 88 | Chiogenes | 292 |
| Burnet | 150 | Carolina Allspice | 162 | Chionanthus | 401 |
| Burning-bush | 116 | Carpel-weed | 97 | Chives | 534 |
| Bur-reed | 481 | CARPINEÆ | 450 | CHLORIDEÆ | 604 |
| Bush-clover | 137 | Carpinus | 457 | Choke-berry | 161 |
| Bush-Honeysuckle | 205 | Carrión Flower | 520 | Chondrilla | 279 |
| Butter-and-eggs | 326 | Carrot | 191 | Chondrosium | 621 |
| Buttercup | 40 | Carya | 448 | Chrysanthemum | 265 |
| Butterfly-Pea | 141 | CARYOPHYLLACEÆ | 87 | Chrysastrum | 239 |
| Butterfly-weed | 397 | Caryophyllata | 152 | Chrysostrum | 248 |
| Butternut | 447 | Cashew Family | 111 | Chrysopsis | 246 |
| Butter-weed | 271 | Cassandra | 294 | Chrysosplenium | 170 |
| Butterwort | 320 | Cassena | 306 | CHICORACEÆ | 221 |
| Button-bush | 211 | Cassia | 144 | Cichorium | 275 |
| Button Snakeroot | 191, 223 | Cassiope | 295 | Cichory | 275 |
| Button-weed | 210 | Castanea | 454 | Cicuta | 196 |
| Buttonwood | 447 | Castilleja | 336 | Cimicifuga | 48 |
| Cabomba | 55 | Catalpa | 321 | CINCHONEÆ | 208 |
| CABOMBEÆ | 54 | Catbrier | 518 | Cinna | 612 |
| Cacalia | 270 | Catchfly | 89 | Cinnamon-Fern | 671 |
| CACTACEÆ | 184 | Catgut | 131 | Cinque-foil | 153 |
| Cactus | 185 | Cat-mint | 353 | Circæa | 176 |
| Cactus Family | 184 | Catnip | 353 | Cirsium | 273 |
| Cænotus | 236 | Cat-tail Family | 480 | Cissus | 113 |
| CÆSALPINIÆ | 125 | Cat-tail Flag | 480 | CISTACEÆ | 80 |
| Cakile | 75 | Cat's-tail Grass | 608 | Citrullus | 186 |
| CAKILINEÆ | 64 | Caulinia | 483 | Cladium | 570 |
| CALAMAGROSTIDEÆ | 603 | Caulophyllum | 53 | Cladrastis | 143 |
| Calamagrostis | 614 | Ceanothus | 115 | Claytonia | 98 |
| Calamintha | 349 | Cedar | 474 | Clearweed | 445 |
| Calaminth | 349 | Cedronella | 354 | Cleavers | 208 |
| Calamovilfa | 616 | Celandine | 59 | CLEMATIDEÆ | 34 |
| Calamus | 478 | Celandine Poppy | 59 | Clematis | 35 |
| Calico-bush | 298 | CELASTRACEÆ | 115 | Clethra | 296 |
| Calla | 476 | Celastrus | 116 | Cliff-brake | 659, 660 |
| Calliastrum | 229 | Celtis | 443 | Climbing Fern | 570 |
| Callicarpa | 341 | Cenchrus | 653 | Climbing Fumitory | 60 |
| Callirrhœ | 100 | Centaurea | 272 | Climbing Hemp-weed | 226 |
| CALLITRICHACEÆ | 427 | Centaurella | 389 | Clinopodium | 349 |
| Callitriche | 428 | Centaur | 385 | Clintonia | 529 |
| Calluna | 297 | Centrosema | 141 | Clitoria | 141 |
| Calomelissa | 349 | Centunculus | 317 | Clothur | 251 |
| Calopogon | 508 | Century-plant | 513 | Cloud-berry | 157 |
| Caltha | 44 | Cephalanthus | 211 | Clover | 126 |
| CALYCANTHACEÆ | 162 | Cephaloxys | 539 | Club-Moss | 673 |
| Calycanthus | 162 | Cerastium | 93 | Club-Moss Family | 672 |
| Calycanthus Family | 162 | Cerasus | 149 | Club-Rush | 560 |
| Calycocarpum | 52 | Ceratophyllaceæ | 427 | Cnicus | 272 |
| Calypso | 508 | Ceratostochæna | 569 | Cnidoseolus | 435 |
| Calystegia | 375 | Cercis | 143 | Cocculus | 51 |
| Camassia | 533 | Chærophyllum | 197 | Cockle | 90 |
| Camelina | 73 | Chætocyperus | 560 | Cocklebur | 251 |
| CAMELINEÆ | 63 | Chaffseed | 336 | Coffee | 208 |
| Camellia | 103 | Chaffweed | 274 | Cohosh | 53 |
| CAMELLIACEÆ | 103 | Chain-Fern | 660 | Colic-root | 515 |
| Campanula | 285 | Chamælium | 527 | Collinsia | 327 |
| CAMPANULACEÆ | 285 | Chamomile | 265 | Collinsonia | 350 |
| Campanula Family | 285 | Charlock | 70 | Colpodium | 610 |
| Campion | 89 | Cheat-Grass | 634 | Coltsfoot | 227 |
| Camptosorus | 663 | Checkerberry | 293 | Columbine | 45 |
| Canary-Grass | 643 | Cheilanthes | 659 | Columbo | 386 |
| | | Chelidonium | 59 | Comandra | 425 |

| | | | | | |
|--------------------|----------|----------------------|----------|-----------------------|----------|
| Comaropsis | 153 | Ctenium | 620 | Dichronema | 567 |
| Comarum | 155 | Cuckoo-flower | 66 | Dicksonia | 669 |
| Comfrey | 361 | Cucumber | 186 | DICOTYLEDONOUS PL. | 33 |
| Commelyna | 546 | Cucumber-tree | 49 | Dictamnus | 110 |
| COMMELYNACEÆ | 546 | Cucumis | 186 | Dielytra | 61 |
| Compass Plant | 249 | Cucurbita | 186 | Diervilla | 205 |
| COMPOSITÆ | 215 | CUCURBITACEÆ | 186 | Digitaria | 645 |
| Composite Family | 215 | Cudweed | 268 | Digraphis | 643 |
| Comptonia | 458 | Culver's Physic | 332 | Dilepyrum | 614 |
| Cone-flower | 254 | Culver's-root | 332 | Diodia | 210 |
| CONIFERÆ | 468 | Cunila | 346 | Dionæa | 83 |
| Conioselinum | 193 | Cuphea | 184 | Dioscorea | 518 |
| Conium | 197 | Cup-plant | 249 | DIOSCOREACEÆ | 518 |
| Conobea | 329 | CUPRESSINÆ | 468 | Diospyros | 308 |
| Conoclinium | 227 | Cupressus | 473 | Diphylleia | 53 |
| Conopholis | 323 | Cupseed | 52 | Diplachne | 623 |
| Conostylis | 514 | CUPULIFERÆ | 449 | Diplocea | 624 |
| Convallaria | 530 | Currant | 164 | Diplopappus | 238 |
| CONVOLVULACEÆ | 374 | Cuscuta | 377 | DIPSACEÆ | 215 |
| CONVOLVULÆ | 374 | CUSCUTINÆ | 374 | Dipsacus | 215 |
| Convolvulus | 375 | Custard-Apple Family | 50 | Dipteracanthus | 339 |
| Convolvulus Family | 374 | Cyanococcus | 291 | Direa | 424 |
| Coprosmanthus | 520 | Cycloloma | 406 | Discopleura | 195 |
| Coptis | 45 | CYNARÆ | 221 | Ditch-Grass | 483 |
| Coral-berry | 203 | Cynodon | 622 | Ditch Stone-crop | 171 |
| Corallorhiza | 509 | Cynoglossum | 365 | Dittany | 346 |
| Coral-root | 509 | Cynthia | 276 | Dock | 419 |
| Cord Grass | 619 | CYPERACEÆ | 550 | Dockmackie | 207 |
| Corema | 440 | CYPERÆ | 551 | Dodder | 377 |
| Coreopsis | 258 | Cyperus | 552 | Dodecatheon | 314 |
| Corispermum | 409 | Cypress | 473 | Dogbane | 393 |
| CORNACEÆ | 199 | Cypress Family | 468 | Dogbane Family | 392 |
| Corn-cockle | 90 | Cypress-Vine | 374 | Dog's-tail Grass | 623 |
| Cornel | 200 | CYPRIPEDIÆ | 498 | Dog's-tooth Violet | 532 |
| Corn Salad | 214 | Cypripedium | 511 | Dogwood | 200 |
| Cornus | 200 | Cystopteris | 667 | Dogwood Family | 199 |
| Corpse-Plant | 304 | Dactylis | 625 | Door-weed | 417 |
| Corydalis | 61 | Dactyloctenium | 622 | Double-bristled Aster | 238 |
| Corylus | 456 | Daffodil | 512 | Draba | 71 |
| Cosmanthus | 369 | Daisy | 239 | Dracocephalum | 353 |
| Cotton-Grass | 565 | Dalea | 129 | Dragon-Arum | 475 |
| Cotton-Rose | 269 | Dalibarda | 153, 156 | Dragon-head | 353 |
| Cotton-Thistle | 274 | Dandelion | 280 | Dragon-root | 476 |
| Cotton-wood | 467 | Dangleberry | 289 | Drop-seed Grass | 610, 613 |
| Couch-Grass | 637 | Danthonia | 639 | Drosera | 82 |
| Cowbane | 192 | Daphne | 424 | ROSERACEÆ | 82 |
| Cowberry | 290 | Darnel | 637 | DRYADEÆ | 147 |
| Cow-Herb | 88 | Dasytoma | 335 | Dryas | 151 |
| Cow-Parsnip | 191 | Date Plum | 308 | Dryopteris | 664 |
| Cowslip | 313 | Datura | 383 | Dubruellia | 445 |
| Cowslips | 44 | Daucus | 191 | Duck's-meat | 478 |
| Cow-wheat | 338 | DAVALLIÆ | 657 | Duckweed | 478 |
| Crab-Apple | 161 | Day-flower | 546 | Duckweed Family | 478 |
| Crab-Grass | 622, 646 | Day-Lily | 535 | Dulichium | 556 |
| Cranberry | 289 | Dead-Nettle | 359 | Dupontia | 624 |
| Cranberry-tree | 207 | Decodon | 183 | Dutchman's Breeches | 61 |
| Crane-fly Orchis | 508 | Deerberry | 290 | Dutchman's Pipe | 404 |
| Cranesbill | 107 | Deer Grass | 181 | Dwarf Dandelion | 275 |
| Crantzia | 190 | Delphinium | 45 | Dyer's Green-weed | 126 |
| CRASSULACEÆ | 171 | Dennstedtia | 669 | Dyer's Rocket | 76 |
| Cratægus | 159 | Dentaria | 65 | Dyer's Weed or Weld | 76 |
| Creeping Snowberry | 292 | Deschampsia | 641 | Dysodia | 262 |
| Crocus | 516 | Desmanthus | 145 | Eatonia | 625 |
| Crossopetalum | 387 | Dcsmodium | 134 | EBENACEÆ | 307 |
| Crotalaria | 126 | Dewberry | 157 | Ebony Family | 307 |
| Croton | 437 | Devil's-bit | 527 | Echinacea | 253 |
| Crotonopsis | 439 | Devil-wood | 401 | Echinochloa | 649 |
| Crowberry | 440 | Deyeuxia | 615 | Echinocystis | 186 |
| Crowberry Family | 440 | Dianthera | 338 | Echinodorus | 492 |
| Crowfoot | 40 | Dianthus | 88 | Echinosperrnum | 365 |
| Crowfoot Family | 34 | Diapensia | 373 | Echium | 361 |
| Crownbeard | 262 | DIAPENSIEÆ | 370 | Eclipta | 252 |
| CRUCIFERÆ | 62 | Diarrhena | 624 | Eel-grass | 483, 496 |
| Crypsis | 609 | Dicentra | 61 | Egg-plant | 380 |
| CRYPTOGAMOUS PL. | 653 | Dichondra | 377 | Egyptian Grass | 622 |
| Cryptogramme | 660 | DICHONDREÆ | 374 | ELÆAGNACEÆ | 424 |
| Cryptotænia | 197 | | | | |

| | | | | | |
|-------------------------|----------|-----------------------|----------|-------------------|----------|
| Elæagnus | 425 | Fagus | 455 | Foxtail Grass | 608, 649 |
| ELATINACEÆ | 86 | Fall Dandelion | 276 | Fragaria | 155 |
| Elatine | 86 | False Acacia | 131 | Frangula | 115 |
| Elder | 205 | False Asphodel | 527 | Frasera | 386 |
| Elecampane | 246 | False Beech-drops | 305 | FRAXINEÆ | 400 |
| Eleocharis | 557 | False Bugbane | 39 | Fraxinella | 110 |
| Eleoegenus | 558 | False Dandelion | 280 | Fraxinus | 401 |
| Elephantopus | 222 | False Dragon-head | 354 | French Marigold | 262 |
| Elephant's-foot | 222 | False Flax | 73 | French Mulberry | 341 |
| Ellisia | 368 | False Foxglove | 335 | Fringed Polygala | 122 |
| Elm | 442 | False Goatsbeard | 167 | Fringe-tree | 401 |
| Elm Family | 441 | False Golden-rod | 239 | Frœlichia | 414 |
| Elodea | 86, 495 | False Gromwell | 362 | Frog's-bit Family | 494 |
| Elodes | 86 | False Hellebore | 525 | Frost-weed | 80 |
| Eleusine | 622 | False Honeysuckle | 299 | Fuirena | 556 |
| Elymus | 639 | False Indigo | 130, 142 | Fumaria | 62 |
| EMPETRACEÆ | 440 | False Lettuce | 281 | FUMARIACEÆ | 60 |
| Empetrum | 440 | False Loosestrife | 180 | Fumitory | 62 |
| ENDOGENOUS PLANTS | 475 | False Mallow | 100 | Fumitory Family | 60 |
| Enchanter's Nightshade | 176 | False Mermaid | 108 | | |
| Enemion | 44 | False Mistletoe | 426 | GALACINEÆ | 305 |
| Engelmannia | 438 | False Mitre-wort | 170 | Galactia | 142 |
| Enslenia | 398 | False Nettle | 445 | Galanthus | 512 |
| Epigæa | 293 | False Pimpernel | 330 | Galax | 305 |
| Epilobium | 177 | False Red-top | 629 | Galax Family | 305 |
| Epipactis | 506 | False Solomon's Seal | 530 | GALEGEÆ | 124 |
| Epiphegus | 322 | False Spikenard | 530 | Galeopsis | 357 |
| EQUISETACEÆ | 653 | False Water-Dropwort | 192 | Galingale | 552 |
| Equisetum | 653 | Farkle-berry | 290 | Galinsoga | 264 |
| Eragrostis | 631 | Featherfoil | 317 | Gall-of-the-Earth | 278 |
| Erechtithes | 269 | Feather Geranium | 407 | Galium | 208 |
| Erianthus | 651 | Feather Grass | 617 | Gama-Grass | 650 |
| ERICACEÆ | 286 | Fedia | 214 | Garden Nasturtium | 105 |
| ERICÆ | 287 | Ferns | 655 | Garget | 405 |
| ERICINEÆ | 287 | Fescue-Grass | 633 | Garlic | 533 |
| Erigenia | 198 | Festuca | 633 | Gaultheria | 293 |
| Erigeridium | 237 | FESTUCINEÆ | 604 | Gaura | 176 |
| Erigeron | 236 | Fetid Horehound | 359 | Gaylussacia | 288 |
| ERIOGONEÆ | 414 | Fetid Marigold | 262 | Geisleria | 438 |
| Eriocaulon | 549 | Fever-bush | 423 | Gelsemium | 391 |
| ERIOCAULONACEÆ | 549 | Feverfew | 266 | Genista | 126 |
| Eriophorum | 565 | Fever-wort | 205 | GENISTEÆ | 123 |
| Erodium | 108 | Fig Family | 441 | Gentian | 387 |
| Erophila | 72 | Figwort | 326 | Gentiana | 387 |
| Ervum | 139 | Figwort Family | 324 | GENTIANACEÆ | 384 |
| Eryngium | 190 | Filago | 269 | GENTIANEÆ | 384 |
| Eryngo | 190 | Filbert | 456 | Gentian Family | 384 |
| Erythraea | 385 | FILICES | 655 | GERANIACEÆ | 105 |
| Erythronium | 532 | Fimbristylis | 566 | GERANIEÆ | 105 |
| Erysinium | 69 | Finger-Grass | 646 | Geranium | 107 |
| ESCALLONIEÆ | 163 | Florin Grass | 612 | Geranium Family | 105 |
| Eubotrys | 294 | Fir | 471 | Gerardia | 334 |
| Eulophus | 198 | Fire-Pink | 89 | GERARDIEÆ | 325 |
| Euonymus | 116 | Fireweed | 269 | Germander | 343 |
| EUPATORIACEÆ | 217 | Five-finger | 153 | Geum | 151 |
| EUPATORIEÆ | 218 | Flax | 104 | Giant Hyssop | 352 |
| Eupatorium | 224 | Flax Family | 104 | Gill | 353 |
| Euphorbia | 430 | Fleabane | 236 | Gillenia | 150 |
| EUPHORBIACEÆ | 430 | Floating Heart | 390 | Ginseng | 198, 199 |
| Euphrasia | 356 | Flœrkea | 108 | Ginseng Family | 198 |
| EUPHRASIEÆ | 325 | Flower-de-Luce | 516 | Girasole | 258 |
| Euphrosyne | 250 | Flowering Fern | 670 | Glade Mallow | 100 |
| Eurhynchospora | 568 | FLOWERING PLANTS | 33 | Gladious | 516 |
| Euthamia | 245 | Flowering Wintergreen | 122 | Glasswort | 409 |
| Eutoca | 369 | FLOWERLESS PLANTS | 653 | Glaucium | 60 |
| Eutriana | 621 | Fly-catch Grass | 607 | Glaux | 316 |
| Eu vaccinium | 290 | Fly-poison | 526 | Glechoma | 353 |
| Euxolus | 412 | Fool's-Parsley | 193 | Gleditschia | 145 |
| Evening Primrose | 178 | Forestiera | 402 | Globe-flower | 44 |
| Evening Primrose Family | 176 | FORESTIEREÆ | 400 | Glyceria | 626 |
| Everlasting | 265, 269 | Forget-me-not | 364 | Glycyrrhiza | 133 |
| Everlasting Pea | 139 | Forked Chickweed | 96 | GNAPHALINEÆ | 220 |
| Evolvulus | 376 | Forsteronia | 393 | Gnaphalium | 268 |
| EXOGENOUS PLANTS | 88 | Fothergilla | 173 | Goat's-Beard | 150 |
| Eyebricht | 336 | Four-o'clock Family | 404 | Goat's Rue | 131 |
| | | Fowl-meadow Grass | 627, 629 | Golden Aster | 246 |
| Fagopyrum | 419 | Foxglove | 335 | Golden-club | 477 |

| | | | | | |
|---------------------|---------------|---------------------------|----------|----------------------|----------|
| Golden-rod | 239 | Hedge Bindweed | 375 | Horse-Mint | 351 |
| Golden Saxifrage | 170 | Hedgehog-Grass | 650 | Horse-Nettle | 381 |
| Gold-thread | 45 | Hedge-Hyssop | 329 | Horseradish | 65 |
| GONOLOBÆ | 394 | Hedge-Mustard | 70 | Horse-Sugar | 310 |
| Gonolobus | 399 | Hedge-Nettle | 358 | Horsetail | 653 |
| Good-King-Henry | 408 | HEDYSARÆ | 124 | Horsetail Family | 653 |
| Goodyera | 503 | Hedysarum | 134 | Hottonia | 317 |
| Gooseberry | 164 | HELENIÆ | 220 | HOTTONIÆ | 313 |
| Goosefoot | 406 | Helenium | 263 | Hound's-Tongue | 365 |
| Goosefoot Family | 405 | Heleochloa | 628 | Houstonia | 212 |
| Goose-Grass | 208, 417 | HELIANTHÆ | 219 | Huckleberry | 288 |
| Gordonia | 104 | Helianthemum | 80 | Hudsonia | 81 |
| Gourd | 186 | Helianthus | 255 | Humulus | 446 |
| Gourd Family | 186 | HELIOPHYTUM | 366 | Huntsman's Cup | 58 |
| GRAMINEÆ | 602 | Heliospis | 253 | Hydrangea | 165 |
| Grape | 112 | HELIOTROPEÆ | 361 | HYDRANGIÆ | 163 |
| Grape-Hyacinth | 534 | Heliotropium (Heliotrope) | 366 | Hydrastis | 47 |
| Graphephorum | 624 | Hellebore | 45 | HYDROCHARIDACEÆ | 494 |
| Grass of Parnassus | 166 | Helleborus | 45 | Hydrocharis | 495 |
| Grass of the Andes | 642 | Helonias | 526 | Hydrocotyle | 189 |
| Grass-wrack | 483 | Helosciadium | 196 | Hydrolea | 370 |
| Gratiola | 329 | Hemerocallis | 535 | HYDROLEÆ | 367 |
| GRATIOLÆ | 324 | Hemianthus | 330 | Hydropeltis | 55 |
| Great Laurel | 300 | Hemicarpha | 556 | HYDROPHYLLACEÆ | 367 |
| Greek Valerian | 371 | Hemlock | 196, 197 | HYDROPHYLLÆ | 367 |
| Greenbrier | 518 | Hemlock-Parsley | 193 | Hydrophyllum | 367 |
| Green Dragon | 426 | Hemlock Spruce | 471 | HYDROPTERIDES | 676 |
| Green Milkweed | 398 | Hemp | 446 | Hymenopappus | 262 |
| Green Violet | 76 | Hemp Family | 442 | Hyoscyamus | 383 |
| Gromwell | 362 | Hemp-Nettle | 357 | HYPERICACEÆ | 83 |
| Grossularia | 164 | Henbane | 383 | Hypericum | 84 |
| GROSSULARIÆ | 163 | Hepatica | 38 | Hypobrychia | 182 |
| Ground Cherry | 381 | Heracleum | 191 | Hypopeltis | 668 |
| Ground Hemlock | 474 | Herb-Robert | 107 | Hypopitys | 304 |
| Ground Ivy | 353 | Hercules's Club | 199 | Hypoxis | 5-3 |
| Ground Laurel | 293 | Herd's Grass | 608, 512 | Hyssop | 346 |
| Ground-nut | 140, 199 | Herpestis | 329 | Hyssopus | 346 |
| Ground Pine | 674 | Hesperis | 68 | Ictodes | 477 |
| Ground Pink | 373 | Heteranthera | 545 | Ilex | 305 |
| Ground Plum | 132 | Heuchera | 169 | ILLECEBREÆ | 87 |
| Groundsel | 270 | HIBISCEÆ | 99 | Ilysanthes | 330 |
| Groundsel-Tree | 247 | Hibiscus | 102 | Impatiens | 108 |
| Guinea-Corn | 652 | Hickory | 448 | Inkberry | 307 |
| Gymnanthena | 499 | Hieracium | 277 | Indian Bean | 321 |
| Gymnocladus | 144 | High-water-shrub | 250 | Indian Chickweed | 97 |
| Gymnopogon | 621 | Hierochloa | 642 | Indian Cucumber-root | 523 |
| GYMNOSPERMEÆ | 468 | HIPPOCASTANÆ | 117 | Indian Currant | 203 |
| Gymnostichum | 639 | Hippuris | 175 | Indian Fig | 185 |
| Gynamblosis | 438 | Hoary Pea | 131 | Indian-Grass | 652 |
| Gyromia | 523 | Hobble-bush | 207 | Indian-Heliotrope | 366 |
| Habenaria | 498 | Hog Pea-nut | 141 | Indian Hemp | 393 |
| Hackberry | 443 | Hog-weed | 251 | Indian Mallow | 101 |
| Hackmatack | 472 | Holcus | 642 | Indian Millet | 652 |
| HÆMODOURACEÆ | 514 | Holly | 305 | Indian Physic | 150 |
| Hair-Grass | 611, 614, 641 | Holly Family | 305 | Indian Pipe | 304 |
| Halenia | 386 | Holosteum | 93 | Indian-pipe Family | 288 |
| Halesia | 310 | Holy Grass | 642 | Indian Plantain | 270 |
| HALORAGEÆ | 174 | Honey-Locust | 145 | Indian Poke | 625 |
| HAMAMELACEÆ | 173 | Honeysuckle | 203 | Indian Rice | 608 |
| HAMAMELÆ | 173 | Honeysuckle Family | 202 | Indian Tobacco | 283 |
| Hamamelis | 173 | Honewort | 197 | Indian Turnip | 475 |
| Hamiltonia | 426 | Honkenya | 92 | Inula | 246 |
| Harbinger-of-Spring | 198 | Hop | 446 | INGLEÆ | 218 |
| Hardhack | 149 | Hopea | 310 | Iodanthus | 68 |
| Harebell | 285 | Hop-Hornbeam | 456 | Ipomœa | 375 |
| Hart's-tongue | 662 | Hop-tree | 110 | Iresine | 413 |
| Hawkbit | 276 | Hordeum | 638 | IRIDACEÆ | 515 |
| Hawkweed | 277 | HORDEINÆ | 605 | Iris | 516 |
| Hawthorn | 159 | Horehound | 357 | Iris Family | 515 |
| Hazel-nut | 456 | Hornbeam | 457 | Iron-weed | 222 |
| Heal-all | 355 | Horned Pondweed | 483 | Iron-wood | 456, 457 |
| Heart's Ease | 79 | Horned Rush | 570 | Isanthus | 344 |
| Heather | 297 | Horn Poppy | 60 | Isoetes | 675 |
| Heath Family | 286 | Hornwort | 427 | Isoplepis | 560 |
| Hedeoma | 350 | Horse-Balm | 350 | Isopyrum | 44 |
| Hedera | 199 | Horsechestnut | 117 | Itea | 165 |
| | | Horse-Gentian | 205 | | |

| | | | | | |
|----------------------|-----|----------------------|----------|-----------------------|----------|
| Iva | 250 | Leatherwood | 424 | Logania Family | 391 |
| Ivy | 199 | Leavvworthia | 65 | Loisclcuria | 301 |
| Ixia | 617 | Lechca | 81 | Lolium | 637 |
| Jacob's Ladder | 371 | Lecontia | 476 | LOMENTACEÆ | 64 |
| Jagged Chickweed | 93 | Ledum | 300 | Long Moss | 515 |
| Jamestown-weed | 383 | Leek | 534 | Lonicera | 203 |
| Jasminum | 400 | Leersia | 607 | LONICEREÆ | 202 |
| Jatropha | 435 | LEGUMINOSEÆ | 123 | Loosestrife | 183, 315 |
| Jeffersonia | 53 | Leiophyllum | 301 | Loosestrife Family | 182 |
| Jerusalem Artichoke | 258 | Lemna | 478 | Lophanthus | 352 |
| Jerusalem Oak | 407 | LEMNACEÆ | 478 | Lophiola | 514 |
| Jerusalem Sage | 360 | LENTIBULACEÆ | 317 | Lopseed | 341 |
| Jessamine | 400 | Leontodon | 276 | LORANTHACEÆ | 426 |
| Jewel-weed | 108 | Leonurus | 359 | Lorinseria | 661 |
| Joe-Pye Weed | 225 | Lepachys | 255 | Lousewort | 337 |
| Jointed Charlock | 75 | LEPIDINEÆ | 63 | Lovage | 194 |
| Joint-Grass | 645 | Lepidium | 74 | Lucerne | 123 |
| Joint-weed | 417 | Lepidanche | 379 | Ludwigia | 180 |
| Jonquil | 512 | Lepigonum | 95 | Lungwort | 364 |
| Judas-tree | 143 | Leptandra | 332 | Lupine | 126 |
| Juglans | 447 | Leptanthus | 545 | Lupinus | 126 |
| JUGLANDACEÆ | 447 | Leptochloa | 623 | Luzula | 536 |
| JUNCACEÆ | 536 | Leptopoda | 263 | Lychnis | 90 |
| JUNCAGINEÆ | 490 | Lepurus | 637 | Lycium | 382 |
| Juncus | 537 | Lespedeza | 137 | Lycopersicum | 380 |
| June-berry | 162 | Lettuce | 280 | LYCOPODIACEÆ | 672 |
| Juniper | 474 | Leucanthemum | 265 | Lycopodium | 673 |
| Juniperus | 473 | Leucojum | 512 | Lycopsis | 361 |
| Jussiaea | 179 | Leucothoë | 294 | Lycopus | 345 |
| | | Lever-wood | 456 | Lygodesmia | 279 |
| | | Liatrix | 223 | Lygodium | 670 |
| Kalmia | 297 | LIGULIFLORÆ | 221, 275 | Lyonia | 296 |
| Kentucky Blue-Grass | 630 | Ligusticum | 194 | Lysimachia | 315 |
| Kentucky Coffee-tree | 144 | Ligustrum | 400 | LYTHRACEÆ | 182 |
| Kidney Bean | 140 | Lilac | 400 | Lythrum | 183 |
| Kinnikinnik | 200 | LILIACEÆ | 520 | | |
| Knapweed | 272 | LILIEÆ | 522 | Madder Family | 208 |
| Knawel | 96 | Lilium (Lily) | 532 | Macrotys | 48 |
| Knotgrass | 417 | Lily of the Valley | 530 | Magnolia | 49 |
| Knotweed | 414 | LIMNANTHEÆ | 106 | MAGNOLIACEÆ | 48 |
| Koeleria | 625 | Limnanthes Family | 106 | Magnolia Family | 48 |
| Kosteletzkya | 102 | Limnanthemum | 390 | Maianthemum | 530 |
| Krigia | 275 | Limnetis | 619 | Maidenhair | 658 |
| Kuhnia | 224 | Limnobiium | 495 | MALAXIDEÆ | 497 |
| Kyllingia | 555 | Limnochloa | 557 | Mallow | 99 |
| | | Limosella | 331 | Mallow Family | 98 |
| LABIATÆ | 341 | LINACEÆ | 104 | Malus | 161 |
| Labrador Tea | 300 | Linaria | 326 | Malva | 99 |
| Lachnanthes | 514 | Linden | 103 | MALVACEÆ | 98 |
| Lachnocaulon | 550 | Linden Family | 103 | Malvastrum | 100 |
| Lactuca | 280 | Lindera | 423 | MALVÆ | 99 |
| Lady's Mantle | 151 | Lindernia | 330 | Mandrake | 54 |
| Lady's Slipper | 511 | Linnaea | 202 | Manna-Grass | 626 |
| Lady's Thumb | 416 | Linum | 104 | Man-of-the-Earth | 375 |
| Ladies-Tresses | 504 | Lion's-foot | 278 | Maple | 118 |
| Lagenaria | 186 | Liparis | 509 | Maple Family | 117 |
| Lambkill | 298 | Lip-Fern | 659 | Mare's Tail | 175 |
| Lamb-Lettuce | 214 | Lipocarpha | 556 | Mariscus | 455 |
| Lamium | 359 | Lippia | 340 | Marrubium | 357 |
| Lampsona | 275 | Liquidambar | 174 | Marshallia | 264 |
| Lapitheia | 385 | Liquorice | 133 | Marsh Elder | 250 |
| Laportea | 445 | Liriodendron | 50 | Marsh-Fleabane | 247 |
| Lappa | 275 | Listera | 506 | Marsh Grass | 619 |
| Larch | 472 | Lithospermum | 362 | Marsh-Mallow | 99 |
| Larix | 472 | Live-for-ever | 172 | Marsh-Marigold | 44 |
| Larkspur | 45 | Liver-leaf | 38 | Marsh-Rosemay | 312 |
| Lastrea | 664 | Lizard's-tail | 427 | Marsh St. John's-wort | 86 |
| Lathyrus | 139 | Lizard's-tail Family | 427 | Marsilia | 677 |
| LAURACEÆ | 422 | LOASACEÆ | 184 | Marsiliaceæ | 676 |
| Laurel | 297 | Loasa Family | 184 | Martynia | 321 |
| Laurel Family | 422 | Lobadium | 112 | Maruta | 264 |
| Laurestinus | 206 | Lobelia | 282 | Marvel of Peru | 404 |
| Lead-Plant | 130 | LOBELIACEÆ | 282 | Masterwort | 191 |
| Leadwort Family | 312 | Lobelia Family | 282 | Matricaria | 266 |
| Leaf-Cup | 247 | Lobolly Bay | 104 | Matrimony-Vine | 382 |
| Leather-Flower | 86 | Locust-tree | 130 | Mayaca | 547 |
| Leather-leaf | 294 | LOGANIACEÆ | 391 | May-Apple | 54 |

| | | | | | |
|---------------------|-----|------------------------|-----|-------------------|-----|
| Mayflower | 293 | Monocera | 620 | Nettle-tree | 443 |
| Maypops | 186 | MONOCOTYLEDONOUS | | New Jersey Tea | 115 |
| May-weed | 264 | PLANTS | 475 | Nicandra | 382 |
| Meadow-Beauty | 181 | Monopetalous Exogenous | | Nicotiana | 383 |
| Meadow-Grass | 628 | Plants | 202 | Niellia | 149 |
| Meadow-Parsnip | 194 | Monotropa | 304 | Nigella | 48 |
| Meadow-Rue | 38 | MONOTROPEÆ | 288 | Nightshade | 380 |
| Meadow Soft-Grass | 642 | Montelia | 413 | Nightshade Family | 380 |
| Meadow-Sweet | 149 | Moonseed | 51 | Nimble Will | 614 |
| Medeola | 523 | Moonseed Family | 51 | Nine-Bark | 149 |
| Medicago (Medick) | 128 | Moonwort | 671 | Nipple-wort | 275 |
| MELAMPODINEÆ | 219 | Moosewood | 424 | Nonesuch | 128 |
| Melampyrum | 338 | Moose-wood | 119 | Nondo | 194 |
| MELANTHIEÆ | 521 | MOREÆ | 441 | Nothoscordum | 534 |
| Melanthium | 524 | Morning-Glory | 375 | Nuphar | 56 |
| MELASTOMACEÆ | 181 | Morocarpus | 408 | Nut-Grass | 553 |
| Melastoma Family | 181 | Morus | 444 | Nut-Rush | 570 |
| Melica | 626 | Moss Campion | 90 | NYCTAGINACEÆ | 404 |
| Melic Grass | 626 | Moss Pink | 373 | Nymphaea | 56 |
| Melilotus (Melilot) | 128 | Motherwort | 359 | NYMPIHEÆCEÆ | 54 |
| Melissa | 350 | Mountain Ash | 161 | Nyssa | 201 |
| Melothria | 187 | Mountain Holly | 307 | | |
| MENISPERMACEÆ | 51 | Mountain Mint | 346 | Oak | 450 |
| Menispermum | 51 | Mountain Rice | 616 | Oakesia | 440 |
| Mentha | 344 | Mountain Sorrel | 419 | Oak Family | 449 |
| Mentzelia | 184 | Mouse-ear | 364 | Oat | 640 |
| Menyanthes | 390 | Mouse-ear Chickweed | 93 | Oat-Grass | 642 |
| MENYANTHEÆ | 384 | Mouse-ear Cress | 70 | Obeliscaria | 255 |
| Menziesia | 298 | Mouse-tail | 44 | Obolaria | 389 |
| Mermaid-weed | 175 | Mud-Plantain | 545 | Enothera | 178 |
| Mertensia | 363 | Mudwort | 331 | Oil-nut | 426 |
| Mexican Poppy | 59 | Mugwort | 267 | Old Witch Grass | 647 |
| Mexican Tea | 408 | Muhlenbergia | 613 | Oldenlandia | 211 |
| Mezereum Family | 424 | Mulberry | 444 | Olea | 401 |
| Micranthemum | 330 | Mulgedium | 281 | OLEACEÆ | 400 |
| Microstylis | 509 | Mullein | 325 | Oleander | 892 |
| Mignonette | 76 | Mullein-Foxglove | 334 | Oleaster Family | 424 |
| Mignonette Family | 76 | Mullugo | 97 | OLEINEÆ | 400 |
| Mikania | 226 | Muscari | 534 | Olive | 401 |
| Milfoil | 265 | Muskit Grass | 620 | Olive Family | 400 |
| Milium | 643 | Muskmelon | 186 | ONAGRACEÆ | 176 |
| Milk-Pea | 142 | Musk-Plant | 329 | Oncostylis | 567 |
| Milkweed | 394 | Musquash-Root | 196 | Onion | 533 |
| Milkweed Family | 394 | Mustard | 62 | Onoclea | 668 |
| Milkwort | 120 | Mustard Family | 62 | Onopordon | 274 |
| Milkwort Family | 120 | Myosotis | 364 | Onosmodium | 362 |
| Milk-Vetch | 132 | Myosurus | 44 | OPHIOGLOSSACEÆ | 657 |
| Millet-Grass | 643 | Myrica | 457 | Ophioglossum | 672 |
| Mimosa Family | 125 | MYRACACEÆ | 457 | OPHYRDEÆ | 497 |
| MIMOSEÆ | 125 | Myriophyllum | 174 | Oplotecha | 414 |
| Mimulus | 328 | | | Opuntia | 185 |
| Mint | 344 | Nabalus | 278 | Orache | 409 |
| Mint Family | 341 | NAIADACEÆ | 482 | Orange Family | 110 |
| Mirabilis | 404 | Naias (Naiad) | 482 | Orange-Grass | 86 |
| Mist-flower | 227 | Naked-beard Grass | 621 | Orange-root | 47 |
| Mistletoe | 426 | Naked Broom-rape | 323 | Orchard Grass | 625 |
| Mistletoe Family | 426 | Napæa | 100 | ORCHIDACEÆ | 497 |
| Mitchella | 211 | Narcissus | 512 | Orchis | 498 |
| Mitella | 169 | Nardosmia | 227 | Orchis Family | 497 |
| Mithridate Mustard | 74 | Narthecium | 535 | Origanum | 348 |
| Mitreola | 392 | Nasturtium | 64 | Oritrophium | 235 |
| Mitrewort | 392 | Naumburgia | 315 | Ornithogalum | 533 |
| Mitre-wort (False) | 170 | Neckweed | 333 | OROBANCHACEÆ | 322 |
| Mocasson Flower | 511 | Negundo | 119 | Orobancha | 323 |
| Mock Bishop-weed | 195 | Nelumbium | 55 | Orontium | 477 |
| Mocker-nut | 449 | Nelumbo | 55 | Orpine | 172 |
| Mock-Orange | 186 | Nelumbo Family | 55 | Orpine Family | 171 |
| Modiola | 101 | NELUMBONEÆ | 55 | Orthomeris | 235 |
| Mœhringia | 91 | Nemopanthus | 307 | ORYZEÆ | 603 |
| Mœnchia | 94 | Nemophila | 368 | Oryzopsis | 616 |
| MOLLUGINEÆ | 88 | NEOTIEÆ | 497 | Osier | 461 |
| Monarda | 351 | Nepeta | 353 | Osmorrhiza | 197 |
| MONARDEÆ | 342 | NEPETEÆ | 342 | OSMUNDACEÆ | 657 |
| Moneses | 303 | Nephrodium | 664 | Osmunda | 670 |
| Moneywort | 316 | Nesæa | 183 | Ostrich-Fern | 667 |
| Monkey-flower | 328 | Nettle | 444 | Ostrya | 456 |
| Monkshood | 46 | Nettle Family | 440 | Oswego Tea | 351 |

| | | | | | |
|-----------------------|----------|----------------------|-----|----------------------------|----------|
| Otophylla | 335 | Persimmon | 308 | Plumeless Thistle | 274 |
| OxALIDÆ | 106 | Perularia | 499 | Pneumonanthe | 387 |
| Oxalis | 109 | Peruvian Bark | 208 | Poa | 628 |
| Ox-eye | 253 | Petaloma | 433 | POACEÆ | 603 |
| Ox-eye Daisy | 265 | Petalostemon | 130 | PODALYRIÆ | 125 |
| Oxybaphus | 404 | Phacelia | 369 | Podophyllum | 54 |
| Oxycoccus | 289 | PHÆNOGAMOUS PL. | 33 | Podostemaceæ | 429 |
| Oxydendrum | 296 | Phalacrolooma | 237 | Podostemon | 429 |
| Oxydendron | 623 | PHALARIDÆ | 606 | Pogonia | 507 |
| Oxyria | 419 | Phalaris | 643 | Poinsettia | 433 |
| Oxytrichium | 236 | Pharbitis | 375 | Poison Dogwood | 111 |
| Oxytropis | 133 | PHASEOLÆ | 125 | Poison Hemlock | 197 |
| | | Phaseolus | 140 | Poison Ivy | 111 |
| Pachysandra | 439 | Phegopteris | 663 | Poison Oak | 111 |
| Padus | 148 | Phelipæa | 323 | Poison Sumach | 111 |
| Pæonia | 48 | Philadelphus | 166 | Pokeweed | 405 |
| Pæpalanthus | 550 | PHLEOIDÆ | 603 | Pokeweed Family | 405 |
| Painted Cup | 336 | Phleum | 608 | Polanisia | 75 |
| Panax | 199 | Phlomis | 360 | POLEMONIACEÆ | 370 |
| Pancreatium | 513 | Phlox | 371 | POLEMONIÆ | 370 |
| PANICEÆ | 606 | Phoradendron | 426 | Polemonium | 371 |
| Panic-Grass | 645 | Phragmites | 636 | Polemonium Family | 370 |
| Panicum | 645 | Phryma | 341 | Polygala | 120 |
| Papaver | 58 | PHRYMÆ | 339 | POLYGALACEÆ | 120 |
| PAPAVERACEÆ | 58 | Phyllanthus | 439 | POLYGONACEÆ | 414 |
| Papaw | 50 | Phyllodoce | 297 | Polygonatum | 531 |
| PAPILIONACEÆ | 123 | Phyllotachys | 573 | Polygonum | 414 |
| Pappoose-root | 53 | Physalis | 381 | Polymnia | 247 |
| Papyrus | 552 | Physocarpus | 149 | Polypetalous Exogenous Pl. | 34 |
| Pardanthus | 517 | Physostegia | 354 | POLYPODIACEÆ | 656 |
| Parietaria | 446 | Phytolacca | 405 | POLYPODIÆ | 656 |
| Parnassia | 166 | PHYTOLACCACEÆ | 405 | Polypodium | 658 |
| Paronychia | 96 | Picea | 471 | Polypody | 618 |
| Parsley Family | 187 | Pickerel-weed | 545 | Polypogon | 652 |
| Parsley Piert | 151 | Pickerel-weed Family | 544 | Polyprenum | 391 |
| Parsnip | 191 | Picrococcus | 290 | Polystichum | 666 |
| Parthenium | 249 | Pieris | 295 | Polytænia | 191 |
| Partridge-berry | 211, 293 | Pigeon-Berry | 405 | POMEÆ | 147 |
| Partridge Pea | 144 | Pig-nut | 449 | Pomme Blanche | 129 |
| PASPALÆ | 606 | Pigweed | 406 | Pomme de Prairie | 129 |
| Paspalum | 644 | Pilea | 445 | Pond Spice | 423 |
| Pasque-flower | 36 | Pilinophytum | 438 | Pondweed | 484 |
| Passiflora | 185 | Pimpernel | 316 | Pondweed Family | 482 |
| PASSIFLORACEÆ | 185 | Pine | 469 | Pontederia | 545 |
| Passion-Flower | 185 | Pine-Apple Family | 515 | PONTEDERIACEÆ | 544 |
| Passion-Flower Family | 185 | Pine-drops | 303 | Poor Man's Weather-glass | 316 |
| Pastinaca | 191 | Pine-Family | 468 | Poplar | 50, 466 |
| Pavia | 118 | Pine-sap | 304 | Poppy | 58 |
| Pea | 139 | Pine-weed | 86 | Poppy Family | 58 |
| Pear | 161 | Pinguicula | 320 | Populus | 466 |
| Pear Family | 147 | Pink | 88 | Porcupine Grass | 617 |
| Pearlwort | 94 | Pink Family | 87 | Portulaca | 97 |
| Pecan-nut | 448 | Pink-root | 392 | PORTULACACEÆ | 97 |
| Pedicularis | 337 | Pinus | 469 | Portuna | 295 |
| Pelargonium | 106 | Pinweed | 81 | Potamogeton | 484 |
| Pellæa | 659 | Pinxter-flower | 299 | Potato | 380 |
| Pellitory | 446 | Pipe-Vine | 404 | Potentilla | 153 |
| Peltandra | 476 | Pipewort | 549 | POTERIEÆ | 147 |
| Pencil-Flower | 138 | Pipewort Family | 549 | Poterium | 150 |
| Pennycress | 73 | Pipissewa | 303 | Poverty-Grass | 618 |
| Pennyroyal | 350 | Piptatherum | 617 | Prairie Clover | 130 |
| Pentalophus | 363 | Pitcher-Plants | 57 | Prairie Dock | 249 |
| Penthorum | 171 | Planera | 443 | Prickly Pear | 185 |
| Pentstemon | 327 | Planer-tree | 443 | Prickly-Poppy | 59 |
| Peplis | 182 | Plane-tree | 447 | Prim | 400 |
| Pepperbush | 296 | Plane-tree Family | 446 | Primrose | 313 |
| Peppergrass | 74 | PLANTAGINACEÆ | 310 | Primrose Family | 313 |
| Pepperidge | 201 | Plantago | 310 | Primula | 313 |
| Peppermint | 344 | Plantain | 310 | PRIMULACEÆ | 313 |
| Pepper-root | 65 | Plantain Family | 310 | PRIMULÆ | 313 |
| Pepperwort | 74 | PLATANACEÆ | 446 | Prince's Feather | 412, 415 |
| Periploca | 400 | Platanthera | 500 | Prince's Pine | 303 |
| PERIFLOCEÆ | 394 | Platanus | 447 | Prinoides | 306 |
| Peristylus | 500 | Platensis | 397 | Prinos | 305, 307 |
| Periwinkle | 392 | Pluchea | 247 | Privet | 400 |
| Persea | 422 | Plum | 147 | Prosartes | 528 |
| Persicaria | 415 | PLUMBAGINACEÆ | 312 | Proserpinaca | 175 |

| | | | | | |
|-------------------------|----------|------------------------|----------|-------------------|----------|
| Prunus | 147 | RHINANTHIDEÆ | 324 | Sand-Myrtle | 301 |
| Pseudo-Polygonella | 417 | Rhinanthus | 337 | Sand-Spurrey | 95 |
| Psilocarya | 668 | Rhododendron | 299 | Sandwort | 90 |
| Psoralea | 128 | Rhodora | 300 | Sanguinaria | 60 |
| PSORALIEÆ | 124 | RHODOREÆ | 287 | Sanguisorba | 150 |
| Psyllophora | 573 | Rhubarb | 414 | Sanicle | 190 |
| Ptelea | 110 | Rhus | 111 | Sanicula | 190 |
| PTERIDEÆ | 656 | Rhynchosia | 142 | SANTALACEÆ | 425 |
| Pteris | 658 | Rhynchospora | 567 | SAPINDACEÆ | 116, 117 |
| Pterospora | 303 | RHYNCHOSPOREÆ | 551 | Saponaria | 88 |
| Puccoon | 382 | Ribes | 164 | SAPOTACEÆ | 308 |
| Pulmonaria | 364 | Ribesia | 165 | Sappodilla Family | 308 |
| Pulsatilla | 36 | Ribgrass | 310 | Sarothra | 86 |
| Pulse Family | 123 | Rice Cut-Grass | 607 | Sarracenia | 57 |
| Pumpkin | 186 | Rich-weed | 350, 445 | SARRACENIACEÆ | 57 |
| Purple Cone-flower | 253 | Ripplegrass | 311 | Sarsaparilla | 198, 199 |
| Purslane | 97 | River-weed | 429 | Sassafras | 483 |
| Purslane Family | 97 | River-weed Family | 429 | Satureia | 349 |
| Putty-root | 510 | Robinia | 130 | SATURIEÆ | 342 |
| Pycnanthemum | 346 | Rock-Cress | 67 | SAURURACEÆ | 427 |
| Pycreus | 552 | Rock-Rose | 80 | Saururus | 427 |
| Pyrola | 301 | Rock-Rose Family | 80 | Savin | 473 |
| PYROLEÆ (Pyrola Family) | 288 | Roman Wormwood | 251 | Savory | 349 |
| Pyrrhopappus | 280 | Rosa (Rose) | 158 | Saxifraga | 167 |
| Pyrularia | 426 | ROSEÆ | 147 | SAXIFRAGACEÆ | 163 |
| Pyrus | 161 | ROSACEÆ | 146 | Saxifrage | 167 |
| Pyxidantha | 373 | Rose-Acacia | 131 | SAXIFRAGEÆ | 163 |
| | | Rose-Bay | 299 | Saxifrage Family | 163 |
| Quaking-Grass | 633 | Rose Family | 146 | Scabious | 215 |
| Quamash | 533 | Rose-Mallow | 102 | Schedonorus | 635 |
| Quamoclit | 374 | Roseroot | 172 | Scheuchzeria | 491 |
| Queen of the Prairie | 149 | Rosin-Plant | 248 | Schizæa | 669 |
| QUERCINEÆ | 450 | Rosin-weed | 249 | SCHIZÆACEÆ | 657 |
| Quercitron | 454 | Roubieva | 408 | Schollera | 545 |
| Quercus | 450 | Rowan Tree | 162 | Schrankia | 145 |
| Queria | 96 | RCBEÆ | 147 | Schwalbea | 336 |
| Quick- or Quitch-Grass | 637 | RUBIACEÆ | 208 | Schweinitzia | 304 |
| Quillwort | 675 | Rubus | 156 | Scilla | 533 |
| | | Rudbeckia | 254 | SCIRPEÆ | 551 |
| Radish | 75 | Rue | 110 | SCIRPINEÆ | 551 |
| Ragiopteris | 668 | Rue-Anemone | 38 | Scirpus | 560 |
| Ragweed | 250 | Ruellia | 339 | Scleranthus | 96 |
| Ragwort | 271 | Rumex | 419 | Scleria | 570 |
| Ram's-head | 511 | Ruppia | 483 | SCLERINEÆ | 551 |
| Ramsted | 326 | Rush | 537 | Sclerochloa | 628 |
| RANUNCULACEÆ | 34 | Rush Family | 536 | Sclerolepis | 222 |
| RANUNCULEÆ | 34 | Rush-Grass | 609 | Scoke | 405 |
| Ranunculus | 40 | Ruta | 110 | Scopolendrium | 662 |
| RAPHANEÆ | 64 | Rutabaga | 71 | Scorpion-grass | 364 |
| Raphanus | 75 | RUTACEÆ | 109 | Scotch Thistle | 274 |
| Raspberry | 156 | Rye-Grass | 637 | Scouring Rush | 653 |
| Rattle-box | 126 | | | Scrophularia | 326 |
| Rattlesnake-Grass | 627 | Sabbatia | 384 | SCROPHULARIACEÆ | 324 |
| Rattlesnake's-Master | 191 | SACCHAREÆ | 607 | Scutch-Grass | 622 |
| Rattlesnake-Plantain | 503 | Sacred Bean | 55 | Scutellaria | 355 |
| Rattlesnake-root | 278 | Sage | 350 | Sea Blite | 410 |
| Rattlesnake-weed | 277 | Sagina | 94 | Sea Lavender | 312 |
| Ray-Grass | 637 | Sagittaria | 492 | Sea Milkwort | 316 |
| Rayless Golden-rod | 245 | St. Andrew's Cross | 83 | Sea Ox-eye | 253 |
| Reboulea | 625 | St. John's-wort | 84 | Sea Purslane | 97 |
| Red Bay | 422 | St. John's-wort Family | 83 | Sea Rocket | 75 |
| Red-bud | 143 | St. Peter's-wort | 83 | Sea Sand-Reed | 616 |
| Red Osier | 200 | SALICACEÆ | 461 | Sea Spear-Grass | 628 |
| Red-root | 115, 514 | Salicornia | 409 | Sedge | 571 |
| Red-top | 612 | Salix | 461 | Sedge Family | 550 |
| Reed | 636 | Salsola | 410 | Sedum | 172 |
| Reed Bent-Grass | 614 | Saltmarsh-Grass | 620 | Seed-box | 180 |
| Reed-mace | 480 | Saltwort | 410 | Selaginella | 674 |
| Reed Meadow-Grass | 627 | Salvia | 350 | Self-heal | 355 |
| Rein-Orchis | 498 | Salvinia | 677 | Senebiera | 74 |
| Rensselæria | 476 | SAMBUCEÆ | 202 | Seneca Grass | 642 |
| Reseda | 76 | Sambucus | 205 | Seneca Snakeroot | 122 |
| RESEDAÇEÆ | 76 | SAMOLEÆ | 313 | Senecio | 270 |
| RHAMNACEÆ | 113 | Samolus | 317 | SENECIONEÆ | 221 |
| Rhamnus | 114 | Samphire | 409 | SENECIONIDEÆ | 219 |
| Rheumatism-root | 54 | Sandalwood Family | 425 | Senna | 144 |
| Rhexia | 181 | Sand-Grass | 624 | Sensitive Brier | 145 |

| | | | | | |
|-----------------------|-------------------------|-----------------------|----------|---------------------|----------|
| Sensitive Fern | 668 | Southern Buckthorn | 308 | STRATIOTIDÆ | 496 |
| Sensitive Joint-Vetch | 134 | Sow Thistle | 282 | Strawberry | 155 |
| Sericocarpus | 228 | Spanish-Bayonet | 635 | Strawberry Blite | 408 |
| Service-berry | 162 | Spanish-Needles | 262 | Strawberry Bush | 116 |
| Sesame-Grass | 650 | Sparanium | 481 | Strawberry Tomato | 382 |
| Sesuvium | 97 | Spartina | 619 | Streptopus | 629 |
| Setaria | 649 | Spatter-Dock | 56 | Striped Dogwood | 119 |
| Seymeria | 334 | Spear-Grass | 628 | Struthiopteris | 667 |
| Shad-bush | 162 | Spearmint | 344 | Sturtia | 104 |
| Shave-Grass | 655 | Spearwort | 41 | Stylipus | 152 |
| Sheep-berry | 206 | Specularia | 286 | Stylisma | 378 |
| Shellflower | 327 | Speedwell | 332 | Stylophorum | 59 |
| Shepherdia | 424 | Spergula | 96 | Stylosanthes | 138 |
| Shepherd's Purse | 73 | Spergularia | 95 | STYRACACEÆ | 309 |
| Shield-Fern | 664 | Spermacoce | 210 | STYRACEÆ | 309 |
| Shin-leaf | 301 | Spice-bush | 423 | Styrax | 309 |
| Shooting-Star | 314 | Spiderwort | 547 | Suæda | 410 |
| Shrubby Althæa | 102 | Spiderwort Family | 546 | Subularia | 73 |
| Shrubby Bitter-sweet | 116 | Spigelia | 392 | Succory | 275 |
| Shrubby Trefoil | 110 | Spike-Grass | 628, 635 | Sugarberry | 443 |
| Shrub Yellow-root | 47 | Spikenard | 199 | Sullivantia | 169 |
| Sibbaldia | 153 | Spike-Rush | 657 | Sumach | 111 |
| STETHORPÆÆ | 324 | Spindle-tree | 116 | Summer Haw | 160 |
| Sickle-pod | 68 | Spiræa | 140 | Summer Savory | 349 |
| Sicyos | 186 | SPIRÆÆ | 146 | Sundew | 82 |
| Sida | 101 | Spiranthes | 504 | Sundew Family | 82 |
| Side-saddle Flower | 57 | Spirodela | 479 | Sundrops | 179 |
| Sieversia | 153 | Spleenwort | 661 | Sunflower | 255 |
| Silene | 88 | Spoonwood | 298 | Supple-Jack | 114 |
| SLENEÆ | 87 | Sporobolus | 610 | Swamp Honeysuckle | 299 |
| SLICULOSÆ | 63 | Spotted Cowbane | 196 | Swamp Loosestrife | 183 |
| SLIQUOSÆ | 63 | Spotted Wintergreen | 303 | Sweet Bay | 49 |
| Silkweed | 394 | Spring-Beauty | 98 | Sweet-Brier | 159 |
| Silphium | 248 | Spruce | 471 | Sweet Cicely | 197 |
| Silver-bell Tree | 310 | Spurge | 430 | Sweet Clover | 128 |
| Silver-Berry | 425 | Spurge Family | 430 | Sweet Coltsfoot | 227 |
| Silver-Weed | 155 | Spurge-Nettle | 436 | Sweet-Fern | 458 |
| Simarubacææ | 111 | Spurred Butterfly-Pea | 141 | Sweet Flag | 478 |
| Sinapis | 70 | Spurred Gentian | 386 | Sweet-Gale Family | 457 |
| SYMBRIÆÆ | 63 | Spurrey | 96 | Sweet-Gum Tree | 174 |
| Sisymbrium | 70 | Squash | 186 | Sweet-leaf | 310 |
| Sisyrinchium | 517 | Squaw-root | 323 | Sweet Pepperbush | 296 |
| Sitolobium | 669 | Squaw-weed | 271 | Sweet Pine-sap | 304 |
| Stium | 196 | Squill | 533 | Sweet-Scented Shrub | 162 |
| Skullcap | 355 | Squirrel Corn | 61 | Sweet Vernal-Grass | 643 |
| Skunk Cabbage | 477 | Squirrel-tail Grass | 638 | Sweet-William | 371 |
| Sloe | 148 | STACHYDEÆ | 343 | Swine-Cress | 74 |
| Smartweed | 416 | Stachys | 358 | Sycamore | 447 |
| SMILACEÆ | 518 | Stagger-bush | 296 | Syena | 547 |
| Smilacina | 530 | Staff-tree | 116 | Symphoricarpus | 203 |
| Smilax | 518 | Staff-tree Family | 115 | Symphytum | 361 |
| Smilax Family | 518 | Staphylea | 117 | Symplocarpus | 477 |
| Snake-head | 327 | STAPHYLEACEÆ | 117 | SYMPLOCINEÆ | 309 |
| Snake-root | 190, 191, 223, 226, 404 | Statice | 312 | Symplocos | 310 |
| | | Star-Cucumber | 186 | Synandra | 354 |
| Snapdragon | 326 | Star Flower | 314 | Syndesmon | 38 |
| Sneeze-weed | 263 | Star-grass | 513, 615 | Synthyris | 331 |
| Sneezewort | 265 | Star-of-Bethlehem | 533 | Syringa | 166, 400 |
| Snowberry | 203 | Star-Thistle | 272 | | |
| Snowdrop | 310, 512 | Starwort | 92, 228 | Tacamahac | 467 |
| Snowflake | 512 | Steeple-Bush | 149 | Tænidia | 195 |
| Soapberry Family | 116 | Steironema | 315 | Tagetes | 262 |
| Soapwort | 88 | Stellaria | 92 | TAGETINEÆ | 220 |
| SOLANACEÆ | 380 | STELLATÆ | 208 | Talinum | 98 |
| Solanum | 380 | Stenactis | 237 | Tall Red-top | 624 |
| Solea | 76 | Stenanthium | 625 | Tamarack | 472 |
| Solidago | 239 | Stickseed | 365 | Tanacetum | 266 |
| Solomon's Seal | 631 | Stillingia | 436 | Tansy | 266 |
| Sonchus | 282 | Stipa | 617 | Tansy-Mustard | 70 |
| SOPHOREÆ | 125 | STIPRÆÆ | 604 | Tape-grass | 496 |
| Sorbus | 181 | Stitchwort | 92 | Taraxacum | 280 |
| Sorghum | 652 | Stone-crop | 172 | TARCHONANTHEÆ | 219 |
| Sorrel | 419 | Stone-root | 350 | Tare | 138 |
| Sorrel Family | 106 | Storax | 309 | TAXINEÆ | 468 |
| Sorrel-tree | 296 | Storax Family | 309 | Taxodium | 473 |
| Sour-Gumtree | 201 | Storksbill | 108 | Taxus | 474 |
| Sour-wood | 296 | Stramonium | 383 | Tea-berry | 293 |

| | | | | | |
|-------------------------|-----|-----------------------|----------|------------------------|----------|
| Tea-Plant | 103 | Trifolium | 126 | Vernonia | 222 |
| Tear-thumb | 418 | Triglochin | 491 | VERNONIACEÆ | 217 |
| Teasel | 215 | TRILLIDÆ | 520 | Veronica | 332 |
| Teasel-Family | 215 | Trillium | 522 | VERONICEÆ | 324 |
| Tecoma | 321 | Triodallus | 286 | Vervain | 339 |
| Telmatopace | 479 | Triosteum | 205 | Vervain Family | 339 |
| Tephrosia | 131 | Triplasis | 624 | Vesicaria | 73 |
| TEPHROSIE | 124 | Triple-awned Grass | 618 | Vetch | 138 |
| TERNSTRÆMIACEÆ | 103 | Tripsacum | 650 | Vetchling | 139 |
| Tetragonotheca | 252 | Tripterella | 496 | Viburnum | 206 |
| Tetranthera | 423 | Trisetum | 640 | Vicia | 138 |
| Teucrium | 343 | Triticum | 637 | VICIEÆ | 124 |
| Thalictrum | 38 | Trollius | 44 | Vigna | 574 |
| Thaspium | 194 | Tropæolum | 105 | Vilfa | 609 |
| Thermopsis | 679 | Troximon | 276 | Vincetoxicum | 399 |
| Thimbleberry | 157 | Trumpet-Creeper | 321 | Vine Family | 112 |
| Thin-Grass | 611 | Trumpet-flower | 321 | Viola | 77 |
| Thistle | 273 | Trumpets | 58 | VIOLACEÆ (Violet Fam.) | 76 |
| Thlaspi | 73 | Trumpet-weed | 225 | Violet | 77 |
| THLASPIDÆ | 63 | Tsuga | 471 | Viper's Bugloss | 361 |
| Thorn | 159 | TUBULIFLORÆ | 216 | Virgaurea | 240 |
| Thorn-Apple | 383 | Tuckermannia | 440 | Virgilia | 143 |
| Thorough-wax | 195 | Tulip-tree | 50 | Virginian Cowslip | 364 |
| Thoroughwort | 224 | Tupelo | 201 | Virginian Creeper | 113 |
| Three-leaved Nightshade | 522 | TUSSILAGINÆ | 218 | Virginia Snakeroot | 404 |
| Three-seeded Mercury | 436 | Tussilago | 227 | Virgin's-Bower | 35 |
| Three-thorned Acacia | 145 | Turnip | 71 | Viscum | 426 |
| Thrift | 312 | Turritis | 68 | VITACEÆ | 112 |
| Thuja | 472 | Turtle-head | 327 | Vitis | 112 |
| Thunbergia | 338 | Twayblade | 506, 509 | Vitis-Idæa | 290 |
| THYMELEACEÆ | 424 | Twig Rush | 570 | Waahoo | 116 |
| Thymus (Thyme) | 348 | Twin-flower | 202 | Wake-Robin | 522, 523 |
| Tiarella | 170 | Twin-leaf | 53 | Waldsteinia | 153 |
| Tickseed | 258 | Twisted Stalk | 529 | Walking-leaf | 663 |
| Tickseed Sunflower | 260 | Typha | 480 | Walnut | 447 |
| Tick-Trefoil | 134 | TYPHACEÆ | 480 | Walnut Family | 447 |
| Tiedemannia | 192 | Udora | 496 | Water Arum | 476 |
| Tiger-flower | 516 | ULMACEÆ | 441 | Water Beech | 457 |
| Tigridia | 516 | Ulmaria | 149 | Water Chinquepin | 55 |
| Tilia | 103 | Ulmus | 442 | Wart-Cress | 64, 74 |
| TILIACEÆ | 103 | UMBELLIFERÆ | 187 | Water-Hemlock | 196 |
| Tillæa | 171 | Umbrella-Grass | 556 | Water-Ilmp | 413 |
| Tillandsia | 515 | Umbrella-leaf | 53 | Water Horehound | 345 |
| Timothy | 608 | Umbrella-tree | 49 | Waterleaf | 367 |
| Tiniaria | 418 | Umbrella-tree | 321 | Waterleaf Family | 367 |
| Tipularia | 508 | Umicorn-plant | 321 | Water-Lily | 56 |
| Tithymalopsis | 433 | Umiola | 635 | Water-Lily Family | 54, 55 |
| Tithymalus | 433 | Uralepis | 624 | Water-Locust | 145 |
| Toad-Flax | 326 | Urtica | 444 | Watermelon | 186 |
| Tobacco | 383 | URTICACEÆ | 440 | Water-Nilfoil | 174 |
| Tofieldia | 527 | URTICEÆ | 441 | Water-Nilfoil Family | 174 |
| Tomato | 380 | Utricularia | 318 | Water-Nymph | 56 |
| Toothache-Grass | 620 | Uvularia | 528 | Water-Parsnip | 196 |
| Toothache-tree | 110 | UVULARIÆ | 521 | Water-Pennywort | 189 |
| Toothwort | 65 | Vaccaria | 88 | Water-pepper | 416 |
| Touch-me-not | 108 | VACCINIEÆ | 287 | Water-Pimpernel | 317 |
| Tovaria | 417 | Vaccinium | 289, 679 | Water-Plantain | 491 |
| Tower-Mustard | 69 | Vahlodea | 641 | Water-Plantain Family | 490 |
| Toxicodendron | 111 | Valeriana (Valerian) | 213 | Water-Purslane | 181 |
| Trachynotia | 619 | VALERIANACEÆ | 213 | Water-Rice | 608 |
| Tradescantia | 547 | Valerianella | 214 | Water-Shield Family | 54 |
| Tragia | 437 | Valerian Family | 213 | Water-Star-grass | 545 |
| Trailing Arbutus | 293 | Vallisneria | 496 | Water-Starworts | 427 |
| Trautvetteria | 39 | VALLISNERIÆ | 495 | Water-Violet | 317 |
| Treacle Mustard | 69 | Vanilla Grass | 642 | Water-weed | 495 |
| Tread-softly | 436 | Vanilla-plant | 224 | Water-Willow | 338 |
| Tree of Heaven | 110 | Velvet-leaf | 101 | Water-wort | 86 |
| Trefoil | 126 | Venus's Fly-trap | 83 | Water-wort Family | 86 |
| Trichelostylis | 567 | Venus's Looking-glass | 286 | Wax-Myrtle | 457 |
| Trichochloa | 614 | Veratrum | 525 | Wax-work | 116 |
| Trichodium | 611 | VERBASCÆ | 324 | Wayfaring-tree | 207 |
| Trichophorum | 565 | Verbascum | 325 | Western Wall-flower | 69 |
| Trichostema | 343 | Verbena | 339 | Whahoo | 443 |
| Tricuspis | 623 | VERBENACEÆ | 339 | Wheat | 637 |
| Tridynia | 315 | VERBENÆ | 339 | Whin | 126 |
| Trientalis | 314 | Verbesina | 262 | White Alder | 296 |
| TRIFOLIEÆ | 123 | | | | |

| | | | | | |
|----------------------|----------|--------------------|----------|--------------------------|-----|
| White Cedar | 472, 473 | Windsoria | 623 | Xanthium | 251 |
| White Daisy | 265 | Winged Pigweed | 406 | Xerophyllum | 526 |
| White Grass | 607 | Winterberry | 307 | Xylosteum | 204 |
| White Lettuce | 278 | Winter-Cress | 69 | XYRIDACEÆ | 547 |
| White Snakeroot | 226 | Wintergreen | 293, 301 | Xyris | 548 |
| White Thorn | 159 | Wire-Grass | 623, 629 | Yam | 518 |
| White-topped Aster | 228 | Wistaria | 131 | Yam Family | 518 |
| White-weed | 265 | Witch-Hazel | 173 | Yard-Grass | 622 |
| Whitlow Grass | 71 | Witch-Hazel Family | 173 | Yarrow | 265 |
| Whitlow-wort | 96 | Withe-rod | 206 | Yaupon | 306 |
| Whortleberry Family | 287 | Woad-Waxen | 126 | Yellow-eyed Grass | 548 |
| Wicopy | 424 | Wolfberry | 203 | Yellow-eyed Grass Family | 547 |
| Wild Allspice | 423 | Wolflia | 480 | Yellow (False) Jessamine | 391 |
| Wild Balsam-apple | 186 | Wolfsbane | 46 | Yellow Pond-Lily | 56 |
| Wild Bean | 140 | Wood Betony | 337, 358 | Yellow Puccoon | 47 |
| Wild Bergamot | 351 | Woodbine | 113, 203 | Yellow-Rattle | 337 |
| Wild Chamomile | 266 | Wood-Fern | 664 | Yellow Rocket | 69 |
| Wild Comfrey | 366 | Wood-Grass | 652 | Yellow-wood | 143 |
| Wild Elder | 199 | Wood-Nettle | 445 | Yew | 474 |
| Wild Ginger | 403 | Wood Reed-Grass | 612 | Yew Family | 468 |
| Wild Hyacinth | 532 | Wood-Rush | 536 | Yucca | 535 |
| Wild Liquorice | 209, 210 | Wood-Sage | 343 | Zannichellia | 483 |
| Wild Marjoram | 348 | Woodsia | 668 | Zanthorhiza | 47 |
| Wild Oat-Grass | 639 | Wood-Sorrel | 109 | Zanthoxylum | 110 |
| Wild Pink | 89 | Woodwardia | 660 | Zapania | 340 |
| Wild Potato-Vine | 375 | Wool-Grass | 565 | Zephyranthes | 513 |
| Wild Sarsaparilla | 198 | Woolly Beard-Grass | 651 | Zizania | 608 |
| Wild Sensitive-Plant | 144 | Worm-grass | 392 | Zizia | 195 |
| Willow | 461 | Wormseed | 408 | Zostera | 483 |
| Willow Family | 461 | Worm-seed Mustard | 69 | Zygadenus (Zygadene) | 524 |
| Willow-herb | 177 | Wormwood | 266 | | |
| Wind-flower | 86 | Woundwort | 358 | | |

EXPLANATION OF THE PLATES.

Genera of Cyperaceæ or Sedges.

TAB. I.

- CYPERUS.** — Small plant of *C. diandrus* (1); a spike magnified (2); a piece of the rhachis with one scale enclosing its flower (3); a separate flower more magnified (4). *C. erythrorhizos*, a spike magnified, the lower scales and flowers have fallen, showing the little internal scales of the section *Papyrus* (5), formed of the winged margins of the joints of the rhachis detached; a separate one more enlarged (6); a flower (7); an achenium (8), cut in two. *C. dentatus*, a piece of the rhachis of a spike with lower part of one scale, showing how it is decurrent on the joint beneath (cut across) to form scale-like wings (9).
- DULICHIMUM.** — Upper part of a plant (1); part of a spike somewhat enlarged (2); piece of rhachis and one scale decurrent on the joint beneath (3); magnified flower (4).
- KYLLINGIA.** — Plant of *K. pumila* (1); one-flowered spike on a piece of the rhachis, enlarged (2); the same more enlarged and open (3); achenium (4), and section of same magnified (5).

TAB. II.

- HEMICARPHA.** — Plant (1), natural size; a spike enlarged, with its bract (2); magnified scale of the same (3); a flower (5), with its single stamen and minute internal scale, magnified; achenium (6), magnified.
- LIPOCARPHA.** — Upper part of plants with spikes (1); diagram of a flower, viz. of ovary between the two internal scales, and single stamen, scale of the spike on one side, axis of the spike on the other (2); scale of spike detached (3); a flower with its two inner scales (4); achenium magnified (5).
- FUIRENA.** — Upper portion of plants (1); scale of spike enclosing a flower (2); open scale of same (3); flower (4); one of the scales and one of the bristles of the flower (5), achenium (6), and section of same (7).

TAB. III.

- ELEOCHARIS.** — Small plant of *E. olivacea* (1); the spike enlarged (2); detached scale (3); flower (4); achenium and bristles (5). *E. quadrangulata*, spike (6); a scale (7); flower (8); achenium and bristles (9). *E. tuberculosa*; the achenium with its great tubercle, and bristles (10).
- SCIRPUS.** — Summit of plant of small *S. debilis* (1); a spike (2); a scale of the same (3), and flower (4); achenium with its bristles (5). *S. (Trichophorum)*

Eriophorum; a small portion of the inflorescence (6); a flower (7); a spike in fruit (8); achenium from the same, with the tortuous bristles much lengthened (9); section of the achenium (10).

ERIPHORUM.—Small plant of *E. alpinum*, in flower (1); spike (2); a scale (3), and (4) a flower from the same; the spike in fruit, the bristles forming a cottony tuft (5); achenium and its bristles (6).

FIMBRISTYLIS.—Summit of a small flowering stem of *F. laxa* (1); a spike of the same (2); a detached scale (3), and (4) a flower of the same; achenium (5). *F. (Trichostylis) autumnalis*; a spike, enlarged (6); flower (7); achenium (8), and (9) section of the same.

TAB. IV.

DICHROMENA.—Head and involucre of *D. latifolia* (1); a scale from one of the spikes (2), and the same cut across (3); a flower (4); achenium with its tubercle (5).

RHYNCHOSPORA.—Upper part of flowering stem of *R. Torreyana* (1); a spike (2); detached flower (3); achenium (4) with short bristles at its base; one of these bristles more magnified (5).

R. (§ PSILOCARYA):—Part of plant (1), enlarged spike (2), detached scale (3), flower (4), and achenium with its beak (5) of *R. (Psilocarya) scirpoides*.

R. (§ CERATOSCHENUS):—Upper part of fruiting plant (1), detached spike (2), flower (3), and beaked achenium with its bristles (4) of *R. (Ceratostchenus) macrostachya*.

TAB. V.

CLADIUM.—Summit of a plant of *C. mariscoides* (1); detached spike (2); same, open, showing a staminate and a perfect flower (3); the nut-like achenium (4), and the longitudinal section of the same (5).

SCLERIA.—Summit of a flowering stem of *S. reticularis* (6); three spikelets from a cluster, the middle one pistillate, the lateral ones staminate (7); staminate spikelet displaying four male flowers, the filaments of two of them have lost their anthers (8); pistillate spikelet displaying a single pistillate flower (9); achenium with the 3-lobed double cup underneath (10).

CAREX.—Plant of *C. pauciflora* (11); a staminate flower with its scale (12); scale (13), and mature pistillate flower, in its perigynium (14); cross section of perigynium and of the contained achenium (15); achenium on its stalk, style and stigmas (16). *C. Steudelii*, upper part of flowering plant (17); the spike enlarged (18); a staminate flower and its scale (19); pistillate flower in its perigynium (20); the same with half the perigynium cut away to show the contained achenium and style (21).

TAB. VI.

CAREX.—*C. trisperma*, upper part of a stem in fruit (1); enlarged spike displayed, with three staminate and two pistillate flowers (2); a scale (3) and a ripe perigynium (4), of the latter; with a section of the perigynium near the base, and of the contained perigynium (5). *C. straminea*, summit of a fruiting plant (6); a spike enlarged (7); scale of a pistillate flower (8); the winged perigynium and the contained achenium cut across (9); detached achenium with persistent style and stigmas (10). *C. umbellata*, whole plant (11); a perigynium and its scale (12); cross-section towards the base of perigynium and its contained achenium (13); detached achenium with its persistent style and stigmas (14). *C. bullata*, upper part of plant in fruit, with one pistillate and two staminate spikes (15); one of its staminate flowers with the scale (16); a pistillate scale (17) and mature perigynium (18); longitudinal section of the latter, showing the achenium and its style (19), and cross section of the same (20).

Genera of Gramineæ or Grasses.

TAB. VII.

- LEERSIA. — Panicle of *L. oryzoides*, reduced in size (1); a branchlet of the same with its spikelets, of the natural size (2); and an open spikelet in flower, enlarged (3).
- ZIZANIA. — A staminate (1) and a pistillate (2) flower or spikelet of *Z. aquatica*; a magnified pistil with a pair of squamulæ or hypogynous scales (3); a grain (4); and a magnified longitudinal section of the lower part of the same, showing the embryo at the outside of the base of the albumen.
- ALOPECURUS. — Part of a plant of *A. geniculatus*, in flower (1); a few spikelets from the spike-like inflorescence, moderately magnified (2); an open spikelet in flower, more magnified (3), and the single lower palet detached (4).
- PHLEUM. — A detached spikelet of *P. pratense*, having the flower with its palets raised above the glumes, magnified.
- CRYPISIS. — Inflorescence (1) of *C. schœnoides*; a separate enlarged spikelet (2); and the same open, in flower (3).
- VILFA. — An enlarged spikelet of *V. vaginæflora* (1); and the same displayed (2).
- SPOROBOLUS. — A spikelet of *S. cryptandrus*, magnified (1); the same with the flower open, the palets raised above the glumes (2); and the fruit (3), more magnified, showing the seed loose in the pericarp (utricle).
- AGROSTIS. — Panicle of *A. vulgaris* (1); with an enlarged open spikelet of the same: also (3) the rough pedicel and glumes of *A. scabra*, with the flower separated, the latter having no upper palet.

TAB. VIII.

- POLYPOGON. — Spike-like contracted panicle of *P. Monspelienis* (1); an enlarged detached spikelet, showing the long awns to the glumes (2); the same open in flower (3); and a separate flower without the glumes (4).
- CINNA. — A magnified spikelet of *C. arundinacea* (1); and the same open, displaying the palets, the single stamen, and the pistil (2).
- MUHLENBERGIA. — A magnified closed spikelet of *M. sylvatica* (1); the same with the open flower raised out of the glumes (2). A magnified spikelet of *M. diffusa* (3); its minute and unequal glumes more magnified (4); and an open spikelet of the same (5).
- BRACHYELYTRUM. — A spikelet of *B. aristatum* enlarged (1); the same displayed (2).
- CALAMAGROSTIS. — An open spikelet of *C. Canadensis*, enlarged, displaying all the parts (1); the same with the flower raised out of the glumes, showing the hairy rudiment behind the upper palet (2).
- ORYZOPSIS. — An open magnified spikelet of *O. asperifolia* (1); and the flower of the same removed from the glumes (2). Notice the remarkably long squamulæ or hypogynous scales, which here nearly equal the palets in length.
- STIPA. — Glumes and flower (a little separated) of *S. avenacea*, enlarged.
- ARISTIDA. — A spikelet of *A. purpurascens*, enlarged.

TAB. IX.

- SPARTINA. — Portion of the inflorescence of *S. stricta*, of the natural size (1); a spikelet enlarged (2); and the same displayed, the flower raised above the glumes (3).
- CTENIUM. — Spike of *C. Americanum* (1); a single spikelet magnified (2); and the same displayed, the glumes separated (3).
- BOUTELLOUA. — A portion of the compound spike, of the natural size (1); and a spikelet displayed and magnified (2), the flowers raised out of the glumes.

- GYMNOPOGON. — Inflorescence of *G. racemosus*, reduced in size (1); and a magnified spikelet with the parts displayed (2).
- CYNODON. — Inflorescence, of digitate spikes (1); a spikelet magnified and displayed, showing a perfect flower and a rudiment (2).
- DACTYLOCTENIUM. — Inflorescence of *D. Ægyptiacum*, of digitate spikes (1); one of the spikelets magnified (2); the fruit magnified (3), showing the seed loose in the thin pericarp (utricle); and (4) the wrinkled seed more magnified.
- ELEUSINE. — One of the spikes from the digitate inflorescence of *E. Indica* (1); a magnified spikelet (2); the same with the flowers more displayed (3); a flower from the last showing its parts (4); the fruit magnified, showing the seed loose in the utricle (5); and the wrinkled seed detached (6).
- LEPTOCHLOA. — Small portion of the inflorescence of *L. fascicularis* (1); one of its spikelets displayed and magnified (2); an open flower of the same (3).

TAB. X.

- TRICUSPIS. — Magnified spikelet of *T. seslerioides* (1); the same displayed and the lowest flower open (2); back view of the lower palets spread out (3).
- GRAPHEPHORUM. — A magnified spikelet of *G. melicoides*, displayed (1); a part of the hairy rhachis and one flower of the same (2).
- DIARRHENA. — A spikelet of *D. Americana*, enlarged (1); the grain and palets (2).
- DACTYLIS. — A spikelet of *D. glomerata* magnified and displayed.
- KÆLERIA. — A magnified spikelet of *K. cristata*, expanded, showing the glumes, the three flowers, and a rudiment (1); lower half of a lower palet, partly spread open (2); it is much more folded and keeled in its natural condition.
- EATONIA. — A magnified spikelet of *E. obtusata*, expanded, showing the glumes, the two flowers, and a rudiment.
- MELICA. — A magnified spikelet of *M. mutica*, expanded, showing the glumes, two perfect flowers, and an abortive one.
- GLYCERIA. — A magnified spikelet of *G. nervata* (1); a separate flower with one joint of the rhachis (2); and (3) the lower half of a lower palet, showing its form (rounded on the back, not keeled.)
- BRIZOPYRUM. — A pistillate spikelet of *B. spicatum*, enlarged (1); a flower from the same (2); and a flower from a staminate spikelet (3).
- POA. — Panicle of *P. compressa*, reduced in size (1); a magnified spikelet (2); a separate flower more magnified (3); a lower palet cut across and somewhat outspread (4).
- ERAGROSTIS. — A spikelet of *E. pilosa*, enlarged (1); the same, from which the glumes and all of six lower flowers except the upper palet have fallen away (2); a magnified flower, open (3); the lower palet of the same outspread (4).
- BRIZA. — A spikelet of *B. media*, enlarged (1); a separate flower (2).
- FESTUCA. — A spikelet of *F. elatior*, enlarged (1); a separate flower (2); lower part of a lower palet, outspread (3).
- BROMUS. — A spikelet of *B. secalinus*, or Chess (1); and a separate flower, enlarged (2).

TAB. XI.

- UNIOLA. — A spikelet of *U. latifolia*, of about the natural size (1); a flower, enlarged (2); empty lower palet of the lowest (sterile) flower (3).
- PHRAGMITES. — A spikelet of *P. communis*, enlarged (1); one of the perfect flowers, enlarged (2); and the lowest flower (3), which has stamens only.
- ARUNDINARIA. — A spikelet of *A. macrosperma* (1); and a separate flower, magnified (2).
- LEPTURUS. — Portion of the spike of *L. paniculatus*, enlarged (1); and a flower magnified (2).

- LOLIUM.** — Portion of the spike of *L. temulentum* (1); and a separate flower, magnified (2).
- TRITICUM.** — Portion of the spike of *T. repens*, or Couch-Grass, of about the natural size (1); a flower, magnified (2).
- HORDEUM.** — The three one-flowered spikelets from one joint of the spike of *H. jubatum*, with their awn-like glumes, the lateral flowers abortive and neutral the middle one alone perfect (1); this perfect flower (with an awn-like rudiment) open and enlarged (2).
- ELYMUS.** — The two spikelets of one joint of the spike of *E. Virginicus*, about the natural size (1); the glumes and the flowers of one spikelet, enlarged and displayed (2); and an open flower, more magnified (3).
- GYMNOSTICHUM.** — A spikelet of *G. Hystrix* (1), and an expanded flower, magnified (2).

TAB. XII.

- AIRA.** — Panicle of *A. flexuosa* (1); a spikelet, magnified, the parts displayed (2); and one of the flowers detached and open (3).
- DANTHONIA.** — Panicle of *D. spicata* (1); a spikelet enlarged (2); and a separate flower from the same (3).
- TRisetum.** — A spikelet of *T. subspicatum*, var. *molle*, expanded and magnified (1); and a separate open flower (2).
- AVENA.** — A spikelet of *A. striata* displayed and magnified (1); and a separate flower (2).
- ARRENATHERUM.** — A spikelet of *A. avenaceum*, displayed and magnified: 1, the glumes; 2, the flowers, the lower one staminate only, the next one perfect, and the third a rudiment.
- HOLCUS.** — A spikelet of *H. lanatus* magnified (1); the same displayed to show the two flowers, the lower one perfect and awnless, the upper staminate and awned (2).

TAB. XIII.

- HIEROCHLOA.** — A spikelet of *H. borealis*, enlarged (1); the same displayed, the flowers separated from the glumes, the two lateral ones with three stamens and no pistil; the middle or terminal one with a pistil and only two stamens (2).
- ANTHOXANTHUM.** — The spike-like inflorescence of *A. odoratum* (1); a spikelet magnified (2); another with the parts displayed, the flowers raised from the glumes, the lateral ones neutral, each of a single and awned palet, the middle one perfect and diandrous (3).
- PHALARIS.** — A spikelet of *P. arundinacea* enlarged (1); the glumes, and the perfect flower with a hairy rudiment on each side of it (2).
- MILIUM.** — Portion of the panicle of *M. effusum* (1); a closed spikelet magnified (2); and the same displayed (3).
- AMPHICARPUM.** — A spikelet from the panicle of *A. Purshii*, magnified (1); the same with the parts displayed (2); and a radical (fertile) spikelet, enlarged (3).
- PASPALUM.** — Inflorescence of *P. læve* (1); a closed spikelet, enlarged (2); the same with the parts displayed (3).
- PANICUM.** — Part of a spike of *P. (Digitaria) sanguinale* (1); one of its spikelets magnified (2); the same with the parts displayed (3); in this the lower flower is neutral and of a single palet. A spikelet of *P. capillare*, magnified (4), and the same displayed (5); the lower flower a single palet. A spikelet of *P. clandestinum*, magnified (6), and the same displayed (7); the lower neutral, of two palets. A spikelet of *P. virgatum*, magnified (8); the same displayed (9); the lower flower of two palets and staminate.
- SETARIA.** — A magnified spikelet of *S. glauca*, with the accompanying cluster of bristles (1); the spikelet displayed, showing the neutral lower flower of two palets and the perfect flower (2).

TAB. XIV.

- CENCHRUS. — Involucre of *C. tribuloides*, in flower, enlarged (1); longitudinal section of the same (2); a spikelet displayed (3): the stigmas should belong to the right-hand flower: the left-hand or lower flower is only staminate.
- TRIPSACUM. — Piece of the spike (of the natural size), pistillate below, staminate above (1); a longitudinal section of one of the pistillate spikelets (2); a pistillate spikelet with its parts displayed (3); a staminate (two-flowered) spikelet, with its parts displayed (4).
- ERIANTHUS. — Part of the hairy inflorescence with two spikelets of *E. alopecuroides*, enlarged (1); one of the spikelets displayed (2).
- ANDROPOGON. — Small portion of the spike of *A. furcatus*, enlarged, with one fertile and awned spikelet, and one staminate and awnless spikelet (1); the fertile spikelet (2); and the staminate spikelet (3) displayed.
- SORGHUM. — A fertile spikelet of *S. nutans*, enlarged, with a sterile pedicel on each side (1); the spikelet displayed (2).

Genera of Filices or Ferns.

TAB. XV.

- POLYPODIUM. — Plant; piece of the frond (1): a magnified sporangium with its stalk, and another bursting and discharging spores, of *P. vulgare*, L.
- STRUTHIOPTERIS. — Pinna of the sterile frond (1) of *S. Germanica*, Willd.; portion of a fertile frond (2); a piece of one pinna cut off to show the manner in which it is rolled up (3); and a portion of the last, magnified, with one side unrolled (4); towards the base of the sporangia all removed, to show how the fruit-dots are borne each on the middle of a vein.
- PELLÆA. — Sterile and fertile plants of *P. gracilis*, and a portion of the fertile frond (1) enlarged, with a piece of the marginal indusium turned back to display the fruit; the sporangia are all removed from the fruit-bearing tips of the two forks of the lower vein.

TAB. XVI.

- PTERIS. — A pinnule of *P. aquilina*, L.; and a piece of one of the lobes, enlarged (2), the marginal indusium rolled back on one side, displaying the fruit; the sporangia all removed from the lower part to show the receptacle that bears them, viz. a cross line connecting the tips of the veins.
- ADIANTUM. — Piece of the frond of *A. pedatum*, L. (1); a pinnule somewhat enlarged (2); and a piece of one (3) more enlarged, with the indusium of one fruit-dot turned back to show the attachment of the fruit.
- CHEILANTHES. — Small plant of *C. vestita* (1); and a fruit-bearing pinnule, enlarged (2).
- WOODWARDIA. — Portion of the sterile (1) and of the fertile frond (2) of *W. angustifolia*; a piece of the latter enlarged (3); piece of the frond of *W. Virginica* (4); and part of a fruiting lobe (5), enlarged.

TAB. XVII.

- CAMPTOSORUS. — Plant of *C. rhizophyllus*, Link.; and a portion of a frond, with fruit-dots, enlarged (1).
- SCOLOPENDRIUM. — Tip of a fertile frond of *S. vulgare*; and (2) a piece enlarged, with two fruit-dots.
- ASPLENIUM. — A pinna of *A. thelypteroides*, Michx. (1); and part of a lobe (2) in fruit, enlarged.
- DICKSONIA. — Pinna of *D. punctilobula*, Hook. (1); portion of a pinnule (2), enlarged; and a fruit-dot in its cup-shaped indusium (3).

TAB. XVIII.

- CYSTOPTERIS.** — Piece of the frond of *C. bulbifera*, *Bernh.* (1); a lobe in fruit (2), enlarged; and a small portion more magnified (3), bearing a fruit-dot with its indusium thrown back.
- WOODSIA.** — Small frond of *W. glabella*, *R. Br.* (1); a part of a fruiting pinna of the same (2), magnified; and a separate indusium (3), more magnified; a piece of a fruitful pinnule of *W. obtusa*, *Torr.* (4), enlarged; and a fruit with the opened indusium beneath (5), more magnified.
- ASPIDIUM.** — Pinna of *A. (Dryopteris) marginale*, *Swartz* (1); and a magnified fruiting portion (2); piece of *A. (Polystichum) acrostichoides* (3); and a small fruiting portion (4), magnified.
- ONOCLEA.** — Sterile and fertile frond of *O. sensibilis*, *L.*; front view of a fruiting contracted pinnule, enlarged (1); and the same laid open and viewed from the other side (2): on one lobe the sporangia are removed from the veins.

TAB. XIX.

- SCHIZEA.** — Plant of *S. pusilla*, *Pursh.*, of the natural size; a fertile pinna with eleven sporangia (1), magnified; and a separate sporangium (2), more magnified.
- LYGODIUM.** — Summit of frond of *L. palmatum*, *Swartz* (1), with fertile and sterile divisions; a fruiting lobe enlarged (2), with two of the lower scales, or indusia, removed, displaying a sporangium under each; and a sporangium more magnified (3).
- OSMUNDA.** — Small piece of the frond of *O. Claytoniana*, *L.* (1), with a fertile and a sterile pinna; a portion of the fruit magnified (2); and one sporangium more magnified (3).
- BOTRYCHIUM.** — Plant of *B. lunarioides*, *Swartz*; and a portion of the fruit (1), with six sporangia, magnified.
- OPHIOGLOSSUM.** — Frond of *O. vulgatum*, *L.*; and a portion of the fruiting spike enlarged (1).

Genera of Equisetaceæ, Lycopodiaceæ, &c.

TAB. XX.


- EQUISETUM.** — Upper part of fertile plant of *E. limosum*, *L.* (1); one of the shield-shaped scales or receptacles of the spike, with the six sporangia underneath (2), enlarged; same seen from below, discharging the spores (3); a magnified spore with the club-shaped filaments spreading (4); and (5) the same with the filaments coiled up.
- LYCOPODIUM.** — Plant of *L. Carolinianum*, *L.*; and (1) a magnified scale of the spike removed, with the sporangium in its axil, discharging powdery spores.
- SELAGINELLA.** — Plant of *S. rupestris*, *Spring*; part of a fertile spike, enlarged (1); scale from the upper part of it (2), with its sporangium, containing innumerable powdery spores; scale from the base (3), with its sporangium containing few large spores; and (4) three large spores.
- ISOETES.** — Plant of *I. lacustris* (1); sporocarp containing the minute spores, cut across (2), enlarged; same divided lengthwise (5); sporocarp with the large spores, divided lengthwise (3); and (4) three large spores more magnified.
- AZOLLA.** — Plant (1); a portion magnified (2), with two kinds of organs; sterile sporocarp, or antheridium, more magnified (3); fertile sporocarp more magnified (4); the same burst open, showing the stalked sporangia (5); one of the latter more magnified (6); another bursting (7); and three spores (8), beset with bristles.







Cyperus


Dulichium

Kyllingia





Гривинъ






of Aug 125






Genera of Grasses.


Tab. VII.




Genera of Grasses


Tab. VIII.


















Hierochloa




Anthoxanthum



Phalaris




Milium




Amphicarpum



Paspalum




Seturia





Panicum



Cenchrus




Tripsacum




Andropogon



Erianthus




Sorghum





Polypodium


Struthiopteris

Pellaea









Genera of Lycopodiaceae, Equisetaceae, &c.

